Clinical Outcomes, Costs, and Cost-effectiveness of Strategies for People Experiencing

Sheltered Homelessness During the COVID-19 Pandemic

Travis P. Baggett, MD, MPH [1, 2, 3]

Justine A. Scott, MPH [4]

Mylinh H. Le, BA [4]

Fatma M. Shebl, MD, PhD [4]

Christopher Panella, BA [4]

Elena Losina, PhD [8, 9,10]

Clare Flanagan, MPH [4]

Jessie M. Gaeta, MD [3, 7]

Anne Neilan, MD, MPH [2, 4, 5, 11]

Emily P. Hyle, MD, MSc [2, 4, 5, 6]

Amir Mohareb, MD [4, 5]

Krishna P. Reddy, MD, MS [2, 4, 12]

Mark J. Siedner, MD, MPH [2, 4, 5, 13]

Guy Harling, ScD, MPH [14, 15, 16, 17]

Milton C. Weinstein, PhD [19]

Andrea Ciaranello, MD, MPH [2, 4, 5, 6]

Pooyan Kazemian, PhD** [18]

Kenneth A. Freedberg, MD, MSc** [1, 2, 4, 5, 6, 19]

**These two authors contributed equally to this work

1.   Division of General Internal Medicine, Massachusetts General Hospital, Boston, MA

2.   Harvard Medical School, Boston, MA

3.   Institute for Research, Quality, and Policy in Homeless Health Care, Boston Health Care

26   for the Homeless Program, Boston, MA

4.   Medical Practice Evaluation Center, Massachusetts General Hospital, Boston, MA

5.   Division of Infectious Diseases, Massachusetts General Hospital, Boston, MA

6.   Harvard University Center for AIDS Research, Boston, MA

7.   Section of General Internal Medicine, Boston University School of Medicine

8.   Department of Biostatistics, Boston University School of Public Health, Boston, MA

9.   Orthopedic and Arthritis Center for Outcomes Research (OrACORe), Department of

33   Orthopedic Surgery, Brigham and Women's Hospital, Boston, MA.

34   10.   Policy and Innovation eValuation in Orthopedic Treatments (PIVOT) Center, Department

35   of Orthopedic Surgery, Brigham and Women's Hospital, Boston, MA.

36   11.   Division of General Academic Pediatrics, Department of Pediatrics, Massachusetts

37   General Hospital, Boston, MA

38   12.   Division of Pulmonary and Critical Care Medicine, Massachusetts General Hospital,

39   Boston, MA

40   13.   Africa Health Research Institute, KwaZulu-Natal, South Africa

41   14.   Department of Epidemiology and Harvard Center for Population & Development Studies,

42   Harvard T.H. Chan School of Public Health, Boston, MA, USA

43   15.   Africa Health Research Institute, KwaZulu-Natal, South Africa

44   16.   Institute for Global Health, University College London, London, UK

45   17.   MRC/Wits Rural Public Health & Health Transitions Research Unit (Agincourt), University

46   of Witwatersrand, South Africa

47   18.   Department of Operations, Weatherhead School of Management, Case Western Reserve

48   University, Cleveland, OH

19.   Department of Health Policy and Management, Harvard T. H. Chan School of Public

Health, Boston, MA

Corresponding author:

Kenneth A. Freedberg, MD MSc   Phone: (617) 724-3341

Medical Practice Evaluation Center   Fax: (617) 726-4120

Massachusetts General Hospital   Email: kfreedberg@mgh.harvard.edu

100 Cambridge Street, Suite 1600

Boston, MA 02114

Word count (Abstract): 376/350 words

Word count (Text): 3,014/3,000 words

Tables: 2

Figures: 3

Key Points

Question: What are the projected clinical outcomes and costs of strategies for reducing

COVID-19 infections among people experiencing sheltered homelessness?

Findings: In this microsimulation modeling study, daily symptom screening with polymerase

chain reaction (PCR) testing of screen-positive individuals, paired with non-hospital care site

management of people with mild to moderate COVID-19, substantially reduced infections and

lowered costs over 4 months compared to no intervention, across a wide range of epidemic

scenarios. In a surging epidemic, adding periodic universal PCR testing to symptom screening

and non-hospital care site management improved clinical outcomes at modestly increased

costs. Periodic universal PCR testing paired with temporary housing further reduced infections

but at much higher cost.

Meaning: Daily symptom screening with PCR testing of screen-positive individuals and use of

alternate care sites for COVID-19 management among sheltered homeless people was

associated with substantially reduced new cases and costs compared to other strategies.

ABSTRACT

Importance: Approximately 356,000 people stay in homeless shelters nightly in the US. They are at high risk for COVID-19.

Objective: To assess clinical outcomes, costs, and cost-effectiveness of strategies for COVID-19 management among sheltered homeless adults.

Design: We developed a dynamic microsimulation model of COVID-19 in sheltered homeless adults in Boston, Massachusetts. We used cohort characteristics and costs from Boston Health Care for the Homeless Program. Disease progression, transmission, and outcomes data were from published literature and national databases. We examined surging, growing, and slowing epidemics (effective reproduction numbers [R_e] 2.6, 1.3, and 0.9). Costs were from a health care sector perspective; time horizon was 4 months, from April to August 2020.

Setting & Participants: Simulated cohort of 2,258 adults residing in homeless shelters in Boston.

Interventions: We assessed daily symptom screening with polymerase chain reaction (PCR) testing of screen-positives, universal PCR testing every 2 weeks, hospital-based COVID-19 care, alternate care sites [ACSs] for mild/moderate COVID-19, and temporary housing, each compared to no intervention.

Main Outcomes and Measures: Cumulative infections and hospital-days, costs to the health care sector (US dollars), and cost-effectiveness, as incremental cost per case prevented of COVID-19.

Results: We simulated a population of 2,258 sheltered homeless adults with mean age of 42.6 years. Compared to no intervention, daily symptom screening with ACSs for pending tests or confirmed COVID-19 and mild/moderate disease led to 37% fewer infections and 46% lower costs (R_e=2.6), 75% fewer infections and 72% lower costs (R_e=1.3), and 51% fewer infections and 51% lower costs (R_e=0.9). Adding PCR testing every 2 weeks further decreased infections; incremental cost per case prevented was $1,000 ($R_e$=2.6), $27,000 ($R_e$=1.3), and $71,000

103 (R_e=0.9). Temporary housing with PCR every 2 weeks was most effective but substantially more

104 costly than other options. Results were sensitive to cost and sensitivity of PCR and ACS

105 efficacy in preventing transmission.

106 **Conclusions & Relevance:** In this modeling study of simulated adults living in homeless

107 shelters, daily symptom screening and ACSs were associated with fewer COVID-19 infections

108 and decreased costs compared with no intervention. In a modeled surging epidemic, adding

109 universal PCR testing every 2 weeks was associated with further decrease in COVID-19

110 infections at modest incremental cost and should be considered during future surges.

112 **Keywords:** Homelessness, COVID-19, cost-effectiveness analysis, simulation model

INTRODUCTION

Over 1.4 million people experience sheltered homelessness annually in the US, including approximately 356,000 each night.[1,2] The crowded circumstances of homeless shelters place this population at increased risk for coronavirus disease 2019 (COVID-19). The United States (US) Centers for Disease Control and Prevention (CDC) issued comprehensive guidance for preventing and mitigating COVID-19 among people experiencing sheltered homelessness, including recommendations for infection control practices in shelters, symptom screening of shelter guests, and dedicated settings for isolation and management of individuals with symptoms or confirmed illness.[3] The high burden of COVID-19 among sheltered homeless populations[4–7] highlights an urgent need to understand the clinical outcomes and costs of CDC-recommended and other prevention and treatment strategies. After a cluster of COVID-19 cases at a single large shelter in Boston, universal polymerase chain reaction (PCR) testing of 408 shelter residents found that 36% had SARS-CoV-2 infection.[4] Eighty-eight percent of these individuals reported no symptoms at the time of testing, raising questions about how to identify COVID-19 disease in this population and the role of non-hospital alternate care sites (ACSs) to isolate those who do not require hospitalization. Our objective was to project the clinical and economic impact of COVID-19 management approaches for adults experiencing sheltered homelessness.

METHODS

Analytic Overview

We developed the Clinical and Economic Analysis of COVID-19 interventions (CEACOV) model, a dynamic microsimulation of the natural history of COVID-19 disease and the impact of prevention, testing, and treatment interventions. We used CEACOV to project the clinical impact, costs, and cost-effectiveness of various COVID-19 management strategies for people experiencing sheltered homelessness, including different combinations of symptom screening,

PCR testing, ACSs, and relocating all shelter residents to temporary housing. Using data from

the early stage of an outbreak among homeless adults in Boston, Massachusetts, we modeled a

cohort of sheltered homeless adults and examined management strategies under various

epidemic scenarios, given evolving and heterogenous epidemic dynamics across the US.[4,8] We

evaluated 3 scenarios over a 4-month time horizon, from April to August 2020, with different

effective reproduction numbers (R_e) representing surging (R_e=2.6), growing (R_e=1.3), and

slowing (R_e=0.9) epidemics. Outcomes included number of infections, utilization of hospital and

intensive care unit (ICU) beds, costs, and cost per COVID-19 case was conducted from a health

care sector perspective. This study was approved by the Partners prevented. The analysis

Human Research Committee.

Model Structure

Disease states and progression

CEACOV is a dynamic microsimulation model of COVID-19 based on an SEIR framework,

including susceptible, exposed, infectious, recovered, and death states.[9] Infected individuals

face daily probabilities of disease progression through 6 COVID-19 states: pre-infectious

latency, asymptomatic, mild/moderate, severe, critical, and recuperation. With mild/moderate

disease, individuals have mild symptoms, such as cough or fever, that generally do not require

inpatient management in a stably housed population. With severe disease, symptoms warrant

inpatient management. With critical disease, patients require ICU care. Recovered individuals

cannot transmit and are assumed immune from repeat infection.[10] eFigure1 displays how

patients move through the model. We describe model validation in the Supplemental Methods.

Transmission

Individuals with COVID-19 transmit to susceptible individuals at health state-stratified rates. We

model a closed cohort, with transmissions occurring between people experiencing sheltered

homelessness. All susceptible people face equal probabilities of contacting infected individuals

and becoming infected (homogenous mixing). The number of projected infections depends on

COVID-19 prevalence, proportion of the population susceptible, transmission rates, and

interventions that change contact rates or infectivity per contact. Transmission rates are

calibrated to achieve the desired R_e, which captures the average number of transmissions per

case. More details can be found in the Supplemental Methods.

Testing and care interventions

Symptom screens or PCR tests are offered at intervals defined in each strategy; test

sensitivities and specificities depend on COVID-19 health state. Care interventions include

hospital care, ACSs, and temporary housing. Since adequate isolation for COVID-19 is not

possible within congregate homeless shelters, care of homeless individuals with mild/moderate

COVID-19 occurs either in hospitals or ACSs, such as large tents or non-hospital facilities with

on-site medical staff.[11,12] ACSs reduce transmission and hospital use for people with

mild/moderate illness. Temporary housing reduces transmission by preemptively moving

everyone from shelters to individual living units (e.g., hotel or dormitory rooms) for the entire

simulation period. Anyone who develops mild/moderate COVID-19 remains in temporary

housing, which offers health monitoring and space for isolation but less intensive staffing and

infection control than ACSs.

Resource use, costs, cost-effectiveness, and budget impact

The model tallies resource utilization, including tests and days in hospital, ICU, ACS, or

temporary housing, and daily costs, including medical supplies and personnel. We included a

budget impact analysis to determine total costs over the 4-month simulation. To understand the

tradeoffs between cost and infections prevented and highlight the relative "return on investment"

for each strategy, we present efficiency frontiers, plotting number of infections prevented against

total cost for each strategy.[13] Since we focus on a cohort relevant to an individual city, and since

overall COVID-19 mortality is low, we report incremental cost per COVID-19 case prevented as

an outcome; $1,000/case prevented is approximately equivalent to $61,000/quality-adjusted life

year (QALY) gained at current case fatality levels (Table 2, notes).

Strategies

We assessed 8 strategies:

1) *NoIntervention*: Only basic infection control practices are implemented in shelters.

2) *SxScreen/PCR/Hospital*: CDC-recommended symptom screening daily in shelters.[14] Screen-

negative individuals remain in shelters. Screen-positive individuals are sent to the hospital for

PCR testing. PCR-positive individuals remain in hospital; PCR-negative individuals return to

shelter.

3) *SxScreen/PCR/ACS*: CDC-recommended symptom screening daily in shelters. Screen-

negative individuals remain in shelters. Screen-positive individuals are sent to an ACS for

people under investigation, where they undergo PCR testing and await results. PCR-positive

individuals with mild/moderate illness are transferred to ACSs for confirmed COVID-19 cases.

PCR-negative individuals return to shelter.

4) *UniversalPCR/Hospital*: Universal PCR testing every 2 weeks in shelters. Those with

symptoms at the time of testing await results at the hospital; individuals without symptoms await

results in shelters. PCR-negative individuals return to or stay in shelters. PCR-positive

individuals, regardless of illness severity, remain in or are sent to the hospital.

5) *UniversalPCR/ACS*: Universal PCR testing every 2 weeks in shelters. Those with symptoms

at the time of testing are sent to an ACS for people under investigation while awaiting results;

individuals without symptoms await results in shelters. PCR-negative individuals return to or

stay in shelters. PCR-positive individuals with mild/moderate illness are transferred to ACSs for

confirmed COVID-19 cases.

6) *UniversalPCR/TempHousing*: All shelter residents are pre-emptively moved to temporary

housing for the duration of the 4-month period. Universal PCR testing occurs every 2 weeks.

PCR-positive individuals with mild/moderate illness remain in temporary housing and are

transferred to the hospital if they progress to severe or critical disease.

7) *Hybrid/Hospital*: This includes the *SxScreen/PCR/Hospital* strategy and adds shelter-based

universal PCR testing every 2 weeks for those without symptoms.

8) *Hybrid/ACS*: This includes the *SxScreen/PCR/ACS* strategy and adds shelter-based

universal PCR testing every 2 weeks for those without symptoms.

In all 8 strategies, people with severe or critical illness are sent to the hospital. Individuals are

eligible for repeat PCR testing after 5 days since their most recent negative test. See eFigure2

for details.

Input Parameters

Cohort characteristics

The simulated cohort represents 2,258 adults living in Boston homeless shelters.[2] 83% are aged

18-59 years, and 17% are $\geq$60 years (Table 1). Initial prevalence of active or past COVID-19 is

assumed to be 2.2%. To reflect symptoms similar to but not due to COVID-19 (e.g., from other

respiratory viruses or seasonal rhinitis), susceptible and recovered individuals have a 0.01%

daily probability of exhibiting mild/moderate COVID-like symptoms.[15–17]

Progression of COVID-19 and transmission

Average duration of each COVID-19 state varies by severity (eTable1). The probabilities of

developing severe or critical disease or dying increase with age.[18,19] Transmission rates are

highest for individuals in asymptomatic and mild/moderate states; individuals in severe and

critical states have fewer infectious contacts due to hospitalization.[19–22]

Testing

We assumed symptom screen sensitivity of 0% for asymptomatic infection, 62% for

mild/moderate COVID-19, and 100% for severe or critical COVID-19.[4] The PCR test is a

nasopharyngeal sample with one-day result delay, 70% sensitivity for people with no symptoms

or mild/moderate symptoms,[23,24] 100% sensitivity for severe or critical illness, and 100%

specificity.

Hospitalization, alternate care sites, and temporary housing

Mortality was decreased with hospitalization among those with critical illness.[18,19] We assumed

hospitalization reduces transmission by 100%, while ACSs reduce transmission by 80% and

temporary housing by 60%. Temporary housing was assumed less effective at reducing

transmission compared to ACSs due to less stringent infection control measures in temporary

housing and potential mixing of uninfected and infected individuals. Length-of-stay at hospitals

and ACSs depends on severity and duration of illness.[18–21,25–28]

Resource use and costs

The nasopharyngeal PCR test costs $51.[29] Hospitalization costs $1,641/day; ICU costs

$2,683/day (Table 1; Supplement).[30–32] ACSs cost $304/day; temporary housing costs $141/day

(data from BHCHP).

Sensitivity Analyses

In one-way sensitivity analyses, we examined: 1) PCR sensitivity, PCR frequency, and symptom

screen sensitivity (eTables2-4); 2) efficacy of ACS and temporary housing in reducing

transmission (eTables5-6); and 3) costs of PCR test, symptom screen, hospital care, ACS, and

temporary housing (eTables7-11). In two-way sensitivity analyses, we varied influential

parameters simultaneously (eTables12-13). To relate these findings to other settings, eTable14

displays outcomes per 1,000 homeless adults and the number of sheltered homeless adults in

select US cities.

RESULTS

Base Case

Surging epidemic (R_e=2.6)

With R_e=2.6, the number of projected COVID-19 cases was highest with *NoIntervention* (1,954)

and lowest with *UniversalPCR/TempHousing* (376) (Table 2; Figure 1). Other than the

temporary housing strategy, strategies that rely on daily symptom screening were more effective

in preventing infections (1,133 to 1,239 cumulative infections) than those with universal PCR

testing every two weeks alone (1,679 to 1,681 cumulative infections). Hybrid strategies involving

daily symptom screening plus universal PCR testing every two weeks performed better than

either alone (967 to 985 cumulative infections).

With R_e=2.6, all ACS-based strategies had lower total costs ($3.27 to $4.14 million) than

hospital-based strategies ($12.20 to $12.91 million) and *NoIntervention* ($6.10 million; Table 2;

Figure 2, eTable15). *UniversalPCR/TempHousing* was most costly ($39.12 million), and

SxScreen/PCR/ACS was least costly ($3.27 million).

Compared with *SxScreen/PCR/ACS*, *Hybrid/ACS* had 20% fewer cases (985 vs 1,239) at

$1,000/case prevented (Table 2). *UniversalPCR/TempHousing*, the most clinically effective

strategy, had an incremental cost of $58,000/case prevented compared to *Hybrid/ACS*. All other

strategies were dominated, or less effective and more costly than another strategy or

combination of strategies (Table 2; Figure 2, eTable15).

Growing epidemic (R_e=1.3)

With R_e=1.3, projected cases ranged from 538 (*NoIntervention*) to 95

(*UniversalPCR/TempHousing*) (Table 2; Figure 1). All strategies had at least 60% fewer

infections than *NoIntervention*. ACS strategies had fewer infections, fewer hospital days, and

lower costs than *NoIntervention*, whereas hospital strategies had higher costs than

NoIntervention (Table 2; Figure 2, eTable15). *SxScreen/PCR/ACS* had 75% fewer infections

than *NoIntervention* and the lowest cost. Compared to *SxScreen/PCR/ACS*, *Hybrid/ACS* yielded

an additional 6% decrease in infections at $27,000/case prevented.

UniversalPCR/TempHousing had the lowest number of infections at $6,854,000/case prevented

(Table 2; Figure 3).

Slowing epidemic (R_e=0.9)

With R_e=0.9, cumulative infections were fewer than in the other scenarios, ranging from 174

(*NoIntervention*) to 71 (*UniversalPCR/TempHousing*) (Table 2; Figure 1). All strategies had at

least 46% fewer infections than *NoIntervention*. *SxScreen/PCR/ACS* had 51% fewer infections

and 51% lower costs than *NoIntervention;* it was the only strategy that cost less than

NoIntervention (Table 2; Figure 2, eTable15). Compared to *SxScreen/PCR/ACS*, *Hybrid/ACS*

yielded an additional 8% decrease in infections at $71,000/case prevented (Table 2; Figure 3).

Sensitivity Analyses

One-way sensitivity analysis

Across the 3 epidemic scenarios, changes in PCR sensitivity, PCR cost, PCR frequency, and

ACS efficacy had the greatest impacts on the incremental cost per case prevented. If PCR

sensitivity increased from 70% to 90% with R_e=2.6, the number of infections with *Hybrid/ACS*

decreased from 985 to 668; incremental cost per case prevented was $100 compared with

SxScreen/PCR/ACS (eTable2). If PCR cost decreased from $51 to $25 in R_e=2.6, the

Hybrid/ACS strategy became cost-saving compared with *SxScreen/PCR/ACS* (eTable7).

Results for higher PCR costs are also shown in eTable7. If ACS efficacy in preventing

transmissions decreased, total cases increased in all ACS-based strategies, and *Hybrid/ACS*

became relatively less effective compared to *SxScreen/PCR/ACS* (eTable5).

With R_e=2.6, *Hybrid/ACS* with universal PCR testing every 7 rather than every 14 days was

associated with 29% fewer infections (incremental cost of $1,000/case prevented compared

with testing every 14 days, eTable16). Every 3-day testing had fewer infections, at $2,000/case

prevented. In other R_e scenarios, the *Hybrid/ACS* strategy did not result in a cost per case

prevented below $20,000 compared with *SxScreen/PCR/ACS*, regardless of universal testing

frequency.

ACS-based management approaches remained less costly than hospital care unless daily ACS

costs began to approach hospital costs. Although *UniversalPCR/TempHousing* had the lowest

number of cases in all scenarios, with R_e=2.6, daily costs of temporary housing needed to be

≤$20/day to have an incremental cost per case prevented of ≤$1,000 compared to *Hybrid/ACS*

(eTable11). In the lower R_e scenarios, *UniversalPCR/TempHousing* had higher costs per case

prevented.

Two-way sensitivity analysis

In two-way sensitivity analysis there were several combinations where *Hybrid/ACS* was cost-

saving or had an incremental cost per case prevented compared to *SxScreen/PCR/ACS* of

$1,000-$3,000 as the sensitivity of PCR increased and PCR cost decreased (eTable12).

DISCUSSION

We developed a microsimulation model to examine the impact of COVID-19 testing and

isolation strategies on infections and health care costs among adults experiencing sheltered

homelessness. Across all epidemic scenarios, daily symptom screening with PCR testing of

screen-positive individuals and ACS-based COVID-19 management was the most efficient

strategy and was cost-saving relative to no intervention.

In all cases, strategies employing ACSs for isolation of symptomatic individuals with pending

tests, and for those with confirmed mild or moderate COVID-19, were associated with

substantially decreased costs compared to analogous strategies relying on hospital-based care

while achieving similar clinical outcomes. ACSs are especially useful for managing COVID-19 in

homeless populations since people with mild to moderate illness cannot be effectively isolated

in shelters. With high levels of SARS-CoV-2 infection among people experiencing

homelessness in Boston and other cities,[4–7,33] ACSs could avert many hospitalizations,

preserving beds for severely ill individuals and reducing costs. Boston created several such

ACSs, ranging from 16-bed tents to a 500-bed field unit in a downtown convention center.[34] In

cities with smaller numbers of sheltered homeless adults (eTable14), using existing facilities

(e.g. hotels/motels) as ACSs would avoid the fixed costs of new ACSs and allow for rapid

implementation of care sites for people with mild to moderate COVID-19.

In a surging epidemic, adding universal PCR testing every 14 days to daily symptom screening

had clinical benefits at an incremental cost of $1,000/case prevented. We selected a 2-week

testing interval because this was deemed by BHCHP clinical staff to be realistic and in line with

practice during the study time period; however, reducing the universal testing interval to every 7

days yielded additional benefits at $1,000/case prevented. In sensitivity analyses, this "hybrid"

approach of daily symptom screening with additional periodic universal PCR testing was less

expensive than daily symptom screening alone when PCR sensitivity increased and PCR cost

decreased. In a growing or slowing epidemic, testing beyond daily symptom screening

prevented a small number of new cases at high incremental costs. If PCR turnaround time was

longer than the 1 day we modeled, all strategies would have more cases and higher costs.

Temporary housing with universal PCR testing every 2 weeks was the most effective strategy

for reducing COVID-19 in all scenarios but was also the most costly, except in sensitivity

analyses where temporary housing costs were reduced below plausible ranges. However, this

analysis does not account for other potential benefits of temporary housing on physical or

mental health.[35] Ultimately, broader policies around supportive housing measures for people

experiencing homelessness should account for more than COVID-19 mitigation, recognizing

that the COVID-19 pandemic is one of many health risks of homelessness.[36]

This study complements the findings of a dynamic transition model of structural interventions for

COVID-19 among people experiencing homelessness in England.[37] In that analysis, single-

room accommodations for people with COVID-19 symptoms and people without symptoms but

at high risk for COVID-19 complications were projected to reduce infections, hospitalizations,

and deaths by 36% to 64%. Our analysis adds to this by examining additional structural

interventions (e.g. ACSs and temporary housing) in a US context, combined with various

COVID-19 diagnostic approaches (e.g. symptom screening, universal PCR testing, and hybrid

strategies), and by adding cost-effectiveness to inform policy and practice.

This analysis has limitations. The findings are specific to individual adults; we excluded adults

experiencing homelessness as part of a family, because family shelters more likely provide

private living quarters.[38] We also excluded unsheltered homeless individuals because disease

transmission dynamics and infection control considerations are distinct for this subpopulation.[39]

We assumed homogeneous mixing of sheltered homeless adults; in reality this population is

spread over numerous shelters. This homogenous mixing assumption may impact the number

of infections projected by our model, but we expect this impact to be small. In the base case, we

did not assume increased comorbidities among homeless adults compared with the general

population.[40] The analysis is based on the possibility that ACSs and PCR tests can be made

available relatively quickly to homeless adults. This may be difficult in some settings because

those responsible for making ACSs and PCR tests available may not be those responsible for

hospital costs, and record-keeping may be challenging. Finally, we focused this analysis on

Boston, which has a 29.7% higher cost of living than the US average.[41] Costs of temporary

housing may be considerably lower in other cities. In sensitivity analysis, however, results were

robust to even large changes in testing, hospital, and housing costs.

In summary, daily symptom screening and use of ACSs for those with pending test results or

mild to moderate COVID-19 was associated with reduced infections and lower costs compared

to no intervention. In a surging epidemic, adding universal PCR testing every 2 weeks was

associated with further reduction in infections at a reasonable cost. Routine symptom screening,

implementation of ACSs, and selective use of universal PCR testing should be implemented for

sheltered homeless populations in the US.

ACKNOWLEDGEMENTS

We thank Elizabeth Lewis, MBA and Agnes Leung, MHA for their assistance with clinical and

cost data from Boston Health Care for the Homeless Program. We also thank Guner Ege

Eskibozkurt, BA and Mary Feser, BA for research assistance. All acknowledged individuals

contributed as part of their institutional roles.

KAF and TPB had full access to all the data in the study and take responsibility for the integrity

of the data and the accuracy of the data analysis.

AUTHOR ROLES

All authors contributed substantively to this manuscript in the following ways: study and model

design (all authors), data analysis (MHL, FMS, EL), interpretation of results (all authors),

drafting the manuscript (KAF, TPB), and critical revision of the manuscript (all authors) and final

approval of submitted version (all authors).

CONFLICTS OF INTEREST AND FINANCIAL DISCLOSURES

The authors have no conflicts of interest or financial disclosures.

FUNDING

This work was supported by the National Institute of Allergy and Infectious Disease [T32

AI007433 to AM] and the National Institute of Arthritis and Musculoskeletal and Skin Diseases

[K24 AR057827 to EL] at the National Institutes of Health, and by the Royal Society and

Wellcome Trust [210479/Z/18/Z to GH].

The funding sources had no role in the study design, data collection, data analysis, data

interpretation, writing of the manuscript, or in the decision to submit the manuscript for

440   publication. The content is solely the responsibility of the authors and does not necessarily

441   represent the official views of the funding sources.

REFERENCES

1. Henry M, Bishop K, de Sousa T, Shivji A, Watt R. *The 2017 Annual Homeless Assessment Report (AHAR) to Congress PART 2: Estimates of Homelessness in the United States.* The U.S. Department of Housing and Urban Development; 2018.

2. Henry M, Watt R, Mahathey A, Ouellette J, Sitler A. *The 2019 Annual Homeless Assessment Report (AHAR) to Congress, Part 1: Point-in-Time Estimates of Homelessness.* The U.S. Department of Housing and Urban Development; 2020.

3. CDC. Interim guidance for homeless service providers to plan and respond to coronavirus disease 2019 (COVID-19). Centers for Disease Control and Prevention. Published February 11, 2020. Accessed July 29, 2020. https://www.cdc.gov/coronavirus/2019-ncov/community/homeless-shelters/plan-prepare-respond.html

4. Baggett TP, Keyes H, Sporn N, Gaeta JM. Prevalence of SARS-CoV-2 infection in residents of a large homeless shelter in Boston. *JAMA.* Published online April 27, 2020. doi:10.1001/jama.2020.6887

5. Testing at Worcester homeless shelter finds 43% positive for coronavirus. WBUR. Published May 20, 2020. Accessed July 29, 2020. https://www.wbur.org/news/2020/04/17/worcester-homeless-population-covid-19-coronavirus

6. Tobolowsky FA, Gonzales E, Self JL, et al. COVID-19 outbreak among three affiliated homeless service sites - King County, Washington, 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69(17):523-526. doi:10.15585/mmwr.mm6917e2

7. Mosites E, Parker EM, Clarke KEN. Assessment of SARS-CoV-2 infection prevalence in homeless shelters — Four U.S. cities, March 27–April 15, 2020. *MMWR Morb Mortal Wkly Rep.* 2020;69. doi:10.15585/mmwr.mm6917e1

8. Johns Hopkins University. COVID-19 United States cases by county. Published 2020. Accessed July 29, 2020. https://coronavirus.jhu.edu/us-map

9.   Martcheva M (Maia). *An Introduction to Mathematical Epidemiology*. 1st ed. 2015. Springer

US : Imprint: Springer; 2015.

10.   Chandrashekar A, Liu J, Martinot AJ, et al. SARS-CoV-2 infection protects against

rechallenge in rhesus macaques. *Science*. Published online May 20, 2020:eabc4776.

doi:10.1126/science.abc4776

11.   CDC. Considerations for alternate care sites. Centers for Disease Control and Prevention.

Published February 11, 2020. Accessed July 29, 2020.

https://www.cdc.gov/coronavirus/2019-ncov/hcp/alternative-care-sites.html

12.   MacKenzie OW, Trimbur MC, Vanjani R. An Isolation Hotel for People Experiencing

Homelessness. *N Engl J Med*. 2020;383(6):e41. doi:10.1056/NEJMc2022860

13.   Gold M, Siegel J, Russell L, Weinstein MC. *Cost-Effectiveness in Health and Medicine*.

Oxford University Press; 1996.

14.   CDC. Screening clients for COVID-19 at homeless shelters or encampments. Centers for

Disease Control and Prevention. Published May 20, 2020. Accessed July 29, 2020.

https://www.cdc.gov/coronavirus/2019-ncov/community/homeless-shelters/screening-

clients-respiratory-infection-symptoms.html

15.   Rui P, Okeyode T. National ambulatory medical care survey: 2016 national summary

tables. Centers for Disease Control and Prevention. Published 2019. Accessed August 4,

2020. https://www.cdc.gov/nchs/data/ahcd/namcs_summary/2016_namcs_web_tables.pdf

16.   Centers for Disease Control and Prevention. Percentage of visits for ILI by age group

reported by a subset of ILINet providers. Published July 24, 2020. Accessed August 4,

2020. https://www.cdc.gov/coronavirus/2019-ncov/covid-data/covidview/07242020/percent-

ili-visits-age.html

17.   CDC. U.S. Outpatient Influenza-like Illness Surveillance Network (ILINet): Overall

percentage of visits for ILI. Centers for Disease Control and Prevention. Published June 5,

2020. Accessed September 25, 2020. https://www.cdc.gov/coronavirus/2019-ncov/covid-

data/covidview/06052020/percent-ili-visits.html

18. Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019

novel coronavirus–infected pneumonia in Wuhan, China. *JAMA*. 2020;323(11):1061-1069.

doi:10.1001/jama.2020.1585

19. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z. Clinical course and risk factors for mortality of

adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *The Lancet*.

2020;395(10229):1054-1062.

20. *Report of the WHO-China Joint Mission on Coronavirus Disease 2019 (COVID-19)*. WHO-

China Joint Mission on Coronavirus Disease 2019 (COVID-19); 2020. Accessed July 29,

2020. https://www.who.int/docs/default-source/coronaviruse/who-china-joint-mission-on-

covid-19-final-report.pdf

21. Hu Z, Song C, Xu C, et al. Clinical characteristics of 24 asymptomatic infections with

COVID-19 screened among close contacts in Nanjing, China. *medRxiv*. Published online

January 1, 2020:2020.02.20.20025619. doi:10.1101/2020.02.20.20025619

22. Liu Y, Gayle AA, Wilder-Smith A, Rocklöv J. The reproductive number of COVID-19 is

higher compared to SARS coronavirus. *Journal of Travel Medicine*. 2020;27(taaa021).

doi:10.1093/jtm/taaa021

23. Yang Y, Yang M, Shen C, et al. Evaluating the accuracy of different respiratory specimens

in the laboratory diagnosis and monitoring the viral shedding of 2019-nCoV infections.

medRxiv. Published online January 1, 2020:2020.02.11.20021493.

doi:10.1101/2020.02.11.20021493

24. Wang W, Xu Y, Gao R, et al. Detection of SARS-CoV-2 in different types of clinical

specimens. *JAMA*. 2020;323(18):1843-1844. doi:10.1001/jama.2020.3786

25.   Li Q, Guan X, Wu P, et al. Early transmission dynamics in Wuhan, China, of novel

518   coronavirus–infected pneumonia. *N Engl J Med*. 2020;382(13):1199-1207.

519   doi:10.1056/NEJMoa2001316

520   26.   He X, Lau EH, Wu P, et al. Temporal dynamics in viral shedding and transmissibility of

521   COVID-19. *medRxiv*. Published online January 1, 2020:2020.03.15.20036707.

522   doi:10.1101/2020.03.15.20036707

523   27.   Linton NM, Kobayashi T, Yang Y, et al. Incubation period and other epidemiological

524   characteristics of 2019 novel coronavirus infections with right truncation: a statistical

525   analysis of publicly available case data. *J Clin Med*. 2020;9(2):538.

526   doi:10.3390/jcm9020538

527   28.   Yu P, Zhu J, Zhang Z, Han Y. A familial cluster of infection associated with the 2019 novel

528   coronavirus indicating possible person-to-person transmission during the incubation

529   period. *The Journal of Infectious Diseases*. 2020;221(11):1757-1761.

530   doi:10.1093/infdis/jiaa077

531   29.   Medicare Administrative Contractor (MAC) COVID-19 test pricing. Published May 19,

532   2020. Accessed July 29, 2020. https://www.cms.gov/files/document/mac-covid-19-test-

533   pricing.pdf

534   30.   Cox C, Rudowitz R, Neuman T, Cubanski J, Rae M. How health costs might change with

535   COVID-19. Health System Tracker. Published April 15, 2020. Accessed July 29, 2020.

536   https://www.healthsystemtracker.org/brief/how-health-costs-might-change-with-covid-19/

537   31.   Rae M, Claxton G, Kurani N, McDermott D, Cox C. Potential costs of COVID-19 treatment

538   for people with employer coverage. Peterson-Kaiser Health System Tracker. Published

539   2020. Accessed July 29, 2020. https://www.healthsystemtracker.org/brief/potential-costs-

540   of-coronavirus-treatment-for-people-with-employer-coverage/

541   32.   *COVID-19: The Projected Economic Impact of the COVID-19 Pandemic on the US*

542   *Healthcare System*. FAIR Health, Inc.; 2020. Accessed July 29, 2020.

https://s3.amazonaws.com/media2.fairhealth.org/brief/asset/COVID-19%20-

%20The%20Projected%20Economic%20Impact%20of%20the%20COVID-

19%20Pandemic%20on%20the%20US%20Healthcare%20System.pdf

33. Stokes S. Atlanta tests more than 2,000 people who are homeless for COVID-19. WABE.

Published April 21, 2020. Accessed July 29, 2020. https://www.wabe.org/atlanta-tests-

more-than-2000-people-who-are-homeless-for-covid-19/

34. Convention centers fill with beds for COVID-19, including 500 for Boston's homeless.

Accessed July 29, 2020. https://www.wbur.org/commonhealth/2020/04/09/convention-

centers-transform-into-field-hospitals-in-boston-and-worcester

35. Padgett DK, Stanhope V, Henwood BF, Stefancic A. Substance use outcomes among

homeless clients with serious mental illness: comparing Housing First with Treatment First

programs. *Community Ment Health J*. 2011;47(2):227-232. doi:10.1007/s10597-009-9283-

7

36. Tsai J, Wilson M. COVID-19: a potential public health problem for homeless populations.

Lancet Public Health. 2020;5(4):e186-e187. doi:10.1016/S2468-2667(20)30053-0

37. Lewer D, Braithwaite I, Bullock M, Eyre MT, Aldridge RW. COVID-19 and homelessness in

England: a modelling study of the COVID-19 pandemic among people experiencing

homelessness, and the impact of a residential intervention to isolate vulnerable people and

care for people with symptoms. *medRxiv*. Published online January 1,

2020:2020.05.04.20079301. doi:10.1101/2020.05.04.20079301

38. Overview of the Department of Housing and Community Development. Mass.gov.

Published August 28, 2019. Accessed October 2, 2020. https://www.mass.gov/info-

details/overview-of-the-department-of-housing-and-community-development#ea-program-

39. CDC. Interim guidance on unsheltered homelessness and coronavirus disease 2019

(COVID-19) for homeless service providers and local officials. Centers for Disease Control

and Prevention. Published May 13, 2020. Accessed July 29, 2020.

https://www.cdc.gov/coronavirus/2019-ncov/community/homeless-shelters/unsheltered-

homelessness.html

40. Brown RT, Hemati K, Riley ED, et al. Geriatric conditions in a population-based sample of

older homeless adults. *The Gerontologist*. 2016;57(4):757-766. doi:10.1093/geront/gnw011

41. Cost of living data series. Missouri Economic Research and Information Center. Published

2020. Accessed September 25, 2020. https://meric.mo.gov/data/cost-living-data-series

42. COVID-19 Dashboard. Massachusetts Department of Public Health. Published 2020.

Accessed July 29, 2020. https://www.mass.gov/doc/covid-19-dashboard-april-20-

2020/download

43. Mizumoto K, Kagaya K, Zarebski A, Chowell G. Estimating the asymptomatic proportion of

coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship,

Yokohama, Japan, 2020. *Euro Surveill*. 2020;25(10):2000180. doi:10.2807/1560-

7917.ES.2020.25.10.2000180

44. Haridy R. CDC director warns 25 percent of COVID-19 cases may present no symptoms.

New Atlas. https://newatlas.com/health-wellbeing/covid-19-cases-contagious-

asymptomatic-presymptomatic-cdc-director/. Published April 1, 2020. Accessed July 29,

2020.

45. Li R, Pei S, Chen B, et al. Substantial undocumented infection facilitates the rapid

dissemination of novel coronavirus (SARS-CoV2). *Science*. Published online March 16,

2020:eabb3221. doi:10.1126/science.abb3221

46. American community survey 1-year estimates. U.S. Census Bureau. Published 2020.

Accessed July 29, 2020. https://censusreporter.org/profiles/04000US25-massachusetts/

47. Richardson S, Hirsch JS, Narasimhan M, et al. Presenting characteristics, comorbidities,

and outcomes among 5700 patients hospitalized with COVID-19 in the New York City

area. *JAMA*. Published online April 22, 2020. doi:10.1001/jama.2020.6775

48. Sanders GD, Neumann PJ, Basu A, et al. Recommendations for conduct, methodological

practices, and reporting of cost-effectiveness analyses: second panel on cost-effectiveness

in health and medicine. *JAMA*. 2016;316(10):1093-1103. doi:10.1001/jama.2016.12195

49. Sullivan PW, Ghushchyan V. Preference-Based EQ-5D index scores for chronic conditions

in the United States. *Med Decis Making*. 2006;26(4):410-420.

doi:10.1177/0272989X06290495

50. Gardner JW, Sanborn JS. Years of Potential Life Lost (YPLL) - What Does it Measure?

Epidemiology. 1990;1(4).

51. Martinez R, Soliz P, Caixeta R, Ordunez P. Reflection on modern methods: years of life

lost due to premature mortality-a versatile and comprehensive measure for monitoring

non-communicable disease mortality. *Int J Epidemiol*. 2019;48(4):1367-1376.

doi:10.1093/ije/dyy254

Table 1. Input parameters for an analysis of management strategies for people experiencing sheltered homelessness during the
COVID-19 pandemic.
609

Parameter	Value				Source
Cohort characteristics					
Cohort size	2,258				[2]
Age distribution, %*					
18-59y	82.9				BHCHP
>60y	17.1				
Natural history					
Probability of COVID-19 severity,[a] stratified by age	Asymptomatic infection	Mild/moderate illness	Severe illness	Critical illness	
18-59y	0.262	0.719	0.012	0.007	Derived from [42–45]
>60y	0.180	0.788	0.001	0.031	
Duration of illness state among hospitalized patients, stratified by COVID-19 severity, mean, days[b]	Asymptomatic infection	Mild/moderate illness	Severe illness	Critical illness	
Pre-infectious latent to asymptomatic state	2.6	2.6	2.6	2.6	Derived from [20,25–27]
Asymptomatic to mild/moderate state	NA	2.0	2.0	2.0	[20,26]
Mild/moderate to severe state	NA	NA	6.5	3.0	[18]
Severe to critical illness state	NA	NA	10.5	7.1	[19]
Critical illness to recuperation state	NA	NA	NA	11.9	[19]
Duration of illness state among non-hospitalized patients, stratified by COVID-19 severity, mean, days[b]	Asymptomatic infection	Mild/moderate illness	Severe illness	Critical illness	
Pre-infectious latent to asymptomatic state	2.6	2.6	2.6	2.6	Derived from [20,25–27]
Asymptomatic to mild/moderate state	NA	2.0	2.0	2.0	[20,26]
Mild/moderate to severe state	NA	NA	6.5	3.0	[18]
Severe to critical illness state	NA	NA	NA	6.5	[19]
Duration of viral shedding, stratified by COVID-19 severity, mean, days[b]	Asymptomatic infection	Mild/moderate illness	Severe illness	Critical illness	
	9.5	12	19	24	[19–21]

Table 1 continued. Input parameters for an analysis of management strategies for people experiencing sheltered homelessness
during the COVID-19 pandemic.

Parameter	Value		Source
Natural history, continued			
Daily probability of mortality in the critical state, stratified by age	18-59y	>60y	
Hospital care	0.004	0.050	Derived from [18,19]
No hospital care	0.166	0.203	Derived from [42,46,47]
Daily probability of onward transmission, stratified by disease state			
Asymptomatic state	0.2394		
Mild/moderate state	0.1948		
Severe state	0.0135		Derived from [19–22]
Critical state	0.0107		
Recuperation state	0.0135		
Persons with other respiratory illnesses exhibiting mild/moderate COVID-like symptoms, daily, %	0.01		[15–17]
Duration of mild/moderate COVID-like symptoms, mean, days	5		Assumed
Intervention			
Reduction in transmission rates,[c] %			
ACS for people with pending PCR test results	80		Assumed
ACS for people with confirmed COVID-19	100		Assumed
Temporary housing	60		Assumed
Hospitalization	100		Assumed

Intervention cost, 2020 USD	Daily cost		Total daily cost	
	Materials	Personnel		
ACS	79	225	304	BHCHP
Temporary housing*	85	56	141	BHCHP
Hospital (non-ICU) bed*	NA	NA	1,641	Derived from [30–32]
ICU bed	NA	NA	2,683	

Table 1 continued. Input parameters for an analysis of management strategies for people experiencing sheltered homelessness
during the COVID-19 pandemic.

Parameter	Value	Source
Testing		
Symptom screen		
Sensitivity, stratified by disease state, %		
Pre-infectious latent	0	Assumed
Asymptomatic state	0	Assumed
Mild/moderate state[d]	62	Derived from [4], Assumed
Severe state	100	Assumed
Critical state	100	Assumed
Result return delay, days	0	Assumed
Unit cost, 2020 USD	0	Assumed
PCR, nasopharyngeal specimen		
Sensitivity, stratified by disease state, %		
Pre-infectious latent	0	Assumed
Asymptomatic state	70	Assumed
Mild/moderate state	70	[23,24]
Severe state	100	Assumed
Critical state	100	Assumed
Specificity, %	100	Assumed
Result return delay, days	1	Assumed
Unit cost, 2020 USD	51	[29]

Abbreviations: ACS, alternate care sites; BHCHP, Boston Health Care for the Homeless Program; COVID-19, coronavirus disease 2019; ICU, intensive care unit; mod., moderate; NA, not applicable; PCR, polymerase chain reaction; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; USD, United States dollars.; y, years.

[a] Severity probability refers to the likelihood that an individual, once infected with SARS-CoV-2, will eventually progress to the specified severity of COVID-19 disease.
[b] Durations of illness state and of viral shedding were derived from model inputs of transition probabilities. See eTable1 for more details.
[c] In ACSs for people with pending PCR test results, there are people without COVID-19 who are susceptible to infection. Transmission in ACSs for people with pending PCR test results is thus not completely reduced. In ACSs for people with confirmed COVID-19, we assumed complete reduction in transmission among sheltered homeless people and did not examine SARS-CoV-2

transmission to healthcare workers. Temporary housing is a less medicalized setting compared to hospitals and ACSs and was
assumed to have a lower reduction in SARS-CoV-2 transmission rates.
[d] The sensitivity of symptom screening for identifying individuals with mild to moderate COVID-19 was derived from an unpublished
reanalysis of data from SARS-CoV-2 testing at a single large shelter in Boston, MA.[4] Among COVID-positive individuals presenting
with mild to moderate symptoms at time of testing, 83% (15/18) would have been identified using symptom screening instrument
concordant with CDC guidelines.[14] To account for the underreporting of symptoms among shelter residents due to stigma and/or fear
of losing shelter accommodations, we estimated that only 75% of those with mild to moderate COVID-19 would report their
symptoms. Thus, we estimated that the symptom screen would identify 62% (0.83*0.75) shelter residents with mild to moderate
COVID-19.
*Data on cohort characteristics and costs of alternative care sites and temporary housing were derived from unpublished data from
the Boston Health Care for the Homeless Program.

Table 2. Results of an analysis of management strategies for people experiencing sheltered homelessness during the COVID-19
pandemic at 4 months (n=2,258).
643

Strategy	Cumulative infections, n	Reduction in cases,[a] %	Peak daily hospital bed use, n	Total hospital days, n	Total cost, [b] 2020 USD	Cost compared with *NoIntervention*, [b] 2020 USD	Incr. cost per case prevented, [b, c] 2020 USD
Effective reproduction number (R$_e$) = 2.6							
SxScreen/PCR/ACS	1,239	36.6	5	394	3,267,000	- 2,831,000	NA
Hybrid/ACS	985	49.6	4	305	3,628,000	- 2,470,000	1,000
UniversalPCR/ACS	1,681	14.0	9	569	4,143,000	- 1,955,000	Dominated
NoIntervention	1,954	NA	64	3,567	6,098,000	NA	Dominated
Hybrid/Hospital	967	50.5	80	6,796	12,202,000	+ 6,104,000	Dominated
SxScreen/PCR/Hospital	1,133	42.0	93	7,656	12,620,000	+ 6,522,000	Dominated
UniversalPCR/Hospital	1,679	14.1	112	7,165	12,914,000	+ 6,816,000	Dominated
UniversalPCR/TempHousing	376	80.8	1	121	39,119,000	+ 33,021,000	58,000
Effective reproduction number (R$_e$) = 1.3							
SxScreen/PCR/ACS	137	74.5	1	48	409,000	- 1,052,000	NA
Hybrid/ACS	103	80.8	1	69	1,325,000	- 136,000	27,000
UniversalPCR/ACS	207	61.5	1	34	1,426,000	- 35,000	Dominated
NoIntervention	538	NA	9	867	1,461,000	NA	Dominated
SxScreen/PCR/Hospital	125	76.7	22	966	1,604,000	+ 143,000	Dominated
Hybrid/Hospital	100	81.4	23	815	2,368,000	+ 907,000	382,000
UniversalPCR/Hospital	207	61.4	19	977	2,631,000	+ 1,170,000	Dominated
UniversalPCR/TempHousing	95	82.3	1	39	38,974,000	+ 37,513,000	6,854,000

645 **Table 2 continued.** Results of an analysis of management strategies for people experiencing sheltered homelessness during the
646 COVID-19 pandemic at 4 months (n=2,258).
647

Strategy	Cumulative infections, n	Reduction in cases,[a] %	Peak daily hospital bed use, n	Total hospital days, n	Total cost,[b] 2020 USD	Cost compared with *NoIntervention,*[b] 2020 USD	Incr. cost per case prevented,[b, c] 2020 USD
Effective reproduction number (R_e) = 0.9							
SxScreen/PCR/ACS	85	51.2	1	30	264,000	- 276,000	NA
NoIntervention	174	0.0	5	318	540,000	NA	Dominated
SxScreen/PCR/Hospital	82	53.2	20	669	1,113,000	+ 573,000	Dominated
UniversalPCR/ACS	94	45.7	1	31	1,226,000	+ 686,000	Dominated
Hybrid/ACS	71	59.1	1	25	1,240,000	+ 700,000	71,000
UniversalPCR/Hospital	95	45.5	19	534	1,901,000	+ 1,361,000	Dominated
Hybrid/Hospital	71	59.4	22	595	2,004,000	+ 1,464,000	Dominated
UniversalPCR/TempHousing	71	59.2	1	29	38,954,000	+ 38,414,000	Dominated

648 Abbreviations: ACS, alternate care site; COVID-19, coronavirus disease 2019; Dominated, less clinically effective and more costly
649 than an alternative strategy, or a combination of two alternative strategies;[48] Incr., incremental; NA, not applicable; *PCR,* polymerase
650 chain reaction; *UniversalPCR*, universal polymerase chain reaction test for everyone; USD, United States dollars; *SxScreen,*
651 symptom screen; *TempHousing,* temporary housing.
652
[a] Reduction in cases are calculated by dividing the number of cases prevented with the use of an alternative strategy by the number
of cumulative cases for *NoIntervention*.
[b] All costs are rounded to the nearest thousands.
[c] Incremental costs per case prevented are calculated by dividing the difference in total costs by the difference in cumulative
infections compared to the next most expensive strategy. All strategies are listed in order of ascending total costs, per convention of
cost-effectiveness analysis.
659
Using 9.50 years of life lost per COVID-19 death from the model, and a mean age-stratified utility of 0.85 for the modeled
population,[42,49–51] a cost per case prevented of $1,000 is equivalent to an incremental cost-effectiveness ratio (ICER) of
$61,000/quality-adjusted life year (QALY) gained. A ratio of $27,000/case prevented is equivalent to $1,728,000/QALY gained. Any
higher cost per case prevented has an even higher ICER.

LEGENDS TO FIGURES

Figure 1. Cumulative infections by management strategy for people experiencing sheltered

homelessness in Boston during the COVID-19 pandemic over a 4-month period.

These panels depict the projected number of cumulative infections over time by management

strategy. Panels A, B, and C show model results for R_e of 2.6, 1.3, and 0.9, respectively. In each

panel, time 0 on the horizontal axis represents the start of model simulation, with SARS-CoV-2

infection prevalence of 2.2%. *UniversalPCR/Hospital* and *UniversalPCR/ACS* are overlapping

lines since they differ only in costs; they are shown separately for clarity. The same is true for

Hybrid/Hospital and *Hybrid/ACS*. The insets in Panels B and C magnify the vertical axis for

clarity. See Methods for strategy definitions.

Abbreviations: ACS, alternate care site; COVID-19, coronavirus disease 2019; *PCR,*

polymerase chain reaction; *UniversalPCR*, universal polymerase chain reaction test for

everyone; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; *SxScreen,* symptom

screen; *TempHousing,* temporary housing.

Figure 2. Health care sector costs of implementing different management strategies for people

experiencing sheltered homelessness in Boston during the COVID-19 pandemic over a 4-month

period.

These panels show the total and component COVID-19-related health care costs, from a health

care sector perspective, associated with different intervention strategies when applied to the

adult sheltered homeless population in Boston. Panels A, B, and C show model results for R_e of

2.6, 1.3, and 0.9, respectively. Costs are derived from model-generated results and are

undiscounted. See Methods for strategy definitions.

Abbreviations: ACS, alternate care site; COVID-19, coronavirus disease 2019; ICU, intensive

care unit; M, millions; *PCR,* polymerase chain reaction; *UniversalPCR*, universal polymerase

chain reaction test for everyone; USD, United States dollars; *SxScreen,* symptom screen;

TempHousing, temporary housing.

Figure 3. Infections averted and costs of management strategies for people experiencing

sheltered homelessness in Boston during the COVID-19 pandemic over a 4-month period.

Panels A, B, and C show model results for R_e of 2.6, 1.3, and 0.9, respectively. The circle

markers represent *NoIntervention* and all strategies with universal PCR testing, which include

UniversalPCR/Hospital, and *UniversalPCR/TempHousing*. The square markers represent

strategies that are based on symptom screening, *SxScreen/PCR/Hospital* and

SxScreen/PCR/ACS. The triangle markers represent strategies that use a combination of

symptom screening and universal PCR testing, including *Hybrid/Hospital* and *Hybrid/ACS*.

Additionally, the inside shading of the markers indicates the presence of ACSs for the isolation

of individuals with symptoms or a positive test result. The dashed line represents the efficient

frontier; strategies below this line are dominated; less clinically effective and more costly, or with

a higher incremental cost per case prevented than an alternative strategy or combination of

strategies. Costs are from model-generated results and are undiscounted. Results for the

UniversalPCR/TempHousing strategy are not shown for R_e of 1.3 and 0.9. In addition to all base

case strategies, Panel A shows the *Hybrid/ACS* strategy with PCR testing every 7 days. See

Methods for strategy definitions.

Abbreviations: ACS, alternate care site; COVID-19, coronavirus disease 2019; *PCR,*

polymerase chain reaction; *UniversalPCR*, universal polymerase chain reaction test for

everyone; USD, United States dollars; *SxScreen,* symptom screen; *TempHousing,* temporary

housing.

Figure 1.

(A) $R_e = 2.6$

(B) $R_e = 1.3$

(C) $R_e = 0.9$

—— NoIntervention
– – UniversalPCR/ACS
– – UniversalPCR/Hospital
– – SxScreen/PCR/ACS
– – SxScreen/Hospital
······ Hybrid/ACS
······ Hybrid/Hospital
—— UniversalPCR/TempHousing

719 **Figure 2.**
720
721 (A) R_e=2.6

722
723

724   (B) R_e=1.3

725
726

727   (C) R_e=0.9

728
729

730 **Figure 3.**

732 (A) R_e=2.6

734　　(B)　　R_e=1.3

736　　(C)　　R_e=0.9