The COVID-19 Pandemic Vulnerability Index (PVI) Dashboard: Monitoring
county-level vulnerability using visualization, statistical modeling, and
machine learning

Authors: Skylar W. Marvel[1], John S. House[2], Matthew Wheeler[2], Kuncheng Song[1], Yihui
Zhou[1], Fred A. Wright[1,3], Weihsueh A. Chiu[4], Ivan Rusyn[4], Alison Motsinger-Reif[2]*, David M.
Reif[1]*

Affiliations:

[1] Bioinformatics Research Center, Department of Biological Sciences, North Carolina State
University, Raleigh, NC 27695, USA.

[2] Biostatistics and Computational Biology Branch, National Institute of Environmental Health
Sciences, Research Triangle Park, NC, 27709, USA.

[3] Department of Statistics, North Carolina State University, Raleigh, NC 27695, USA

[4] Veterinary Integrative Biosciences, College of Veterinary Medicine and Biomedical Sciences,
Texas A&M University, College Station, TX 77845, USA.

*Correspondence to:
Alison Motsinger-Reif
alison.motsinger-reif@nih.gov
111 TW Alexander Dr, Rall Building, Research Triangle Park, NC 27709 USA
AND
David M. Reif
dmreif@ncsu.edu
Box 7566, 1 Lampe Drive, Raleigh NC 27695, USA

Declaration of competing financial interests:

The authors declare they have no actual or potential competing financial interests.

26
27

Abstract

Background: While the COVID-19 pandemic presents a global challenge, the U.S. response
places substantial responsibility for both decision-making and communication on local health
authorities, necessitating tools to support decision-making at the community level.

Objectives: We created a Pandemic Vulnerability Index (PVI) to support counties and
municipalities by integrating baseline data on relevant community vulnerabilities with dynamic
data on local infection rates and interventions. The PVI visually synthesizes county-level
vulnerability indicators, enabling their comparison in regional, state, and nationwide contexts.

Methods: We describe the data streams used and how these are combined to calculate the PVI,
detail the supporting epidemiological modeling and machine-learning forecasts, and outline the
deployment of an interactive web Dashboard. Finally, we describe the practical application of the
PVI for real-world decision-making.

Results: Considering an outlook horizon from 1 to 28 days, the overall PVI scores are
significantly associated with key vulnerability-related outcome metrics of cumulative deaths,
population adjusted cumulative deaths, and the proportion of deaths from cases. The modeling
results indicate the most significant predictors of case counts are population size, proportion of
black residents, and mean $PM_{2.5}$. The machine learning forecast results were strongly predictive
of observed cases and deaths up to 14 days ahead. The modeling reinforces an integrated concept
of vulnerability that accounts for both dynamic and static data streams and highlights the drivers
of inequities in COVID-19 cases and deaths. These results also indicate that local areas with a
highly ranked PVI should take near-term action to mitigate vulnerability.

Discussion: The COVID-19 PVI Dashboard monitors multiple data streams to communicate
county-level trends and vulnerabilities and facilitates decision-making and communication
among government officials, scientists, community leaders, and the public to enable effective
and coordinated action to combat the pandemic.

Introduction

Defeating the COVID-19 pandemic requires well-informed, data-driven decisions at all
levels of government, from federal and state agencies to county health departments. Numerous
datasets are being collected in response to the pandemic, enabling the development of predictive
models and interactive monitoring applications (Wynants et al. 2020; ESRI 2020). However, this
multitude of data streams—from disease incidence to personal mobility to comorbidities—is
overwhelming to navigate, difficult to integrate, and challenging to communicate. Synthesizing
these disparate data is crucial for decision-makers, particularly at the state and local levels, to
prioritize resources efficiently, identify and address key vulnerabilities, and evaluate and
implement effective interventions. To address this situation, we developed a COVID-19
Pandemic Vulnerability Index (PVI) Dashboard (https://covid19pvi.niehs.nih.gov/) for
interactive monitoring that features a county-level Scorecard to visualize key vulnerability
drivers, historical trend data, and quantitative predictions to support decision-making at the local
level (Figure 1).

We assembled U.S. county- and state-level datasets into 12 key indicators across four
major domains: current infection rates (infection prevalence, rate of increase), baseline
population concentration (daytime density/traffic, residential density), current interventions
(social distancing, testing rates), and health and environmental vulnerabilities (susceptible
populations, air pollution, age distribution, comorbidities, health disparities, and hospital beds).
These 12 indicators (some of which combine multiple datasets) are integrated at the county level
into an overall PVI score, employing methods previously used for geospatial prioritization and
profiling (Bhandari et al. 2020; Marvel et al 2018). The individual data streams comprising these
indicators measure either well-established, general vulnerability factors for public health
disasters or emerging factors relevant to the COVID-19 pandemic (Centers for Disease Control
and Prevention 2015).

In developing the PVI, we performed rigorous statistical modeling of the underlying data
to enable quantitative analysis and monitoring and provide short-term predictions of cases and
deaths. Our modeling efforts directly address the discussion raised by Chowkwanyun and Reed
(2020) about racial disparities in COVID-19 case and death rates. By contextualizing factors
such as these racial disparities, correcting for socioeconomic factors, health resource allocation,
and co-morbidities, and highlighting place-based risks and resource deficits, the PVI can help
explain differences in the spatial distribution of cases. Specifically, we performed three types of
modeling efforts, all of which are regularly updated. First, epidemiological modeling on
cumulative case- and death-related outcomes provides insights into the epidemiology of the
pandemic. Second, dynamic time-dependent modeling provides similar outcome estimates as
national-level models but with county-level resolution. Finally, a Bayesian machine learning
approach provides data-driven, short-term forecasts. Herein, we describe the development of the
PVI, including the epidemiological modeling and machine-learning forecasts, and its use in an
interactive web Dashboard.

Methods

Data Streams Included in the Pandemic Vulnerability Index

To the best of our knowledge, we have assembled the most extensive set of community-
level data streams related to COVID-19. These data streams span four major domains, namely
infection rate, population concentration, intervention measures, and healthcare vulnerability.
The specific components (i.e., datasets) comprising the current PVI model are provided in a
dedicated Details page linked from the Dashboard. Supplementary Table 1 describes each
component, outlines the rationale for its inclusion, and provides a link to the associated data
source. To empower additional modeling efforts, the complete time series of all daily PVI scores
and the source data are publicly available at https://github.com/COVID19PVI/data. The software
used to generate PVI scores and profiles from these data is freely available at https://toxpi.org
(Marvel et al. 2018).

These data streams comprise both static and dynamic data, including static measures of
population concentration and healthcare vulnerabilities. Many of the data streams are from the
CDC's Social Vulnerability Index (SVI), which was developed by the Agency for Toxic
Substances and Disease Registry (ATSDR's) Geospatial Research, Analysis and Services

Program (GRASP). GRASP creates and maintains databases that help emergency response
planners and public health officials identify and map the communities most likely to need
support before, during, and after a disaster or hazard event such as the current pandemic. The
SVI has been successfully used in a variety of emergency response scenarios, including mapping
fire outbreaks to determine vulnerability metrics (Lue & Wilson 2017) and hazard mitigation
planning studies (Horney et al. 2017). The CDC's SVI uses U.S. Census data to rank each census
tract's social vulnerability based on 15 factors, including poverty, vehicle access, and housing
crowding. Additional data streams from the 2020 County Health Rankings that summarize the
prevalence of important co-morbidities and risk factors at the county level are also included
(Bhandari et al. 2020). Regarding interventions, it has been established that increased testing
rates and the implementation of social distancing are effective interventions to slow the spread of
COVID-19 and fatalities due to the disease (Wu et al. 2020). Testing rates are from the COVID
Tracking Project (The COVID Tracking Project 2020), and daily ordinal grades of social
distancing adherence are from Unacast (2020), which analyzes relative mobility compared to the
same period during the previous year using mobile device data. By anchoring movement to pre-
COVID-19 activity, this measure is applicable in both rural and urban settings, which have
different mobility patterns in general. Dynamic measures of disease spread and the number of
transmissible cases are estimated from John Hopkins University data (Dong, Du, & Gardner
2020).

PVI Calculation

For each county, the PVI, a dimensionless index score, is calculated as a weighted
combination of all data sources and represents a formalized, rational integration of information
from various domains. The score is calculated using the Toxicological Prioritization Index
(ToxPi) framework for data integration, as described in Reif et al. (2010). Briefly, the individual
factors are ranked for each county by scaling the raw value from 0 to 1. Factors for which lower
values represent higher risk (percent testing and social distancing) are reverse-scaled so that
higher values represent higher vulnerability. This allows all factors to be expressed on the same
scale (0 to 1) and removes the difficulties associated with applying different units to different
factors. The overall PVI is then calculated using a weighted sum. The choice of factors used in
the creation of the PVI score and the weights applied to them were informed by our
epidemiological modeling (described in subsequent section) as well as general knowledge of
contributors to general health morbidities.

The PVI profiles translate numerical results into visual representations as component
slices of a radar chart, with each slice representing one piece (or related pieces) of information.
For each profile, the radial length of a slice represents its rank relative to all other entities (i.e.,
counties), with a longer radius indicating higher concern or risk. The relative width (e.g.,
fraction of a full, 360° circle) of a slice indicates the contribution of its score to the overall
model. These visual profiles provide a risk assessment of the strength, relative contribution, and
robustness of the multiple data sources used in the model. Figure 2 illustrates the PVI workflow
and the results for two example counties. This type of data integration framework has been
proven effective for communicating risk prioritization and profiling information among
scientists, regulators, stakeholders, and the general public and has been featured in publications

by the U.S. National Academy of Sciences, Engineering, and Medicine (2017) and the World
Health Organization's International Agency for Research on Cancer (Loomis et al. 2018).

<u>Epidemiological Modeling</u>

The diverse array of data assembled for the epidemiological modeling that informs the
COVID-19 PVI Dashboard represents an advance over the ever-increasing number of models
related to COVID-19. To provide context and ensure that the data streams provide conclusions
and priority rankings that are broadly consistent with other epidemiological models, we
performed cross-sectional analysis of cumulative (i) cases, (ii) deaths, (iii) deaths as a proportion
of the population, and (iv) deaths as a proportion of reported cases using data current as of
8/24/2020. We emphasize that the PVI is not intended to be an epidemiological modeling tool
per se as it does not explicitly distinguish between factors of vulnerability for cases vs. deaths.
Our modeling described here is intended to anchor the components of the PVI and provide
context within the larger field of COVID-19-related epidemiological modeling. Additionally, this
modeling is not intended to provide forecasts, which are the primary focus of projection models,
as discussed in the subsequent section (see <u>Forecasting</u>).

As the initial analyses displayed evidence of count overdispersion, we performed
generalized linear modeling in R version 3.5 with the gam() procedure using a negative binomial
model with observed cumulative counts as the response (see Supplementary Tables 2–5) (R Core
Team 2018). For analyses (i), (ii), and (iv), we used log(population size) values as predictors
with estimated coefficients. For analysis (iii), we used the "offset" command to model the death
rate. Similarly, for analysis (iv), we used log(cumulative cases) as an offset to model the death
rate among cases, which may produce biased results due to regional variation in reporting rates.
It should be noted that a constant underreporting bias across counties would be absorbed into the
intercept and would otherwise produce valid coefficient estimates for the predictors. Analysis
(iv) may provide important clues about the death risk as including cases in the denominator
removes a large portion of the stochastic variation. Moreover, for all analyses, we used the
proportion of the state population that has been tested as a predictor to account for additional
sources of bias.

To anchor our efforts to previous work, we included as additional fixed predictors those
from Wu et al. (2020), who focused primarily on the effects of a $PM_{2.5}$ air pollution index using
an analysis analogous to our model (iii). Before analysis, we removed predictors with pairwise
correlation with any other predictor greater than 0.85 and predictors that would be collinear with
a series of predictors, such as the overall proportion of minority residents. For pairs exceeding
the correlation threshold, we favored predictors with the lower missingness rate (if any) or those
that are reported in other work. Dynamic predictors (i.e., those that changed substantially over
the modeled period) were incorporated using simple county averages over the March-August
period covered by the PVI. With over 3,100 counties (according to FIPS codes), most with >0
cases and deaths, the analysis can easily support the 27 to 28 final predictors used. To facilitate
comparison with previous sources, we used predictors as they are given in their source.
Accordingly, in some instances, predictors are represented as proportions and, in other instances,
they are represented as percentages.

To provide additional context, we also performed negative binomial modeling (R version
3.5 bam() with "REML" fitting) (R Core Team 2018) of daily cases up to 6/11/2020
(Supplementary Table 6), using the fixed county predictors as well as unaveraged dynamic
predictors. Due to the nature of the model, we included the two-week-lagged cumulative number
of cases as an additional predictor, as well as a smoothing spline time-dependent term to reflect a
nationwide component of risk. Although it is formally a fixed-effects model, we refer to this
model as dynamic and treat each day outcome as an independent realization, with the rate
determined by the predictors. To account for potential time-dependent latent correlation
structures, we determined standard errors for the coefficients by bootstrapping, treating each
county across all dates as an observational unit for bootstrap resampling. We also built a
dynamic version of the generalized linear model for cases and deaths as a proportion of the
population to further investigate the effects of social distancing and other predictors that change
daily. Final significance testing was based on bootstrapping to account for potential time-
dependent correlation structures.

Forecasting

For the accurate prediction of future COVID-19 cases and deaths, it is necessary to
account for the fluid nature of the data streams comprising the PVI. Accordingly, we developed a
Bayesian spatiotemporal random-effects model that jointly describes the log-observed and log-
death counts to build local forecasts. Log-observed cases for a given day are predicted using
known covariates (e.g., population density, social distancing metrics), a spatiotemporal random-
effect smoothing component, and the time-weighted average number of cases for these counts.
This smoothed time-weighted average is related to a Euler approximation of a differential
equation; it provides modeling flexibility while approximating potential mechanistic models of
disease spread. The smoothed case estimates are used in a similar spatiotemporal model that
predicts future log-death counts based on a geometric mean estimate of the estimated number of
observed cases for the previous seven days as well as the other data streams. The Dashboard
shows the resulting county-level predictions and corresponding confidence intervals (Fig. 1).
Details of the model are provided in the Supplemental Information.

Dashboard Technical Details

We used the ArcGIS JavaScript API (v4.13) (ESRI 2020) and custom PHP and JavaScript
files to build the Dashboard web application. The API is used to overlay county borders,
COVID-19 count data, and PVI model images on a Basemap or custom WebMap. County
boundary data is from a feature service, while all other data is stored in an SQL database. PVI
model images are base64-encoded scalable vector graphics (SVGs) rendered as inline images.
The custom scripts controlling the Dashboard's functionality optimize the efficiency of HTTP
requests and other computational overhead to promote real-time interactivity. The Dashboard is
hosted by the National Institute of Environmental Health (NIEHS) Office of Scientific
Computing, which provides high-availability HTTPS load balanced with NGINX and a secure
environment for web applications. Automated data updates are pushed to the public servers
daily, and the daily update process is paralleled on a private server to permit independent data
integrity assessment.

Complete, continually-updated documentation is available from a link on the Dashboard to
a Quick Start Guide that introduces the PVI and Dashboard tools
(https://www.niehs.nih.gov/research/programs/coronavirus/covid19pvi). A Details page provides
additional in-depth information
(https://www.niehs.nih.gov/research/programs/coronavirus/covid19pvi/details/index.cfm). The
Supplemental Information and Figure S2 explain in detail the main features. The Dashboard is
also mirrored as an iFrame on the CDC's COVID Data Tracker, under the Unique Populations
tab (https://covid.cdc.gov/covid-data-tracker/).

Results

PVI and Vulnerability

The summarization and communication goals of the PVI and the corresponding scorecards
are human-centric and designed to convey and distill high-dimensional complex data. Our PVI
model communicates an integrated concept of vulnerability that accounts for both dynamic
(infection rate and interventions) and static (community population and healthcare
characteristics) drivers. To gauge the association of daily PVI scores versus observed death
outcomes, we assessed the rank-correlation between the overall PVI and the key vulnerability-
related outcome metrics of cumulative deaths (Figure S1A), population adjusted cumulative
deaths (Figure S1B), and the proportion of deaths from cases (Figure S1C). The Spearman Rho
values for the PVI (from March 15 to August 12th) versus outcomes 1, 7, 14, 21, and 28 days
ahead of a given day are displayed. All daily rank-correlation estimates were highly
significant (p-values < 5.1E-14). The mean Rho values increase with a longer time horizon (blue
text on Supplementary Figures 1A, 1B, 1C) and thus a highly-ranked PVI provides evidence that
local actions should be taken to mitigate undesirable outcomes.

Epidemiological Modeling

Supplementary Tables 2-7 display the regression coefficients produced with generalized linear
modeling in a cross-sectional analysis of county cases and deaths up to 8/24/2020. As expected,
the most significant predictor for the case count is population size (p<1E-300). The next most
significant predictors associated with case counts are the proportion of black residents (p=1.28E-
61) and mean $PM_{2.5}$ (p=9.08E-32), followed by Insurance percent coverage (positively
associated, p=1.51E-27) and proportion of Hispanic residents (p=6.92E-20) (Supplementary
Table 2). In addition, the proportion of the population tested for SARS-Cov-2 infection is
associated with case counts (p=3.39E-13), which we attribute to statewide responses to emerging
infection clusters. In this cumulative analysis, social distancing and travel-related predictors were
significant even though they represent aggregate values per county. For deaths as a proportion of
the county population (Supplementary Table 4), the same predictors are highly significant,
although the Insurance percent coverage is much less significant than for cases (p=4.48E-05) .
We note that cases and deaths per county population represent overall societal risk, for which
vulnerability measures are relevant. The rank correlation coefficients for the PVI vs. the fitted
values for the number of cases and the death rate are 0.54 and 0.55, respectively ($p<10^{-16}$ for
both).

Our analysis of the proportion of deaths per cases is enlightening, despite the previously
noted caveat regarding potential bias due to testing variation. We note that a true case fatality
rate potentially involves very different predictors than a case rate model, and deaths per county
population are also closely tied to case rates. Our modeling (Supplementary Table 5) shows that
after multiple test corrections, state testing rates are no longer significant in comparison to
multiple-testing thresholds (p=0.036), which is consistent with the hypothesis that testing
uncovers cases but does not predict case fatalities. The proportion of black residents (p=2.64E-
12) and mean $PM_{2.5}$ (p=1.92E-04) were significant, but less so than in the deaths/population
model. Deaths/reported cases were found to be associated with the proportion of owner-occupied
residences (p=9.31E-07) and inversely associated with median house value (p=0.000863). Both
measures tend to be associated with wealth, but the relationship is complicated by the fact that
high housing prices impede home ownership.

The results of the dynamic model for cases (Supplementary Table 6) with bootstrapped p-
values had much stronger significance than the results of the cumulative case model, which we
attribute to the dynamic model's ability to account for additional sources of variation due to the
use of lagged case counts, a smooth time-dependent term to account for national trend, and the
inclusion of daily dynamic predictors. Again, the most significant predictors are the population
size ($p<1E-300$), the proportion of black residents ($p<1E-300$), the two-week-lagged cumulative
number of cases as a predictor of current cases ($p<1E-300$), and mean $PM_{2.5}$ ($p<1E-300$). We
also ran the analogous model for deaths/population size (Supplementary Table 7) and the same
predictors were found to be highly significant. In summary, the dynamic versions of the
generalized linear model reinforce and amplify the conclusions from the previous cumulative
models. However, the models are not designed to perform forecasting, which can be viewed as
essentially a machine learning exercise. For forecasting, careful cross-validation approaches can
be used to assess the accuracy of the results.

The most consistent significant predictors for COVID-19 related case rates and mortality
are the proportion of black residents and the mean $PM_{2.5}$, reinforcing conclusions from previous
reports (Dong, Du, & Gardner 2020). A one-percentage-point increase in the proportion of black
residents is associated with a 2.9% increase in the COVID-19 death rate. The effect of a 1 g/m^3
increase in $PM_{2.5}$ is associated with an approximately 14.5% increase in the COVID-19 death
rate, which is at the high end of a previously reported confidence interval from a report in late
April 2020 (Wu et al. 2020) when deaths had reached 38% of the total as of June 2020. We find
that these effects persist when including numerous additional predictors and correcting for
factors such as socioeconomic status, housing density, and comorbidities. Moreover, the effects
persist for a range of response values, including cumulative (i) cases, (ii) deaths, (iii) deaths as a
proportion of the population, and (iv) deaths as a proportion of reported cases (Supplemental
Tables 2-5). These results strongly suggest the important role of structural variation by location,
which results in drastic health disparities. The results of the dynamic version of the generalized
linear model (Supplemental Tables 6-7) support the importance of social vulnerability indicators
and may be viewed as a sensitivity test that the impact of social distancing and other dynamic
measures do not alter the significance of many of the social vulnerability indicators.

Functional Power-Adjusted Relative Rate Model Forecasting

Data-driven machine learning was implemented for near term predictions of case and death
outcomes at a local level. The resulting county-level predictions and corresponding confidence
intervals for the next seven days are shown in the Dashboard 'Predictions' element. Additional
details of the model and implementation are included in the Supplemental Information. The
accuracy of the predictions were assessed by calculating the Pearson Correlation (Rho) of the
predicted values versus the observed from June 3 through August 5, 2020. To avoid weekend-
related reporting variation in cases and deaths, both predictions and observed cases/deaths were
summed to weeks based on Wednesday through Tuesday. The accuracy of county-level
predictions of Covid19-cases and -deaths were assessed by calculating the Pearson Correlation
(Rho) of weekly predicted values versus weekly actual values across U.S. counties. For the 10
forecasts of Covid19 cases made each Wednesday from June 3rd through August 5th, the median
Rho for 1-week out was 0.96. For deaths, the 1-week out median Rho for these 10 periods was
0.88. Summary Rho distributions are shown in Supplemental Figure 3, and scatterplots for all
counties for the most recent analysis week are shown in Supplemental Figure 4.

Dashboard Features

The interactive visualization within the PVI Dashboard communicates factors underlying
vulnerability and empowers community action. On loading, the Dashboard displays the top 250
PVI profiles (by rank) for the current day. The data, PVI scores, and predictions are updated
daily, and users can scroll through historical PVI and county outcome data. Individual profiles
are an interactive map layer with numerous display options and filters that allow sorting by
overall score and combinations of slice scores, clustering by profile similarity (i.e., vulnerability
"shape"), and searching for counties by name or state. Any user-selected county overlays the
summary Scorecard and populates the surrounding panels with county-specific information
(Figure 1). Scrollable panels on the left include plots of vulnerability drivers relative to their
nationwide distribution across all U.S. counties, with the location of the selected county
delineated. The panels across the bottom of the Dashboard report cumulative county numbers of
cases and deaths; timelines of cumulative cases, deaths, PVI scores, and PVI ranks; daily
changes in cases and deaths for the most recent 14-day period (a measure commonly used in
reopening guidelines); and predicted cases and deaths for a seven-day forecast horizon.

Taken together, the Dashboard features support the interactive evaluation and visualization
of current data for localities while providing context with respect to all U.S. counties. Full time
series of case, death, and PVI trends enable the examination of the track records of counties of
interest as well as the comparison of trajectories for peer, or comparable, counties in terms of
varying success with specific interventions. For example, using Orleans County (home to New
Orleans, LA) as an exemplar, we employed the multi-criteria filtering capabilities in the
Dashboard to find a peer county for comparison. By bounding the PVI to similar ranges of
vulnerability drivers (i.e., slices) for population mobility, residential density, and population
demographics, we identified a subset of candidate counties and ultimately chose Clayton County,
GA to illustrate the effects of dramatic differences in public action/interventions. Figure 3 shows
detailed results for the two counties, which have similar baseline vulnerabilities but implemented
divergent interventions at the outset of the pandemic. Specifically, pronounced differences in

intervention measures (social distancing and testing) are associated with varying dynamics of the
infection rates in these counties, as visualized through the considerable differences in magnitude
of the blue (intervention-related) and red (infection-related) slices over time. Note that all data
streams are scaled so that a larger slice indicates increased vulnerability (e.g., the larger blue
slices represent less adherence to social distancing and lower testing rates). As visualized in
Figure 3, the PVI rank for Orleans County improves over time (i.e., follows a downward
rank/percentile path), effectively blunting the curve caused by the accelerated increase in the
number of cases through early interventions. There is no similar positive change for Clayton
County, reflecting differences due to varying interventions in the two areas. In this way, the PVI
Dashboard enables customized empirical comparisons and evaluations across peer counties.

Discussion

Numerous expert groups have coalesced around a general roadmap to address the current
COVID-19 pandemic that comprises (i) reducing the spread through social distancing, (ii)
gradually easing restrictions while monitoring for resurgence and healthcare overcapacity, and
(iii) eventually moving to pharmaceutical interventions. However, the responsibility for
navigating the COVID-19 response falls largely on state and local officials, who require data at
the community level to make equitable decisions on allocating resources, caring for vulnerable
sub-populations, and enhancing/relaxing social distancing measures. The goal of the COVID-19
PVI Dashboard is to empower informed actions to combat the pandemic from the local to the
national levels on multiple time scales. The Dashboard accomplishes this goal by combining
underlying COVID-19-specific structural vulnerabilities with dynamic infection and intervention
data at the county level to produce an integrated concept of vulnerability that can inform
decision-making on actions at the local level.

Furthermore, the general public must embrace interventions for them to be effective, and
interactive visualization is a proven approach to communication among diverse audiences. The
PVI Dashboard provides interactive, visual profiles of vulnerability atop an underlying statistical
framework that enables the comparison of counties by clustering and the evaluation of the PVI's
sensitivity to component data. The Dashboard's county-level Scorecards illustrate both overall
vulnerability and the components driving it. A key utility of a public-facing, interactive
dashboard is that decision-makers can point to it for support, thus promoting transparency and
public buy-in for actions taken in the interest of public health. Example use cases include the
priority distribution of medical resources such as hospital beds, targeted community outreach
activities, and the establishment of contact-tracing mechanisms. Eventually, the PVI could be
used to support the priority distribution of vaccines to highly-vulnerable communities.

The modeling efforts presented here support decision-making in multiple ways. The
epidemiological modeling enables testing the impact of changes in dynamic interventions, such
as changes in social distancing, and the forecasting efforts support short-term resource allocation
decisions, such as hospital staffing and the distribution of supplies. These forecasts also help
communicate the trends that are part of the CDC's reopening criteria (Centers for Disease
Control and Prevention 2020), such as whether interventions and local government actions
translate into flattened curves. The PVI score itself constitutes an integrated indicator of
vulnerability that is strongly associated with mortality outcomes in the near-to-medium term.

The overall PVI score highlights highly-ranked counties that should consider taking local actions
or receive targeted help to mitigate undesirable outcome trends. The *Timelines* panel of the
Dashboard illustrates observed county-level changes over time to answer questions such as
"Have we flattened the curve?" while the *Predictions* panel presents statistically robust forecasts
that consider all of the data to answer the question, "What's next?". Overall, the COVID-19 PVI
Dashboard can help facilitate decision-making and communication among government officials,
scientists, community leaders, and the public to enable more effective and coordinated action to
combat the pandemic.

A growing number of risk factors have been highlighted in the rapidly expanding scientific
studies on COVID-19. Infectious disease epidemiology is rapidly evolving, and new risk factors
and environmental variables (e.g., weather conditions) are continually being discovered. From
the early identification of older individuals' vulnerability (Verity et al. 2020) to the dramatic
racial disparities that have been more recently highlighted (Coughlin et al. 2020), there is
mounting evidence that socially marginalized populations are suffering disproportionally. While
this is not unique to the ongoing pandemic, it is clear that pandemic vulnerability and response
are dynamic and differ across communities (Quinn & Kumar 2020). The COVID-19 PVI
Dashboard provides contextualized local summaries of differences in vulnerability and highlights
racial disparities, even when adjusting for multiple covariates. The Dashboard's presentation of
information in a relative sense enables the fair comparison of communities of different sizes to
support prioritization decisions. Further, the PVI visualization is a human-centric and
communicates how particular communities' drivers (i.e., slices) differ in terms of their
contribution to overall vulnerability. This visualization promotes transparency by clarifying the
judgments and trade-offs entering into such an assessment while maintaining a direct link to the
underlying quantitative data.

COVID-19 will continue to present public health challenges into the foreseeable future. By
integrating vulnerability information (both historical and forward-looking), the Dashboard
supports key decision-making for managing the ongoing pandemic. We will continue to update
the data streams combined to calculate the PVI and will add additional variables as evidence of
new risk factors and potential drivers of vulnerability emerge and supported by publicly
available data. We will also continually develop software tools so that people can actively build
their own models and will update the modeling efforts as needed. Combating endemic diseases
requires long-range thinking, informed action, and political will, and we offer the COVID-19
PVI Dashboard as an interactive monitoring tool to support these sustained efforts.

Acknowledgments

We would like to thank the IT and web services staff at NIEHS for their help and support,
as well as JK Cetina and DJ Reif for useful technical input and advice. **Funding:** This work was
supported by P42 ES027704, P30 ES029067, P42 ES031009, and P30 ES025128 and by
intramural funds from the National Institute of Environmental Health Sciences (Z ES103352-01).
Author contributions: *Concept and design:* SM, FW, WC, IR, AMR, DR. *Acquisition,*
analysis, or interpretation of data: All authors. *Drafting of the manuscript:* All authors. *Critical*
revision of the manuscript for important intellectual content: All authors. *Statistical*
analysis: JH, MW, KS, YZ, AMR, DR. *Obtained funding:* WC, IR, DR, AMR. **Competing**

interests: Authors declare no competing interests. **Data and materials availability:** All data is
available through links in the main text or the supplementary materials.

References

Bhandari S, Lewis P, Craft E, Marvel SW, Reif DM, Chiu WA. 2020. HGBEnviroScreen:
enabling community action through data integration in the Houston–Galveston–Brazoria Region.
Int J Environ Res Public Health 17(4):1130, PMID: 32053902, https:// doi:
10.3390/ijerph17041130.

Centers for Disease Control and Prevention (CDC). 2015. Planning for an emergency: strategies
for identifying and engaging at-risk groups. A guidance document for emergency managers.
Atlanta, GA: CDC.
https://svi.cdc.gov/Documents/Publications/SVI_Community_Materials/atriskguidance.pdf
[accessed 31 August 2020].

Centers for Disease Control and Prevention. 2020. Reopening Guidance for cleaning and
disinfecting public spaces, workplaces, businesses, schools, and homes.
https://www.cdc.gov/coronavirus/2019-ncov/community/reopen-guidance.html [accessed on 31
August 2020].

Chowkwanyun M, Reed AL. 2020. Racial health disparities and Covid-19–caution and context.
N Engl J Med 383:202-203, PMID: 32374952, http://doi: 10.1056/NEJMp2012910.

Coughlin SS, Moore JX, George V, Johnson JA, Hobbs J. 2020. COVID-19 among African
Americans: from preliminary epidemiological surveillance data to public health action. Am J
Public Health 110(8):1157-1159, PMID: 32639918, https://doi:10.2105/AJPH.2020.305764.

Dong E, Du H, Gardner L. An interactive web-based dashboard to track COVID-19 in real time.
Lancet Infect Dis 20(5):533-534, PMID: 32087114, https://doi:10.1016/S1473-3099(20)30120-1.

Esri. 2020. ArcGIS Javascript API, version 4.13. https://www.esri.com/en-us/home [accessed on
31 August 2020].

Esri. 2020. COVID-19 GIS Hub. https://coronavirus-resources.esri.com [accessed 31 August
2020].

Horney J, Nguyen M, Salvesen D, Dwyer C, Cooper J, Berke P. 2017. Assessing the quality of
rural hazard mitigation plans in the southeastern United States. J Plann Educ Res 37(1):56-65,
PMID: 24180063, https://doi: 10.5055/jem.2013.0138.

Loomis D, Guyton K, Grosse Y, Ghissasi FE, Bouvard V, Benbrahim-Tallaa L, et al. 2015.
Carcinogenicity of lindane, DDT, and 2,4-dichlorophenoxyacetic acid. Lancet Oncol. 16(8):891-
892, PMID: 26111929, https://doi:10.1016/S1470-2045(15)00081-9.
482

483   Lue E, Wilson JP. 2017. Mapping fires and American Red Cross aid using demographic
484   indicators of vulnerability. Disasters 41(2):409-26, PMID: 27170545, https://doi:
485   10.1111/disa.12198.
486
487   Marvel SW, To K, Grimm FA, Wright FA, Rusyn I, Reif DM. 2018. ToxPi Graphical User
488   Interface 2.0: Dynamic exploration, visualization, and sharing of integrated data models. BMC
489   Bioinf 19(1):1-7, PMID: 29506467, https://doi: 10.1186/s12859-018-2089-2.
490
491   National Academy of Sciences, Engineering, and Medicine. 2017. Using 21st century science to
492   improve risk-related evaluations. Washington, DC: National Academies Press.
493
494   Quinn SC, Kumar S. 2014. Health inequalities and infectious disease epidemics: a challenge for
495   global health security. Biosecur Bioterror 12(5):263-273, PMID: 25254915,
496   https://doi:10.1089/bsp.2014.0032.
497
498   R Core Team. *R: A language and environment for statistical computing*. 2019, R Foundation for
499   Statistical Computing: Vienna, Austria. https://www.R-project.org/ [accessed on 31 August
500   2020].
501
502   Reif DM, Martin MT, Tan SW, Houck KA, Judson RS, Richard, AM et al. 2010. Endocrine
503   profiling and prioritization of environmental chemicals using ToxCast data. Environ Health
504   Perspect 118(12):1714-1720, PMID: 20826373, https://doi:10.1289/ehp.1002180.
505
506   Sattar N, McInnes IB, McMurray JJV. 2020. Obesity is a risk factor for severe COVID-19
507   infection: multiple potential mechanisms. Circulation 142(1):4-6, PMID: 32320270,
508   https://doi:10.1161/circulationaha.120.047659.
509
510   The COVID Tracking Project. 2020. https://covidtracking.com [accessed 31 August 2020] .
511
512   Unacast. 2020. Social Distancing Scoreboard. https://www.unacast.com/covid19/social-
513   distancing-scoreboard [accessed on 31 August 2020].
514
515   Verity R, Okell LC, Dorigatti I, Winskill P, Whittaker C, Imai I, et al. 2020. Estimates of the
516   severity of coronavirus disease 2019: a model-based analysis. Lancet Infect Dis 20(6):669-677,
517   PMID: 32240634, https://doi:10.1016/S1473-3099(20)30243-7.
518
519   Wu J, Tang B, Bragazzi NL, Nah K, McCarthy Z. 2020a. Quantifying the role of social
520   distancing, personal protection and case detection in mitigating COVID-19 outbreak in Ontario,
521   Canada. J Math Ind 10(1):15, PMID: 32501416, https:// doi:10.1186/s13362-020-00083-3.
522
523   Wu X, Nethery RC, Sabath BM, Braun D, Dominici F. 2020b. Exposure to air pollution and
524   COVID-19 mortality in the United States. medRxiv preprint, PMID: 32511651,
525   https://doi.org/10.1101/2020.04.05.20054502.
526

527   Wynants L, Van Calster B, Bonten MM, Collins GS, Debray TP, De Vos M, et al. 2020.
528   Prediction models for diagnosis and prognosis of covid-19 infection: systematic review and
529   critical appraisal. BMJ 369:m1328, PMID: 32265220, https://dx.doi.org/10.1136/bmj.m1328.
530
531

538
539

Figure 1. Dashboard displaying map view with PVI Scorecard and associated data for a selected county. The Dashboard allows U.S.-wide navigation to area(s) of interest. The filter is set to display the top 250-ranked (i.e., most vulnerable) PVI profiles for the selected date (displayed in the upper left panels for each data layer). The Scorecard displayed shows the contribution of each indicator (slice) for Florence, SC, which is in a cluster of high-PVI counties across the rural Southeast U.S. The scorecard summarizes the overall PVI score and rank compared to all 3,142 U.S. counties. In the graphical profile, longer slices indicate higher vulnerability driven by a particular indicator, with corresponding indicator-wise scores (0 = minimum; 1 = maximum) provided in the lower portion of the Scorecard. The scrollable score distributions at left compare the selected county PVI to the distributions of overall and slice-wise scores across the U.S. The panels below the map are populated with county-specific information on observed trends in cases and deaths, observed numbers for the selected date, historical timelines (for cumulative cases, cumulative deaths, PVI, and PVI rank), daily case and death counts for the most recent 14-day period, and a 7-day forecast of predicted cases and deaths. The information displayed for both observed COVID-19 data and PVI layers is scrollable back through March 2020. Documentation of additional features and usage, including advanced options (accessible via the collapsed menu at the upper left), is provided in a Quick Start Guide (linked at the upper right corner).

Figure 2. Translation of data into PVI profiles. Information from all 3,142 U.S. counties is
translated into PVI slices. The illustration shows how *Air Pollution* data (average density of
PM$_{2.5}$ per county) are compared for two example counties. The county (County Y) with the
higher relative measurement has a longer *Air Pollution* slice than the county (County X) with a
lower measurement. This procedure is repeated for all slices, resulting in an integrated, overall
PVI profile.

569
Figure 3. PVI data from the PVI Dashboard are shown for Orleans County, LA (left) and
Clayton County, GA (right). The PVI profiles from March 15th, 2020 are shown above the
timelines for each county. Comparing Orleans Parish (County) to Clayton County, they have
similar ranks for Population Concentration (green slices) and Health and Environment (purple
slices). For Intervention Practices (blue slices), differences can be seen; Orleans County's
adherence to social distancing measures and increased COVID-19 testing (reverse-scored and
indicated by smaller blue slices) is visualized compared to Clayton. The changes in overall PVI
rank across the timeline of the pandemic are shown. While the trajectory of new cases was
blunted in Orleans County, it continued increasing in Clayton. Note that both counties observed
spikes in the Cases trajectory as Social Distancing measures were relaxed at the end of June.
580