1   **Examining the Interplay between Face Mask Usage, Asymptomatic Transmission, and Social**

2   **Distancing on the Spread of COVID-19**

3   **Short title: Face mask impact on prevention of COVID-19**

Adam Catching[1,2 §], Sara Capponi[3,4 §], Ming Te Yeh[1], Simone Bianco[3,4 *], Raul Andino[1 *]

[1] Department of Microbiology and Immunology, University of California, San Francisco, San Francisco,

CA 94158

[2] Graduate Program in Biophysics, University of California, San Francisco, San Francisco, CA 94158

[3] Industrial and Applied Genomics, AI and Cognitive Software, IBM Almaden Research Center, San Jose,

CA, 95120

[4] Center for Cellular Construction, San Francisco, CA 94158

24 § authors contributed equally to the manuscript

26 * corresponding authors: sbianco@us.ibm.com, Raul.Andino@ucsf.edu

Abstract

The COVID-19 global crisis is facilitated by high virus transmission rates and high percentages of

asymptomatic and presymptomatic infected individuals. Containing the pandemic hinged on combinations

of social distancing and face mask use. Here we examine the efficacy of these measures, using an agent-

based modeling approach that evaluates face masks and social distancing in realistic confined spaces

scenarios. By explicitly considering different fractions of asymptomatic individuals, as well as a realistic

hypothesis of face mask protection during inhaling and exhaling, we find that face masks are more

effective than social distancing in curbing the infection. Importantly, combining face masks with even

moderate social distancing provides optimal protection. The finding that widespread usage of face masks

limits COVID-19 outbreaks can inform policies to reopening of social functions.

53   Keywords: epidemic, virus, agent-based simulation

Author summary:

The COVID-19 outbreak has created an enormous burden on the worldwide population. Among the various ways of preventing the spread of the virus, face masks have been proposed as a main way of reducing transmission. Yet, the interplay between the usage of face mask and other forms of Non-Pharmaceutical Intervention is still not completely clear. In this paper we introduce a stochastic individual-based model which aims at producing realistic scenarios of disease spread when mask wearing with different inward and outward efficacy and social distancing are enforced. The model elucidates the conditions which make the two forms of intervention synergistic in preventing the spread of the disease.

Introduction

The COVID-19 outbreak has caused a catastrophic mortality and economic damage around the world. The causative agent of COVID-19, SARS-CoV-2, is an airborne pathogen that can be transmitted between humans through droplets and aerosols that can travel 1–8 meters[1]. The virus is transmitted by both symptomatic and asymptomatic individuals. COVID-19 causes severe symptoms that require hospitalization, as well as severe long-term sequels and death. Accordingly, the outbreak has seriously impacted healthcare systems around the world[2]. One of the major difficulties to contain the COVID-19 pandemic has been detection of infected asymptomatic or pre-symptomatic individuals, who are estimated to be responsible for as much as 95% of all transmissions[3][4]. As these individuals carry and spread the virus without manifesting any sign of the disease, they represent a crucial variable in managing the outbreak.

In the absence of an effective vaccine or antiviral, most countries have implemented non-pharmaceutical interventions (NPIs) to curb the spread of COVID-19[5]. These include closure of schools, workplaces, churches, offices, factories and other social venues, while encouraging preventative measures ranging from maintaining social distancing (SD, usually 2 m/6 ft between individuals) to total quarantine and societal lockdowns. These measures aim to reduce the effective contact rate of the population, which in turn decreases the disease reproductive number R_0. These NPIs limit the epidemic, but they present important drawbacks. Total lockdown can only be implemented for short periods, due to

its severe impact on the social fabric and economy of a country. Meanwhile, essential workers remain vulnerable to infection and transmission due to the frequent encounters with infected, often asymptomatic, individuals. Because of their social and economic impact, lockdowns and SD measures have been lifted in some countries, leading to reactivation of virus spread and ensuing increased morbidity and mortality.

Face masks covering the nose and mouth area also provide a level of filtration that blocks virus transmission to a certain extent[6-8]. Masks prevent the spread of droplets and aerosols generated by an infected individual[1], reducing viral transmission by 95%. Uninfected individuals wearing a mask are about 85% protected against infection[9]. Masks may be more effective to control the source of infectious virus because they prevent the larger expelled droplets from being converted into smaller droplets that can travel farther. Accordingly, face masks reduce the spread of influenza[10] and coronaviruses[11,12].

In the past, several papers have used theoretical models to study how efficacious mask wearing is in avoiding the spread of airborne viruses. In 2010 at least two studies were focused on the effect of face mask usage to contrast the diffusion of Novel Influenza A (H1N1)[13,14]. Although these works are not on COVID-19 or another Coronavirus disease, they underline the necessity of developing more accurate models to describe similar diseases and reinforce their results in the light of new diseases. Specifically on COVID-19, the efficacy of mask wearing has recently been studied using an ordinary differential equations (ODE) model[15], which considered also a varying percentage of asymptomatic individuals, compliance with mandate to wear masks, and a different inward and outward efficacy of protection. However, the authors consider face masks as the sole preventative, excluding additional NPIs like SD or shelter in place.

Additionally, the problem has been studied in several papers using ABMs, each being focuses on a specific part of the problem. In some cases, the presence of asymptomatic infected individuals was not considered[16,17]. In other studies, the efficacy of wearing masks was not analyzed when combined with other NPIs[18]. Finally, the difference in inward and outward protection given by a face mask was often neglected or parametrized with a single value[19]. A rather comprehensive data-driven investigation of all these effects has been performed by Hoertel et al.[20]. However, their results are specific to the country of

France, and the high dimensionality of the parameter space makes it difficult to disentangle the effect of

the various interventions.

In our current study, we analyzed the relative efficacy of wearing face masks and/or exercising

SD to reducing the spread of COVID-19 in the presence of asymptomatic individuals. Face masks have

distinct inward and outward protection, parameterized using a Gamma function (see Methods). Through

stochastic computer simulations of infection spread, we modeled realistic outbreak scenarios and found

that SD only yields beneficial effects if accompanied by a widespread population lockdown. In contrast,

wearing face masks is a highly effective strategy to reduce the spread of infection. Our results are

general, and suggest that, even when a large fraction of infected individuals is asymptomatic, mask

wearing is the most effective strategy to control virus spread and alleviate the impact of COVID-19

outbreak, particularly when combined with conditions of partial SD compatible with the function of society.

Results

Stochastic model description and calibration. We developed an agent-based model (ABM) to examine

the effectiveness of wearing masks and SD on the rate of infections and viral spread during the

pandemic. Unlike ODE models of the spread of COVID-19 disease[15,21], ABMs are stochastic models that

allow the description of non-homogeneous distributions of agents that act individually[12,22]. In ABMs, each

individual behaves dynamically and independently in response to environmental changes[23] according to

rules that describe their interactions. To model the COVID-19 pandemic, we used a SEAIR system in

which each agent represents an individual who can be susceptible (S), exposed (E), asymptomatically

infected (A), symptomatically infected (I), or recovered (R). To obtain a realistic model of virus spread, we

chose parameters that describe the spread of SARS-CoV2: transmission events occur through contacts

made between susceptible and infectious individuals in close proximity (distance $\leq 2r$, Fig. 1A), and

exposed individuals undergo an incubation period of 5.1 days to become infectious (Fig. 1, spheres with

red border). This incubation time represents the interval required to increase viral loads to levels sufficient

for transmission[3][4,24]. We assume that recovered individuals (Fig. 1, spheres with black border), who

resolve the infection, cannot be re-infected or infect others, which is a reasonable assumption for the

duration of our models (45 days).

To define the probabilities of infection, we used reported COVID-19 parameters[24]. The probability

of transmission follows a Γ (*gamma)* distribution (see Methods) whose shape is described by a constant

(α), estimated to be 0.25. Wearing masks reduces this probability (Fig. 1, probability of infection). To

estimate the protective effect of masks, we used parameters determined for FDA-approved surgical

masks, whose efficacy has been experimentally verified to inhibit virus transmission[10]. Based on previous

studies (Leung et al.)[9], we assumed that, if an infectious individual wears a mask, the effective probability

of transmission is reduced by a factor of 0.05. If a susceptible individual wears a mask, α is reduced by a

factor of 0.15 (Fig. 1A). Our assumption that wearing masks is more effective to reduce transmission than

to prevent getting infected is supported by experimental data[25]. However, if both infectious and

susceptible individuals wear masks the probability of transmission is the product of these probabilities

(0.0075) and, thus, sufficiently low such that transmission is effectively null. We calibrated our model by

running simulations without any individuals wearing a face mask or practicing SD and considering that

50% of infected individuals are asymptomatic (Movie 1). In this way, we determined the simulation

parameters, such as velocity and density of individuals, to obtain a value of $R_0 = 2.5$, consistent with what

was reported early on as the infection rate of the epidemic in Wuhan[9].

Percentage of population wearing masks determines the daily infection incidence and cumulative

number of cases. We started each simulation with one individual being infected and all others

susceptible and assuming that 50% of COVID-19 infections are asymptomatic (or pre-symptomatic). Non-

infected and asymptomatic individuals circulate in the population without any restriction (i.e., they do not

isolate themselves or become hospitalized) (see Methods). In contrast, symptomatic individuals no longer

move after 12 hours of the symptom onset, simulating hospitalization or self-isolation. Thus, symptoms

are assumed to manifest after the incubation time of 5.1 days (Fig. 2B). Using these assumptions and

model calibration, we carried out a set of simulations in which we gradually increased the percentage of

individuals in the population that wear masks. Individuals, who are assigned randomly to wear a mask at

the beginning of the simulation, keep on the mask for the entire duration of the simulation. Increasing the

fraction of the population wearing face masks has a highly significant effect on the spread of the virus

(Fig. 1B, 1C and Movie 2). Mask wearing reduces the cumulative number of infected individuals at the

end of the simulation (Fig. 1B). Strikingly, this effect negatively correlates with the fraction of the

population using masks without any other intervention, suggesting that masks are effective for reducing

the overall number of cases (Fig. 1B). The description of the dynamics of infection generated by this

model is consistent with previous clinical studies[9,25] and highlights the benefit of wearing masks (Fig. 1C,

Movie 1).

If the daily incidence surpasses the treatment capacity, it will overwhelm the healthcare system

with detrimental consequences for medical care of infected individuals and increased mortality and

morbidity. Thus, we examined the effects of face masks on the daily incidence of infection over time (Fig.

2B). If all individuals move freely and randomly interact with others (i.e., 0% SD), the rate of daily infection

through the population depends on the percentage of individuals wearing masks. The average number of

new daily infection varies considerably, according to the number of individuals wearing face masks. When

0% of individuals wear face masks (Fig. 2b, red line), the number of daily infections peaks sharply at day

27 with a maximum of 54.8 ± 18.1 infections per day for a population of 500 individuals. No additional

infections are observed after day 39, when the entire population (100%) has been infected. Thus, without

any intervention, the infection quickly reaches every individual through an epidemic characterized by a

very sharp infection peak. Note that, in these original models, we assume all individuals recover after ~11

days post-infection and cannot be re-infected. When 40% of all individuals wear masks, the number of

individuals infected at any given day is reduced by approximately 30% (maximum of infected individuals

35.1 ± 12.8, Fig. 2A, blue line), both flattening the curve and extending the duration of the outbreak by

more than 10 days, with a peak maximum at day 32. By day 52, an average of 472 individuals has been

infected, and none of the individuals is infectious or exposed any longer. Even more significantly, if 80%

of the population wear masks, we observed a significant flattening of the curve, with a substantial

reduction in the maximum number of infected individuals per day, 5.9 ± 6.8 (Fig. 2A, green line), and the

number of new infected individuals reached zero by day 57.8 ± 35.0. Thus, the shape of the outbreak

changes from a curve characterized by a sharp peak when no intervention is considered to a broader

peak when 80% of the individuals wear masks. By increasing the percentage of the individuals wearing

masks, the number of newly infected individuals per day substantially decreases, which will reduce

mortality and morbidity. Moreover, since the use of masks eliminates the sharp peak that characterizes

SARS CoV2 epidemics, the overall impact of the outbreak on the health system is alleviated. These

results highlight the importance of widespread mask wearing as an effective intervention that can be

implemented as soon as the first cases are reported.

Effect of social distancing on viral infection spread. Next, we evaluated the effect of SD in shaping

the spread of infection. Practicing SD does not affect the probability of infection but reduces the chances

of encounters leading to transmission. Thus, SD was introduced into the model by limiting the proportion

of individuals in the population that move freely in the field. This simulates a scenario in which a given

proportion of individuals in the population quarantine or shelter-in-place, thereby reducing the probability

of contacts and transmission (Fig. 2A). As reported, if a percentage of individuals practice SD, even

without any other non-pharmacological intervention, the number of infections and daily infection rate are

reduced (compare Fig. 2B, SD 0, 40, and 80%, red line). However, if no one wears masks but 40% of

individuals practice SD, we observed a very small effect in the shape of the infection curve (Fig. 2B, the

peak maximum decreases from 51.0 ± 13.3 to 45.7 ± 5.9). When 80% of the population practices SD, a

more significant reduction in the number of new daily infections was observed (Fig. 2B, 80%). Notably,

the shape of the new daily infection curve broadens considerably upon increasing the percentage of

individuals wearing face masks (Fig. 2B, compare blue and green curves with red curve). For instance, if

80% of the population wears masks with 40% SD, the peak maximum decreases to one-tenth (from 51.0

± 13.3 to 5.7 ± 5.9) and is slightly delayed (Fig. 2B). At 80% SD, the peak of new daily infections is no

longer observed, and the number of new infected individuals per day averages 1.6 ± 1.3 (Fig. 2B). Thus,

the effects of individuals wearing face masks and practicing SD are synergistic, with the most pronounced

effects occurring when 60% or 80% of individuals wear masks.

Effectiveness of combining mask wearing and social distancing to control infection in populations

with high proportions of pre-symptomatic and asymptomatic cases. Asymptomatic or pre-

symptomatic SARS-CoV-2 infection is emerging as possibly the most common clinical manifestation of

COVID-19[26-30]. This finding could only be revealed once mass testing campaigns were performed,

regardless of symptoms (e.g., universal testing campaigns). One of the earliest studies documenting

clinical manifestations in a testing campaign (which still focused testing mainly on symptomatic

individuals) was on the *Diamond Princess* cruise ship, where the rate of asymptomatic infection was

18%[31,32]. In a mass testing campaign in Iceland, where testing was offered to a segment of the general

population (regardless of symptoms), 43% of individuals were asymptomatic at the time of testing[33]. Thus,

the actual percentage of pre-symptomatic and asymptomatic cases is currently unknown, but it is clear

that a large number of SARS CoV2 new infections derive from undetected infections. Controlling the

outbreaks by isolation or even by increased population testing is a big challenge and may be difficult to

implement. We thus determined the efficacy of mask wearing and SD in the context of different

proportions of asymptomatic incidence. We assume that asymptomatic individuals are less infectious than

symptomatic individuals: asymptomatic individuals may have lower viral loads and reduced coughing,

sneezing, and nasal secretions, all of which may facilitate transmission[34-36]. The model considers that

symptomatic infected individuals isolated themselves 12 hours after the onset of symptoms, because they

get hospitalized or self-isolated. In contrast, asymptomatic individuals remain infectious, circulate and

transmit for a period of 7 days (Fig. 3A). Because some individuals can be pre-symptomatic (i.e.,

symptoms emerge later after the initial incubation time (Fig. 2B, *t= 5.1* days)), we modelled infectivity of

asymptomatic/pre-symptomatic as linearly declining until the individual is no longer infectious, after 7

days[37]. Using these assumptions, we carried out simulations, varying the percent of asymptomatic/pre-

symptomatic cases to determine the efficiency of non-pharmacological interventions in populations with

25, 50, and 75% pre-symptomatic and asymptomatic infected individuals (Fig. 3).

Our simulations demonstrate that the total number of infected individuals increases linearly with

the increase of the percentage of asymptomatic individuals in the population (Fig. 3B). To compare each

condition, we normalized the peak of new daily infected individuals to conditions in which no one

practiced SD or wore masks. Strikingly, increasing the percentage of individuals wearing masks linearly

reduces the normalized peak number of infected individuals per day (Fig. 3B). For instance, when 40% of

the individuals wear face masks, the peak number of infected decreases similarly independently of the

percent of asymptomatic cases considered (to 0.6–0.7, Fig. 3B). Furthermore, the linear relationship

between the decrease in peak number of newly infected individuals per day and increase in percent

individuals wearing masks is independent of the proportion of asymptomatic individuals in the population

(Fig. 3B). This finding indicates that, when the fraction of asymptomatic cases is high, as in the case for

COVID-19, wearing face masks is as effective to reduce the peak number of infected as when a low

percentage of individuals are asymptomatically infected.

In contrast, SD was only effective in populations with high incidence of asymptomatic infections

when a very high fraction of the population practice SD (more than 60%) (Fig. 3B). Furthermore, the low

efficacy of SD as a containment strategy is more pronounced when the proportion of asymptomatic

individuals increase (compare 25, 50, and 75%). Of note, if the number of individuals wearing face masks

is high, increasing the number of individuals practicing SD has negligible effects on the daily number of

infected individuals (see supplementary materials). One noteworthy observation from our model is that, at

high rates of SD (e.g. 80 %), there is more variability.

Our analysis uncovers a linear relationship between the fraction of a population wearing masks

and the reduction in infection rate. In contrast, we find SD requires a high fraction of compliance to be

effective. These findings indicate that having a high percentage of individuals wearing face masks is more

beneficial in preventing virus spread and reducing the peak number of infected individuals than having a

high percent of people practicing SD. Notably, the benefit of wearing masks is not affected if the percent

of pre-symptomatic and asymptomatic increases.

Interplay of face masks and social distancing for controlling infection spread and protecting from

COVD-19. Next, we determined the average cumulative incidence as a function of the percentage of

individuals wearing masks and practicing SD (standard deviation described in Fig. SM3-A). When neither

SD nor masks were used our simulations indicate that up to 99% of the population will end up infected,

leading to unacceptable levels of mortality and morbidity. When the proportion of the population wearing

masks was increased, no significant effect on the total number of infected people was noted until more

than 40% of the population wore masks. However, if 80% of people use masks, the average cumulative

incidence of infection decreased to around 35%. Therefore, wearing face masks alone would greatly limit

the spread of the virus. In contrast, SD alone does not have a significant effect. Increasing SD

compliance to 80% only reduced total infections by 8%, and 87% of the population was eventually

infected. Importantly, the combination of wearing masks and practicing SD by a high proportion of the

population dramatically reduces the total number of infected individuals to 10% of the population (Fig. 4A;

and Fig. SM2-C).

To gain additional insights into the characteristics of the epidemic in response to these mitigation

strategies, we studied the shape of the epidemic curve by calculating the peak and full-width half-

maximum (FWHM) of the peak of the new daily infections (Methods). While the peak reports on the

maximum number of daily infections at the height of the epidemic, the FWHM is a simple way to represent

the duration of the characteristic peak of new daily infections. Indeed, FWHM reports on the extinction

time of the infection within the population (number of days until no individual is still infected) (Fig. 4A, and

Fig. SM3-B standard deviation). A larger FWHM corresponds to a flatter but longer epidemic curve. A

flatter epidemic curve enables better management of the epidemic, as the healthcare system is not

overwhelmed by the number of cases at a given time. Initially, the FWHM was calculated assuming no

individual practice SD. FWHM was ~8.3 days if none of the individuals wears masks, ~12.8 if 40% of

individuals wear masks, and but increases dramatically to 85.4 when 80% of the population wears masks

(Fig. 2B). Our data indicate that wearing masks has a more profound effect than SD on flattening the

epidemic curve. For instance, when 80% of the population wears masks, the epidemic curve is eight

times flatter than without any non-pharmacological intervention (Fig. 4B). In contrast, if 80% of the

population practices SD, the flattening of the curve is less than threefold. The most dramatic effect on

flattening of the curve is observed when wearing masks is combined with SD; for instance, if 80% of the

population wears masks and 80% practices SD, the curve is flattened over 10-fold, compared to no

intervention.

We also find that the reduction in the number of infected individuals per day correlates with a

lengthening of the outbreak. Broadening of the peak affects the extinction time of the infection (Fig. 4C).

For instance, when the percentage of individuals wearing face masks rises from 0 to 80% of the

population, the extinction time increases from ~39.0 to ~58 days (Fig. 4C). Similarly, as a higher

proportion of individuals practice SD, the time to extinction of the infection also increases (Fig. 4C, ~39 to

~54 days for 0% to 80% of SD). Importantly, even though the time to epidemic extinction is extended, the

total number of infected individuals dramatically decreases (Fig. 4A). If a higher proportion of the

population (80%) wears face masks and practices SD, the time to epidemic extinction is reduced (Fig. 4C)

because the total number of infected individuals is dramatically reduced (Fig. 4A). Thus, stochastic effects

dominate, and this can be appreciated by the wider error bars (Fig. 4F).

To relate the impact of these interventions to their societal impact, we determined the number of

deaths per million after each mitigation strategy, assuming a mortality rate of ~3%[38]. This analysis

illustrates the heavy cost of lives of the virus, but also demonstrates that a high level of mask wearing

compliance is the most effective non-pharmacological approach to protect human lives, particularly when

combined with even moderate SD measures (Fig. 4D). In contrast, SD, without masks wearing is not

effective to reduce mortality (Fig. 4E). Finally, our simulations predict that increasing the proportion of the

population wearing masks will increase the time to outbreak extinction (from ~40 to ~60 days) (Fig. 4F).

Together with the broadening of the peak (Fig. 4C), this shows an effective flattening of the curve.

Importantly, with 80% mask wearing, we observed an increase in the statistical distribution from the

average time to extinction (Fig. 4F, see 95% confidence intervals). Thus, a generally low disease

incidence triggers stochastic events leading to extinction of the infection. Our simulations represent real

outbreak scenarios and reveal that as the outbreak approaches its extinction there is an increase in the

uncertainty of whether or not the infection has been completed eliminated, which argue to be prudent

before society reopening can be done safety.

Discussion

Here we use realistic simulations rooted in experimentally measured parameters of SARS-Cov2 spread,

contagion mode and mortality, to evaluate two available NPIs that reduce the spread of a respiratory

infection, such as COVID-19. In our simulation, we assumed proper use of FDA-approved face masks.

We showed that a high degree of compliance in the use of masks, regardless of whether the wearer

displays symptoms, slows the spread of infection. Face masks substantially reduce the transmission of

respiratory droplets and aerosols containing viral particles[9-11]. Increasing the fraction of the population

wearing face masks reduces the number of new infected individuals per day and flattened the curve of

total individuals infected (Fig. 2A and 4A). These two effects should reduce mortality and morbidity,

alleviate the current stress on healthcare systems, and enable a more effective management of severe

cases. However, solely wearing masks cannot entirely prevent an outbreak from occurring. It cannot by

itself extinguish the virus, since as long as a small fraction of the population is non-compliant, the virus

can persist in the population. Our models show that combining proper use of masks with practices such

as SD, indisputably decreases the number of new infected individuals per day (Fig. 2). This conclusion

agrees with that of ODE models[39]. In addition, our model was demonstrated to be scalable and able to

describe realistic situations such as interactions between 5000 agents (see supplementary materials).

Our analysis provides guidance for policies to protect the population from COVID-19. Optimizing

the use of masks with SD practices effectively limits the virus spread and reduces several parameters in

the epidemic, including cumulative incidence, shape of the peak, and the extinction rate (Fig. 3). In

particular, we observed that wearing masks is more effective than SD. Even in a population with a high

number of asymptomatic infections, increasing the use of masks up to 80% results in a significant

reduction in infection (Fig. 4A). Meanwhile, even 80% of individuals practicing SD has only a marginal

effect (Fig. 4D). This result can be understood in terms of contact rate, since we assume that

asymptomatic infectious individuals have higher mobility than symptomatic infectious ones. If the vast

majority of the population is asymptomatic, then high compliance with face mask use is a key factor for

curbing the epidemic.

Our simulations also provide insights into how enforcing different mitigation practices affects the

length of the epidemic. Assuming a homogenous population, the trajectory of epidemic extinction lasts

50–60 days, when 80% of the population either wears masks or practices SD (Fig. 4C). However, when

80% of the population is wearing masks and 0% of the population is practicing SD, the cumulative

incidence is reduced three times (~35%), and the peak is very broad (FWHM of ~ 85 days). In contrast,

when the population solely practices SD (80%), the majority of the population (93%) will end up infected,

and the peak of daily infected individuals will be sharper (FWHM of ~ 20 days).

Our model indicates that the synergistic utilization of face mask wearing and social distancing

practice is most effective in controlling SARS-CoV-2 spread. We observed that wearing masks in

combination with some degree of SD relaxes the need for a complete lockdown, leading to a potential

suggestion for an intervention policy based on a mix of the two measures. The effectiveness of mask

wearing to control virus spread is not reduced if a large fraction of the population is asymptomatic. This

suggests that, in the absence of universal testing, widespread use of face masks is necessary and

sufficient to prevent a large outbreak. Our results are supported by the real data of Korea[33] and Taiwan[34],

where an early mandate to requiring face mask usage, in combination with SD, severely limited the

spread of the virus. While more work is necessary to specifically assess the impact of other variables

shaping COVID-19 outbreaks, such as increased mobility, age stratification, testing a fraction of the

population, our study can accurately inform strategies to reduce the spread of the virus. In particular, our

results may be highly relevant toward informing specific realistic situations, such as the spread of the

disease in a confined space, where effective SD may not be easily achievable (e.g., schools, essential

businesses, correctional facilities, public transportation, hospitals). These strategies, if effectively

implemented, will save countless lives from the SARS-Cov2 infection. According to our model, if the

United States (330 million people) does not implement any NPIs, then ~627,000 people are expected to

die. Contrastingly, if 80% of the population wear face masks, that number would significantly shrink to

about 250,000. If both face masks and SD were practiced by 80% of the population from the start of a

pandemic, the mortality rate decreases to 65,600 people.

Materials and Methods

We developed and used our python codes using the NumPy library version 1.15.4[40] to carry out ABM and

describe the dynamic evolution of a SEAIR system affected by COVID-19 disease, in which each

individual can be in a susceptible (S), exposed (E), asymptomatic infected (A), symptomatic infected (I),

or recovered (R) status. In addition, each individual can wear a face mask (M) or can practice social

distancing (SD), where wearing face masks or practicing SD are independent binary values of an

individual. In the simulations, each individual was represented by a position in a 2-D lattice of 21x21

dimension with periodic boundary conditions. Individual initial positions are assigned randomly by

NumPy's random module, and the simulations start with all susceptible individuals and only one exposed.

During the simulation, each individual move along a randomly oriented trajectory at a constant velocity,

moves the same distance between each time step of the simulation, and interacts with individuals whose

position is within a fixed diameter of another individual. The data from each individual are saved as

vectors of attributes, including position, velocity, state (S, E, A, I, R, M, SD), and number of individuals

they come into contact with at every simulation time step. A time step corresponds to 1 hour. We

implemented the state of M by defining sub-routine during the interaction between two individuals where

the probability of infection was reduced by an amount corresponding to which of the two individuals is

wearing a mask. For the SD state, we assigned the individual to be stationary and not follow along a

randomly assigned trajectory.

We refer to Ferguson et al. for defining the rules governing the interactions among individuals[24].

For the interaction between symptomatic infected and susceptible individuals, the infection probability

was randomly sampled from a gamma distribution with a mean of 1 and a shape of 2.5, whereas for that

between asymptomatic infected and susceptible individuals, the infection probability is reduced by 33 %.

We assume this reduction of infectivity for the asymptomatic, based on the absence of transmission-

aiding symptoms, such as coughing, sneezing, and a runny nose[24]. The time between a susceptible

individual being exposed to being in the infectious infected state is 5.1 days[24]. Pre-symptomatic agents

are defined as infected agents during the incubation period. Symptomatic infected individuals are

infectious for 12 hours before self-isolating. After this time, we assume that these individuals no longer

infect those around them because they are hospitalized or self-isolating. Asymptomatic infected

individuals are infectious for 7 days, and the infectivity linearly declines until the individual is no longer

infectious on day 7[37]. If an infected, a susceptible, or both individuals are wearing a mask their probability

of another individual being infected is reduced to 5, 15, or 0%, respectively, from the original gamma

distribution.

First, we carried out a simulation with 500 individuals with 0% of individuals wearing face masks

and 0% practicing SD to optimize parameters as velocities and individual density to obtain a basic

reproduction number, R_0, equal to 2.5, based on reported values. The basic reproduction number was

computed from the mean number of symptomatic cases, resulting from a single symptomatic individual.

Then, we carried out four more sets of simulations in which we increased the percentage of individuals

wearing masks by the 20, 40, 60, and 80%. Finally, for each of these simulation sets we carried out four

more sets in which we increased the percentage of individuals practicing SD by the 20, 40, 60, and 80%.

Simulations are stochastic, and we simulated each condition 100 times to increase the statistical power of

the analysis. In total, we simulated 25 different conditions for a total of 2500 simulations, 100 for each

condition. In all these simulations, the probability of a new infection being asymptomatic was 50%.

Simulations were run until there were no individuals either infected or exposed. Summary values were

computed by methods in the NumPy module. Error bars were calculated by the standard deviation

method in NumPy and for figure 3 and 4 the error bars were calculated by 95% confidence interval. The

full-width half-maximum (FWHM) was calculated for each simulation from the newly infected per day data

by taking the maximum and using it to find the days that intersect the value of half the maximum value.

The first and last days of this intersection were then used to calculate the number of days of the FWHM.

From the set of FWHM for each simulation, an average and standard deviation were calculated using the

NumPy module. FWHM error bars are large due to a large variation in how flat the curves are. Graphs

were created by using the Matplotlib[41] and Seaborn[42] modules in python.

We followed the same protocol described above to perform simulations, varying the probability of

a new infection being asymptomatic from 25 to 75%. Finally, to monitor the reproducibility of our results

with a larger population and contrast with real data, we ran four sets of 30 simulations with 5000

individuals in the combinations of 0 or 80% SD and 0 or 80% wearing masks. All 5000 individual

simulations were done with 50% asymptomatic infection rates.

As the position, trajectory, and state of each individual in the simulation are explicitly known,

using the matplotlib library's animation package, we converted each time step of the simulation to an

image in a movie, representing 1 day as a second spanning the length of the simulation.

For each simulation, we saved in a .csv in tidy data format the summary values of the

simulations, which we used for the analysis. The simulation was written in python and all scripts are

available on GitHub at https://github.com/adamcatching/SARS_SEIR_Simulation, simulations were

generated using script mask_single_sim.py.

Acknowledgments

We acknowledge the use of the UCSF Computing Cluster Wynton
(https://wynton.ucsf.edu/hpc/index.html). We thank Judith Frydman, Alma Andino-Frydman, and Patrick
Dolan for their suggestions on the manuscript.

Author contributions

441 A. C., S. C., and M. T. Y. conceived the study. A. C. and S. C. designed the simulations. A. C. wrote and
442 executed the simulations. A. C., S. C., and S. B. analyzed the data. S. B. and R. A. provided guidance
443 and support. All authors wrote the manuscript.

Competing interest statement

The authors declare no competing interests.

References

1   Bourouiba, L. Turbulent Gas Clouds and Respiratory Pathogen Emissions: Potential

Implications for Reducing Transmission of COVID-19. *Jama* **323**, 1837,

doi:10.1001/jama.2020.4756 (2020).

2   Miller, I. F., Becker, A. D., Grenfell, B. T. & Metcalf, C. J. E. Disease and healthcare

burden of COVID-19 in the United States. *Nature Medicine*, 1, doi:10.1038/s41591-020-

0952-y (2020).

3   Bai, Y. *et al.* Presumed Asymptomatic Carrier Transmission of COVID-19. *Jama* **323**,

1406, doi:10.1001/jama.2020.2565 (2020).

4   Wölfel, R. *et al.* Virological assessment of hospitalized patients with COVID-2019.

Nature **581**, 465, doi:10.1038/s41586-020-2196-x (2020).

5   Schwartz, I. B., Kaufman, J., Hu, K. & Bianco, S. Predicting the impact of asymptomatic

transmission, non-pharmaceutical intervention and testing on the spread of COVID19.

6   Feng, S. *et al.* Rational use of face masks in the COVID-19 pandemic. *The Lancet*

Respiratory Medicine **8**, 434, doi:10.1016/s2213-2600(20)30134-x (2020).

7   Ng, K. *et al.* COVID-19 and the Risk to Health Care Workers: A Case Report. *Annals of*

Internal Medicine, doi:10.7326/l20-0175 (2020).

8   Suess, T. *et al.* The role of facemasks and hand hygiene in the prevention of influenza

transmission in households: results from a cluster randomised trial; Berlin, Germany,

2009-2011. *BMC Infectious Diseases* **12**, 1, doi:10.1186/1471-2334-12-26 (2012).

471 9 Leung, N. H. L. *et al.* Respiratory virus shedding in exhaled breath and efficacy of face

472  masks. *Nature Medicine* **26**, 676, doi:10.1038/s41591-020-0843-2 (2020).

473 10 Offeddu, V., Yung, C. F., Low, M. S. F. & Tam, C. C. Effectiveness of Masks and

474  Respirators Against Respiratory Infections in Healthcare Workers: A Systematic Review

475  and Meta-Analysis. *Clinical Infectious Diseases* **65**, 1934, doi:10.1093/cid/cix681

476  (2017).

477 11 Konda, A. *et al.* Aerosol Filtration Efficiency of Common Fabrics Used in Respiratory

478  Cloth Masks. *ACS Nano* **14**, 6339, doi:10.1021/acsnano.0c03252 (2020).

479 12 Howard, J. *et al.* Face Masks Against COVID-19: An Evidence Review.

480  doi:10.20944/preprints202004.0203.v1 (2020).

481 13 Tracht, S. M., Valle, S. Y. D. & Hyman, J. M. Mathematical Modeling of the

482  Effectiveness of Facemasks in Reducing the Spread of Novel Influenza A (H1N1). *Plos*

483  *One* **5**, doi:10.1371/journal.pone.0009018 (2010).

484 14 Brienen, N. C. J., Timen, A., Wallinga, J., Steenbergen, J. E. V. & Teunis, P. F. M. The

485  Effect of Mask Use on the Spread of Influenza During a Pandemic. *Risk Analysis* **30**,

486  1210, doi:10.1111/j.1539-6924.2010.01428.x (2010).

487 15 Ngonghala, C. N. *et al.* Mathematical assessment of the impact of non-pharmaceutical

488  interventions on curtailing the 2019 novel Coronavirus. *Mathematical Biosciences* **325**,

489  doi:10.1016/j.mbs.2020.108364 (2020).

490 16 Mahdizadeh Gharakhanlou, N. & Hooshangi, N. Spatio-temporal simulation of the novel

491  coronavirus (COVID-19) outbreak using the agent-based modeling approach (case study:

492  Urmia, Iran). *Informatics in Medicine Unlocked* **20**, doi:10.1016/j.imu.2020.100403

493  (2020).

17   Chang, S. L., Harding, N., Zachreson, C., Cliff, O. M. & Prokopenko, M. Modelling

transmission and control of the COVID-19 pandemic in Australia. (2020).

18   Kai, D., Goldstein, G.-P., Morgunov, A., Nangalia, V. & Rotkirch, A. Universal Masking

is Urgent in the COVID-19 Pandemic: SEIR and Agent Based Models, Empirical

Validation, Policy Recommendations. *arXiv:2004.13553 [physics, q-bio]* (2020).

19   Silva, P. C. L. *et al.* COVID-ABS: An agent-based model of COVID-19 epidemic to

simulate health and economic effects of social distancing interventions. *Chaos, Solitons*

& Fractals **139**, doi:10.1016/j.chaos.2020.110088 (2020).

20   Hoertel, N. *et al.* A stochastic agent-based model of the SARS-CoV-2 epidemic in

France. *Nature Medicine* **26**, 1417, doi:10.1038/s41591-020-1001-6 (2020).

21   Kuniya, T. Prediction of the Epidemic Peak of Coronavirus Disease in Japan, 2020.

Journal of Clinical Medicine **9**, doi:10.3390/jcm9030789 (2020).

22   Bauer, A. L., Beauchemin, C. A. A. & Perelson, A. S. Agent-based modeling of host–

pathogen systems: The successes and challenges. *Information Sciences* **179**, 1379,

doi:10.1016/j.ins.2008.11.012 (2009).

23   Rahmandad, H., Hu, K., Tebbens, R. J. D. & Thompson, K. M. Development of an

individual-based model for polioviruses: implications of the selection of network type

and outcome metrics. *Epidemiology and Infection* **139**, 836,

doi:10.1017/s0950268810001676 (2011).

24   Ferguson, N. *et al.* Impact of non-pharmaceutical interventions (NPIs) to reduce COVID-

19 mortality and healthcare demand.

25　Liang, M. *et al.* Efficacy of face mask in preventing respiratory virus transmission: A

systematic review and meta-analysis. *Travel Medicine and Infectious Disease*,

doi:10.1016/j.tmaid.2020.101751 (2020).

26　Oran, D. P. & Topol, E. J. Prevalence of Asymptomatic SARS-CoV-2 Infection. *Annals*

of Internal Medicine, doi:10.7326/m20-3012 (2020).

27　Ing, A. J., Cocks, C. & Green, J. P. COVID-19: in the footsteps of Ernest Shackleton.

Thorax **75**, 693, doi:10.1136/thoraxjnl-2020-215091 (2020).

28　Arons, M. M. *et al.* Presymptomatic SARS-CoV-2 Infections and Transmission in a

Skilled Nursing Facility. *New England Journal of Medicine* **382**, 2081,

doi:10.1056/NEJMoa2008457 (2020).

29　Day, M. Covid-19: four fifths of cases are asymptomatic, China figures indicate. *Bmj*

369, doi:10.1136/bmj.m1375 (2020).

30　Cline, S. *Cases at seafood plant cause spike in Oregon COVID numbers | News Break*,

<https://www.newsbreak.com/oregon/salem/news/1581086288035/cases-at-seafood-

plant-cause-spike-in-oregon-covid-numbers> (

31　Moriarty, L. F. Public Health Responses to COVID-19 Outbreaks on Cruise Ships —

Worldwide, February–March 2020. *MMWR. Morbidity and Mortality Weekly Report* **69**,

doi:10.15585/mmwr.mm6912e3 (2020).

32　Mizumoto, K., Kagaya, K., Zarebski, A. & Chowell, G. Estimating the asymptomatic

proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond

Princess cruise ship, Yokohama, Japan, 2020. *Euro Surveillance: Bulletin Europeen Sur*

Les Maladies Transmissibles = European Communicable Disease Bulletin **25**,

doi:10.2807/1560-7917.es.2020.25.10.2000180 (2020).

33   Gudbjartsson, D. F. *et al.* Spread of SARS-CoV-2 in the Icelandic Population. *New*

England Journal of Medicine **382**, 2302, doi:10.1056/NEJMoa2006100 (2020).

34   Buitrago-Garcia, D. C. *et al.* Asymptomatic SARS-CoV-2 infections: a living systematic

review and meta-analysis. *medRxiv*, doi:10.1101/2020.04.25.20079103 (2020).

35   Sakurai, A. *et al.* Natural History of Asymptomatic SARS-CoV-2 Infection. *New*

England Journal of Medicine **0**, doi:10.1056/NEJMc2013020 (2020).

36   Fernandez, M. *WHO: Data suggests it's "very rare" for coronavirus to spread through*

asymptomatics, <https://www.axios.com/who-coronavirus-asymptomatic-patients-

08d84e31-1846-44d8-8fe1-2343ff850764.html> (

37   He, X. *et al.* Temporal dynamics in viral shedding and transmissibility of COVID-19.

Nature Medicine, 1, doi:10.1038/s41591-020-0869-5 (2020).

38   *Mortality Risk of COVID-19 - Statistics and Research*,

<https://ourworldindata.org/mortality-risk-covid> (

39   Kai, D., Goldstein, G.-P., Morgunov, A., Nangalia, V. & Rotkirch, A. Universal Masking

is Urgent in the COVID-19 Pandemic: SEIR and Agent Based Models, Empirical

Validation, Policy Recommendations. (2020).

40   Van Der Walt, S., Colbert, S. C. & Varoquaux, G. The NumPy array: a structure for

efficient numerical computation. *Computing in Science & Engineering* **13**, 22,

doi:10.1109/mcse.2011.37 (2011).

41   *Matplotlib: A 2D Graphics Environment - IEEE Journals & Magazine*,

<https://ieeexplore.ieee.org/document/4160265> (

42   *mwaskom/seaborn: v0.8.1 (September 2017) | Zenodo*,

<https://zenodo.org/record/883859> (

Figures

Figure 1. Individual states and rules of interactions. Using the same coloring system used in animated simulations, we show the different states an individual can be in during a simulation. Individuals keep the masked or not masked attribute assigned at initialization during the course of the simulation. **A**) Spread of infection is caused by interactions of overlapping trajectories between infected and susceptible individuals. If the distance between individuals, d, is less than the radius of the two individuals, r, then an interaction occurs. Interactions initiate the generation of a random number that determines if transmission occurs. The chance of infection is randomly generated, based on a gamma distribution with shape parameter, alpha, of 0.25. This probability is further modulated by which individuals in the interactions are wearing face masks.

B) Agents are shown at their positions and states during snapshots of the simulation. Representative

simulations of are shown of 0, 40, or 80% of the population wearing face masks. Snapshots are collected

at days 0, 20, and 40 of the simulations. **C**) Progression of the outbreak, the trajectory of the number of

new daily infected individuals (Currently infected) and cumulatively infected from a representative

simulation are graphed in light blue and black, respectively.

Figure 2. Average new infections per day when varying the population wearing masks and

practicing social distancing. A) SD is implemented by assigning agents to not move along a trajectory

during the simulation. This lack of movement reduces the number of trajectories overlaps between agents

and subsequently reduces the number of transmission events. **B)** The number of new infections per day

for 0, 40, or 80% of a population wearing face-masks are displayed by red, blue, or green trajectories,

respectively, for 0% of the population practicing SD, for the 40% and for the 80%. Simulations were

repeated 100 times for each condition, the curves and the highlighted regions around the curves

represent the mean value +/- one standard deviation.

Figure 3. Effects of the asymptomatic population on the infected peak number. A) Schematic

representation of the asymptomatic transmission during the simulations. After 5.1 days of coming into

contact with an infected individual, infectious agents can develop symptoms and after 12 hours will be

isolated. On the contrary, asymptomatic individuals don't develop symptoms of the disease so they are

not isolated but keep circulating thus contributing to spread the infection for 7 more days after the initial

contact with an infectious agent. **B**) The normalized peak of new daily infected is represented as a

function of the percentage of the population wearing mask (left panels, green solid circles) and as a

function of increasing the percentage of asymptomatic individuals (25% top left panel, 50% central left

panel, 75% bottom left panel). The normalized peak of new daily infected is represented as a function of

the percentage of the population practicing social distance (right panels, blue solid circles) and as a

function of increasing the percentage of asymptomatic individuals (25% top right panel, 50% central right

panel, 75% bottom right panel). Error bars represent 95% confidence intervals calculated.

Andino 20.05D

Figure 4. Summary values from 2500 simulations of varying percentages of population wearing masks or social distancing. A) Proportion of infected population. The average cumulative incidence is represented as a function of the population practicing SD or wearing a mask, which are given by the x and y axis, respectively. **B**) The shape of the daily new infection curve. Full-width half-maximum (FWHM), denoting the number of days between the first day and last day of cases that have half the peak number of infected individuals. **C**) Extinction rate of the epidemic estimated as the average number of days for which the simulation reports no new infected individuals. In the figure, the numbers represent the mean value calculated over 100 of simulations carried out for each condition. For clarity, we reported the standard deviation of the mean for each value reported in this figure in Fig. SM5. **D**) and **E**) Number of death people. Extrapolated for one million individuals from the cumulative incidence of infections estimated from our model at a mortality rate of 3%. **F)** Number of days from the start of infection spread to

618   the last infected agent recovering. Error bars represent 95% confidence intervals calculated for **D**), **E**) and

619   **F**).