1 **Examining face-mask usage as an effective strategy to control COVID-19 spread**

Adam Catching[1,2][§], Sara Capponi[3,4][§], Ming Te Yeh[1], Simone Bianco[3,4][*], Raul Andino[1][*]

[1] Department of Microbiology and Immunology, University of California, San Francisco, San Francisco, CA
94158

[2] Graduate Program in Biophysics, University of California, San Francisco, San Francisco, CA 94158

[3] Industrial and Applied Genomics, AI and Cognitive Software, IBM Almaden Research Center, San Jose,
CA, 95120

[4] Center for Cellular Construction, San Francisco, CA 94158

22 [§] authors contributed equally to the manuscript

24 [*] corresponding authors: sbianco@us.ibm.com, Raul.Andino@ucsf.edu

Abstract

The COVID-19 global crisis is facilitated by high virus transmission rates and high percentages of asymptomatic and presymptomatic infected individuals. Containing the pandemic hinged on combinations of social distancing and face mask use. Here we examine the efficacy of these measures, using an agent-based modeling approach that evaluates face masks and social distancing in realistic confined spaces scenarios. We find face masks are more effective than social distancing. Importantly, combining face masks with even moderate social distancing provides optimal protection. The finding that widespread usage of face masks limits COVID-19 outbreaks can inform policies to reopening of social functions.

Keywords: epidemic, virus, individual-based simulation

Introduction

The COVID-19 outbreak has caused a catastrophic mortality and economic damage around the world. The causative agent of COVID-19, SARS-CoV-2, is an airborne pathogen that can be transmitted between humans through droplets and aerosols that can travel 1–8 meters[1]. The virus is transmitted by both symptomatic and asymptomatic individuals. COVID-19 causes severe symptoms that require hospitalization, as well as severe long-term sequels and death. Accordingly, the outbreak has seriously impacted healthcare systems around the world[2]. One of the major difficulties to contain the COVID-19 pandemic has been detection of infected asymptomatic or pre-symptomatic individuals, who are estimated to be responsible for as much as 95% of all transmissions[3][4]. As these individuals carry and spread the virus without manifesting any sign of the disease, they represent a crucial variable in managing the outbreak.

In the absence of an effective vaccine or antiviral, most countries have implemented non-pharmaceutical interventions (NPIs) to curb the spread of COVID-19[5]. These include closure of schools, workplaces, churches, offices, factories and other social venues, while encouraging preventative measures ranging from maintaining social distancing (SD, usually 2 m/6 ft between individuals) to total quarantine and societal lockdowns. These measures aim to reduce the effective contact rate of the population, which in turn decreases the disease reproductive number R_0. These NPIs limit the epidemic, but they present important drawbacks. Total lockdown can only be implemented for short periods, due to its severe impact on the social fabric and economy of a country. Meanwhile, essential workers remain vulnerable to infection and transmission due to the frequent encounters with infected, often asymptomatic, individuals. Because of their social and economic impact, lockdowns and SD measures have been lifted in some countries, leading to reactivation of virus spread and ensuing increased morbidity and mortality.

Face masks covering the nose and mouth area also provide a level of filtration that blocks virus transmission to certain extend[6-8]. Masks prevent the spread of droplets and aerosols generated by an infected individual[1], reducing viral transmission by 95%. Uninfected individuals wearing a mask are about 85% protected against infection[9]. Masks may be more effective to control the source of infectious virus because they prevent the larger expelled droplets from being converted into smaller droplets that can travel farther. Accordingly, face masks reduce the spread of influenza[10] and coronaviruses[11,12].

In our current study, we analyzed the relative efficacy of wearing face masks and/or exercising SD to reducing the spread of COVID-19. Through stochastic computer simulations of infection spread, we modeled realistic outbreak scenarios and found that SD only yields beneficial effects if accompanied by a widespread population lockdown. In contrast, wearing face masks is a highly effective strategy to reduce the spread of infection. Our results suggest that, even when a large fraction of infected individuals is asymptomatic, mask wearing is the most effective strategy to control virus spread and alleviate the impact of COVID-19 outbreak, particularly when combined with conditions of partial SD compatible with the function of society.

Results

Stochastic model description and calibration. We developed an agent-based model (ABM) to examine the effectiveness of wearing masks and SD on the rate of infections and viral spread during the pandemic. Unlike other computational approaches to model the spread of COVID-19 disease[13,14], ABMs are stochastic models that allow the description of non-homogeneous distributions of agents that act individually[12,15]. In ABMs, each individual behaves dynamically and independently in response to environmental changes[16] according to rules that describe their interactions. To model the COVID-19 pandemic, we used a SEAIR system in which each agent represents an individual who can be susceptible (S), exposed (E), asymptomatically infected (A), symptomatically infected (I), or recovered (R). To obtain a realistic model of virus spread, we chose parameters that describe the spread of SARS-CoV2: transmission events occur through contacts made between susceptible and infectious individuals in close proximity (distance ≤ 2r, Fig. 1A), and exposed individuals undergo an incubation period of 5.1 days to become infectious (Fig. 1, spheres with red border). This incubation time represents the interval required to increase viral loads to levels sufficient for transmission (REF). We assume that recovered individuals (Fig. 1, spheres with black border),

who resolve the infection, cannot be re-infected or infect others, which is a reasonable assumption for the
duration of our models (45 days).

To define the probabilities of infection, we used reported COVID-19 parameters[17]. The probability
of transmission follows a Γ (*gamma)* distribution (see Methods) whose shape is described by a constant
(α), estimated to be 0.25. Wearing masks reduces this probability (Fig. 1, probability of infection). To
estimate the protective effect of masks, we used parameters determined for FDA-approved surgical masks,
whose efficacy has been experimentally verified to inhibit virus transmission[10]. Based on previous studies
(Ferguson et al.), we assumed that, if an infectious individual wears a mask, the effective probability of
transmission is reduced by a factor of 0.05. If a susceptible individual wears a mask, α is reduced by a
factor of 0.15 (Fig. 1A). Our assumption that wearing masks is more effective to reduce transmission than
to prevent getting infected is supported by experimental data[18]. However, if both infectious and susceptible
individuals wear masks the probability of transmission is the product of these probabilities (0.0075) and,
thus, sufficiently low such that transmission is effectively null. We calibrated our model by running
simulations without any individuals wearing a face mask or practicing SD and considering that 50% of
infected individuals are asymptomatic (Movie 1). In this way, we determined the simulation parameters,
such as velocity and density of individuals, to obtain a value of R_0 = 2.5, consistent with what was reported
early on as the infection rate of the epidemic in Wuhan[9].

Percentage of population wearing masks determines the daily infection incidence and cumulative
number of cases. We started each simulation with one individual being infected and all others susceptible
and assuming that 50% of COVID-19 infections are asymptomatic (or pre-symptomatic). Non-infected and
asymptomatic individuals circulate in the population without any restriction (i.e., they do not isolate
themselves or become hospitalized) (see Methods). In contrast, symptomatic individuals no longer move
after 12 hours of the symptom onset, simulating hospitalization or self-isolation. Thus, symptoms are
assumed to manifest after the incubation time of 5.1 days plus 12 hours (Fig. 2B). Using these assumptions
and model calibration, we carried out a set of simulations in which we gradually increased the percentage
of individuals in the population that wear masks. Individuals assigned randomly to wear a mask keep on
the mask for the entire duration of the simulation. Increasing the fraction of the population wearing face
masks has a highly significant effect on the spread of the virus (Fig. 1B, 1C and Movie 2). Mask wearing
reduces the cumulative number of infected individuals at the end of the simulation (Fig. 1B). Strikingly, this
effect negatively correlates with the fraction of the population using masks without any other intervention,
suggesting that masks are effective for reducing the overall number of cases (Fig. 1B). The description of
the dynamics of infection generated by this model is consistent with previous clinical studies[9,18] and
highlights the benefit of wearing masks (Fig. 1C, Movie 1).

If the daily incidence surpasses the treatment capacity, it will overwhelm the healthcare system
with detrimental consequences for medical care of infected individuals and increased mortality and
morbidity. Thus, we examined the effects of face masks on the daily incidence of infection over time (Fig.
2B). If all individuals move freely and randomly interact with others (i.e., 0% SD), the rate of daily infection
through the population depends on the percentage of individuals wearing masks. The average number of
new daily infection varies considerably, according to the number of individuals wearing face masks. When
0% of individuals wear face masks (Fig. 2b, red line), the number of daily infections peaks sharply at day
27 with a maximum of 54.8 ± 18.1 infections per day for a population of 500 individuals. No additional
infections are observed after day 39, when the entire population (100%) has been infected. Thus, without
any intervention, the infection quickly reaches every individual through an epidemic characterized by a very
sharp infection peak. Note that, in these original models, we assume all individuals recover after ~11 days
post-infection and cannot be re-infected. When 40% of all individuals wear masks, the number of individuals
infected at any given day is reduced by approximately 30% (maximum of infected individuals 35.1 ± 12.8,
Fig. 2A, blue line), both flattening the curve and extending the duration of the outbreak by more than 10
days, with a peak maximum at day 32. By day 52, an average of 472 individuals has been infected, and
none of the individuals is infectious or exposed any longer. Even more significantly, if 80% of the population
wear masks, we observed a significant flattening of the curve, with a substantial reduction in the maximum
number of infected individuals per day, 5.9 ± 6.8 (Fig. 2A, green line), and the number of new infected
individuals reached zero by day 57.8 ± 35.0. Thus, the shape of the outbreak changes from a curve
characterized by a sharp peak when no intervention is considered to a broader peak when 80% of the
individuals wear masks. By increasing the percentage of the individuals wearing masks, the number of

newly infected individuals per day substantially decreases, which will reduce mortality and morbidity.
Moreover, since the use of masks eliminates the sharp peak that characterizes SARS CoV2 epidemics, the
overall impact of the outbreak on the health system is alleviated. These results highlight the importance of
widespread mask wearing as an effective intervention that can be implemented as soon as the first cases
are reported.

Effect of social distancing on viral infection spread. Next, we evaluated the effect of SD in shaping the
spread of infection. Practicing SD does not affect the probability of infection but reduces the chances of
encounters leading to transmission. Thus, SD was introduced into the model by limiting the proportion of
individuals in the population that move freely in the field. This simulates a scenario in which a given
proportion of individuals in the population quarantine or shelter-in-place, thereby reducing the probability of
contacts and transmission (Fig. 2A). As reported, if a percentage of individuals practice SD, even without
any other non-pharmacological intervention, the number of infections and daily infection rate are reduced
(compare Fig. 2B, SD 0, 40, and 80%, red line). However, if no one wears masks but 40% of individuals
practice SD, we observed a very small effect in the shape of the infection curve (Fig. 2B, the peak maximum
decreases from 51.0 ± 13.3 to 45.7 ± 5.9). When 80% of the population practices SD, a more significant
reduction in the number of new daily infections was observed (Fig. 2B, 80%). Notably, the shape of the new
daily infection curve broadens considerably upon increasing the percentage of individuals wearing face
masks (Fig. 2B, compare blue and green curves with red curve). For instance, if 80% of the population
wears masks with 40% SD, the peak maximum decreases to one-tenth (from 51.0 ± 13.3 to 5.7 ± 5.9) and
is slightly delayed (Fig. 2B). At 80% SD, the peak of new daily infections is no longer observed, and the
number of new infected individuals per day averages 1.6 ± 1.3 (Fig. 2B). Thus, the effects of individuals
wearing face masks and practicing SD are synergistic, with the most pronounced effects occurring when
60% or 80% of individuals wear masks.

Effectiveness of combining mask wearing and social distancing to control infection in populations
with high proportions of pre-symptomatic and asymptomatic cases. Asymptomatic or pre-
symptomatic SARS-CoV-2 infection is emerging as possibly the most common clinical manifestation of
COVID-19[19-23]. This finding could only be revealed once mass testing campaigns were performed,
regardless of symptoms (e.g., universal testing campaigns). One of the earliest studies documenting clinical
manifestations in a testing campaign (which still focused testing mainly on symptomatic individuals) was on
the *Diamond Princess* cruise ship, where the rate of asymptomatic infection was 18%[24,25]. In a mass testing
campaign in Iceland, where testing was offered to a segment of the general population (regardless of
symptoms), 43% of individuals were [[infected but? If you don't include this, it would imply that 57% were
showing symptoms.]] asymptomatic at the time of testing[26]. Thus, the actual percentage of pre-symptomatic
and asymptomatic cases is currently unknown, but it is clear that a large number of SARS CoV2 new
infections derive from undetected infections. Controlling the outbreaks by isolation or even by increased
population testing is a big challenge and may be difficult to implement. We thus determined the efficacy of
mask wearing and SD in the context of different proportions of asymptomatic incidence. We assume that
asymptomatic individuals are less infectious than symptomatic individuals: asymptomatic individuals may
have lower viral loads and reduced coughing, sneezing, and nasal secretions, all of which may facilitate
transmission[27-29]. However, the model considers that symptomatic infected individuals isolated themselves
12 hours after the onset of symptoms, because they get hospitalized or self-isolated. In contrast,
asymptomatic individuals remain infectious, circulate and transmit for a period of 7 days (Fig. 3A). Because
some individuals can be pre-symptomatic (i.e., symptoms emerge later after the initial incubation time (Fig.
2B, *t= 5.1* days)), we modelled infectivity of asymptomatic/pre-symptomatic as curve that declines linearly
until the individual is no longer infectious, after 7 days[30]. Using these assumptions, we carried out
simulations, varying the percent of asymptomatic/pre-symptomatic cases to determine the efficiency of non-
pharmacological interventions in populations with 25, 50, and 75% pre-symptomatic and asymptomatic
infected individuals (Fig. 3).

Our simulations demonstrate that the total number of infected individuals increases linearly with
the increase of the percentage of asymptomatic individuals in the population (Fig. 3B). To compare each
condition, we normalized the peak of new daily infected individuals to conditions in which no one practiced
SD or wore masks. Strikingly, increasing the percentage of individuals wearing masks linearly reduces the
normalized peak number of infected individuals per day (Fig. 3B). For instance, when 40% of the individuals
wear face masks, the peak number of infected decreases similarly independently of the percent of

asymptomatic cases considered (to 0.6–0.7, Fig. 3B). Furthermore, the linear relationship between the
decrease in peak number of newly infected individuals per day and increase in percent individuals wearing
masks is independent of the proportion of asymptomatic individuals in the population (Fig. 3B). This finding
indicates that, when the fraction of asymptomatic cases is high, as in the case for COVID-19, wearing face
masks is as effective to reduce the peak number of infected as when a low percentage of individuals are
asymptomatically infected.

In contrast, SD was only effective in populations with high incidence of asymptomatic infections
when a very high fraction of the population practice SD (more than 60%) (Fig. 3C). Furthermore, the low
efficacy of SD as a containment strategy is more pronounced when the proportion of asymptomatic
individuals increase (compare 25, 50, and 75% Fig. 3C). Of note, if the number of individuals wearing face
masks is high, increasing the number of individuals practicing SD has negligible effects on the daily number
of infected individuals (not shown). One noteworthy observation from our model is that, at high rates of
asymptomatic infection (e.g. 75 %), there is more variability in the protective effect of SD (Figure 3C, note
larger 95% confidence intervals if 80% practice SD). This suggests that stochastic nature of epidemics
affects the efficiency of SD more significantly than the efficiency of mask wearing.

Our analysis uncovers a linear relationship between the fraction of a population wearing masks and
the reduction in infection rate. In contrast, we find SD requires a high fraction of compliance to be effective.
These findings indicate that having a high percentage of individuals wearing face masks is more beneficial
in preventing virus spread and reducing the peak number of infected individuals than having a high percent
of people practicing SD. Notably, the benefit of wearing masks is not affected if the percent of pre-
symptomatic and asymptomatic increases.

Interplay of face masks and social distancing for controlling infection spread and protecting from
COVD-19. Next, we determined the average cumulative incidence as a function of the percentage of
individuals wearing masks and practicing SD (standard deviation described in Fig. SM3-A). When neither
SD nor masks were used our simulations indicate that up to 95% of the population will end up infected,
leading to unacceptable levels of mortality and morbidity. When the proportion of the population wearing
masks was increased, no significant effect on the total number of infected people was noted until more than
60% of the population wore masks. However, if 80% of people use masks, the average cumulative
incidence of infection decreased to around 35%. Therefore, wearing face masks alone would greatly limit
the spread of the virus. In contrast, SD alone does not have a significant effect. Increasing SD compliance
to 80% only reduced total infections by 8%, and 87% of the population was eventually infected. Importantly,
the combination of wearing masks and practicing SD by a high proportion of the population dramatically
reduces the total number of infected individuals to 1–10% of the population (Fig. 4A; and Fig. SM2-C).

To gain additional insights into the characteristics of the epidemic in response to these mitigation
strategies, we studied the shape of the epidemic curve by calculating the peak and full-width half-maximum
(FWHM) of the peak of the new daily infections (Methods). While the peak reports on the maximum number
of daily infections at the height of the epidemic, the FWHM is a simple way to represent the duration of the
characteristic peak of new daily infections. Indeed, FWHM reports on the extinction time of the infection
within the population (number of days until no individual is still infected) (Fig. 4A, and Fig. SM3-B standard
deviation). A larger FWHM corresponds to a flatter but longer epidemic curve. A flatter epidemic curve
enables better management of the epidemic, as the healthcare system is not overwhelmed by the number
of cases at a given time. Initially, the FWHM was calculated assuming no individual practice SD. FWHM
was ~8.3 days if none of the individuals wears masks, ~12.8 if 40% of individuals wear masks, and but
increases dramatically to 85.5 when 80% of the population wears masks (Fig. 2B). Our data indicate that
wearing masks has a more profound effect than SD on flattening the epidemic curve. For instance, when
80% of the population wears masks, the epidemic curve is eight times flatter than without any non-
pharmacological intervention (Fig. 4B). In contrast, if 80% of the population practices SD, the flattening of
the curve is less than threefold. The most dramatic effect on flattening of the curve is observed when
wearing masks is combined with SD; for instance, if 80% of the population wears masks and 80% practices
SD, the curve is flattened over 10-fold, compared to no intervention.

We also find that the reduction in the number of infected individuals per day correlates with a
lengthening of the outbreak. Broadening of the peak affects the extinction time of the infection (Fig. 4C).
For instance, when the percentage of individuals wearing face masks rises from 0 to 80% of the population,

the extinction time increases from ~39.0 to ~53 days (Fig. 4C). Similarly, as a higher proportion of
individuals practice SD, the time to extinction of the infection also increases (Fig. 4C, ~39 to 51 days for
0% to 80% of SD). Importantly, even though the time to epidemic extinction is extended, the total number
of infected individuals dramatically decreases (Fig. 4A). However, if a higher proportion of the population
(80%) wears face masks and practices SD, the time to epidemic extinction is reduced because the total
number of infected individuals is dramatically reduced (Fig. 4A).

To relate the impact of these interventions to their societal impact, we determined the number of
deaths per million after each mitigation strategy, assuming a mortality rate of ~3%[31]. This analysis illustrates
the heavy cost of lives of the virus, but also demonstrates that a high level of mask wearing compliance is
the most effective non-pharmacological approach to protect human lives, particularly when combined with
even moderate SD measures (Fig. 4D). In contrast, SD, without masks wearing is not effective to reduce
mortality (Fig. 4E). Finally, our simulations predict that increasing the proportion of the population wearing
masks will increase the time to outbreak extinction (from ~40 to ~60 days) (Fig. 4F). Together with the
broadening of the peak (Fig. 4C), this shows an effective flattening of the curve. Importantly, with 80% mask
wearing, we observed an increase in the statistical distribution from the average time to extinction (Fig. 4F,
see 95% confidence intervals). Thus, a generally low disease incidence triggers stochastic events leading
to extinction of the infection. Our simulations represent real outbreak scenarios and reveal that as the
outbreak approaches its extinction there is an increase in the uncertainty of whether or not the infection has
been completed eliminated, which argue to be prudent before society reopening can be done safety.

Discussion

Here we use realistic simulations rooted in experimentally measured parameters of SARS-Cov2 spread,
contagion mode and mortality, to evaluate two available NPIs that reduce the spread of a respiratory
infection, such as COVID-19. In our simulation, we assumed proper use of FDA-approved face masks. We
showed that a high degree of compliance in the use of masks, regardless of whether the wearer displays
symptoms, slows the spread of infection. Face masks substantially reduce the transmission of respiratory
droplets and aerosols containing viral particles[31]. Increasing the fraction of the population wearing face
masks reduces the number of new infected individuals per day and flattened the curve of total individuals
infected (Fig. 2A and 4A). These two effects should reduce mortality and morbidity, alleviate the current
stress on healthcare systems, and enable a more effective management of severe cases. However, solely
wearing masks cannot entirely prevent an outbreak from occurring. It cannot by itself extinguish the virus,
since as long as a small fraction of the population is non-compliant, the virus can persist in the population.
Our models show that combining proper use of masks with practices such as SD, indisputably decreases
the number of new infected individuals per day (Fig. 2). This conclusion is in agreement with that of ODE
models[32].

Our analysis provides guidance for policies to protect the population from COVID-19. Optimizing
the use of masks with SD practices effectively limits the virus spread and reduces several parameters in
the epidemic, including cumulative incidence, shape of the peak, and the extinction rate (Fig. 3). In
particular, we observed that wearing masks is more important than SD. Even in a population with a high
number of asymptomatic infections, increasing the use of masks up to 80% results in a significant reduction
in infection (Fig. 4A). Meanwhile, even 80% of individuals practicing SD has only a marginal effect (Fig.
4D). This result can be understood in terms of contact rate, since we assume that asymptomatic infectious
individuals have higher mobility than symptomatic infectious ones. If the vast majority of the population is
asymptomatic, then high compliance with face mask use is a key factor for curbing the epidemic.

Our simulations also provide insights into how enforcing different mitigation practices affects the
length of the epidemic. Assuming a homogenous population, the trajectory of epidemic extinction lasts 50–
60 days, when 80% of the population either wears masks or practices SD (Fig. 4C). However, when 80%
of the population is wearing masks and 0% of the population is practicing SD, the cumulative incidence is
reduced three times (~35%), and the peak is very broad (FWHM of ~ 85 days). In contrast, when the
population solely practices SD (80%), the majority of the population (93%) will end up infected, and the
peak of daily infected individuals will be sharper (FWHM of ~ 20 days).

Our model indicates that the synergistic utilization of face masks wearing and social distancing
practice is most effective controlling SARS-CoV-2 spread. We observed that wearing masks in combination
with some degree of SD relaxes the need for a complete lockout, leading to a partial reopening of the
society. The effectiveness masks wearing to control virus spread is not reduced if a large fraction of the
population is asymptomatic. This suggests that, in the absence of universal testing, widespread use of face
masks is necessary and sufficient to prevent a large outbreak. Our results are supported by the real data
of Korea[33] and Taiwan[34], where an early mandate to requiring face mask usage, in combination with SD,
severely limited the spread of the virus. While more work is necessary to specifically assess the impact of
other variables shaping COVID-19 outbreaks, such as increased mobility, age stratification, testing a
fraction of the population, our study can accurately inform strategies to reduce the spread of the virus. In
particular, our results may be highly relevant toward informing specific realistic situations, such as the
spread of the disease in a confined space, where effective SD may not be easily achievable (e.g., schools,
essential businesses, correctional facilities, public transportation, hospitals). These strategies, if effectively
implemented, will save countless lives from the SARS-Cov2 infection. According to our model, if the United
States (330 million people) does not implement any NPIs, then ~627,000 people are expected to die.
Contrastingly, if 80% of the population wear face masks, that number would significantly shrink to about
250,000. If both face masks and SD were practiced by 80% of the population from the start of a pandemic,
the mortality rate decreases to 65,600 people.

Materials and Methods

We developed and used our python codes using the NumPy library version 1.15.4[36] to carry out ABM and
describe the dynamic evolution of a SEAIR system affected by COVID-19 disease, in which each individual
can be in a susceptible (S), exposed (E), asymptomatic infected (A), symptomatic infected (I), or recovered
(R) status. In addition, each individual can wear a face mask (M) or can practice social distancing (SD),
where wearing face masks or practicing SD are independent binary values of an individual. In the
simulations, each individual was represented by a position in a 2-D lattice of 21x21 dimension. Individual
initial positions are assigned randomly by NumPy's random module, and the simulations start with all
susceptible individuals and only one exposed. During the simulation, each individual move along a
randomly oriented trajectory at a constant velocity, moves the same distance between each time step of
the simulation, and interacts with individuals whose position is within a fixed diameter of another individual.
The data from each individual are saved as vectors of attributes, including position, velocity, state (S, E, A,
I, R, M, SD), and number of individuals they come into contact with at every simulation time step. A time
step corresponds to 1 hour. We implemented the state of M by defining sub-routine during the interaction
between two individuals where the probability of infection was reduced by an amount corresponding to
which of the two individuals are wearing a mask. For the SD state, we assigned the individual to be
stationary and not follow along a randomly assigned trajectory.

We refer to Ferguson et al. for defining the rules governing the interactions among individuals[17].
For the interaction between symptomatic infected and susceptible individuals, the infection probability was
randomly sampled from a gamma distribution with a mean of 1 and a shape of 2.5, whereas for that between
asymptomatic infected and susceptible individuals, the infection probability is reduced by 33 %. We assume
this reduction of infectivity for the asymptomatic, based on the absence of transmission-aiding symptoms,
such as coughing, sneezing, and a runny nose[17]. The time between a susceptible individual being exposed
to being in the infectious infected state is 5.1 days[17]. Symptomatic infected individuals are infectious for 12
hours before self-isolating. After this time, we assume that these individuals no longer infect those around
them because they are hospitalized or self-isolating. Asymptomatic infected individuals are infectious for 7
days, and the infectivity linearly declines until the individual is no longer infectious on day 7[37]. If an infected,
a susceptible, or both individuals are wearing a mask their probability of another individual being infected
is reduced to 5, 15, or 0%, respectively, from the original gamma distribution.

First, we carried out a simulation with 500 individuals with 0% of individuals wearing face masks
and 0% practicing SD to optimize parameters as velocities and individual density to obtain a basic
reproduction number, R_0, equal to 2.5, based on reported values. The basic reproduction number was
computed from the mean number of symptomatic cases, resulting from a single symptomatic individual.
Then, we carried out four more sets of simulations in which we increased the percentage of individuals
wearing masks by the 20, 40, 60, and 80%. Finally, for each of these simulation sets we carried out four
more sets in which we increased the percentage of individuals practicing SD by the 20, 40, 60, and 80%.

Simulations are stochastic, and we simulated each condition 100 times to increase the statistical power of
the analysis. In total, we simulated 25 different conditions for a total of 2500 simulations, 100 for each
condition. In all these simulations, the probability of a new infection being asymptomatic was 50%.
Simulations were run until there were no individuals either infected or exposed. Summary values were
computed by methods in the NumPy module. Error bars were calculated by the standard deviation method
in NumPy. The full-width half-maximum (FWHM) was calculated for each simulation from the newly infected
per day data by taking the maximum and using it to find the days that intersect the value of half the maximum
value. The first and last days of this intersection were then used to calculate the number of days of the
FWHM. From the set of FWHM for each simulation, an average and standard deviation were calculated
using the NumPy module. FWHM error bars are large due to a large variation in how flat the curves are.
Graphs were created by using the Matplotlib[38] and Seaborn[39] modules in python.

We followed the same protocol described above to perform simulations, varying the probability of
a new infection being asymptomatic from 25 to 75%. Finally, to monitor the reproducibility of our results
with a larger population and contrast with real data, we ran four sets of 30 simulations with 5000 individuals
in the combinations of 0 or 80% SD and 0 or 80% wearing masks. All 5000 individual simulations were
done with 50% asymptomatic infection rates.

As the position, trajectory, and state of each individual in the simulation are explicitly known, using
the matplotlib library's animation package, we converted each time step of the simulation to an image in a
movie, representing 1 day as a second spanning the length of the simulation.

For each simulation, we saved in a .csv in tidy data format the summary values of the simulations,
which we used for the analysis. The simulation was written in python and all scripts are available on GitHub
at https://github.com/adamcatching/SARS_SEIR_Simulation, simulations were generated using script
mask_single_sim.py.

Acknowledgments

394 We acknowledge the use of the UCSF Computing Cluster Wynton (https://wynton.ucsf.edu/hpc/index.html).
395 We thank Judith Frydman, Alma Andino-Frydman and Patrick Dolan for his encouraging discussions.

Author contributions

398 A. C., S. C., and M. T. Y. conceived the study. A. C. and S. C. designed the simulations. A. C. wrote and
399 executed the simulations. A. C., S. C., and S. B. analyzed the data. S. B. and R. A. provided guidance and
400 support. All authors wrote the manuscript.

Competing interest statement

403 The authors declare no competing interests.

References

Bourouiba, L. Turbulent Gas Clouds and Respiratory Pathogen Emissions: Potential Implications for Reducing Transmission of COVID-19. *Jama* **323**, 1837, doi:10.1001/jama.2020.4756 (2020).

Miller, I. F., Becker, A. D., Grenfell, B. T. & Metcalf, C. J. E. Disease and healthcare burden of COVID-19 in the United States. *Nature Medicine*, 1, doi:10.1038/s41591-020-0952-y (2020).

Bai, Y. *et al.* Presumed Asymptomatic Carrier Transmission of COVID-19. *Jama* **323**, 1406, doi:10.1001/jama.2020.2565 (2020).

Wölfel, R. *et al.* Virological assessment of hospitalized patients with COVID-2019. *Nature* **581**, 465, doi:10.1038/s41586-020-2196-x (2020).

Schwartz, I. B., Kaufman, J., Hu, K. & Bianco, S. Predicting the impact of asymptomatic transmission, non-pharmaceutical intervention and testing on the spread of COVID19.

Feng, S. *et al.* Rational use of face masks in the COVID-19 pandemic. *The Lancet Respiratory Medicine* **8**, 434, doi:10.1016/s2213-2600(20)30134-x (2020).

Ng, K. *et al.* COVID-19 and the Risk to Health Care Workers: A Case Report. *Annals of Internal Medicine*, doi:10.7326/l20-0175 (2020).

Suess, T. *et al.* The role of facemasks and hand hygiene in the prevention of influenza transmission in households: results from a cluster randomised trial; Berlin, Germany, 2009-2011. *BMC Infectious Diseases* **12**, 1, doi:10.1186/1471-2334-12-26 (2012).

Leung, N. H. L. *et al.* Respiratory virus shedding in exhaled breath and efficacy of face masks. *Nature Medicine* **26**, 676, doi:10.1038/s41591-020-0843-2 (2020).

Offeddu, V., Yung, C. F., Low, M. S. F. & Tam, C. C. Effectiveness of Masks and Respirators Against Respiratory Infections in Healthcare Workers: A Systematic Review and Meta-Analysis. *Clinical Infectious Diseases* **65**, 1934, doi:10.1093/cid/cix681 (2017).

Konda, A. *et al.* Aerosol Filtration Efficiency of Common Fabrics Used in Respiratory Cloth Masks. *ACS Nano* **14**, 6339, doi:10.1021/acsnano.0c03252 (2020).

Howard, J. *et al.* Face Masks Against COVID-19: An Evidence Review. doi:10.20944/preprints202004.0203.v1 (2020).

Kuniya, T. Prediction of the Epidemic Peak of Coronavirus Disease in Japan, 2020. *Journal of Clinical Medicine* **9**, doi:10.3390/jcm9030789 (2020).

Ngonghala, C. N. *et al.* Mathematical assessment of the impact of non-pharmaceutical interventions on curtailing the 2019 novel Coronavirus. *Mathematical Biosciences* **325**, doi:10.1016/j.mbs.2020.108364 (2020).

15 Bauer, A. L., Beauchemin, C. A. A. & Perelson, A. S. Agent-based modeling of host–
pathogen systems: The successes and challenges. *Information Sciences* **179**, 1379,
doi:10.1016/j.ins.2008.11.012 (2009).

16 Rahmandad, H., Hu, K., Tebbens, R. J. D. & Thompson, K. M. Development of an
individual-based model for polioviruses: implications of the selection of network type and
outcome metrics. *Epidemiology and Infection* **139**, 836, doi:10.1017/s0950268810001676
(2011).

17 Ferguson, N. *et al.* Impact of non-pharmaceutical interventions (NPIs) to reduce COVID-
19 mortality and healthcare demand.

18 Liang, M. *et al.* Efficacy of face mask in preventing respiratory virus transmission: A
systematic review and meta-analysis. *Travel Medicine and Infectious Disease*,
doi:10.1016/j.tmaid.2020.101751 (2020).

19 Oran, D. P. & Topol, E. J. Prevalence of Asymptomatic SARS-CoV-2 Infection. *Annals of
Internal Medicine*, doi:10.7326/m20-3012 (2020).

20 Ing, A. J., Cocks, C. & Green, J. P. COVID-19: in the footsteps of Ernest Shackleton.
Thorax **75**, 693, doi:10.1136/thoraxjnl-2020-215091 (2020).

21 Arons, M. M. *et al.* Presymptomatic SARS-CoV-2 Infections and Transmission in a Skilled
Nursing Facility. *New England Journal of Medicine* **382**, 2081,
doi:10.1056/NEJMoa2008457 (2020).

22 Day, M. Covid-19: four fifths of cases are asymptomatic, China figures indicate. *Bmj* **369**,
doi:10.1136/bmj.m1375 (2020).

23 Cline, S. *Cases at seafood plant cause spike in Oregon COVID numbers | News Break*,
<https://www.newsbreak.com/oregon/salem/news/1581086288035/cases-at-seafood-
plant-cause-spike-in-oregon-covid-numbers> (

24 Moriarty, L. F. Public Health Responses to COVID-19 Outbreaks on Cruise Ships —
Worldwide, February–March 2020. *MMWR. Morbidity and Mortality Weekly Report* **69**,
doi:10.15585/mmwr.mm6912e3 (2020).

25 Mizumoto, K., Kagaya, K., Zarebski, A. & Chowell, G. Estimating the asymptomatic
proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess
cruise ship, Yokohama, Japan, 2020. *Euro Surveillance: Bulletin Europeen Sur Les
Maladies Transmissibles = European Communicable Disease Bulletin* **25**,
doi:10.2807/1560-7917.es.2020.25.10.2000180 (2020).

26 Gudbjartsson, D. F. *et al.* Spread of SARS-CoV-2 in the Icelandic Population. *New
England Journal of Medicine* **382**, 2302, doi:10.1056/NEJMoa2006100 (2020).

27 Buitrago-Garcia, D. C. *et al.* Asymptomatic SARS-CoV-2 infections: a living systematic
review and meta-analysis. *medRxiv*, doi:10.1101/2020.04.25.20079103 (2020).

28 Sakurai, A. *et al.* Natural History of Asymptomatic SARS-CoV-2 Infection. *New England
Journal of Medicine* **0**, doi:10.1056/NEJMc2013020 (2020).

29 Fernandez, M. *WHO: Data suggests it's "very rare" for coronavirus to spread through*
asymptomatics, <https://www.axios.com/who-coronavirus-asymptomatic-patients-
08d84e31-1846-44d8-8fe1-2343ff850764.html> (

30 *Advice on the use of masks in the community, during home care and in healthcare settings*
in the context of the novel coronavirus (COVID-19) outbreak,
<https://www.who.int/publications-detail-redirect/advice-on-the-use-of-masks-in-the-
community-during-home-care-and-in-healthcare-settings-in-the-context-of-the-novel-
coronavirus-(2019-ncov)-outbreak> (

31 *Mortality Risk of COVID-19 - Statistics and Research*,
<https://ourworldindata.org/mortality-risk-covid> (

32 Grover, C. Efficacy of face masks depends on spatial relation between host and recipient
and who is being protected. *Bmj* **369**, doi:10.1136/bmj.m2016 (2020).

33 Stutt, R. O. J. H., Retkute, R., Bradley, M., Gilligan, C. A. & Colvin, J. A modelling
framework to assess the likely effectiveness of facemasks in combination with 'lock-down'
in managing the COVID-19 pandemic. *Proceedings of the Royal Society A: Mathematical,*
Physical and Engineering Sciences **476**, doi:10.1098/rspa.2020.0376 (2020).

34 Cheng, V. C.-C. *et al.* The role of community-wide wearing of face mask for control of
coronavirus disease 2019 (COVID-19) epidemic due to SARS-CoV-2. *Journal of*
Infection, doi:10.1016/j.jinf.2020.04.024 (2020).

35 Yuan, E. J. *et al.* Where to buy face masks? Survey of applications using Taiwan's open
data in the time of COVID-19. *Journal of the Chinese Medical Association* **Latest Articles**,
doi:10.1097/jcma.0000000000000325 (2020).

36 Van Der Walt, S., Colbert, S. C. & Varoquaux, G. The NumPy array: a structure for
efficient numerical computation. *Computing in Science & Engineering* **13**, 22,
doi:10.1109/mcse.2011.37 (2011).

37 He, X. *et al.* Temporal dynamics in viral shedding and transmissibility of COVID-19.
Nature Medicine, 1, doi:10.1038/s41591-020-0869-5 (2020).

38 *Matplotlib: A 2D Graphics Environment - IEEE Journals & Magazine*,
<https://ieeexplore.ieee.org/document/4160265> (

39 *mwaskom/seaborn: v0.8.1 (September 2017) | Zenodo*,
<https://zenodo.org/record/883859> (

511
512
513
514

Figures

Figure 1. Individual states and rules of interactions. Using the same coloring system used in animated simulations, we show the different states an individual can be in during a simulation. Individuals keep the masked or not masked attribute assigned at initialization during the course of the simulation. **A**) Spread of infection is caused by interactions of overlapping trajectories between infected and susceptible individuals. If the distance between individuals, d, is less than the radius of the two individuals, r, then an interaction occurs. Interactions initiate the generation of a random number that determines if transmission occurs. The chance of infection is randomly generated, based on a gamma distribution with shape parameter, alpha, of 0.25. This probability is further modulated by which individuals in the interactions are wearing face masks.

 B) Agents are shown at their positions and states during snapshots of the simulation. Representative simulations of are shown of 0, 40, or 80% of the population wearing face masks. Snapshots are collected at days 0, 20, and 40 of the simulations. **C**) Progression of the outbreak, the trajectory of the number of new daily infected individuals (Currently infected) and cumulatively infected from a representative simulation are graphed in light blue and black, respectively.

Figure 2. Average new infections per day when varying the population wearing masks and practicing social distancing. A) SD is implemented by assigning agents to not move along a trajectory during the simulation. This lack of movement reduces the number of trajectories overlaps between agents and subsequently reduces the number of transmission events. **B)** The number of new infections per day for 0, 40, or 80% of a population wearing face-masks are displayed by red, blue, or green trajectories, respectively, for 0% of the population practicing SD, for the 40% and for the 80%. Simulations were repeated 100 times for each condition, the curves and the highlighted regions around the curves represent the mean value +/- one standard deviation.

Figure 3. Effects of the asymptomatic population on the infected peak number. A) The peak number
of infected individuals is represented as a function of the percentage of the population wearing face masks
and as a function of the percentage of asymptomatic individuals. 25% of asymptomatic individuals is colored
gray, 50% bright blue, 75% dark blue as the legend reports. Sensitivity analysis reports how the peak
number of infected individuals is affected by the 25% (**B**), 50% (**C**), and 75% (**D**) of asymptomatic individuals
varying the percentage of the population wearing face masks and the percentage of the population
practicing SD. The legend reports the percentage of individuals practicing SD: 0% is colored gray, 20%
bright blue, 40% dark blue, 60% purple, 80% red. Error bars represent 95% confidence intervals calculated
for both B) and C).

Figure 4. Summary values from 2500 simulations of varying percentages of population wearing
masks or social distancing. A) Proportion of infected population. The average cumulative incidence is
represented as a function of the population practicing SD or wearing a mask, which are given by the x and
y axis, respectively. **B**) The shape of the daily new infection curve. Full-width half-maximum (FWHM),
denoting the number of days between the first day and last day of cases that have half the peak number of
infected individuals. **C**) Extinction rate of the epidemic estimated as the average number of days for which
the simulation reports no new infected individuals. In the figure, the numbers represent the mean value
calculated over 100 of simulations carried out for each condition. For clarity, we reported the standard
deviation of the mean for each value reported in this figure in Fig. SM5. **D**) and **E**) Number of death people.
Extrapolated for one million individuals from the cumulative incidence of infections estimated from our
model at a mortality rate of 3%. **F**) Number of days from the start of infection spread to the last infected
agent recovering. Error bars represent 95% confidence intervals calculated for **D**), **E**) and **F**).