Newborn dried blood spots for serological surveys of COVID-19

Feimei Liu, BS*[1,2], Mytien Nguyen[1,3], MS*, Pavithra Vijayakumar[3], BS*, Alanna Kaplan[3], PhD,

Amit Meir[4], PhD, Yile Dai[1], BS, Eric Wang[1], BS, Hannah Walsh[6], BS, Aaron M. Ring[1], MD PhD,

Saad B. Omer[5,6,7], MBBS PhD, and Shelli F. Farhadian, MD PhD[6]

*These authors contributed equally

[1] Department of Immunobiology, Yale School of Medicine, New Haven, CT

[2] Yale School of Engineering and Applied Science, New Haven, CT

[3] Yale School of Medicine, New Haven, CT

[4] Boyer Center for Molecular Medicine, Department of Microbial Pathogenesis, Yale University

[5] Yale Institute for Global Health
14
15 [6] Department of Internal Medicine (Infectious Diseases), Yale School of Medicine, New Haven,
16 Connecticut.
17
18 [7] Department of Epidemiology of Microbial Diseases, Yale School of Public Health, New Haven,
19 Connecticut.
20

Corresponding author:

Shelli F. Farhadian MD PhD

135 College St, New Haven, CT 06510

shelli.farhadian@yale.edu

Abstract

There is an urgent need for inexpensive, population-wide surveillance testing for COVID-19. We

tested newborn dried blood spots (DBS) anti-SARS-COV-2 antibodies for all infants born at Yale

from March-May 2020, and found that newborn DBS serologies reflect maternal and

population-wide infection rates during the study period. This suggests a role for DBS in COVID-

19 surveillance in areas where viral testing is limited.

Introduction

Population seroprevalence surveys are important tools for COVID-19 public health decision-

making. However, there are significant hurdles to implementing large-scale serological surveys,

especially in resource poor settings. Newborn dried blood spot (DBS) is an easy, inexpensive,

and widely used method to collect and store newborn blood specimens, and, in cases where

newborn antibody results reflect maternal seropositivity, DBS is an appealing tool to measure

COVID-19 population seroprevalence trends among pregnant women.[1] Here, we evaluated DBS

as a potential COVID-19 surveillance tool.

Methods

Discarded newborn DBS was obtained for all infants born in the Yale New Haven hospital

system from February 19-May 26, 2020, a time period that began two weeks prior to the first

confirmed case in Connecticut (March 8). Routine SARS-CoV-2 PCR testing of all women who

presented to labor and delivery began on April 1, 2020. We adapted a standardized ELISA

Serology Assay[2] for detecting anti-SARS-CoV-2 antibody using newborn DBS. Anti-SARS-CoV-2

IgG was measured in all samples, and IgM was measured where the IgG result was positive. The

threshold for a positive result was set at the 99% confidence interval, at which 1% of true

negatives would be reported as false positives. Banked cord blood from infants born prior to

2019 was run as negative controls. Maternal demographic and clinical features were examined

in association with the likelihood of an antibody positive newborn using multivariable logistic

regression performed in STATA 16.1.

This study was deemed exempt from human subjects regulations by the Yale institutional

review board.

Results

The first positive DBS in the study was observed from an infant born in Connecticut on February

18, 2020 **(Figure 1).** Overall 182/3048 (5.9%) DBS tested were positive for anti-SARS-CoV-2 IgG.

None were positive for anti-SARS-CoV-2 IgM. Of the 182 infants with positive antibody results,

134 had mothers who underwent testing for COVID-19 by viral PCR prior to delivery, of which

65 (49%) had a positive test and 69 (51%) had a negative PCR test. All pre-2019 cord blood

samples were negative for SARS-CoV-2 IgG by this assay.

We examined the inter-relationship between daily newborn IgG antibody positive rates to

maternal and statewide COVID-19 infection rates. The daily positive proportion of DBS

specimens were predicted strongly by both maternal SARS-CoV-2 infection rate ($p<0.001$) and

statewide COVID-19 daily positive test counts per 100,000 people ($p=0.010$). Further

investigations using cross-correlation analysis demonstrated that daily newborn IgG antibody

positive rate strongly correlated with maternal SARS-CoV-2 infection rate at zero lag time

(r=0.47), and statewide COVID-19 positive test counts at a lag-time of 15-days (r=0.45).

We examined maternal demographics and clinical factors that may be associated with the

likelihood of an IgG+ newborn. Of the 92 mothers who tested positive for SARS-CoV-2 during

the universal screening period, 90 had complete demographic and clinical data. When

controlled for body mass index (BMI), hypertension, diabetes, COVID-19 symptoms, gestational

age, and time between maternal COVID-19 diagnosis and delivery (lag time), infants that were

antibody positive for COVID-19 were more likely to be born later during the study period

(adjusted OR, 1.05; 95% CI, 1.01-1.10, p=0.01), and to mothers with older maternal age

(adjusted OR, 1.13; 95% CI, 1.02-1.25, p=0.01). No significant associations were found for the

other factors.

Discussion

Our findings demonstrate utility of the newborn DBS SARS-CoV-2 antibody assay to detect past

maternal infection and suggest a use for DBS to measure population-level trends of COVID-19,

as well as a way to monitor for resurgence of this disease. The detection of seropositive

newborns prior to the availability of viral testing in CT indicates that DBS surveillance may be a

useful tool for COVID-19 surveillance where viral testing is limited. Most, but not all, mothers

who screened positive for SARS-CoV-2 during the study period delivered a newborn with

detectable anti-SAR-CoV-2 IgG antibody, a finding that may reflect the lag time in development

of detectable antibodies after infection.[3]

This study has limitations. Most mothers were screened for SARS-CoV-2 at the time of

hospitalization for delivery, but we were unable to determine the true date of maternal

infection. Because we did not have access to the date of infection or serum antibody testing

results for all mothers, we could not determine whether mothers who screened positive for

SARS-CoV-2 by PCR but delivered a seronegative newborn had poor antibody responses

themselves, had a remote infection with antibody responded that waned prior to delivery, or

whether there was inefficient transplacental transfer of antibody in these cases. However,

given that the majority of PCR positive mothers delivered seropositive newborns, this does not

appear to diminish the utility of DBS as a surveillance tool.

In this study, we demonstrate that levels of IgG in DBS reflect overall population-level trends in

case incidence, with a lag that is consistent with the time to development of detectable

antibodies after infection, making DBS antibody testing an attractive option for large-scale

population surveillance during the COVID-19 pandemic. As dried blood spots are routinely

collected from newborns, no additional sample collection is required, and specimens can be

stored for later assessment. Using DBS as a surveillance tool may therefore be particularly

advantageous in resource-poor settings, where innovative tools of field epidemiology will be

required in order to control the spread of the virus.

Author Contributions: *SFF had full access to all of the data in the study and takes responsibility*
for the integrity of the data and the accuracy of the data analysis.

Concept and design: SFF and SBO

Acquisition, analysis, or interpretation of data: All authors.

Drafting of the manuscript: SFF, MN, PV, FL

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: MN, PV, FL

Administrative, technical, or material support: AR, SFF

Supervision: SFF

Additional contributions: We acknowledge Dr. Wade Schulz who assisted with data acquisition.

References

1.   Bjorkesten J, Enroth S, Shen Q, et al. Stability of Proteins in Dried Blood Spot Biobanks.
Mol Cell Proteomics. 2017;16(7):1286-1296.
2.   Amanat F, Stadlbauer D, Strohmeier S, et al. A serological assay to detect SARS-CoV-2
seroconversion in humans. *Nat Med.* 2020.
3.   Long QX, Liu BZ, Deng HJ, et al. Antibody responses to SARS-CoV-2 in patients with
COVID-19. *Nature medicine.* 2020;26(6):845-848.
136

138　**Figure 1: Rate of newborn COVID-19 positive antibody and cases in Connecticut from February to May**

139　**2020.** Daily positive rate (bars) and 7-day moving average (lines) of newborn IgG antibody (blue) in

140　relation to SARS-CoV-2 PCR in mothers (pink), and positive cases per 100,000 people in Connecticut

141　State (orange). Connecticut statewide data were retrieved from Hughes *et al.* (2020) and rate per

142　100,000 calculated using U.S. Census estimates.