1

# Role of IgM and IgA Antibodies in the Neutralization of SARS-CoV-2

Jéromine Klingler[1,2], Svenja Weiss[1,2], Vincenza Itri[1], Xiaomei Liu[1,2], Kasopefoluwa Y.

Oguntuyo[3], Christian Stevens[3], Satoshi Ikegame[3], Chuan-Tien Hung[3], Gospel Enyindah-

Asonye[1], Fatima Amanat[3,4], Ian Baine[5], Suzanne Arinsburg[5], Juan C. Bandres[2], Erna Milunka

Kojic[6], Jonathan Stoever[7], Denise Jurczyszak[3,4], Maria Bermudez-Gonzalez[3], Viviana Simon[1,3,8],

Arthur Nádas[9], Sean Liu[1,3], Benhur Lee[3], Florian Krammer[3], Susan Zolla-Pazner[1,3*], Catarina E.

Hioe[1,2,3*]

[1]Division of Infectious Diseases, Department of Medicine, Icahn School of Medicine at Mount

Sinai, New York, NY USA.

[2]James J. Peters VA Medical Center, Bronx, NY, USA.

[3]Department of Microbiology, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

[4]Graduate School of Biomedical Sciences, Icahn School of Medicine at Mount Sinai, New York,

NY, USA.

[5]Department of Pathology, Icahn School of Medicine at Mount Sinai, New York, NY, USA.

[6]Division of Infectious Diseases, Department of Medicine, Mount Sinai West and Morningside,

NY, USA.

[7]Pulmonary and Critical Care Medicine, Mount Sinai West, NY, USA.

[8]Global Health Emerging Pathogens Institute, Icahn School of Medicine at Mount Sinai, NY,

USA.

[9]Department of Environment Medicine, NYU School of Medicine, New York, NY, USA

25  *Co-corresponding author

26  Contact: catarina.hioe@mssm.edu, catarina.hioe@va.gov

## Abstract

SARS-CoV-2 has infected millions of people globally. Virus infection requires the receptor-

binding domain (RBD) of the spike protein. Although studies have demonstrated anti-spike and -

RBD antibodies to be protective in animal models and convalescent plasma as a promising

therapeutic option, little is known about immunoglobulin (Ig) isotypes capable of blocking

infection. Here, we studied spike- and RBD-specific Ig isotypes in convalescent and acute

plasma/sera. We also determined virus neutralization activities in plasma/sera, and purified Ig

fractions. Spike- and RBD-specific IgM, IgG1, and IgA1 were produced by all or nearly all

subjects at variable levels and detected early after infection. All samples also displayed

neutralizing activity. Regression analyses revealed that IgM and IgG1 contributed most to

neutralization, consistent with IgM and IgG fractions' neutralization potency. However, IgA also

exhibited neutralizing activity at a lower potency. Together, IgG, IgM and IgA are critical

components of convalescent plasma used for COVID-19 treatment.

## Keywords

43  SARS-CoV-2, COVID-19, antibody isotypes, neutralization, convalescent plasma

## Background

In December 2019, the first patients with coronavirus disease 2019 (COVID-19), caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), were identified in Wuhan's city of Hubei Province, China[1]. Since then, the epidemic has rapidly spread to most regions of the world, infecting millions of people[2]. Effective therapeutics and vaccines against SARS-CoV-2 are urgently needed. Convalescent plasma transfusion showed promising results in patients with severe to life-threatening COVID-19[3–7] and clinical trials to evaluate the efficacy of convalescent plasma treatment for ambulatory and hospitalized COVID-19 patients are underway[8–11]. To this end, more information about the Ig isotypes present in the plasma of COVID-19 convalescent individuals and their antiviral activities are needed. It is also unclear which of the immunoglobulin (Ig) isotypes present in convalescent plasma are protective. The data would likewise inform vaccine development, as more than 100 vaccine candidates are in different stages of preclinical development, and many are now in phase 2 and 3 clinical trials[12]. Although using different strategies[13], most vaccines are based on the SARS-CoV-2 spike protein[14,15], which is a membrane-anchored protein present with two others (membrane and envelope proteins) on the virus envelope surface and contains the receptor-binding domain (RBD) required for binding to and entry into the cells[16–22]. These vaccines aim to protect by inducing neutralizing antibodies (Abs) capable of blocking viral infection.

Monomeric IgG constitutes approximately 75% of the Abs found in serum and exists as four subtypes: IgG1 (~66% of IgG), IgG2 (~23% of IgG), IgG3 (~7% of IgG) and IgG4 (~4% of IgG)[23,24]. IgM Abs represent 10% of total serum Abs and are the first to arise in response to new antigens[24,25]. Although IgM Abs do not undergo extensive somatic hypermutation to increase their affinity as do IgG and IgA Abs, their higher valency due to oligomerization enhances their

avidity and potency against pathogens[24,26,27]. IgA Abs exist as two subtypes: IgA1 and IgA2,

representing 15% of total serum Abs. IgA1 is the primary IgA subtype in serum while IgA2

predominates in mucosal secretions[24]. These two IgA subtypes are dimeric in the mucosa, but in

the circulation, they are monomeric.

SARS-CoV-2 spike-, RBD- and nucleocapsid-specific serum/plasma Abs of IgM, IgG,

and IgA isotypes are found in most COVID-19 patients[28–35], with Ab neutralizing activities

reported developing within the first two weeks of infection and decline over time[31,33,36–38].

However, the neutralizing titers appear to vary greatly[31,33,36–38], and they correlate with Ab

binding levels against RBD, spike, and/or nucleocapsid, and with age, duration of symptoms, and

symptom severity[31,33,37]. Several RBD-specific monoclonal Abs of IgG isotype with potent

antiviral activities have been generated from individuals with high neutralization titers, and these

confer protection in animal models[31,36,39,40]. Moreover, a monoclonal Ab of IgA isotype capable

of recognizing both the SARS-CoV-1 and SARS-CoV-2 spike proteins and blocking ACE2

receptor binding was recently described[41]. However, no direct evidence is available regarding the

neutralizing capacity of plasma IgM and IgA Abs from COVID-19 patients.

Studies on other respiratory viruses such as influenza show that, in addition to IgG, IgA

could also mediate virus neutralization, and their relative contribution depends on the

physiologic compartment in which they are found, with IgA contributing to the protection of

mostly the upper respiratory tract while IgG was protective of the lower respiratory tract[42–45]. Of

note, an anti-hemagglutinin monoclonal polymeric IgA has been demonstrated to mediate more

potent antiviral activities against influenza when compared to a monoclonal IgG against the same

epitope[46]. Interestingly, an IgM Ab with potent antiviral activities targeting influenza B's

receptor binding site has also been described[47]. In addition, respiratory syncytial virus (RSV)-

specific mucosal IgA neutralizing Abs are a better correlate of protection than RSV-specific

serum IgG neutralizing Abs[48]. In the case of SARS-CoV-1, high titers of mucosal IgA in the

lungs correlated with reduced pathology upon viral challenge in animal models[49]. Whether IgA

in the blood and in the respiratory tract's mucosa offers protection against SARS-CoV-2

infection or COVID-19 disease remains an open question. Moreover, few data are available

regarding the contribution of IgM to neutralization and protection against viruses, including

SARS-CoV-2. Of note, in terminally ill COVID-19 patients, systemic SARS-CoV-2 infection

affecting multiple organs including the heart, kidneys, and brain, is evident from autopsy

studies[50–52]. Thus, the capacity of plasma Ig to suppress virus spread to these organs is critical for

effective convalescent plasma therapy against severe COVID-19 disease.

We recently described a multiplex bead Ab binding assay using the Luminex technology

to detect total Ig against spike and RBD[53]. Based on this assay, we characterized the Ig isotype

profiles against spike and RBD in the plasma and serum from acutely infected or convalescent

individuals using this Luminex assay which detects antigen-specific IgM, IgG1-4, IgA1, and

IgA2. Using a pseudovirus assay[54], we also measured neutralizing activities in plasma, serum,

and Ig isotype plasma fractions to determine the neutralizing capacity of IgM, IgA, and IgG. The

data indicated a high prevalence of spike- and RBD-specific IgM and IgA Abs, similar to that of

IgG1, in plasma and serum from COVID-19 patients, and their contributions to virus neutralizing

activities. By testing purified IgG, IgM and IgA Abs from plasma of convalescent COVID-19

subjects, this study presents the first direct evidence that plasma IgG, IgM, and IgA all contribute

to SARS-CoV-2 neutralization.

**Results**

**Levels of Ig isotypes against the SARS-CoV-2 spike and RBD vary in convalescent**

**individuals.** A total of 29 serum (P#5-8) and plasma (TF#1-25) specimens from COVID-19-

convalescent individuals was tested. TF#1-25 were collected between March 26 and April 7,

2020, about 4-8 weeks after the initial outbreak in North American, and used for transfusion into

hospitalized COVID-19 patients[3]. Ten plasma from COVID-negative contemporaneous blood

bank donors (N#4-13) were included for comparison. Sera or plasma from 12 uninfected

individuals banked prior to the COVID-19 outbreak (N#1-3 and N#14-22) were used to establish

background values. The specimens were initially titrated for total Ig against spike and RBD (**Fig.**

**1**). All 29 COVID-19 positive specimens exhibited titration curves of total Ig Abs against spike,

while none of the negative controls did. Similar results were observed with RBD, except that one

contemporaneous COVID-19-negative sample had low levels of RBD-specific Ig (N#10).

Overall the background MFI values were higher for RBD than spike. To assess the

reproducibility of the assay, the samples were tested in at least two separate experiments run on

different days, and a strong correlation was observed between the MFI values from these

independent experiments (**Supplementary Fig. 1**). The areas under the curves (AUCs) highly

correlated with the MFI values from specimens diluted 1:200 ($p < 0.0001$; **Supplementary Fig.**

**2**); consequently, all samples were tested for isotyping at this dilution. At the 1:200 dilution we

were able to discern a diverse range of Ig isotype levels among individual samples (**Fig. 2**). To

evaluate for the presence of spike-specific and RBD-specific total Ig, IgM, IgG1, IgG2, IgG3,

IgG4, IgA1 and IgA2, the specificity and strength of the secondary Abs used to detect the

different isotypes were first validated with Luminex beads coated with myeloma proteins of

known Ig isotypes (IgG1, IgG2, IgG3, IgG4, IgA1, IgA2, and IgM). All eight secondary Abs

were able to detect their specific Ig isotypes with MFI values reaching $> 60,000$

(**Supplementary Fig. 3**).

All 29 convalescent individuals had anti-spike and anti-RBD total Ig (**Fig. 2**), but the Ig

levels were highly variable, with MFI values ranging from 36,083 to 190,150. In addition, all 29

convalescent individuals also displayed IgM Abs against spike at varying levels, and 93% were

positive for anti-RBD IgM when evaluated using cut-off values calculated as mean + 3 standard

deviation (SD) of the 12 pre-pandemic samples (**Fig 2b, c**). An IgG1 response was detected

against both spike and RBD in 97% of the convalescent subjects, with MFI values that ranged

from 1,013 to 59,880. In contrast, IgG2, IgG3, and IgG4 Abs against spike and RBD were

detected in only a small fraction of the subjects, and the levels were very low (MFI values <

1,300) (**Fig. 2**). Surprisingly, almost all individuals produced IgA1 Abs against spike (97%) and

RBD (93%), while 17% exhibited IgA2 against spike, and 48% exhibited IgA2 against RBD

(**Fig. 2**). Low levels, slightly above cut-off, of spike- and RBD-binding total Ig, IgM, and IgG1,

and IgA1 were also detected sporadically in contemporaneous COVID-19 samples, such as N#8,

N#10, and N#11. The responses against spike and RBD were highly correlated for every isotype

(**Supplementary Fig. 4**). Overall, these data demonstrate that IgM, IgG1, and IgA1 Abs were

induced against spike and RBD in all or almost all COVID-19 convalescent individuals (**Fig. 2**).

The levels, however, were highly variable among individuals. No statistical significance in the

levels of total Ig, IgM, IgG1, and IgA1 was observed between female and male individuals

(**Supplementary Fig. 5**).

In **Fig. 3**, regression analyses to assess the impact of individual isotypes on the total Ig

binding showed that IgG1 had the highest $r^2$ values (0.83 and 0.70 for spike- and RBD-binding

IgG1, respectively) with p <0.0001, indicating that IgG1 is the major isotype induced by SARS-

CoV-2 infection against spike and RBD (**Fig. 3a,b**). IgG2 Abs against RBD had an $r^2$ value of

0.55 with p < 0.0001, but IgG2 levels were very low. For all other isotypes, including IgM, the $r^2$

values were less than 0.40 (**Fig. 3c**). Thus, despite the presence of many isotypes in sera and

plasma, as expected, the major isotype is IgG1.

Specimens from two patients (P#1 and P#2) were drawn during the acute phase of the

infection. Serial specimens from these patients were tested to determine the isotypes of Abs

present early in infection. The earliest samples from both patients, drawn at 7 or 8 days after

symptom onset were already positive for total Ig, IgG1, IgA1and IgM Abs against spike

(**Supplementary Fig. 6a**) and RBD (**Supplementary Fig. 6b**), and these levels increased over

the following three to seven days. On the contrary, IgA2 Ab levels were near or below

background on days 7-8 and remained unchanged over two weeks post-onset. IgG4 Abs also

remained low or near background, whereas IgG2 and IgG3 Abs increased slightly to above

background after 10-15 days.

**Neutralizing activities are detected in all convalescent COVID-19 individuals.** We

subsequently tested the ability of samples from convalescent subjects to neutralize a VSV$\Delta$G

pseudovirus bearing the SARS-CoV-2 spike protein (COV2pp). This pseudovirus assay

demonstrated a strong positive correlation with neutralization of the authentic SARS-CoV-2

virus (p = 0.82 and p < 0.0001 for $IC_{50}$ correlation)[54]. The results, shown in **Fig. 4**, demonstrate

the percentages of COV2pp neutralizing activity in serum or plasma specimens from 28 COVID-

19-convalescent individuals and 11 COVID-19-negative individuals over a range of seven serial

four-fold dilutions. A soluble recombinant RBD (sRBD) protein capable of blocking virus

infection was tested in parallel as a positive control.

All specimens from COVID-19-convalescent individuals were able to neutralize the virus

at levels above 50% (**Fig. 4a**). For 26 of 28 specimens, neutralization reached more than 90%

(**Fig. 4a**). The sample with the lowest titer (reciprocal $IC_{50}$ titer = 37) reached a neutralization

plateau of only ~60%. Of note, one sample (TF#11) demonstrated highly potent neutralization

with a reciprocal $IC_{50}$ titer > 40,960, and neutralization was still 75% at the highest dilution

tested. None of the samples from COVID-19-negative individuals reached 50% neutralization

(**Fig. 4b**), while the sRBD positive control demonstrated potent neutralization with an $IC_{50}$ of

0.06 µg/mL (**Fig. 4c**), similar to that recently reported[54].

The samples were also tested for neutralization against a COV2pp bearing spike with a

D614 mutation (D614G mutant), as the D614G variant has become the most prevalent

circulating strain in the global pandemic[55]. Similar to the WT COV2pp, all COVID-19-

convalescent samples had neutralizing activity reaching >50% (**Fig. 4d**), while none of the

negative samples did (**Fig. 4e**). The $IC_{90}$ titers against WT and D614 mutant differed on average

by only 1.7-fold and correlated strongly with each other (**Fig. 4f**).

**IgM and IgG1 contribute most to SARS-CoV-2 neutralization.** Given our observation

that Ab isotype levels and neutralization titers varied tremendously among convalescent COVID-

19 individuals (**Figs. 2 and 5**), we investigated the relative contribution of each Ab isotype to the

neutralizing activities. Regression analyses were performed on 27 COVID-19-convalescent

samples (TF#11 was excluded due to its outlier neutralization titer). As expected, relatively high

$r^2$ values (0.32 – 0.62) and significant p values were observed with total Ig, IgM and IgG1; in

each case, $r^2$ values were higher for spike than for RBD (**Fig. 6a**). For other isotypes, significant

p values were sporadically achieved, but $r^2$ values were weak (**Fig. 6a,b**).

**Neutralizing activities are mediated by IgM, IgG, and IgA fractions.** To assess

directly the capacity of different isotypes to mediate neutralization, we evaluated the

neutralization activities of IgM, IgG, and IgA fractions purified from the plasma of five COVID-

19-convalescent individuals (RP#1-5). The enrichment of IgM, IgG1, and IgA1 Abs reactive

with spike and RBD was validated using the isotyping method used above (**Supplementary Fig.**

**7** and not shown). These IgM, IgG, and IgA fractions were then evaluated for neutralizing

activity along with the original plasma (**Fig. 7**). The RP#1-5 plasma neutralizing reciprocal $IC_{50}$

titers ranged from 35 to 690 (**Fig. 7a,b**). Purified IgM and IgG fractions from RP#1-5 all

mediated neutralization reaching more than 50%. Unexpectedly, plasma IgA fractions also

displayed neutralizing activity, although not to the same potency as IgM and IgG (**Fig 7c,d**). In

contrast, IgM, IgG, and IgA fractions from the negative control (RN#1) showed no neutralization

(**Fig. 7c,d**).

## Discussion

Our study demonstrated that IgG1, IgA1 and IgM Abs against spike and RBD were

highly prevalent in the plasma samples of convalescent COVID-19 patients approximately one to

two months after infection. The presence of these isotypes was detected within 7-8 days after the

onset of symptoms. Importantly, all three isotypes show the capacity to mediate virus

neutralization. While regression analyses indicate the strongest contribution of IgM and IgG1

Abs to neutralizing activity, direct testing of purified isotype fractions showed that IgA also

contributed to neutralizing activity, indicating the protective potential of all three major Ig

isotypes. These data carry important implications for the use of convalescent plasma and

hyperimmunoglobulin as COVID-19 therapeutic modalities, suggesting that the selection be

based on the measurement of all of these Ig isotypes.

While all COVID-19 convalescent individuals exhibited plasma/serum neutralization

activities reaching 50% neutralization, and 26 of 28 specimens attained 90% neutralization,

neutralization levels were highly variable with reciprocal $IC_{50}$ and $IC_{90}$ titers ranging over three

orders of magnitude. The titers were comparable against the initial Wuhan strain and the

currently prevalent D614G strain. Similarly, the levels of spike- and RBD-binding total Ig and Ig

isotypes varied greatly.

A trend toward higher levels of total Ig and each Ig isotype was seen in female compared

to male subjects, as reported in another study[56]. Moreover, except for TF#11 (a male elite

neutralizer), the median neutralizing reciprocal $IC_{90}$ titer was higher in females than males,

although the difference did not reach significance (data not shown). Sex differences in Ab

induction have been observed following vaccination against influenza in humans and mice and

were shown to result from the impact of sex steroids[57,58]. Whether and to what extent this

contributes to sex differences seen in clinical outcomes of COVID-19[59] remains to be

investigated. Other studies have shown that the Ab levels were associated with multiple factors,

including time from disease onset[60] and disease severity[30]. However, other than sex, clinical data

are not available for the subjects studied here, limiting our analysis only to neutralization and Ig

isotypes.

One remarkable finding from our study is that although neutralization titers correlated

with binding levels of IgM and IgG1 and not with those of IgA1 or IgA2, purified IgA fractions

from convalescent COVID-19 patients exhibited significant neutralizing activities. The

importance of this finding is underscored by the data showing that IgA1 was the prominent

isotype in some plasma samples such as TF#7 and TF#24 and that IgA1 could be detected early,

within a week after symptom onset. Data from other studies also supported IgA's significance in

that purified IgA fractions exhibited more or as potent neutralizing activities compared to

purified IgG and that RBD-binding IgA correlated as strongly as the IgG equivalent with micro-

neutralization titers[61,62]. The presence of IgA was also detected in the saliva and bronchoalveolar

lavage samples from COVID-19 patients[62,63]. Nonetheless, Wang *et al.* reported that plasma IgA

monomers were less potent than the plasma IgG and secretory IgA counterparts[64]. In our study,

neutralization activities detected in the IgA fractions were mediated mainly by IgA1, the

predominant IgA isotype in the plasma, and the $IC_{50}$ potency of the IgA fraction was ~4-fold

lower than those of IgM and IgG1 fractions. This difference cannot be explained entirely by

lower amounts of spike-specific IgA1 in the tested fractions, as estimations using spike-specific

monoclonal antibodies of the respective isotypes yielded similar concentrations in IgA1 and IgM

fractions (median of 2 and 2.5 µg/mL respectively). Fine epitope specificity and affinity may

differ for IgA, IgM, and IgG to impact neutralization potency, but are yet to be evaluated.

In addition to neutralization, non-neutralizing Ab activities have been implicated in

protection from virus infection through potent Fc-mediated functions such as antibody-dependent

cellular cytotoxicity (ADCC), antibody-dependent cellular phagocytosis (ADCP), and

complement-mediated lysis; this is reported for HIV, influenza, Marburg, and Ebola viruses[44,65–

68]. The Fc activities were not evaluated in our study, and their contribution to protection against

infection and disease progression in humans is yet unclear[15,69–71]. Interestingly, a recent study

demonstrated enrichment of spike-specific IgM and IgA1 Abs and spike-specific phagocytic and

antibody-dependent complement deposition (ADCD) activity in plasma of individuals who

recovered from a SARS-CoV-2 infection, while nucleocapsid-specific IgM and IgA2 responses

and nucleocapsid-specific ADCD activity were features enriched in deceased patients[72]. DNA

vaccines expressing full-length and truncated spike proteins could curtail SARS-CoV-2 infection

in the respiratory tract by varying degrees in rhesus macaques. Virus reduction correlated with

levels of neutralization and also Fc-mediated effector functions such as ADCD[15]. Interestingly,

these DNA vaccines elicited spike- and RBD-specific IgG1, IgG2, IgG3, IgA, and IgM Abs, and

similar to our findings, neutralization correlated most strongly with IgM. Adenovirus serotype 26

vaccine vectors encoding seven different SARS-CoV-2 spike variants showed varying protection

levels, and virus reduction correlated best with neutralizing Ab titers together with IgM binding

levels, FcγRII-binding, and ADCD responses[73]. Defining the full potential of Abs against SARS-

CoV-2 that includes neutralizing, non-neutralizing and enhancing activities are vital for

determining the optimal convalescent Ab treatment against COVID-19 and assessing the

potential efficacy of COVID-19 vaccine candidates.

When we examined plasma specimens collected within 7-8 days after COVID-19

symptom onset, we detected IgG and IgA against spike and RBD, as well as IgM. This is

consistent with published reports showing that 100% of COVID-19-infected individuals

developed IgG within 19 days after symptom onset and that seroconversion for IgG and IgM

occurred simultaneously or sequentially[30]. IgA was also found early after infection (4-6 days

after symptom onset) and increased over time in several other studies[28,35,63,74]. These studies

suggest that measuring total Ig, rather than IgG, would provide a better outcome for early disease

diagnosis. Indeed, we found no correlation between the levels of different isotypes examined in

our study (data not shown). This lack of correlation may result from their asynchronous,

sequential induction (IgM first, then IgG, and finally IgA). Still, IgA's presence early during

acute infection also suggests the potential contribution of natural IgA, which, similar to natural

IgM, arises spontaneously from innate B1 cells to provide the initial humoral responses before

the induction, maturation, and class-switching of adaptive classical B cells[75,76].

In summary, this study demonstrates that spike- and RBD-specific IgM, IgG1, and IgA1

Abs were present in serum and plasma of all or almost all analyzed COVID-19 convalescent

subjects and were detected at very early stages of infection. The plasma of convalescent

individuals also displayed neutralization activities mediated by IgM, IgG, and IgA1, although

neutralization titers correlated more strongly with IgM and IgG levels. The contribution of IgM,

IgG, and IgA Abs to the neutralizing activities against SARS-CoV-2 demonstrates their

importance in the efficacy of convalescent plasma used for COVID-19 treatment.

**Methods**

**Recombinant proteins.** The recombinant spike and RBD proteins were produced as

previously described[77] in Expi293F cells (ThermoFisher) by transfections of purified DNA using

an ExpiFectamine Transfection Kit (ThermoFisher). The soluble version of the spike protein

included the protein ectodomain (amino acids 1-1213), a C-terminal thrombin cleavage site, a T4

foldon trimerization domain, and a hexahistidine tag. The protein sequence was also modified to

remove the polybasic cleavage site (RRAR to A) and two stabilizing mutations (K986P and

V987P, wild type numbering). The RBD (amino acids 319-541) included the signal peptide

(amino acids 1–14) and a hexahistidine tag. Supernatants from transfected cells were harvested

three days after the transfection by centrifugation of the culture at 4000 g for 20 minutes. The

supernatant was then incubated with 6 mL Ni-NTA agarose (Qiagen) for one to two hours at

room temperature. Next, gravity-flow columns were used to collect the Ni-NTA agarose, and the

protein was eluted. Each protein was concentrated in Amicon centrifugal units (EMD Millipore)

and resuspended in phosphate buffered saline (PBS).

**Human samples.** Twenty-five citrated plasma of convalescent COVID-19 individuals

destined for transfusion to SARS-CoV-2-infected individuals (TF#1-25, collected between

March 26th and April 7th 2020) and ten specimens derived from the blood bank (N#4-13),

representing contemporary COVID-19-negative blood bank donors, were obtained from the

Division of Transfusion Medicine of the Department of Pathology, Molecular and Cell-Based

Medicine (Mount Sinai Hospital System, IRB #20-03574). Four additional de-identified serum

specimens from individuals with COVID-19 (P#5-8) were provided by the Clinical Pathology

Division of the Department of Pathology, Molecular and Cell-Based Medicine at the Icahn

School of Medicine at Mount Sinai. Serum and plasma samples were also obtained from study

participants enrolled in IRB-approved protocols at the Icahn School of Medicine at Mount Sinai

(Icahn School of Medicine at Mount Sinai IRB #16-00772, #16-00791, #17-01243) and the

James J. Peter VA Medical Center (IRB #BAN-1604). Samples from these protocols included

sera from seven participants with documented SARS-CoV-2 infection (P#1 d8, d11, and d15

after symptom onset, P#2 d7 and d10 after symptom onset, and RP#1-5 after convalescence), and

sera from twelve healthy donors (N#1-3, N#14-22) collected prior to the spread of SARS-CoV-2

in the USA. All study participants provided written consent at enrollment and agreed to sample

banking and future research use of their banked biospecimens. All samples were heat-inactivated

and/or treated with 0.05% Triton X-100 prior to use.

**Ig fractionation.** IgA was first isolated from plasma by mixing 1:2 diluted plasma with

peptide M agarose beads (600 µL/28 mL plasma, InvivoGen #GEL-PDM) for 1.5 hours at room

temperature. Beads were then collected on a column and washed with PBS until protein reading

(280 nm) by Nanodrop reached background. IgA was eluted from beads with a pH 2.8 elution

buffer (Thermo Scientific #21004) and neutralized with pH 9 Tris buffer. The pass-through

plasma sample was collected for IgG enrichment using protein G agarose beads (InvivoGen

#GEL-AGG) using the same protocol as above and subsequently for IgM isolation using a

HiTrap IgM column (G.E. Healthcare #17-5110-01) according to the manufacturer's instruction.

An additional purification step was performed using Protein A Plus mini-spin columns to

separate IgG from IgM. Protein concentrations were determined with Nanodrop prior to use in

Luminex and neutralization experiments.

**Multiplex bead Ab binding assay.** The SARS-CoV-2 antigens used in this assay

consisted of a soluble recombinant trimerized form of the spike protein and a recombinant RBD

protein[78]. Antigens were coupled to beads as previously described, with minor changes[53]. Each

antigen was covalently coupled individually to a uniquely labeled fluorochrome carboxylated

xMAP bead set at 2.0 μg protein/million beads using a two-step carbodiimide reaction with the

xMAP Ab Coupling (AbC) Kit following to the manufacturers' instructions (Luminex, Austin,

TX). The coupled beads were pelleted, resuspended at $5 \times 10^6$ beads/mL in storage buffer (PBS,

0.1% bovine serum albumin (BSA), 0.02% Tween-20, and 0.05% sodium azide, pH 7.4), and

stored at -80°C. Three to five million beads per batch were prepared in a 1.5 mL conical tube.

Before each experiment, the beads needed for a single run (2,500 beads/well x number of

wells) were pelleted and resuspended in assay buffer (PBS, 0.1% B.S.A., 0.02% Tween-20) to

deliver 2,500 beads in a volume of 50 μL/well in a 96-well plate. Sera/plasma samples were

serially titrated (1:50 to 1:6400 final dilution) or diluted in assay buffer to 1:100 (for a final

dilution of 1:200). The samples were then added as 50 μL/well to the wells containing the beads

and incubated at room temperature for 1 hour at 600 rpm. After two washes in assay buffer, 100

μL/well of biotinylated antibodies specific for total Ig, IgG1, IgG2, IgG3, IgG4, IgA1, IgA2, or

IgM was added and incubated for 30 minutes at room temperature on a plate shaker; these

antibodies were rabbit biotinylated-anti-human total Ig (Abcam, catalog #ab97158) at 2 μg/mL,

mouse biotinylated-anti-human IgG1 Fc (Invitrogen #MH1515) at 4 μg/mL, mouse biotinylated-

anti-human IgG2 Fc (Southern Biotech #9060-08) at 1 μg/mL, mouse biotinylated-anti-human

IgG3 Hinge (Southern Biotech #9210-08) at 3 μg/mL, mouse biotinylated-anti-human IgG4 Fc

(Southern Biotech #9200-08) at 4 μg/mL, mouse biotinylated-anti-human IgA1 Fc (Southern

Biotech #9130-08) at 4 μg/mL, mouse biotinylated-anti-human IgA2 Fc (Southern Biotech

#9140-08) at 4 µg/mL or goat biotinylated-anti-human IgM (Southern Biotech #2020-08) at 3

µg/mL. After two washes, 100 µL/well of Streptavidin-Phycoerythrin (P.E.) at 1 µg/mL was

added (BioLegend #405204) followed by a 30 minutes incubation at room temperature on a plate

shaker. After two additional washes, 100 µL of assay buffer/well was added and put on a shaker

to resuspend the beads. Each plate was read with a Luminex Flexmap 3D instrument. Specimens

were tested in duplicate, and the results were recorded as mean fluorescent intensity (MFI).

**COV2pp production and titration.** The SARS-CoV-2 pseudoviruses (COV2pp) with

wild-type (WT) or D614G mutated spike proteins were produced as previously described[54].

Briefly, 293T cells were transfected to overexpress SARS-CoV-2 glycoproteins. For background

entry with particles lacking a viral surface glycoprotein, pCAGG empty vector was transfected

into 293T cells. Around 8 hours post-transfection, cells were infected with the VSVΔG-rLuc

reporter virus for 2 hours and then washed with PBS. Two days post-infection, supernatants were

collected and clarified by centrifugation at 1250 rpm for 5 minutes. At the time of collection, a

small batch of VSVΔG-rLuc particles bearing the CoV2pp was treated with TPCK-treated

trypsin (Sigma-Aldrich #T1426-1G) at room temperature for 15 minutes prior to inhibition with

soybean trypsin inhibitor (Fisher Scientific #1707502). Finally, particles were aliquoted prior to

storage in -80°C.

The pseudoviruses were titrated on 20,000 Vero-CCL81 cells seeded in a 96-well black

plate with clear bottom 24 hours before infection. At 18 to 22 hours post-infection, the infected

cells were washed with PBS and processed for detection of Renilla luciferase activity with

Renilla-Glo™ Luciferase Assay System (Promega #E2720). A Cytation3 (BioTek) instrument

was used to read luminescence.

**COV2pp neutralization.** The day before infection, 20,000 Vero-CCL81 cells per well

were seeded in a 96-well black plate with clear bottom. On the assay day, the COV2pp WT or

D614G virus was diluted based on the titration results, and 82.5 µL added to all wells, except six

for cell control, of a 96-well V-bottom plate. The seven serial dilutions of the samples were then

prepared in another 96-well V-bottom plate. Samples of the COVID-19-infected individuals and

COVID-19-negative individuals were tested at 4-fold dilutions from 1:10 to 1:40,960. Purified

IgM, IgG, and IgA fractions and negative control fractions were tested at 4-fold dilutions from

500 to 0.02 µg/mL. 27.5 µL per well of the diluted sample was then added to the plate with the

pseudovirus. For each plate, six wells were kept with the virus only, as virus control, and six

wells with media only, as cell control. The plates were then incubated for 30 minutes at room

temperature. 100 µL of the virus/sample mix or virus or medium alone was then added to the

cells and spinoculated by centrifugation at 1250 rpm for 1 hour at room temperature. After

incubation for 18 to 22 hours at $37^{\circ}$C, the measurement of infection/neutralization was performed

as described for the COV2pp titration.

The percentage of neutralization was calculated with the formula: 100-((sample's R.L.U.

– cell control R.L.U.) / virus control R.L.U.) *100). The inhibitory concentration 50% ($IC_{50}$) and

90% ($IC_{90}$) were respectively defined as the reciprocal sample dilution or purified Ig fraction

concentration achieving 50% and 90% neutralization.

**Statistical analysis.** Statistical tests that included two-tailed Mann-Whitney test,

Spearman rank-order correlation test, and simple linear regressions were performed as

designated in the figure legends. All statistical tests were performed using GraphPad Prism 8.

**Data availability.**

The raw data that support the findings of this study are available from the corresponding

author upon request.

**References.**

1.        Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al. A Novel Coronavirus from

Patients with Pneumonia in China, 2019. N Engl J Med. 2020 20;382(8):727–33.

2.        WHO Coronavirus Disease (COVID-19) Dashboard [Internet]. [cited 2020 Jun 29].

Available from: https://covid19.who.int

3.        Liu STH, Lin H-M, Baine I, Wajnberg A, Gumprecht JP, Rahman F, et al. Convalescent

plasma treatment of severe COVID-19: a propensity score-matched control study. Nat Med.

2020 Sep 15;

4.        Martinez-Resendez MF, Castilleja-Leal F, Torres-Quintanilla A, Rojas-Martinez A,

Garcia-Rivas G, Ortiz-Lopez R, et al. Initial experience in Mexico with convalescent plasma in

COVID-19 patients with severe respiratory failure, a retrospective case series. medRxiv. 2020

Jul 20;2020.07.14.20144469.

5.        Altuntas F, Ata N, Yigenoglu TN, Bascı S, Dal MS, Korkmaz S, et al. Convalescent

plasma therapy in patients with COVID-19. Transfus Apher Sci Off J World Apher Assoc Off J

Eur Soc Haemapheresis. 2020 Sep 19;102955.

6.        Zeng H, Wang D, Nie J, Liang H, Gu J, Zhao A, et al. The efficacy assessment of

convalescent plasma therapy for COVID-19 patients: a multi-center case series. Signal Transduct

Target Ther. 2020 Oct 6;5(1):1–12.

7.        Ibrahim D, Dulipsingh L, Zapatka L, Eadie R, Crowell R, Williams K, et al. Factors

Associated with Good Patient Outcomes Following Convalescent Plasma in COVID-19: A

Prospective Phase II Clinical Trial. Infect Dis Ther. 2020 Sep 20;1–14.

8.    Janssen M, Schäkel U, Fokou CD, Krisam J, Stermann J, Kriegsmann K, et al. A

Randomized Open label Phase-II Clinical Trial with or without Infusion of Plasma from Subjects

after Convalescence of SARS-CoV-2 Infection in High-Risk Patients with Confirmed Severe

SARS-CoV-2 Disease (RECOVER): A structured summary of a study protocol for a randomised

controlled trial. Trials. 2020 Dec;21(1):1–4.

9.    NIH expands clinical trials to test convalescent plasma against COVID-19 [Internet].

National Institutes of Health (NIH). 2020 [cited 2020 Oct 7]. Available from:

https://www.nih.gov/news-events/news-releases/nih-expands-clinical-trials-test-convalescent-

plasma-against-covid-19

10.    Convalescent Plasma to Limit SARS-CoV-2 Associated Complications - Full Text View

- ClinicalTrials.gov [Internet]. [cited 2020 Nov 2]. Available from:

https://clinicaltrials.gov/ct2/show/NCT04373460

11.    Convalescent Plasma in Outpatients With COVID-19 - Full Text View -

ClinicalTrials.gov [Internet]. [cited 2020 Nov 2]. Available from:

https://clinicaltrials.gov/ct2/show/NCT04355767

12.    Le TT, Andreadakis Z, Kumar A, Román RG, Tollefsen S, Saville M, et al. The COVID-

19 vaccine development landscape. Nat Rev Drug Discov. 2020 Apr 9;19(5):305–6.

13.    Amanat F, Krammer F. SARS-CoV-2 Vaccines: Status Report. Immunity. 2020 Apr 3;

14.    Mulligan MJ, Lyke KE, Kitchin N, Absalon J, Gurtman A, Lockhart SP, et al. Phase 1/2

Study to Describe the Safety and Immunogenicity of a COVID-19 RNA Vaccine Candidate

(BNT162b1) in Adults 18 to 55 Years of Age: Interim Report. medRxiv. 2020 Jul

1;2020.06.30.20142570.

15.    Yu J, Tostanoski LH, Peter L, Mercado NB, McMahan K, Mahrokhian SH, et al. DNA

vaccine protection against SARS-CoV-2 in rhesus macaques. Science. 2020 May 20;

16.    Yan R, Zhang Y, Li Y, Xia L, Guo Y, Zhou Q. Structural basis for the recognition of

SARS-CoV-2 by full-length human ACE2. Science. 2020 27;367(6485):1444–8.

17.    Hoffmann M, Kleine-Weber H, Krüger N, Müller M, Drosten C, Pöhlmann S. The novel

coronavirus 2019 (2019-nCoV) uses the SARS-coronavirus receptor ACE2 and the cellular

protease TMPRSS2 for entry into target cells. bioRxiv. 2020 Jan 31;2020.01.31.929042.

18.    Wang K, Chen W, Zhou Y-S, Lian J-Q, Zhang Z, Du P, et al. SARS-CoV-2 invades host

cells via a novel route: CD147-spike protein. bioRxiv. 2020 Mar 14;2020.03.14.988345.

19.    Zhou P, Yang X-L, Wang X-G, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak

associated with a new coronavirus of probable bat origin. Nature. 2020;579(7798):270–3.

20.    Lu R, Zhao X, Li J, Niu P, Yang B, Wu H, et al. Genomic characterisation and

epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding.

Lancet Lond Engl. 2020 22;395(10224):565–74.

21.    Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, Erichsen S, et al.

SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically

Proven Protease Inhibitor. Cell. 2020 16;181(2):271-280.e8.

22.    Walls AC, Park Y-J, Tortorici MA, Wall A, McGuire AT, Veesler D. Structure,

Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. Cell. 2020 16;181(2):281-

292.e6.

23.    Vidarsson G, Dekkers G, Rispens T. IgG Subclasses and Allotypes: From Structure to

Effector Functions. Front Immunol [Internet]. 2014 Oct 20 [cited 2018 Jun 12];5. Available

from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4202688/

24.    Schroeder HW, Cavacini L. Structure and function of immunoglobulins. J Allergy Clin

Immunol. 2010 Feb;125(2 Suppl 2):S41-52.

25.    Boes M. Role of natural and immune IgM antibodies in immune responses. Mol

Immunol. 2000 Dec;37(18):1141–9.

26.    Müller R, Gräwert MA, Kern T, Madl T, Peschek J, Sattler M, et al. High-resolution

structures of the IgM Fc domains reveal principles of its hexamer formation. Proc Natl Acad Sci.

2013 Jun 18;110(25):10183–8.

27.    Brekke OH, Sandlie I. Therapeutic antibodies for human diseases at the dawn of the

twenty-first century. Nat Rev Drug Discov. 2003 Jan;2(1):52–62.

28.    Ma H, Zeng W, He H, Zhao D, Jiang D, Zhou P, et al. Serum IgA, IgM, and IgG

responses in COVID-19. Cell Mol Immunol. 2020 Jul;17(7):773–5.

29.    Grifoni A, Weiskopf D, Ramirez SI, Mateus J, Dan JM, Moderbacher CR, et al. Targets

of T Cell Responses to SARS-CoV-2 Coronavirus in Humans with COVID-19 Disease and

Unexposed Individuals. Cell. 2020 Jun 25;181(7):1489-1501.e15.

30.    Long Q-X, Liu B-Z, Deng H-J, Wu G-C, Deng K, Chen Y-K, et al. Antibody responses

to SARS-CoV-2 in patients with COVID-19. Nat Med. 2020 Jun;26(6):845–8.

31.    Robbiani DF, Gaebler C, Muecksch F, Lorenzi JCC, Wang Z, Cho A, et al. Convergent

antibody responses to SARS-CoV-2 in convalescent individuals. Nature. 2020 Jun 18;

32.    Zhao J, Yuan Q, Wang H, Liu W, Liao X, Su Y, et al. Antibody responses to SARS-

CoV-2 in patients of novel coronavirus disease 2019. Clin Infect Dis Off Publ Infect Dis Soc

Am. 2020 Mar 28;

33.    Okba NMA, Müller MA, Li W, Wang C, GeurtsvanKessel CH, Corman VM, et al.

Severe Acute Respiratory Syndrome Coronavirus 2-Specific Antibody Responses in Coronavirus

Disease Patients. Emerg Infect Dis. 2020 Jul;26(7):1478–88.

34.      Guo L, Ren L, Yang S, Xiao M, Chang D, Yang F, et al. Profiling Early Humoral

Response to Diagnose Novel Coronavirus Disease (COVID-19). Clin Infect Dis Off Publ Infect

Dis Soc Am. 2020 Mar 21;

35.      Padoan A, Sciacovelli L, Basso D, Negrini D, Zuin S, Cosma C, et al. IgA-Ab response

to spike glycoprotein of SARS-CoV-2 in patients with COVID-19: A longitudinal study. Clin

Chim Acta. 2020 Aug 1;507:164–6.

36.      Ju B, Zhang Q, Ge J, Wang R, Sun J, Ge X, et al. Human neutralizing antibodies elicited

by SARS-CoV-2 infection. Nature. 2020 May 26;

37.      Wu F, Wang A, Liu M, Wang Q, Chen J, Xia S, et al. Neutralizing antibody responses to

SARS-CoV-2 in a COVID-19 recovered patient cohort and their implications. medRxiv. 2020

Apr 20;2020.03.30.20047365.

38.      Prévost J, Gasser R, Beaudoin-Bussières G, Richard J, Duerr R, Laumaea A, et al. Cross-

Sectional Evaluation of Humoral Responses against SARS-CoV-2 Spike. Cell Rep Med. 2020

Oct 20;1(7):100126.

39.      Rogers TF, Zhao F, Huang D, Beutler N, Burns A, He W, et al. Isolation of potent SARS-

CoV-2 neutralizing antibodies and protection from disease in a small animal model. Science

[Internet]. 2020 Jun 15 [cited 2020 Jul 16]; Available from:

https://science.sciencemag.org/content/early/2020/06/15/science.abc7520

40.      Zost SJ, Gilchuk P, Case JB, Binshtein E, Chen RE, Nkolola JP, et al. Potently

neutralizing and protective human antibodies against SARS-CoV-2. Nature. 2020 Jul 15;1–10.

41.      Ejemel M, Li Q, Hou S, Schiller ZA, Wallace AL, Amcheslavsky A, et al. IgA MAb

blocks SARS-CoV-2 Spike-ACE2 interaction providing mucosal immunity. BioRxiv Prepr Serv

Biol. 2020 May 15;

42.    Renegar KB, Small PA, Boykins LG, Wright PF. Role of IgA versus IgG in the control of

influenza viral infection in the murine respiratory tract. J Immunol Baltim Md 1950. 2004 Aug

1;173(3):1978–86.

43.    Suzuki T, Kawaguchi A, Ainai A, Tamura S, Ito R, Multihartina P, et al. Relationship of

the quaternary structure of human secretory IgA to neutralization of influenza virus. Proc Natl

Acad Sci U S A. 2015 Jun 23;112(25):7809–14.

44.    Krammer F. The human antibody response to influenza A virus infection and vaccination.

Nat Rev Immunol. 2019;19(6):383–97.

45.    Reynolds HY. Immunoglobulin G and its function in the human respiratory tract. Mayo

Clin Proc. 1988 Feb;63(2):161–74.

46.    Muramatsu M, Yoshida R, Yokoyama A, Miyamoto H, Kajihara M, Maruyama J, et al.

Comparison of antiviral activity between IgA and IgG specific to influenza virus hemagglutinin:

increased potential of IgA for heterosubtypic immunity. PloS One. 2014;9(1):e85582.

47.    Shen C, Zhang M, Chen Y, Zhang L, Wang G, Chen J, et al. An IgM antibody targeting

the receptor binding site of influenza B blocks viral infection with great breadth and potency.

Theranostics. 2019;9(1):210–31.

48.    Habibi MS, Jozwik A, Makris S, Dunning J, Paras A, DeVincenzo JP, et al. Impaired

Antibody-mediated Protection and Defective IgA B-Cell Memory in Experimental Infection of

Adults with Respiratory Syncytial Virus. Am J Respir Crit Care Med. 2015 May 1;191(9):1040–

9.

49.    Du L, He Y, Zhou Y, Liu S, Zheng B-J, Jiang S. The spike protein of SARS-CoV--a

target for vaccine and therapeutic development. Nat Rev Microbiol. 2009 Mar;7(3):226–36.

50.    Schurink B, Roos E, Radonic T, Barbe E, Bouman CSC, de Boer HH, et al. Viral

presence and immunopathology in patients with lethal COVID-19: a prospective autopsy cohort

study. Lancet Microbe. 2020 Sep 25;

51.    Lindner D, Fitzek A, Bräuninger H, Aleshcheva G, Edler C, Meissner K, et al.

Association of Cardiac Infection With SARS-CoV-2 in Confirmed COVID-19 Autopsy Cases.

JAMA Cardiol. 2020 Jul 27;

52.    Remmelink M, De Mendonça R, D'Haene N, De Clercq S, Verocq C, Lebrun L, et al.

Unspecific post-mortem findings despite multiorgan viral spread in COVID-19 patients. Crit

Care Lond Engl. 2020 12;24(1):495.

53.    Weiss S, Klingler J, Hioe C, Amanat F, Baine I, Kojic EM, et al. A High Through-put

Assay for Circulating Antibodies Directed against the S Protein of Severe Acute Respiratory

Syndrome Corona virus 2. medRxiv. 2020 Apr 17;2020.04.14.20059501.

54.    Oguntuyo KY, Stevens CS, Hung C-T, Ikegame S, Acklin JA, Kowdle SS, et al.

Quantifying absolute neutralization titers against SARS-CoV-2 by a standardized virus

neutralization assay allows for cross-cohort comparisons of COVID-19 sera. medRxiv. 2020

Aug 15;2020.08.13.20157222.

55.    Korber B, Fischer WM, Gnanakaran S, Yoon H, Theiler J, Abfalterer W, et al. Tracking

Changes in SARS-CoV-2 Spike: Evidence that D614G Increases Infectivity of the COVID-19

Virus. Cell. 2020 Aug 20;182(4):812-827.e19.

56.    Zeng F, Dai C, Cai P, Wang J, Xu L, Li J, et al. A comparison study of SARS-CoV-2 IgG

antibody between male and female COVID-19 patients: a possible reason underlying different

outcome between gender. medRxiv. 2020 Mar 27;2020.03.26.20040709.

57.    Fink AL, Engle K, Ursin RL, Tang W-Y, Klein SL. Biological sex affects vaccine

efficacy and protection against influenza in mice. Proc Natl Acad Sci U S A. 2018

04;115(49):12477–82.

58.    Potluri T, Fink AL, Sylvia KE, Dhakal S, Vermillion MS, vom Steeg L, et al. Age-

associated changes in the impact of sex steroids on influenza vaccine responses in males and

females. Npj Vaccines. 2019 Jul 12;4(1):1–12.

59.    Jin J-M, Bai P, He W, Wu F, Liu X-F, Han D-M, et al. Gender Differences in Patients

With COVID-19: Focus on Severity and Mortality. Front Public Health [Internet]. 2020 [cited

2020 Aug 6];8. Available from:

https://www.frontiersin.org/articles/10.3389/fpubh.2020.00152/full

60.    Long Q-X, Tang X-J, Shi Q-L, Li Q, Deng H-J, Yuan J, et al. Clinical and

immunological assessment of asymptomatic SARS-CoV-2 infections. Nat Med. 2020 Jun 18;1–

5.

61.    Mazzini L, Martinuzzi D, Hyseni I, Lapini G, Benincasa L, Piu P, et al. Comparative

analyses of SARS-CoV-2 binding (IgG, IgM, IgA) and neutralizing antibodies from human

serum samples. bioRxiv. 2020 Aug 10;2020.08.10.243717.

62.    Sterlin D, Mathian A, Miyara M, Mohr A, Anna F, Claer L, et al. IgA dominates the early

neutralizing antibody response to SARS-CoV-2. medRxiv. 2020 Jun 11;2020.06.10.20126532.

63.    Isho B, Abe KT, Zuo M, Jamal AJ, Rathod B, Wang JH, et al. Persistence of serum and

saliva antibody responses to SARS-CoV-2 spike antigens in COVID-19 patients. Sci Immunol

[Internet]. 2020 Oct 8 [cited 2020 Oct 9];5(52). Available from:

https://immunology.sciencemag.org/content/5/52/eabe5511

64.    Wang Z, Lorenzi JCC, Muecksch F, Finkin S, Viant C, Gaebler C, et al. Enhanced

SARS-CoV-2 Neutralization by Secretory IgA in vitro. bioRxiv. 2020 Sep 9;2020.09.09.288555.

65.     Horwitz, Bar-On Y, Lu C-L, Fera D, Lockhart AAK, Lorenzi JCC, et al. Non-

neutralizing Antibodies Alter the Course of HIV-1 Infection In Vivo. Cell. 2017 Jul 26;

66.     Chung AW, Ghebremichael M, Robinson H, Brown E, Choi I, Lane S, et al.

Polyfunctional Fc-effector profiles mediated by IgG subclass selection distinguish RV144 and

VAX003 vaccines. Sci Transl Med. 2014 Mar 19;6(228):228ra38.

67.     Ilinykh PA, Huang K, Santos RI, Gilchuk P, Gunn BM, Karim MM, et al. Non-

neutralizing Antibodies from a Marburg Infection Survivor Mediate Protection by Fc-Effector

Functions and by Enhancing Efficacy of Other Antibodies. Cell Host Microbe. 2020

10;27(6):976-991.e11.

68.     Gunn BM, Yu W-H, Karim MM, Brannan JM, Herbert AS, Wec AZ, et al. A Role for Fc

Function in Therapeutic Monoclonal Antibody-Mediated Protection against Ebola Virus. Cell

Host Microbe. 2018 08;24(2):221-233.e5.

69.     Zohar T, Alter G. Dissecting antibody-mediated protection against SARS-CoV-2. Nat

Rev Immunol. 2020;20(7):392–4.

70.     Pinto D, Park Y-J, Beltramello M, Walls AC, Tortorici MA, Bianchi S, et al. Cross-

neutralization of SARS-CoV-2 by a human monoclonal SARS-CoV antibody. Nature.

2020;583(7815):290–5.

71.     Natarajan H, Crowley AR, Butler SE, Xu S, Weiner JA, Bloch EM, et al. SARS-CoV-2

antibody signatures robustly predict diverse antiviral functions relevant for convalescent plasma

therapy. MedRxiv Prepr Serv Health Sci. 2020 Sep 18;

72.     Atyeo C, Fischinger S, Zohar T, Slein MD, Burke J, Loos C, et al. Distinct Early

Serological Signatures Track with SARS-CoV-2 Survival. Immunity [Internet]. 2020 Jul 30

[cited 2020 Aug 12]; Available from:

http://www.sciencedirect.com/science/article/pii/S1074761320303277

73. Mercado NB, Zahn R, Wegmann F, Loos C, Chandrashekar A, Yu J, et al. Single-shot

Ad26 vaccine protects against SARS-CoV-2 in rhesus macaques. Nature. 2020 Jul 30;

74. Yu H-Q, Sun B-Q, Fang Z-F, Zhao J-C, Liu X-Y, Li Y-M, et al. Distinct features of

SARS-CoV-2-specific IgA response in COVID-19 patients. Eur Respir J. 2020;56(2).

75. Meyer-Bahlburg A. B-1 cells as a source of IgA. Ann N Y Acad Sci. 2015

Dec;1362:122–31.

76. Rodriguez-Zhurbenko N, Quach TD, Hopkins TJ, Rothstein TL, Hernandez AM. Human

B-1 Cells and B-1 Cell Antibodies Change With Advancing Age. Front Immunol [Internet].

2019 [cited 2020 Aug 6];10. Available from:

https://www.frontiersin.org/articles/10.3389/fimmu.2019.00483/full

77. Amanat F, Stadlbauer D, Strohmeier S, Nguyen THO, Chromikova V, McMahon M, et

al. A serological assay to detect SARS-CoV-2 seroconversion in humans. Nat Med. 2020 May

12;1–4.

78. Stadlbauer D, Amanat F, Chromikova V, Jiang K, Strohmeier S, Arunkumar GA, et al.

SARS-CoV-2 Seroconversion in Humans: A Detailed Protocol for a Serological Assay, Antigen

Production, and Test Setup. Curr Protoc Microbiol. 2020;57(1):e100.

## Funding

This study was supported by the Microbiology Laboratory Clinical Services at the Mount Sinai Health System and the Mount Sinai Health System Translational Science Hub, National Institutes of Health [grant U54TR001433]; the Personalized Virology Initiative supported by institutional funds and philanthropic donations (to V.S.); the Department of Medicine of the Icahn School of Medicine at Mount Sinai Department of Medicine (to S.Z-P., C.E.H.); the Department of Microbiology and the Ward-Coleman estate for endowing the Ward-Coleman Chairs at the Icahn School of Medicine at Mount Sinai (to B.L.), the National Institute of Allergy and Infectious Diseases Centers of Excellence for Influenza Research and Surveillance [contract HHSN272201400008C] (to F.K., V.S.); the Department of Veterans Affairs [Merit Review Grant I01BX003860] (to C.E.H.) and [Research Career Scientist Award 1IK6BX004607] (to C.E.H.); the National Institute of Allergy and Infectious Diseases [grant AI139290] to C.E.H., [grant AI136916] (to V.S.), [grants R01 AI123449, R21 AI1498033] to B.L. K.Y.O. and C.S. were supported by Viral-Host Pathogenesis Training Grant T32 AI07647; K.Y.O. was additionally supported by F31 AI154739. S.I. and C-T. H. were supported by postdoctoral fellowships from CHOT-SG (Fukuoka University, Japan) and the Ministry of Science and Technology (MOST, Taiwan), respectively.

## Acknowledgments

We thank all the donors for their contribution to research.

## Competing interests

660     The authors declare no competing interests.

## Author contributions

J.K., S.W., G.E-A., S.Z-P., and C.E.H. wrote and edited the manuscript. S.W., J.K., C.E.H., and

S.Z-P. designed the experiments. J.K., S.W., V.I., X.L. performed the experiments and collected

the data. J.K., A.N., S.Z-P. and C.E.H. analyzed the data. K.Y.O., C.S., S.I., C-T.H., F.A., B.L.,

and F.K. provided protocols, antigens, cells and pseudovirus stocks. G.E-A., I.B., S.A., J.C.B.,

E.M.K., J.S., S.L., D.J., M.B-G., and V.S. provided samples. All authors read and approved the

final manuscript.

 **Figure legends**

**Fig. 1. Titration of SARS-CoV-2 spike and RBD total Ig in plasma or serum samples from**

**COVID-19 convalescent individuals.** Titration of (a) spike-specific or (b) RBD-specific total Ig

from 29 COVID-19-convalescent individuals, two acute COVID-19 patients with longitudinal

samples, and 13 COVID-19-negative individuals. Specimens were diluted at 2-fold dilutions

from 1:50 to 1:6,400.

**Fig. 2. Levels of Ig isotypes against the SARS-CoV-2 spike and RBD vary in plasma or**

**serum samples from COVID-19 convalescent individuals.** Detection of total Ig, IgM, IgG1,

IgG2, IgG3, IgG4, IgA1 and IgA2 against (a) spike and (b) RBD in specimens from 29 COVID-

19-convalescent individuals, 13 COVID-19-negative contemporary samples, and pre-pandemic

controls. The samples were tested at a dilution of 1:200 and data are shown as mean MFI +

standard deviation (SD) of duplicate measurements from at least two independent experiments.

The pre-pandemic controls are shown as mean MFI + SD of 12 samples (Pre, black bar). The

horizontal red dotted line represents the cut-off value determined as the mean + 3 SD of 12 pre-

pandemic samples for each of the isotypes. (c) Percentages of responders above the cut-off for

each spike- or RBD-specific Ig isotype.

**Fig. 3. IgG1 is the dominant isotype response induced in COVID-19 convalescent**

**individuals.** Simple linear regression of (a) spike-specific or (b) RBD-specific total Ig levels

versus IgM, IgG1 or IgG2 levels or versus (c) spike-specific and RBD-specific IgG3, IgG4,

IgA1, and IgA2 levels from the 29 COVID-19-convalescent individuals from Fig. 1. The dash

lines represent 95% confidence intervals.

**Fig. 4. Neutralization activities are detected in all COVID-19 convalescent individuals.**

Neutralization of COV2pp with (a,b,c) WT or (d,e) D614G mutated spike proteins by samples

693 from (a,d) 28 COVID-19-convalescent individuals and (b,e) 11 COVID-19-negative individuals,

694 compared to (c) a recombinant soluble RBD (sRBD) control. Each plasma or sera specimen was

695 tested at 4-fold dilutions from 1:10 to 1:40,960, and sRBD was tested at 4-fold dilutions from

696 100 to 0.02 µg/mL. The data are shown as mean percentage of neutralization + SD of triplicate.

697 The extrapolated titration curves were generated using a nonlinear regression model in GraphPad

698 Prism (Inhibitor versus response – variable slope (four parameters), least squares regression).

699 The dotted horizontal lines highlight 50% neutralization. (f) Spearman correlation between the

700 $IC_{90}$ titers against COV2pp WT versus D614G.

701 **Fig. 5. Summary of relative Ig isotype levels and neutralization titers.** Table showing sex

702 (purple, F: female, M: male), relative levels of spike-specific (green) and RBD-specific (blue) Ig

703 isotypes (+: bottom quartile, ++: second quartile, +++: third quartile, ++++: top quartile, -: non-

704 responder) and reciprocal $IC_{50}$ and $IC_{90}$ neutralization titers against WT pseudovirus (orange)

705 and D614G pseudovirus (red) of 29 plasma samples from COVID-19-convalescent individuals

706 (nd: not done).

707 **Fig. 6. IgM and IgG1 contribute most to SARS-CoV-2 neutralization.** Simple linear

708 regression of reciprocal $IC_{90}$ neutralization titers of 27 COVID-19-convalescent individuals

709 versus (a) spike-specific or (b) RBD-specific total Ig, IgM, IgG1 and IgA1 Ab levels. The black

710 dash line shows 95% confidence interval. The dotted vertical red line represents the cut-off

711 (mean of 12 pre-pandemic samples + 3 SD) for each isotype from Fig. 1. (b) Simple linear

712 regression of reciprocal $IC_{90}$ neutralization titers of 27 COVID-19-convalescent individuals

713 versus spike-specific or RBD-specific total IgG2-4 and IgA2 Ab levels.

714 **Fig. 7. Purified IgM, IgG, and IgA fractions display neutralizing activities against SARS-**

715 **CoV-2.** (a) Neutralization of COV2pp by five COVID-19-infected individual plasma samples

(RP#1-5) compared to a COVID-19-negative sample (RN#1). Plasma samples were tested at 4-

fold dilutions from 1:10 to 1:40,960 or 1:20 to 1:81,920. Data are shown as the mean percentage

of neutralization. The dotted horizontal lines highlight 50% neutralization. (b) Reciprocal $IC_{50}$

and $IC_{90}$ neutralization titers of RP#1-5 plasma samples (c) Neutralization of COV2pp by

purified IgM, IgG, and IgA fractions from five COVID-19-infected individuals (RP#1-5)

compared to a control Ig fraction. The fractions were tested at 4-fold dilutions from 500 to 0.02

µg/mL. Data are shown as the mean percentage of neutralization. The dotted horizontal lines

highlight 50% neutralization. (d) $IC_{50}$ of purified IgM, IgG, and IgA fractions from RP#1-5. The

statistical significance was determined by a two-tailed Mann-Whitney test (*: $p < 0.05$, **: $p <$

0.01).

**Supplementary Fig. 1.** Spearman correlations of (a) spike-specific or (b) RBD-specific total Ig

MFI values from two independent experiments to show the degree of assay reproducibility.

**Supplementary Fig. 2.** Spearman correlations of the area under the curves (AUCs) of (a) spike-

or (b) RBD-specific total Ig versus total Ig MFI values at a 1:200 dilution.

**Supplementary Fig. 3.** Isotyping validation was performed by coating Luminex beads with

IgG1, IgG2, IgG3, IgG4, IgA1, IgA2, and IgM myeloma proteins and detecting each specifically

with eight different secondary Abs against (a) total Ig, (b) IgM, (c) IgG1, (d) IgG2, (e) IgG3, (f)

IgG4, (g) IgA1 and (h) IgA2. The data are shown as mean MFI + SD of duplicate.

**Supplementary Fig. 4.** Spearman correlations between spike-specific versus RBD-specific total

Ig, IgM, IgG1, IgG2, IgG3, IgG4, IgA1, or IgA2 MFI values.

**Supplementary Fig. 5.** Violin plots of (a) spike-specific or (b) RBD-specific total Ig, IgM,

IgG1, and IgA1 levels from nine COVID-19 convalescent female (F) and 15 male (M) subjects.

The statistical significance was determined by a two-tailed Mann-Whitney test (ns: non-

significant: p > 0.05).

**Supplementary Fig. 6. Induction of IgA1 and IgG1 along with IgM early after disease**

**onset**. Kinetics of induction of spike-specific (left panel) or RBD-specific (right panel) (a) total

Ig, (b) IgM, (c) IgG1, (d) IgG2, (e) IgG3, (f) IgG4, (g) IgA1, and (h) IgA2 from two COVID-19

patients. Longitudinal samples from each patient were tested at a dilution of 1:200 in parallel

with all negative samples and data are shown as mean MFI + SD of duplicate measurements

from at least two experiments. The dotted red line represents the cut-off value calculated as the

mean of 12 pre-pandemic samples + 3 SD from Fig. 1.

**Supplementary Fig. 7. Enrichment of spike-specific (a) IgM, (b) IgG, and (c) IgA in**

**purified fractions from RP#1-5 and RN#1.** Each fraction was measured for the presence of

IgM, IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2 Abs using the isotyping method validated in

Supplementary Fig. 3.

**a**

Spike

**b**

RBD

| | |
|---|---|
| P#1a | TF#8 |
| P#1b | TF#9 |
| P#1c | TF#10 |
| P#2a | TF#11 |
| P#2b | TF#12 |
| P#3 | TF#13 |
| P#4 | TF#14 |
| P#5 | TF#15 |
| P#6 | TF#16 |
| P#7 | TF#17 |
| P#8 | TF#18 |
| TF#1 | TF#19 |
| TF#2 | TF#20 |
| TF#3 | TF#21 |
| TF#4 | TF#22 |
| TF#5 | TF#23 |
| TF#6 | TF#24 |
| TF#7 | TF#25 |

COVID-19-positive subjects

COVID-19-negative subjects

**Fig. 1. Titration of SARS-CoV-2 spike and RBD total Ig in plasma or serum samples from COVID-19 convalescent individuals.** Titration of (a) spike-specific or (b) RBD-specific total Ig from 29 COVID-19-convalescent individuals, two acute COVID-19 patients with longitudinal samples, and 13 COVID-19-negative individuals. Specimens were diluted at 2-fold dilutions from 1:50 to 1:6,400.

| | Spike | RBD |
|---|---|---|
| Total Ig | 100% | 100% |
| IgM | 100% | 93% |
| IgG1 | 97% | 97% |
| IgG2 | 0% | 0% |
| IgG3 | 7% | 17% |
| IgG4 | 7% | 3% |
| IgA1 | 97% | 93% |
| IgA2 | 17% | 48% |

**Fig. 2. Levels of Ig isotypes against the SARS-CoV-2 spike and RBD vary in plasma or serum samples from COVID-19 convalescent individuals.** Detection of total Ig, IgM, IgG1, IgG2, IgG3, IgG4, IgA1 and IgA2 against (a) spike and (b) RBD in specimens from 29 COVID-19-convalescent individuals, 13 COVID-19-negative contemporary samples, and pre-pandemic controls. The samples were tested at a dilution of 1:200 and data are shown as mean MFI + standard deviation (SD) of duplicate measurements from at least two independent experiments. The pre-pandemic controls are shown as mean MFI + SD of 12 samples (Pre, black bar). The horizontal red dotted line represents the cut-off value determined as the mean + 3 SD of 12 pre-pandemic samples for each of the isotypes. (c) Percentages of responders above the cut-off for each spike- or RBD-specific Ig isotype.

| | Linear regression spike MFI | | | Linear regression RBD MFI | | |
|---|---|---|---|---|---|---|
| | r² | p | | r² | p | |
| Total vs IgG3 | 0.35 | 0.0008 | *** | 0.07 | 0.15 | ns |
| Total vs IgG4 | 0.07 | 0.15 | ns | 0.24 | 0.007 | ¨** |
| Total vs IgA1 | 0.06 | 0.20 | ns | 0.23 | 0.009 | *** |
| Total vs IgA2 | 0.06 | 0.20 | ns | 0.13 | 0.05 | ns |

**Fig. 3. IgG1 is the dominant isotype response induced in COVID-19 convalescent individuals.** Simple linear regression of (a) spike-specific or (b) RBD-specific total Ig levels versus IgM, IgG1 or IgG2 levels or versus (c) spike-specific and RBD-specific IgG3, IgG4, IgA1, and IgA2 levels from the 29 COVID-19-convalescent individuals from Fig. 1. The dash lines represent 95% confidence intervals.

**Fig. 4. Neutralization activities are detected in all COVID-19 convalescent individuals.**
Neutralization of COV2pp with (a,b,c) WT or (d,e) D614G mutated spike proteins by samples from (a,d) 28 COVID-19-convalescent individuals and (b,e) 11 COVID-19-negative individuals, compared to (c) a recombinant soluble RBD (sRBD) control. Each plasma or sera specimen was tested at 4-fold dilutions from 1:10 to 1:40,960, and sRBD was tested at 4-fold dilutions from 100 to 0.02 μg/mL. The data are shown as mean percentage of neutralization + SD of triplicate. The extrapolated titration curves were generated using a nonlinear regression model in GraphPad Prism (Inhibitor versus response – variable slope (four parameters), least squares regression). The dotted horizontal lines highlight 50% neutralization. (f) Spearman correlation between the $IC_{90}$ titers against COV2pp WT versus D614G.

| | Sex | Spike | | | | | | | | RBD | | | | | | | | Neutralization | | Neutralization D614G mutant | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Total Ig | IgM | IgG1 | IgG2 | IgG3 | IgG4 | IgA1 | IgA2 | Total Ig | IgM | IgG1 | IgG2 | IgG3 | IgG4 | IgA1 | IgA2 | IC$_{50}$ | IC$_{90}$ | IC$_{50}$ | IC$_{90}$ |
| P#5 | unknown | ++ | + | + | - | - | - | ++ | - | + | - | + | - | - | - | + | + | 36.6 | 2.3 | nd | nd |
| P#6 | unknown | +++ | + | ++ | - | - | + | + | - | ++ | + | + | - | + | - | + | + | 561.0 | 25.6 | nd | nd |
| P#7 | unknown | + | ++ | + | - | - | - | + | - | ++ | ++ | + | - | - | - | + | - | nd | nd | nd | nd |
| P#8 | unknown | ++ | + | ++ | - | - | + | + | - | + | ++ | + | - | - | - | + | - | 376.2 | 10.2 | nd | nd |
| TF#1 | F | + | + | + | - | - | - | + | - | ++ | + | ++ | - | - | - | + | - | 418.8 | 21.5 | 254.8 | 28.6 |
| TF#2 | M | ++ | + | ++ | - | - | - | + | - | ++ | + | + | - | - | - | + | - | 177.5 | 11.8 | 59.6 | 12.7 |
| TF#3 | M | ++ | + | + | - | - | - | + | - | ++ | + | + | - | - | - | + | + | 164.7 | 19.8 | 206.2 | 87.5 |
| TF#4 | F | + | + | + | - | - | - | + | - | ++ | + | + | - | - | - | + | - | 998.6 | 39.9 | 328.7 | 62.1 |
| TF#5 | M | ++++ | ++ | ++++ | - | - | - | ++ | - | ++++ | +++ | +++ | - | - | - | ++ | + | 2344.7 | 515.9 | 4138.0 | 730.1 |
| TF#6 | F | ++ | + | + | - | - | - | + | - | ++ | + | ++ | - | + | - | + | - | 976.6 | 51.9 | 551.8 | 99.7 |
| TF#7 | F | +++ | + | ++ | - | + | - | ++++ | + | ++++ | + | ++ | - | - | - | ++++ | + | 5788.5 | 90.7 | 452.5 | 81.2 |
| TF#8 | M | + | + | + | - | - | - | + | - | ++ | + | + | - | - | - | + | - | 2840.0 | 142.1 | 700.1 | 129.1 |
| TF#9 | F | +++ | ++ | ++ | - | - | - | +++ | + | +++ | ++ | ++ | - | + | - | ++ | + | 4044.1 | 116.5 | 664.2 | 122.8 |
| TF#10 | M | ++ | + | + | - | - | - | + | - | +++ | + | ++ | - | - | - | + | + | 1060.0 | 72.7 | 572.1 | 75.0 |
| TF#11 | M | ++ | ++++ | + | - | - | - | + | - | ++ | ++++ | + | - | - | - | + | - | > 40960 | 7200.0 | > 40960 | 2344.0 |
| TF#12 | F | + | + | + | - | - | - | + | - | ++ | ++ | + | - | - | - | + | - | 285.5 | 30.0 | 325.7 | 54.12 |
| TF#13 | M | ++ | ++ | + | - | - | - | +++ | + | ++ | +++ | + | - | - | - | + | + | 1697.6 | 144.9 | 1573.0 | 156.1 |
| TF#14 | M | +++ | + | ++ | - | - | - | + | - | ++ | + | + | - | + | - | + | - | 17078.5 | 201.4 | 675.2 | 151.1 |
| TF#15 | M | +++ | + | ++ | - | - | - | - | - | +++ | ++ | ++ | - | - | - | - | - | 2006.2 | 29.6 | 548.5 | 67.9 |
| TF#16 | M | +++ | + | ++ | - | - | - | + | + | +++ | ++ | ++ | - | - | - | + | + | 1330.7 | 106.2 | 1961.0 | 233.0 |
| TF#17 | M | ++ | + | + | - | - | - | + | - | +++ | + | ++ | - | - | - | + | - | 198.0 | 27.2 | 185.9 | 39.8 |
| TF#18 | F | ++ | + | ++ | - | - | - | ++ | - | +++ | + | ++ | - | - | - | + | + | 1122.1 | 56.9 | 453.8 | 64.4 |
| TF#19 | M | ++ | + | + | - | - | - | + | - | ++ | + | + | - | - | - | - | - | 326.2 | 17.7 | 201.4 | 31.8 |
| TF#20 | F | ++ | + | ++ | - | - | - | ++ | - | +++ | +++ | ++ | - | + | - | + | + | 1535.3 | 72.4 | 471.0 | 75.5 |
| TF#21 | M | ++++ | ++ | +++ | - | + | - | ++ | - | ++++ | +++ | ++++ | - | - | - | + | + | 5375.7 | 389.7 | 3231.0 | 411.8 |
| TF#22 | M | ++ | + | ++ | - | - | - | + | - | + | - | ++ | - | - | - | + | + | 623.2 | 4.8 | 46.65 | 11.5 |
| TF#23 | unknown | ++ | + | - | - | - | - | ++ | - | +++ | +++ | - | - | - | - | + | - | 3881.9 | 107.8 | 563.7 | 170.9 |
| TF#24 | M | + | + | + | - | - | - | ++ | + | ++ | + | + | - | - | + | ++ | + | 813.6 | 24.4 | 214.8 | 70.7 |
| TF#25 | F | +++ | + | ++ | - | - | - | + | - | ++++ | + | ++++ | - | - | - | + | - | 958.6 | 78.7 | 605.0 | 134.0 |

**Fig. 5. Summary of relative Ig isotype levels and neutralization titers.** Table showing sex (purple, F: female, M: male), relative levels of spike-specific (green) and RBD-specific (blue) Ig isotypes (+: bottom quartile, ++: second quartile, +++: third quartile, ++++: top quartile, -: non-responder) and reciprocal $IC_{50}$ and $IC_{90}$ neutralization titers against WT pseudovirus (orange) and D614G pseudovirus (red) of 29 plasma samples from COVID-19-convalescent individuals (nd: not done).

**a**

Spike

$r^2 = 0.53$
$p < 0.0001$
****

Spike

$r^2 = 0.62$
$p < 0.0001$
****

Spike

$r^2 = 0.52$
$p < 0.0001$
****

Spike

$r^2 = 0.07$
$p = 0.2$
ns

**b**

RBD

$r^2 = 0.37$
$p = 0.0008$
***

RBD

$r^2 = 0.34$
$p = 0.002$
***

RBD

$r^2 = 0.32$
$p = 0.002$
**

RBD

$r^2 = 0.06$
$p = 0.22$
ns

**c**

|  | Linear regression IC$_{90}$ versus Spike MFI | | | Linear regression IC$_{90}$ versus RBD MFI | | |
|---|---|---|---|---|---|---|
|  | $r^2$ | p |  | $r^2$ | p |  |
| **IgG2** | 0.14 | 0.05 | ns | 0.33 | 0.002 | ** |
| **IgG3** | 0.30 | 0.003 | ** | 0.02 | 0.46 | ns |
| **IgG4** | 0.00 | 0.99 | ns | 0.06 | 0.21 | ns |
| **IgA2** | 0.02 | 0.50 | ns | 0.03 | 0.45 | ns |

**Fig. 6. IgM and IgG1 contribute most to SARS-CoV-2 neutralization.** Simple linear regression of reciprocal $IC_{90}$ neutralization titers of 27 COVID-19-convalescent individuals versus (a) spike-specific or (b) RBD-specific total Ig, IgM, IgG1 and IgA1 Ab levels. The black dash line shows 95% confidence interval. The dotted vertical red line represents the cut-off (mean of 12 pre-pandemic samples + 3 SD) for each isotype from Fig. 1. (b) Simple linear regression of reciprocal $IC_{90}$ neutralization titers of 27 COVID-19-convalescent individuals versus spike-specific or RBD-specific total IgG2-4 and IgA2 Ab levels.

**Fig. 7. Purified IgM, IgG, and IgA fractions display neutralizing activities against SARS-CoV-2.** (a) Neutralization of COV2pp by five COVID-19-infected individual plasma samples (RP#1-5) compared to a COVID-19-negative sample (RN#1). Plasma samples were tested at 4-fold dilutions from 1:10 to 1:40,960 or 1:20 to 1:81,920. Data are shown as the mean percentage of neutralization. The dotted horizontal lines highlight 50% neutralization. (b) Reciprocal $IC_{50}$ and $IC_{90}$ neutralization titers of RP#1-5 plasma samples (c) Neutralization of COV2pp by purified IgM, IgG, and IgA fractions from five COVID-19-infected individuals (RP#1-5) compared to a control Ig fraction. The fractions were tested at 4-fold dilutions from 500 to 0.02 µg/mL. Data are shown as the mean percentage of neutralization. The dotted horizontal lines highlight 50% neutralization. (d) $IC_{50}$ of purified IgM, IgG, and IgA fractions from RP#1-5. The statistical significance was determined by a two-tailed Mann-Whitney test (*: $p < 0.05$, **: $p < 0.01$).

**a** Spike

Spearman r = 0.9638
p < 0.0001
****

**b** RBD

Spearman r = 0.9439
p < 0.0001
****

**Supplementary Fig. 1.** Spearman correlations of (a) spike-specific or (b) RBD-specific total Ig MFI values from two independent experiments to show the degree of assay reproducibility.

**Supplementary Fig. 2.** Spearman correlations of the area under the curves (AUCs) of (a) spike- or (b) RBD-specific total Ig versus total Ig MFI values at a 1:200 dilution.

**Supplementary Fig. 3.** Isotyping validation was performed by coating Luminex beads with IgG1, IgG2, IgG3, IgG4, IgA1, IgA2, and IgM myeloma proteins and detecting each specifically with eight different secondary Abs against (a) total Ig, (b) IgM, (c) IgG1, (d) IgG2, (e) IgG3, (f) IgG4, (g) IgA1 and (h) IgA2. The data are shown as mean MFI + SD of duplicate.

**Supplementary Fig. 4.** Spearman correlations between spike-specific versus RBD-specific total Ig, IgM, IgG1, IgG2, IgG3, IgG4, IgA1, or IgA2 MFI values.

**Supplementary Fig. 5.** Violin plots of (a) spike-specific or (b) RBD-specific total Ig, IgM, IgG1, and IgA1 levels from nine COVID-19 convalescent female (F) and 15 male (M) subjects. The statistical significance was determined by a two-tailed Mann-Whitney test (ns: non-significant: $p > 0.05$).

**Supplementary Fig. 6. Induction of IgA1 and IgG1 along with IgM early after disease onset.** Kinetics of induction of spike-specific (left panel) or RBD-specific (right panel) (a) total Ig, (b) IgM, (c) IgG1, (d) IgG2, (e) IgG3, (f) IgG4, (g) IgA1, and (h) IgA2 from two COVID-19 patients. Longitudinal samples from each patient were tested at a dilution of 1:200 in parallel with all negative samples and data are shown as mean MFI + SD of duplicate measurements from at least two experiments. The dotted red line represents the cut-off value calculated as the mean of 12 pre-pandemic samples + 3 SD from Fig. 1.

**Supplementary Fig. 7. Enrichment of spike-specific (a) IgM, (b) IgG, and (c) IgA in purified fractions from RP#1-5 and RN#1.** Each fraction was measured for the presence of IgM, IgG1, IgG2, IgG3, IgG4, IgA1, and IgA2 Abs using the isotyping method validated in Supplementary Fig. 3.