

Environmental risk factors of airborne viral transmission: Humidity, Influenza and SARS-CoV-2 in the Netherlands

Edsard Ravelli*, Rolando Gonzales Martinez**

* Muiderberg, The Netherlands. Address: Scholekster 18, Postalcode: 1399 KT. Contact: research@ravelli.net

** University of Agder (Kristiansand, Norway). ORCID: <https://orcid.org/0000-0001-5401-780X>

Abstract

Objective The relationship between specific humidity and influenza/SARS-CoV-2 in the Netherlands is evaluated over time and at regional level.

Design Parametric and non-parametric correlation coefficients are calculated to quantify the relationship between humidity and influenza, using five years of weekly data. Bayesian spatio-temporal models—with a Poisson and a Gaussian likelihood—are estimated to find the relationship between regional humidity and the daily cases of SARS-CoV-2 in the municipalities and provinces of the Netherlands.

Results An inverse (negative) relationship is observed between specific humidity and the incidence of influenza between 2015 and 2019. The space-time analysis indicates that an increase of specific humidity of one gram of water vapor per kilogram of air (1 g/kg) is related to a reduction of approximately 5% in the risk of COVID-19 infections.

Conclusions The increase in humidity during the outbreak of the SARS-CoV-2 in the Netherlands helped to reduce the risk of regional COVID-19 infections. Public policies that promote higher levels of specific humidification—above 6 g/Kg—can lead to significant reductions in the spread of respiratory viruses, such as influenza and SARS-CoV-2.

Keywords: Humidity; influenza; SARS-CoV-2; spatio-temporal modelling

1. Introduction

Previous experimental evidence indicates that the aerosolization of secretions lubricating the vocal cords can be a major source of microscopic droplet (microdroplet) virus infections [1]. Evidence of airborne spread of the severe acute respiratory syndrome virus (SARS) is provided by [2]. Recent studies also suggest the possibility of airborne transmission of SARS-CoV-2 through respiratory air droplets and aerosols [3].

Weather factors—such as humidity and temperature—are suspected of playing a role in the transmission of viral particles through aerosolized droplet nuclei or aerosols. Particularly, differences in humidity are suspected to provide an explanation for the observed variability of influenza and its transmission [4]. Weather conditions can have a similar effect on SARS-CoV-2 (COVID-19), as airborne inactivation of the human coronavirus 229E is affected by temperature and relative humidity [5], and in experimental settings both temperature and absolute humidity affect the

environmental survival of surrogates of mammalian coronaviruses [6].

Based on observational data, the relationship between SARS-CoV-2 and weather has been recently analyzed in published studies [7-12] and pre-prints [13-18]. All of these studies find a negative correlation between absolute humidity and the spread of SARS-CoV-2, indicating that low levels of humidity increase the risk of COVID-19 cases. In [18], for example, weather conditions are investigated worldwide, and [18] conclude that the community transmission of SARS-CoV-2 is consistent with average temperatures of 5° to 11°C, combined with low specific humidity between 3 to 6 g/kg. Humidity seems to be the most important factor in viral spreading, while temperature play a less important role [13].

Our study provides new evidence about the relationship between specific humidity and the risk of influenza and COVID-19 cases. Five years of hourly weather data from 34 regional weather stations in the Netherlands is analyzed against five years of weekly data of influenza

cases, as well as daily regional data of COVID-19 cases, hospitalizations, and deaths. The Netherlands has accurate and easily accessible data that allows for a high-resolution and precise estimation of the relationship between weather conditions and the spread of viral diseases. As time trends of viral transmission can vary substantially in time and at sub-national level, the combination of high spatial and temporal resolution captures the heterogeneity of viral spreading across time and space.

In our study, parametric and non-parametric correlations are calculated for specific humidity and influenza/SARS-CoV-2 at country-level. An inverse (negative) relationship is observed between specific humidity and the incidence of influenza between 2015 and 2019, but a positive correlation is found between specific humidity and the reported cases of COVID-19. Since the positive correlation between specific humidity and COVID-19 may be caused by the use of aggregated data at national level—besides the lack of herd immunity—a Bayesian spatio-temporal disease model for the Netherlands is estimated at municipality and province level. This model allows to quantify the relation of COVID-19 cases with the specific humidity in the Netherlands at regional sub-national levels and over time.

The results of the spatio-temporal disease model indicate that the increase of specific humidity during the outbreak of the SARS-CoV-2 helped to reduce the risk of regional COVID-19 cases in the Netherlands. Specifically, an increase of specific humidity of one gram of water vapor per kilogram of air (1 g/kg) is related to a reduction of approximately 5% in the risk of COVID-19 cases.

2. Data

Weekly data of influenza reports were collected for the years 2015 to 2019 from the Nivel Primary Care Database (Nivel Zorgregistraties Eerste Lijn). Daily COVID-19 data (Reported Infections, Hospitalizations and Mortality) was provided by The Dutch National Institute for Public Health and the Environment (RIVM). RIVM is the government agency focused as Centre for Infectious Disease Control (CDC), under supervision of the Dutch Government. The COVID-19 data covers the twelve provinces and the 355 municipalities in the Netherlands, from March 13th to July 9th.

Specific humidity (q) was calculated with the data from 34 weather stations, spread across The Netherlands. This data was provided by The Royal Netherlands

Meteorological Institute (KNMI), the Dutch national weather service. Since KNMI only reports relative humidity (h), specific humidity q was calculated using the Rotronic formula¹, which is based on the UK National Physical Laboratory guide for the measurement of humidity [19]:

$$q = \xi \frac{h d_w}{(d_m - d_w)} \quad (1)$$

In (1), q is specific humidity of air vapor mixture (expressed in g/Kg), h is relative humidity (in percentage), d_w is the density of water vapor (kg/m³), d_m is density of the moist or humid air (kg/m³) and $\xi = .622$. Specific humidity is analyzed instead of relative humidity because relative humidity is affected by the surrounding temperature. Specific humidity in contrast remains the same in indoor and outdoor conditions². The hourly weather data provided by the KNMI was aggregated at daily and weekly levels using the average values during the time periods.

Additional population data was used in the spatial-time models to calculate the incidence of SARS-CoV-2 at intra-regional level. This data was provided by the national statistical office, Statistics Netherlands (CBS) and is also used by the RIVM.

3. Methods

Parametric and non-parametric correlation coefficients were calculated between specific humidity q and the incidence of influenza and SARS-CoV-2. Spatio-temporal disease models were estimated with the daily data of specific humidity (q_{it}) and SARS-CoV-2 (y_{it}) in the Netherlands. Spatial dependence in the latent component of the disease mapping is modelled by specifying neighborhood relationships among the area-level risks [20-21].

In the space-time analysis, the Besag-York-Mollié (BYM) spatio-temporal model [22] is used to estimate the impact of specific humidity on the risk of SARS-CoV-2 (COVID-19) cases at province and municipality level. In the BYM model, the number of SARS-CoV-2 cases follows a Poisson stochastic process for each

¹ The Rotronic technical note about humidity definitions is freely available at: https://www.rotronic.com/media/productattachments/files/h/u/humidity_definition_weba.pdf?_ga=2.192612136.342104324.1597340506-398495933.1597340506

² For example, weather stations may report a temperature of 5°C and a relative humidity of 35% for the outside environment (equal to a specific humidity of 1.87g/kg). However, inside the households, the temperature can be 21°C with a relative humidity of 12% due to heating systems, but specific humidity will remain the same (1.87g/kg) regardless of the temperature.

area i and time points t , $y_{it} \sim \mathcal{P}(\lambda_{it})$, as in [23]. The parameter λ_{it} is defined by the ecological regression:

$$\begin{cases} \lambda_{it} = E_{it} p_{it} \\ p_{it} = \exp(\alpha + v_i + u_i + (\beta_t + \tau_i)t + \beta_q q_{it}) \end{cases} \quad (2)$$

where E_{it} is the expected number of cases in each area, at each point in time, p_{it} is the ratio between the number of observed cases y_{it} and the number of expected cases, α is the average rate of cases in all the areas, v_i is the unstructured area-specific effect, which follows a Gaussian prior $v_i \sim \mathcal{N}(0, \sigma_v^2)$, and u_i is the spatially structured area-specific effect, modelled with an intrinsic conditional autoregressive prior that takes into account the number of j -areas which share boundaries with the i -areas ($j \neq i$), i.e. the neighbors of each province/municipality [24].

In (2), the temporal component t is modelled with a parametric approach [25], where β_t represents the global time effect, and $\tau_i \sim \mathcal{N}(0, 1/\sigma_\tau^2)$ is the differential trend that captures the interaction between time and space. Specific humidity levels (q_{it}) were included as a risk factor in the spatio-temporal model (1) to evaluate their impact on the risk of SARS-CoV-2 cases, measured by the parameter β_q . When the incidence of SARS-CoV-2 per 100,000 population is used as the dependent variable, a Gaussian likelihood is used instead of the Poisson model in equation (2)³.

4. Results

Figure 1 shows the weekly historical patterns of specific humidity q and the incidence of influenza in the Netherlands, from 2015 to 2019. Specific humidity q during the winter periods is between 3 and 6 g/kg, while in the summer specific humidity is above 8 g/kg. The highest incidence of influenza infection is observed in weeks with a specific humidity of less than 6 g/kg. Particularly, a higher incidence is observed in children of less than 4 years and individuals with 65 years or more (Figure 2a). Above 6 g/kg, a significant drop in infections is observed. The opposite is also true: the rate of influenza cases is 5.91 higher when specific humidity levels are approximately 2 g/kg compared to the average cases at 8 g/kg (Table 1).

Figure 2b illustrates the inverse non-linear relationship between the levels of specific humidity and the weekly incidence of influenza. The ordinary Pearson

correlation of specific humidity with influenza cases is negative and equal to -0.6986 (t-value: -15.713 , p-value: 0.0000). The value of the non-parametric Spearman correlation is equal to -0.8083 (t-statistic: -22.0993 , p-value: 0.0000).

In the case of SARS-CoV-2, there is a positive correlation between humidity levels and the daily COVID-19 cases in the Netherlands (Figure 3). The Pearson correlation is equal to $.7016$ (t-value: 10.651 , p-value: 0.0000), and the Spearman correlation is 0.8389 (t-statistic: 16.6734 , p-value: 0.0000). This positive correlation can be explained by the lack of herd immunity—due to the novelty of the SARS-CoV-2—, the increasing number of tests, and the use of aggregate data at country level which does not capture the dissimilarities of specific humidity at sub-national levels in the Netherlands.

In order to take into account the differential spatial and temporal patterns of specific humidity across the Netherlands during the outbreak of the COVID-19, a Bayesian spatial-time model based on a Poisson stochastics process was estimated for the number of reported cases of SARS-CoV-2, at municipality and province level. The increase in testing, which increases reported COVID-19 cases, was controlled by relativizing the number of COVID-19 cases with the population at municipality and province level, i.e. estimating an additional Gaussian space-time model for the incidence of COVID-19 cases per 100,000 people⁴.

Table 2 shows the results of estimating the fixed-effects $\{\alpha, \beta_t, \beta_q\}$ of the spatio-temporal model. The low standard deviation of the posterior means in all the models indicates a good level of accuracy in the integrated nested Laplace approximation of the approximate Bayesian inference. The value of the exponentiated estimate of β_t implies a 1.5% daily rate of infection of SARS-CoV-2 in the Netherlands from March to July of 2020, with a 95% credibility interval ranging from 1.47% to 1.62%. The negative sign of the mean estimate of β_q —as well as the sign of the low and upper limit of the 95% credible interval—suggests that

³ The data and R codes to replicate the results with R-INLA (www.r-inla.org) are freely available upon request.

⁴ The Netherlands increased COVID-19 testing from 55,000 weekly tests in March to 110,000 tests per week in July 2020. The testing was only focused on care workers in March, but now the entire population is subject to testing. Due to the possible bias caused by testing, space-time models were also estimated using data of the number of hospitalizations, and similar results were obtained as those for reported cases. The results for hospitalization are available upon request.

regions with higher specific humidity levels have, on average, lower number of COVID-19 cases in the Netherlands. Specifically, the estimated values of $\hat{\beta}_q = -0.048$ at province level and $\hat{\beta}_q = -0.053$ at municipality level in the Poisson space-time model indicate that an increase of one gram of water vapor per kilogram of air (1 g/kg) is related to a reduction of around 5% in the risk of Covid-19 cases.

Table 1. Average incidence of influenza cases at different intervals of specific humidity

Specific humidity	Average incidence of influenza per 100000 people
Below 2 g/kg	147.3
Between 2 and 3	135.4
Between 3 and 4	110.9
Between 4 and 5	79.7
Between 5 and 6	54.4
Between 6 and 7	33.2
Between 7 and 8	29.5
Between 8 and 9	24.9
Between 9 and 10	17.2
Above 10	15.8

Figure 1. Specific humidity and incidence of influenza in the Netherlands
Weekly historical observations per year




Figure 2. Specific humidity and incidence of influenza for age categories in the Netherlands.
a: Weekly historical observations (left), b: bivariate scatterplot (right)





Figure 3. Specific humidity and SARS-CoV-2 (Covid-19) cases in the Netherlands

a: Daily historical observations (left), b: bivariate scatterplot (right)



Similar results are obtained using a Gaussian space-time model for the incidence of SARS-CoV-2 cases per 100,000 population in the Netherlands. On average, regions with higher levels of specific humidity showed a lower incidence of COVID-19 cases per 100,000 people, on a magnitude of at province level and at municipality level.

Figure 4 shows the area-specific relative risks of COVID-19 in the Netherlands—at province and municipality level—calculated with the random-effects estimated with the spatio-temporal models. The regions located in the North of the country,


where specific humidity tends to be higher, show a lower risk of COVID-19 cases. In contrast the regions with less specific humidity, in the South East of the country, display higher levels of relative risk of SARS-CoV-2 cases.

Interestingly, municipalities with the highest concentration of conservative orthodox Calvinist Protestants in the country, i.e. those in the 'Bible belt', have a higher risk of SARS-CoV-2, compared to other regions in the Netherlands. The identification of this type of regions with a high risk of COVID-19 cases can help to allocate resources during the SARS-CoV-2 pandemic [26].

Table 2. Estimation results: Besag-York-Mollie spatio-temporal disease model of daily () COVID-19 cases and specific humidity () in the Netherlands

Parameter	Area's break	Mean estimate	Standard deviation	Credible interval	
				2.5%	97.5%
Number of cases of SARS-CoV-2 (COVID-19)	Province	6.770375	0.092920	6.585284	6.955255
	Municipality	3.407532	0.043961	3.321127	3.493834
	Province	0.015331	0.000354	0.014625	0.016037
	Municipality	0.015307	0.000035	0.015238	0.015375
	Province	-0.047696	0.000361	-0.048404	-0.046988
	Municipality	-0.053141	0.000364	-0.053855	-0.052427
Incidence of cases of SARS-CoV-2 (COVID-19) per 100,000h	Province	67.440261	2.982071	61.584535	73.291041
	Municipality	87.629212	3.374872	81.003201	94.249693
	Province	2.448832	0.049035	2.352544	2.545038
	Municipality	2.866350	0.013162	2.840509	2.892170
	Province	-6.231110	0.706851	-7.619118	-4.844281
	Municipality	-8.399374	0.194467	-8.781178	-8.017889

Figure 4. Area-specific relative risk* of SARS-CoV-2 in the Netherlands



(*) Relative risk: residual risk after regional specific humidity () levels are taken into account.
Area's break: province level (left) and municipality level (right).

5. Discussion

Higher levels of specific humidity () were found to be related to a lower number of reported cases of respiratory viruses in the Netherlands. An inverse non-linear relationship between specific humidity and influenza is observed historically between 2015 and 2019. A similar inverse (negative) relationship between specific humidity and SARS-CoV-2 cases is found using spatio-temporal models that consider the differential patterns of specific humidity () and SARS-CoV-2 cases at municipality and province levels in the Netherlands. Given the seasonal dynamics of humidity in the last 5 years, an uptake (second wave) of SARS-CoV-2 infections can be expected during the weeks 43 to 45 of 2020, if low levels of humidity—below 6 g/Kg—are observed during those periods.

The results are in line with previous studies [7-18]⁵, as our risk analysis shows that higher levels of humidity may result in a slower spread of COVID-19 [10, 12]. The findings are consistent with the hypothesis that suggests that SARS-CoV-2 spreads through airborne aerosolization [28], since lower levels of specific humidity lead to a higher rate of infections due to the

effects of water activity on virus survivability, surface inactivation, salt toxicity, or changes in pH induced by evaporation that modify the surface glycoproteins of enveloped viruses and subsequently compromise their infectivity [29]. Lower levels of humidity can particularly lead to higher aerosol concentration when a dry air increases the evaporation of respiratory droplets, i.e. when droplets smaller in diameter than a few micrometers—referred to as ‘droplet nuclei’—evaporate to about half their initial size [30]. In the case of influenza, for example, the probabilities of exposure and infection risk of aerosol and droplet transmission are within the same order of magnitude, but intranasal inoculation leads to about 20 times lower infectivity that when the virus is delivered in an inhalable aerosol [31].

In practice, the findings suggest that public policies that promote higher levels of specific humidification—above 6 g/Kg—can lead to significant reductions in the spread of respiratory viruses, such as influenza and SARS-CoV-2. For example, the deployment of hygrometers aimed at measuring specific humidity can guide the public behavior in the face of a warning of an increased risk of infection⁶. The results also imply that the use of air

⁵ Compared to these previous studies, we use more detailed data, with humidity levels, per municipality, and allow social gatherings in outside spaces (a temporal and spatial resolution, in order to reduce the possibility of the ecological fallacy that may arise in the analysis of large spatial units that neglect localized patterns of SARS-CoV-2 [27].

⁶ For example, smart social distancing regulations may take into account local humidity levels, per municipality, and allow social gatherings in outside spaces (a temporal and spatial resolution, in order to reduce the possibility of the ecological fallacy that may arise in the analysis of large spatial units that neglect localized patterns of SARS-CoV-2 [27].

conditioning (which decreases the specific humidity in a cooled area), must be complemented with humidification during the fall and winter periods, particularly when specific humidity q decreases below 6 g/kg.

Future studies can study the interaction effects of sunlight with humidity, as sunlight radiation may act as an additional environmental factor in the reduction and prevention of the risk of SARS-CoV-2 [32], since solar radiation plays an important role in vitamin D production, which seems to play a role in reducing COVID-19 infections [33-34].

Acknowledgements

The authors are grateful to Drs. Maurice de Hond, social geographer with a specialization in statistics, and Pierre Capel, professor emeritus in experimental immunology at the Utrecht University in the Netherlands, for providing us with valuable research papers and feedback about our study. Rami Alkhatib, assistant professor of mechatronics at Rafik Hariri University, did a parallel analysis on the same dataset and got similar results, supporting our findings. The data used in the study was obtained with the help of staff members at RIVM, KNMI, CBS, Rotronic, and Mariëtte Hooiveld, J Hendriksen and Korevaar JC from Nivel data (www.nivel.nl/surveillance).

References

1. Morawska, L. J. G. R., Johnson, G. R., Ristovski, Z. D., Hargreaves, M., Mengersen, K., Corbett, S., ... & Katoshevski, D. (2009). Size distribution and sites of origin of droplets expelled from the human respiratory tract during expiratory activities. *Journal of Aerosol Science*, 40(3), 256-269.
2. Yu, I. T., Li, Y., Wong, T. W., Tam, W., Chan, A. T., Lee, J. H., ... & Ho, T. (2004). Evidence of airborne transmission of the severe acute respiratory syndrome virus. *New England Journal of Medicine*, 350(17), 1731-1739.
3. Morawska, L., & Cao, J. (2020). Airborne transmission of SARS-CoV-2: The world should face the reality. *Environment International*, 105730.
4. Shaman, J., & Kohn, M. (2009). Absolute humidity modulates influenza survival, transmission, and seasonality. *Proceedings of the National Academy of Sciences*, 106(9), 3243-3248.
5. Ijaz, M. K., Brunner, A. H., Sattar, S. A., Nair, R. C., & Johnson-Lussenburg, C. M. (1985). Survival characteristics of airborne human coronavirus 229E. *Journal of General Virology*, 66(12), 2743-2748.
6. Casanova, L. M., Jeon, S., Rutala, W. A., Weber, D. J., & Sobsey, M. D. (2010). Effects of air temperature and relative humidity on coronavirus survival on surfaces. *Applied and environmental microbiology*, 76(9), 2712-2717.
7. Gupta, S., Raghuwanshi, G. S., & Chanda, A. (2020). Effect of weather on COVID-19 spread in the US: a prediction model for India in 2020. *Science of The Total Environment*, 138860.
8. Ma, Y., Zhao, Y., Liu, J., He, X., Wang, B., Fu, S., & Luo, B. (2020). Effects of temperature variation and humidity on the death of COVID-19 in Wuhan, China. *Science of The Total Environment*, 138226.
9. Carta, M. G., Scano, A., Lindert, J., Bonanno, S., Rinaldi, L., Fais, S., & Orrù, G. (2020). Association between the spread of COVID-19 and weather-climatic parameters. *European review for medical and pharmacological sciences*, 24(15).
10. Bukhari, Q., Massaro, J. M., D'Agostino, R. B., & Khan, S. (2020). Effects of Weather on Coronavirus Pandemic. *International Journal of Environmental Research and Public Health*, 17(15), 5399.
11. Guo, X., Zhang, H. & Zeng, Y. Transmissibility of COVID-19 in 11 major cities in China and its association with temperature and humidity in Beijing, Shanghai, Guangzhou, and Chengdu. *Infect Dis Poverty* 9, 87 (2020).
12. Qi, H., Xiao, S., Shi, R., Ward, M. P., Chen, Y., Tu, W., ... & Zhang, Z. (2020). COVID-19 transmission in Mainland China is associated with temperature and humidity: A time-series analysis. *Science of the Total Environment*, 138778.
13. Crema, E. (2020). The SARS-COV-2 outbreak around the Amazon rainforest: the relevance of the airborne transmission. *medRxiv*.
14. Bukhari, Q., & Jameel, Y. (2020). Will coronavirus pandemic diminish by summer? Available at SSRN 3556998.
15. Liu, Y., Ning, Z., Chen, Y., Guo, M., Liu, Y., Gali, N. K., ... & Liu, X. (2020). Aerodynamic characteristics and RNA concentration of SARS-CoV-2 aerosol in Wuhan hospitals during COVID-19 outbreak. *BioRxiv*.
16. Oliveiros, B., Caramelo, L., Ferreira, N. C., & Caramelo, F. (2020). Role of temperature and humidity in the modulation of the doubling time of COVID-19 cases. *medRxiv*.

implemented. If increased humidification is not possible, air purification with HEPA filters and ionization can be alternative solutions to deactivate viral aerosols.

17. Raines, K. S., Doniach, S., & Bhanot, G. (2020). The transmission of SARS-CoV-2 is likely comodulated by temperature and by relative humidity. medRxiv.
18. Sajadi, M. M., Habibzadeh, P., Vintzileos, A., Shokouhi, S., Miralles-Wilhelm, F., & Amoroso, A. (2020). Temperature and latitude analysis to predict potential spread and seasonality for COVID-19. Available at SSRN 3550308.
19. National Physical Laboratory (1996). A Guide to the Measurement of Humidity. The Institute of Measurement and Control, United Kingdom. https://rotronic.com/media/productattachments/files/n/p/npl_guide_to_humidity.pdf
20. Blangiardo, M., Bouliari, A., Diggle, P., Piel, F. B., Shaddick, G., & Elliott, P. (2020). Advances in spatiotemporal models for non-communicable disease surveillance. *International Journal of Epidemiology*, 49(Supplement_1), i26-i37.
21. Kim, S., & Castro, M. C. (2020). Spatiotemporal pattern of COVID-19 and government response in South Korea (as of May 31, 2020). *International Journal of Infectious Diseases*.
22. Besag, J., York, J., & Mollié, A. (1991). Bayesian image restoration, with two applications in spatial statistics. *Annals of the institute of statistical mathematics*, 43(1), 1-20.
23. Blangiardo, M., Cameletti, M., Pirani, M., Corsetti, G., Battaglini, M., Baio, G. (2020). Estimating weekly excess mortality at sub-national level in Italy during the COVID-19 pandemic.
24. Blangiardo, M., & Cameletti, M. (2015). *Spatial and spatio-temporal Bayesian models with R-INLA*. John Wiley & Sons.
25. Bernardinelli, L., Clayton, D., Pascutto, C., Montomoli, C., Ghislandi, M., & Songini, M. (1995). Bayesian analysis of space-time variation in disease risk. *Statistics in medicine*, 14(21-22), 2433-2443.
26. Cordes, J., & Castro, M. C. (2020). Spatial analysis of COVID-19 clusters and contextual factors in New York City. *Spatial and Spatio-temporal Epidemiology*, 100355.
27. Konstantinoudis, G., Padellini, T., Bennett, J. E., Davies, B., Ezzati, M., & Blangiardo, M. (2020). Long-term exposure to air-pollution and COVID-19 mortality in England: a hierarchical spatial analysis. medRxiv.
28. Fineberg, H. V., & National Research Council. (2020). Rapid expert consultation on the possibility of bioaerosol spread of SARS-CoV-2 for the COVID-19 pandemic (April 1, 2020). In *The National Academies Press NRC. The National Academies Press, National Research Council, Washington, DC.*
29. Yang, W., & Marr, L. C. (2012). Mechanisms by which ambient humidity may affect viruses in aerosols. *Applied and environmental microbiology*, 78(19), 6781-6788.
30. Brien, N. C., Timen, A., Wallinga, J., Van Steenbergen, J. E., & Teunis, P. F. (2010). The effect of mask use on the spread of influenza during a pandemic. *Risk Analysis: An International Journal*, 30(8), 1210-1218.
31. Teunis, P. F., Brien, N., & Kretzschmar, M. E. (2010). High infectivity and pathogenicity of influenza A virus via aerosol and droplet transmission. *Epidemics*, 2(4), 215-222.
32. Rosario, D. K., Mutz, Y. S., Bernardes, P. C., & Conte-Junior, C. A. (2020). Relationship between COVID-19 and weather: Case study in a tropical country. *International Journal of Hygiene and Environmental Health*, 113587.
33. Grant, W. B., Lahore, H., McDonnell, S. L., Baggerly, C. A., French, C. B., Aliano, J. L., & Bhattoa, H. P. (2020). Evidence that vitamin D supplementation could reduce risk of influenza and COVID-19 infections and deaths. *Nutrients*, 12(4), 988.
34. Whittemore, P. B. (2020). COVID-19 fatalities, latitude, sunlight, and vitamin D. *American Journal of Infection Control*