Immunogenicity and safety of fractional doses of yellow fever vaccines: a randomized, double-blind, non-inferiority trial

Aitana Juan-Giner[1,*], Derick Kimathi[2,3*], Kyra H. Grantz[4,5,6], Mainga M. Hamaluba[2], Patrick Kazooba[7], Patricia Njuguna[2], Gamou Fall[8], Moussa Dia[8], Ndeye S. Bob[8], Thomas P. Monath[9], Alan D. Barrett[10], Joachim Hombach[11], Edgar M. Mulogo[12], Immaculate Ampeire[13], Henry K. Karanja[2], Dan Nyehangane[7], Juliet Mwanga-Amumpaire[7], Derek A.T. Cummings[4,5], Philip Bejon[2,3], George M. Warimwe[2,3,#], Rebecca F. Grais[1,#].

[1] Epicentre, Paris, France
[2] Kenya Medical Research Institute – Wellcome Trust Research Programme, Kilifi, Kenya
[3] Centre for Tropical Medicine & Global Health, University of Oxford, UK
[4] Department of Biology, University of Florida
[5] Emerging Pathogens Institute, University of Florida
[6] Department of Epidemiology, Johns Hopkins Bloomberg School of Public Health
[7] Epicentre Mbarara Research Centre, Uganda
[8] Institut Pasteur Dakar, Senegal
[9] Crozet BioPharma LLC, Devens MA, USA
[10] Sealy Institute for Vaccines Sciences and Department of Pathology, University of Texas Medical Branch, USA
[11] Immunization, Vaccines & Biologicals, World Health Organization, Geneva, Switzerland
[12] Department of Community Health, Mbarara University of Science & Technology, Uganda
[13] Expanded Program on Immunization, Ministry of Health, Uganda

*Co-first author; #Co-senior author

Corresponding author: Rebecca F Grais; Address: 34 Avenue Jean Jaurès, 75019 Paris, France; Tel: +33 (0) 1 40 21 54 75; Email: rebecca.grais@epicentre.msf.org

Word Count: 3,571

Abstract

Background

Yellow fever vaccine stocks have been insufficient to cover exceptional demands for outbreak response. Fractional dosing evidence is limited to the 17DD substrain vaccine. We assessed the immunogenicity and safety of one-fifth fractional dose compared to standard dose of each of the four WHO-prequalified yellow fever vaccines produced from three substrains.

Methods

We conducted a randomized, double-blind, non-inferiority trial in Mbarara, Uganda and Kilifi, Kenya. 960 participants aged 18-59 years without previous yellow fever vaccination or infection were recruited from communities and randomized to receive one of four vaccines and to standard or fractional dosage. Vaccine was administered subcutaneously by unblinded nurse. Other study personnel and participants were blinded to vaccine allocation. Primary immunogenicity outcome, seroconversion, was measured in the per-protocol population; safety outcomes included all vaccinated participants. We defined non-inferiority as no more than 10% decrease in seroconversion in fractional compared to standard dose arms 28 days post-vaccination. Seroconversion was defined as ≥4-fold rise in neutralizing antibody titers measured by 50% plaque reduction neutralization test ($PRNT_{50}$).

ClinicalTrials.gov Identifier: NCT02991495, following participants.

Findings

Between 6[th] November 2017 and 21[st] February 2018, 960 participants, total sample goal, were randomized. The primary per-protocol analysis includes 899 participants, with 110 to 117 participants per arm. The absolute difference in seroconversion between fractional and standard doses by vaccine was 1.71% (95%CI: -2.60, 5.28), -0.90% (95%CI: -4.24, 3.13), 1.82% (95%CI: -2.75, 5.39), 0.0% (95%CI: -3.32, 3.29). Fractional doses from all four vaccines met the non-inferiority criterion. There were no safety concerns.

Interpretation

These results support fractional dosing of all WHO-prequalified yellow fever vaccines for the general adult population for outbreak response in situations of vaccine shortage.

Funding

The study was funded by Médecins Sans Frontières Foundation, Wellcome Trust (grant no. 092654) and the UK Department for International Development. Vaccines were donated in kind.

Research in context

Evidence before this study

In July 2016, following major yellow fever (YF) outbreaks in two countries, WHO published a secretariat information paper including a review of studies assessing the immunogenicity of fractional doses of YF vaccines and recommended consideration of fractional doses to manage a vaccine shortage. Following this, fractional doses of YF vaccine produced by Bio-Manguinhos/Fiocruz (17DD substrain) were given to approximately 7.5 million non-pregnant adults and children ≥2 years of age in Kinshasa, Democratic Republic of Congo. The evidence base to support this action was limited to a single vaccine substrain and to a specific context. To broaden and simplify recommendations, WHO called for additional research to be conducted. We designed a trial to assess non-inferiority in seroconversion of fractional (one-fifth dose) versus standard dose for each of the four WHO-prequalified YF vaccines at 28 days post-vaccination in an adult population in Kenya and Uganda. We selected vaccine batches as close as possible to each manufacturer's minimum release specification.

Added value of this study

This is the first randomized controlled trial assessing all four WHO-prequalified YF vaccines, providing information on the immunogenicity and safety of fractional doses of the different vaccine substrains at 10 days, 28 days and one year post-vaccination. The results show that, at 28 days post-vaccination, most participants had high levels of neutralizing antibodies and that seroconversion rates in the fractional dose arms were non-inferior to standard dose for each of the four vaccines. Seroconversion rates and neutralizing antibodies remained high up to one year post-vaccination for both fractional and standard doses for all vaccines. These results are aligned with previous studies using the 17DD substrain vaccine but extend the evidence to randomized comparisons of all four vaccines and to a sub-Saharan Africa context. To our knowledge, this is the first trial assessing immunogenicity of fractional doses at 10 days post-vaccination.

Implications of all the available evidence

Our study supports the use of one-fifth fractional doses of all four WHO-prequalified yellow fever vaccines for the general adult population and fills a critical knowledge gap to support WHO policy on the use of fractional dosing of yellow fever vaccine for outbreak response. The immunogenicity and safety of fractional dosing in children and specific populations, such as those living with HIV, is yet to be determined. Long-term studies are warranted to confirm the duration of protection.

Introduction

Yellow fever is a mosquito-borne viral disease endemic in 44 countries with periodic outbreaks (1). Four live attenuated yellow fever virus vaccines (derived from 17D-204, 17D-213 and 17DD substrains) are World Health Organization (WHO)-prequalified (Bio-Manguinhos/Fiocruz (Brazil), Federal State Unitary Enterprise of Chumakov Institut of Poliomyelitis and Viral Encephalitides (Russia), Institut Pasteur de Dakar (Senegal), and Sanofi Pasteur (France)). All have been widely used and are considered safe and effective (1). WHO recommends routine vaccination in all countries where the disease is endemic, vaccination of travelers to those areas, and mass vaccination for prevention or control of outbreaks. A stockpile of 2 million doses was reserved for outbreak response in 2000 and increased to 6 million in 2014 to facilitate response to major multi-country, but unpredictable outbreaks (2).

An outbreak of yellow fever in 2016 in Angola raised major concerns about the adequacy of vaccine supply. Routine vaccination was suspended in some African countries to meet demand during the Angola outbreak (3) and a subsequent outbreak in Democratic Republic of Congo (DRC) further contributed to a global shortage (4). In response, WHO reviewed the evidence on fractional dosing of yellow fever vaccine as a dose-sparing option and exceptionally recommended considering off-label use of fractional doses, administered by the standard subcutaneous or intramuscular route, to extend supply in response to ongoing outbreaks (4,5). Fractional doses (defined as one-fifth of the standard dose) of the yellow fever vaccine produced by Bio-Manguinhos/Fiocruz (17DD substrain) were given to approximately 7.5 million non-pregnant adults and children ≥2 years of age in Kinshasa, DRC in August, 2016 (6). Fractional dosing was again used in response to outbreaks in 2017-2018 in the states of São Paulo, Rio de Janeiro and Bahia, Brazil to vaccinate almost 17 million people (7). Four studies have assessed the immunogenicity of fractional doses of yellow fever vaccine. One of these studies (8) used an old vaccine formulation no longer produced and another focused on the intradermal administration (9). The evidence base to support WHO recommendations was primarily from a dose-finding study of the 17DD substrain vaccine amongst healthy male army recruits in Brazil that showed seroconversion rates greater than 97% with doses as low as 587 IU/dose (10) and similar virological and immunological kinetics with doses down to 3,013 IU/dose compared to the released dose of 27,476 IU/dose (11). Based on historic data, standard doses of yellow fever vaccines should contain not less

than 1,000 IU/dose (12). However, release specifications vary by manufacturer, related to the

stability of the vaccine, and are generally several-fold higher than the minimum specification

(13). WHO recommended that the minimal dose administered, standard or fractional, should

preferentially contain 3,000 IU/dose, hence the decision on the dose fractioning (e.g. one-half

or one-fifth of the standard dose) should consider the potency of the vaccine batch (4). The

dose-finding study conducted in Brazil (10,11) provides information for only one of the

WHO-prequalified vaccines produced, that of the 17DD substrain. During the fractional dose

campaign in Kinshasa, DRC, a study confirmed the immunogenicity of fractional doses of the

17DD substrain vaccine in a large-scale campaign (14). As vaccines are produced from three

substrains and differ in viral concentration per dose, the results of these studies are not

generalizable, and it remains important to obtain data for all WHO-prequalified vaccines.

We assessed the immunogenicity and safety of fractional doses of all four WHO-prequalified

yellow fever vaccines. This is the first clinical trial where all four WHO-prequalified vaccines

have been used in the same study, providing information on the different substrains. This

study fills a critical gap to provide evidence to simplify and broaden recommendations for

fractional dosing of yellow fever vaccines globally (4).

Methods

Study design

We conducted a two-center, double-blind, individually-randomized trial in Mbarara, Uganda

and Kilifi, Kenya (15). The study took place at the Epicentre Mbarara Research Centre in

Mbarara, Uganda and the Kenya Medical Research Institute (KEMRI)-Wellcome Trust

Research Programme clinical trials facility in Kilifi, Kenya. Mbarara district is located in

proximity to Masaka, Rukungiri, and Kalangala districts which registered confirmed yellow

fever cases in 2016 (16).

The study protocol was approved by the ethics committee of the World Health Organization

(Switzerland); Scientific & Ethics Review Unit, Kenya Medical Research Institute (Kenya);

Oxford Tropical Research Ethics Committee (United Kingdom); Mbarara University of

Science and Technology Research Ethics Committee (Uganda); and the Uganda National

Council of Sciences and Technology (Uganda). Approval was obtained from the national

regulatory authorities in Uganda and Kenya. The trial was conducted in accordance with

Good Clinical Practice guidelines.

Participants

Participants were recruited from rural communities in the Mbarara municipality, Mbarara

district, Uganda and Kilifi County, Kenya. Communities were informed about the trial using

locally adapted strategies. Individuals interested in participating were invited to the study

sites. Written informed consent was required to participate.

Individuals were eligible to participate if they were 18 to 59 years of age; had no

contraindications for vaccination; were not pregnant or lactating; had no history of previous

yellow fever vaccination or infection; did not require yellow fever vaccination for travel; and

were able to comply with study procedures.

Randomization and masking

Participants were individually randomized to one of 8 arms corresponding to the four

prequalified yellow fever vaccines at standard or fractional dose. Unique allocation numbers

were prepared by an independent statistician (DiagnoSearch LifeSciences, Mumbai, India)

using a computer-generated random number list with non-disclosed fixed blocks of size 10

with equal allocation to dosage within a block and to a given manufacturer by site. The

allocation sequence was concealed using pre-prepared, sequentially numbered scratch-off

booklets stored under key. After enrollment, the vaccination nurse scratched off the

randomization code indicating the allocated vaccine and dosage for the participant. Vaccines

were reconstituted and administered in a private room not accessible to other study staff.

Participants were blinded by masking the volume of the syringe with opaque tape. The

vaccination nurse and supervisor overseeing vaccination were aware of allocation; personnel

assessing outcomes and investigators were blinded to vaccine and dosage throughout.

Procedures

A batch of yellow fever vaccine of standard 10-dose vials was selected from each

manufacturer with potency at time of release closest to internal minimum specification.

Vaccine potencies were independently measured at the National Institute for Biological

Standards and Control (United Kingdom) (Table 1). The freeze-dried preparations and

diluents were kept at 2-8°C until administration. Each day, the needed vaccine vial(s)

was(were) reconstituted using each manufacturer's diluent to prepare fractional and standard

doses prior to vaccination of the first randomized participant. Any remaining reconstituted vaccine was discarded after 6 hours. Fractional doses consisted of one-fifth of the standard dose, as defined for each manufacturer. Vaccine was administered subcutaneously in the deltoid region using 0.5ml auto-disable syringes (needle size 25G x 3/4") with a 45° injection angle for the standard 0.5ml dose and 0.1ml auto-disable syringes (needle size 26G x 3/8") at 90° injection angle for the 0.1ml fractional dose.

Participants were followed at 10 ($\pm$1) days, 28 ($\pm$3) days, and 365 ($\pm$14) days post-vaccination. At each study visit, participants had a medical consultation and a blood sample was obtained at the initial visit prior to vaccination and subsequently at each scheduled follow-up visit.

Outcomes

Serum samples were separated and aliquoted within 4 hours of collection and stored in -80°C freezers at each site. Serum samples were analyzed at the Institut Pasteur de Dakar (Senegal), where neutralizing antibody titers against yellow fever were assessed by 50% plaque reduction neutralization test ($PRNT_{50}$) as primary outcome and $PRNT_{90}$ as supplementary (17,18). Laboratory personnel were blinded to study dose allocation.

The attenuated yellow fever vaccine strain 17D-204 sourced by Institut Pasteur de Dakar was used as challenge strain for the neutralization assays. Viral stock preparation was in C6/36 cells and titrated by plaque assay method previously described (19). Briefly, a defined virus concentration of 10^3 plaque forming units per ml and diluted and heat-inactivated sera were titrated in serial two-fold dilutions from 1:10 to 1:20480. Serum and virus were mixed and incubated before overlaying with 0.6% carboxy-methyl-cellulose and 3% fetal calf serum in L-15 Leibovitz's media. Following 4 to 5 days of incubation at 37°C, plaques were counted, and antibody titer was determined as the dilution that reduced observed plaques by 50% and 90% compared to the number observed in control wells.

Serious adverse events (SAEs), defined as any new health-related problem that resulted in death, was life-threatening, necessitated hospitalization or prolongation of existing hospitalization, or resulted in disability or incapacity, were assessed for the duration of follow-up. Adverse events (AEs) included all untoward medical events and were evaluated in the 30 minutes following vaccination and up to 28 days post-vaccination. At the 10- and 28-

days post-vaccination visits, a clinician asked for the presence of local reaction, headache,

fatigue, muscle pain, fever, gastrointestinal problem and any other symptom since the

previous visit. Participants were also asked to report any other symptoms or concerns during

and outside scheduled visits. The nature, relatedness, severity and outcome of each AE were

recorded (20,21). All events were coded using MedDRA dictionary, version 20.0.

Statistical analysis

Weighing the public health consequence of loss of protection against an increase in available

vaccine doses and coverage, non-inferiority of the fractional dose was defined as

seroconversion rate at 28 days post-vaccination no more than 10% lower than the standard

dose. We assumed 95% seroconversion with standard doses and considered that a loss of

protection to 85% seroconversion with fractional doses would still ensure protection above

80% necessary to interrupt local transmission (22).The non-inferiority margin is supported by

a modelling study where fractional doses were shown beneficial in high-transmission areas if

80 to 90% efficacious (23). To detect a non-inferiority margin of 10% in seroconversion 28

days post-vaccination, with a 2.5% level of significance for a one-sided test, 90% power, and

accounting for 5% loss to follow-up and 15% baseline yellow fever seropositivity, a sample

size of 120 per manufacturer and dose was required, for a total of 240 per manufacturer. In

total, 960 participants were to be recruited (480 per study site).

The primary endpoint was seroconversion, defined as $\geq$4-fold rise in $PRNT_{50}$ titer between

baseline pre-vaccination and 28 days post-vaccination for fractional compared to standard

dose for each WHO-prequalified vaccine. Secondary endpoints include assessment of

geometric mean $PRNT_{50}$ titers (GMT) and geometric mean fold increase (GMFI) at 28 days

post-vaccination, and seroconversion, GMT and GMFI at 10 days and one year after

vaccination. Analysis using $PRNT_{90}$ titers are presented in the supplementary material.

Analysis consisted of pairwise comparisons of fractional versus standard dose of the same

manufacturer's vaccine. No comparisons were made of vaccines from different

manufacturers. Comparison of baseline characteristics were done using χ^2 tests (Fisher's exact

test for k<5) for categorical variables and Student's t-test for continuous variables. Any

$PRNT_{50}$ titer reported as seronegative (below the Limit of Quantification (LOQ), <1:10) was

converted to LOQ/2. Any $PRNT_{50}$ titer >20480 was assigned as 20480.

The number and percentage of participants who seroconverted are presented by study arm (i.e.

manufacturer and dose) with two-sided exact 95% confidence interval (CI) based on the

Clopper-Pearson method. Non-inferiority for the primary outcome was assessed by

constructing a two-sided 95% CI using the Wilson score interval of the point difference

between seroconversion rates in the fractional and standard dose arms. Fractional doses were

considered non-inferior if the lower bound of the CI for difference in seroconversion was

greater than -10%.

Two-sided 95% CIs of the mean difference between log GMT and log GMFI between the

standard and fractional dose of each vaccine were generated using the t-distribution.

Immunogenicity outcomes were assessed in the per-protocol population (PP) and the intent-

to-treat population (ITT). The PP population included all participants for whom the eligibility

criteria were appropriately applied, who were seronegative at baseline ($PRNT_{50}$ below LOQ),

and who had a $PRNT_{50}$ titer at 28 days post-vaccination. The ITT population included any

vaccinated participant with at least one $PRNT_{50}$ result post-vaccination.

AEs and SAEs were summarized as number and percentage by study arm. Safety outcomes

were assessed in all vaccinated participants.

Data analysis was conducted in R, version 3.6.1. A data safety monitoring board (DSMB)

regularly reviewed study data.

ClinicalTrials.gov Identifier: NCT02991495

Role of the funding source

The funder of the study had no role in study design, data collection, data analysis, data

interpretation, or writing of the report. The corresponding author had full access to all data in

the study and had final responsibility for the decision to submit for publication.

Results

From 6[th] November 2017 to 21[st] February 2018, 1,029 participants were assessed for

eligibility, 69 were ineligible, and 960 were randomized to a vaccine manufacturer and dosage

(120 per study arm) and vaccinated (Figure 1). Nearly all (99.2%) participants completed the

day 28 post-vaccination visit and 96.8% completed the one year post-vaccination visit. The most frequent reasons for discontinuation were migration out of study area (n=13), loss to follow-up (n=10) and non-compliance with study visits (n=4). There were 2 discontinuations due to protocol violation: inclusion of a 65-year-old man and re-vaccination of a participant attending the day 28 post-vaccination visit.

The primary analysis (PP) included 110 to 117 participants per arm, 93.6% of the randomized participants. There were 49 participants with detectable $PRNT_{50}$ at baseline (included in ITT analysis) and 11 participants with missing $PRNT_{50}$ results at baseline or 28 days follow-up (Figure 1).

The mean age of participants was 35.7 years of age and 55.1% were females. Participant characteristics were similar at baseline between study arms (Table 2).

At 28 days post-vaccination, seroconversion rates were high in all arms, with at least 98.2% of participants seroconverting (Table 3). The difference in seroconversion between fractional and standard dose arms was 1.71% (95%CI: -2.60, 5.28) for the vaccine produced by Bio-Manguinhos/Fiocruz with the 17DD substrain, -0.90% (95%CI: -4.24, 3.13) for the vaccine produced by Chumakov IPVE with the 17D-213 substrain, 1.82% (95%CI: -2.75, 5.39) for the vaccine produced by Institut Pasteur Dakar and 0.0% (95%CI: -3.32, 3.29) for the vaccine produced by Sanofi Pasteur, both derived from the 17D-204 substrain. The lower bound of the 95% confidence interval for the difference in seroconversion between fractional and standard dose arms for each vaccine manufacturer excluded the defined non-inferiority margin of -10%, indicating non-inferiority of the fractional dose for each vaccine (Table 3 and Figure 2). Non-inferiority is also met considering $PRNT_{90}$ results (Table S1). Results for the ITT population at 28 days post-vaccination were similar, with a lower bound of the confidence interval for the difference in seroconversion between fractional and standard dose arms for each vaccine manufacturer of -2.58%, -6.11%, -2.11% and -3.10% (Table S2). Seroconversion was high even among those adults with baseline seropositivity to yellow fever (Table S3).

At 10 days post-vaccination, the percentage of participants that seroconverted in each arm varied from 52.5% to 78.2%, with no differences between fractional and standard doses by vaccine manufacturer (Table 3). Seroconversion rates remained high one year after

vaccination, with between 96.5% and 100% of those who received a fractional dose and

>99% of those who received the standard dose maintaining a ≥4-fold rise in $PRNT_{50}$ titer

from baseline (Table 3).

GMTs of neutralizing antibodies 28 days post-vaccination were high in each arm, with titers

ranging from 3939 to 5874 in the fractional dose arms and 2576 to 5817 in the standard dose

arms. For one vaccine GMTs were higher (Table 3). At 10 days post-vaccination, GMTs were

lower compared to 28 days post-vaccination and were, on average, lower in the fractional

dose compared to the standard dose arm for three vaccines, though the confidence intervals

for the difference are wide and overlap 0 for all but one vaccine (Table 3). At one year post-

vaccination, GMTs remained high in all study arms and were consistently higher in the

fractional dose arms (range: 2862-5339) compared to standard dose arms (range: 2269-3876).

There was some evidence for a slow decline in GMT from 28 days to one year post-

vaccination in all arms (Figure S5).

GMFIs showed an increase in neutralizing antibody titers from baseline at each time point for

all vaccine arms, with GMFIs reaching highest point 28 days post-vaccination (Tables S7,

S11 and S12).

AEs were common with between 58% to 63% of participants per arm reporting at least one

AE and 32 to 49% reporting an AE classified as vaccine-related within 28 days of

vaccination. There were similar number of events across all arms. The most common related

AEs were headache, fatigue, muscle pain, and self-reported fever (Table S14).

In total, 9 SAEs, including 3 deaths, were reported. All were classified as not related to the

study vaccines (Tables S18 and S19).

Discussion

This study shows that one-fifth fractional doses of the four WHO-prequalified yellow fever

vaccines met the main outcome of non-inferiority in seroconversion by $PRNT_{50}$ compared to

the standard dose. While not firmly established, non-human studies suggest that neutralizing

titers ≥40-50 protect against lethal yellow fever infection, and these levels have been used as

the cut-off for protection from infection in humans (24). At 28 days post-vaccination, almost

all participants showed high levels of neutralizing antibodies far in excess of the assumed

protective threshold. Seroconversion rates and GMT remain high up to one year post-

vaccination for both fractional and standard doses for all vaccines. There were no major safety

concerns with either the standard or fractional dose of each of the four WHO-prequalified

vaccines.

The results of our trial are aligned with previous studies on fractional dosing using the 17DD

substrain vaccine in other contexts showing that nearly all vaccinated individuals seroconvert

within 28 days of vaccination (10,14). We now extend the evidence base to randomized

comparisons for all WHO-prequalified vaccines and to a general adult population in sub-

Saharan Africa. Previous studies indicate that the immune response generated by fractional

doses is long-lived (14,25,26), consistent with our findings of a robust immunogenic response

at one year post-vaccination, though longer-term studies are warranted.

We assessed, for the first time in a randomized trial including all four WHO-prequalified

yellow fever vaccines, seroconversion rates at 10 days post-vaccination in all participants.

These results show overall lower seroconversion and GMTs in the fractional dose arms,

suggesting a possible delayed immune response of the fractional doses. However, previous

studies have shown that 80-90% of standard dose vaccine recipients have protective levels of

neutralizing antibodies 10 days after vaccination (27). This proportion is lower in our study,

even among participants that received a standard dose of vaccine. Our study was performed in

an African population, which may have lower B and T cell responses to yellow fever

vaccination than European populations (28). Given the likely use of fractional doses in

outbreak responses, the low rates of seroconversion at 10 days are concerning and emphasize

the need for early vaccination campaigns in outbreak response. Implications for the current

International Health Regulations, which allow for travel 10 days post-vaccination, may

require further study.

In this study, we used vaccines as close as possible to each manufacturer's internal minimum

specification for potency. Vaccines titers were, however, high, with potencies between 6 and

43 times the minimum specification established by WHO. All fractional doses still contained

potencies above the minimum specification of 1000 IU. Vaccines also differed in expiry dates

and had between 3 and 16 months of remaining shelf-life (i.e. between 1/8[th] and less than 1/5[th]

remaining shelf-life) at the time of testing at NIBSC (Table 1). Current yellow fever vaccines

were developed more than 80 years ago (1) following a production process that has not

significantly changed since and lacking clinical data to support minimum specifications (13).

Historically, vaccines have been released at higher titers than the recommended minimum,

partly to account for the loss of potency during shelf-life (13). Average doses of currently

produced WHO-prequalified vaccines vary between 12,874 and 43,651 IU and consequently,

titers of fractional doses would in principle still exceed WHO's minimum specification (29).

Given the longstanding practice of releasing batches with high potency, and the dependence

on historical precedent for dose requirements, there was significant uncertainty regarding the

use of fractional dosing, even for doses above the minimum specifications. This is reflected in

the WHO recommendations for fractional dosing that the minimal dose administered should

preferentially contain 3000 IU/dose, three times the minimum specification for vaccine

potency (4). Our data suggest that, even at the lower end of the range of vaccine potency and

with minimal remaining shelf-life, fractional doses can be confidently recommended. We

consider that the batches selected for this study represent the WHO-prequalified vaccines as

currently produced by the different manufacturers and that results should be generalizable to

other batches as long as there is not a significant change in the manufacturing process.

Therefore, our study provides additional confidence in using fractional doses for the vaccines

as currently produced.

These results support the use of one-fifth fractional doses of the four WHO-prequalified

yellow fever vaccines for the general adult population when there are insufficient standard

doses to protect the population at risk during an outbreak, thereby filling a critical knowledge

gap to support WHO policy on the use of fractional dosing of yellow fever vaccine (5). The

use of fractional dosing will expand the outbreak stockpile up to five-fold and therefore will

be a critical back-up tool in case of vaccine supply shortage during yellow fever outbreak

response. We are currently conducting sub-studies to assess immunological non-inferiority

and safety of fractional doses among children from 9 months to 5 years of age and adults

living with HIV

Contributors. AJG, PB, KHG, DC, JH, TM, AB and RFG designed the study; PK, DK and GW collected the data; KHG conducted the statistical analysis, GF, NSM and MD conducted the laboratory analyses; and AJG prepared the first draft of the manuscript. AJG and RFG vouch for the accuracy and completeness of the data and analyses reported. All authors contributed to the interpretation of data, critically reviewed the manuscript and decided to publish the paper.

Declaration of interests. We declare no competing interests

Disclaimer. JH is staff of the World Health Organization. The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the World Health Organization

Acknowledgements.
The study was funded by Médecins Sans Frontières Foundation, Wellcome Trust (grant no. 092654) and the UK Department for International Development. The vaccines were donated in kind by Bio-Manguinhos/Fiocruz (Brazil), Federal State Unitary Enterprise of Chumakov Institut of Poliomyelitis and Viral Encephalitides (Russia), Institut Pasteur de Dakar (Senegal), and Sanofi Pasteur (France). Epicentre receives core funding from Médecins Sans Frontières.
We wish to thank study participants and the Scientific Committee, the DSMB and the National Institute for Biological Standards and Control (United Kingdom) for critical support to this project. We also thank the YEFE teams in KEMRI-Wellcome Trust Research Programme, Kilifi, Kenya and in Epicentre Mbarara Research Centre, Mbarara, Uganda.

454   Table 1. Characteristics of study vaccines

Manufacturer	Substrain	Expiry date	Log$_{10}$ IU/dose Mean (SD)
Bio-Manguinhos/Fiocruz (Bio-Manguinhos)	17DD	31/10/2018	4.59 (0.10)
Federal Stata Unitary Enterprise of Chumakov Institute of Poliomyelitis and Viral Encephalitides (Chumakov IPVE)	17D-213	28/02/2018	4.64 (0.05)
Institut Pasteur de Dakar (IPD)	17D-204	31/03/2019	3.83 (0.24)
Sanofi Pasteur	17D-204	28/02/2019	4.22 (0.10)

Note: Standard doses of the vaccines were tested at the National Institute for Biological Standards and
Control (NIBSC), UK, December 2017. The Log$_{10}$ IU/dose is the mean of 3 assays; SD: Standard
deviation.

Figure 1. Trial profile

Note: number analyzed refers to the number included in the immunogenicity analysis at each time

point. Differences with number of completed visits are due to missing PRNT result.

Table 2. Baseline demographic and clinical characteristics

	Bio-Manguinhos		Chumakov IPVE		IPD		Sanofi Pasteur	
	Fractional (N=120)	Standard (N=120)	Fractional (N=119)	Standard (N=120)	Fractional (N=120)	Standard (N=120)	Fractional (N=120)	Standard (N=120)
Age at enrolment, Mean (SD)	34.5 (11.5)	35.0 (11.1)	37.2 (11.4)	35.9 (11.8)	35.5 (11.4)	35.4 (11.7)	35.3 (11.7)	36.8 (12.9)
Female sex, no (%)	73 (60.8)	73 (60.8)	58 (48.7)	62 (51.7)	64 (53.3)	72 (60.0)	61(50.8)	66 (55.0)
HIV+, no (%)	4(3.3)	1 (0.8)	0 (0)	0 (0)	1 (0.8)	1 (0.8)	1 (0.8)	2 (1.7)
Seropositive to YF at baseline, no (%)	6* (5.0)	1 (0.8)	7** (5.9)	4* (3.4)	7 (5.8)	9 (7.5)	7 (5.8)	8 (6.7)

464   *N=119; **N=118 due to missing PRNT titer at baseline

Table 3. Non-inferiority of seroconversion, GMT and GMFI by $PRNT_{50}$ in fractional vs. standard dose of yellow fever vaccine at Day 10, Day 28 and Day 365 post-vaccination, by vaccine manufacturer

| Manufacturer | Time | Dosage | N | Seroconversion | | | GMT | |
				n	% (95% CI)	Difference, % (95% CI)	GMT (95% CI)	Difference GMT (95% CI)
Bio-Manguinhos	D10	Fractional	116	71	61.2 (51.7, 70.1)	1.54 (-10.7, 13.9)	51.1 (35.7, 73.1)	1.11 (0.67, 1.85)
		Standard	119	71	59.7 (50.3, 68.6)		46.0 (31.9, 66.3)	
	D28	Fractional	111	111	100.0 (96.7, 100.0)	1.71 (-2.60, 5.28)	3939 (2812, 5516)	0.97 (0.60, 1.58)
		Standard	115	117	98.3 (94.0, 99.8)		4064 (2850, 5794)	
	D365	Fractional	117	116	99.1 (95.3, 100.0)	0.04 (-4.02, 3.94)	3225 (2284, 4555)	1.17 (0.71, 1.93)
		Standard	112	111	99.1 (95.1, 100.0)		2757 (1914, 3973)	
Chumakov IPVE	D10	Fractional	118	62	52.5 (43.1, 61.8)	-8.80 (-21.4, 3.38)	36.8 (25.5, 53.3)	0.66 (0.40, 1.09)
		Standard	119	73	61.3 (52.0, 70.1)		56.1 (39.5, 79.6)	
	D28	Fractional	110	111	99.1 (95.1, 100.0)	-0.90 (-4.24, 3.13)	5874 (4162, 8289)	1.01 (0.64, 1.59)
		Standard	114	114	100.0 (96.8, 100.0)		5817 (4297, 7876)	
	D365	Fractional	114	110	96.5 (91.3, 99.0)	-2.65 (-7.06, 2.57)	3801 (2667, 5417)	1.08 (0.66, 1.77)
		Standard	116	115	99.1 (95.3, 100.0)		3514 (2486, 4966)	
IPD	D10	Fractional	120	74	61.7 (52.4, 70.4)	-6.71 (-18.8, 5.10)	49.0 (34.2, 70.1)	0.54 (0.31, 0.93)
		Standard	117	80	68.4 (59.1, 76.7)		91.1 (60.2, 137.9)	
	D28	Fractional	112	112	100.0 (96.8, 100.0)	1.82 (-2.75, 5.39)	4279 (3182, 5753)	1.66 (1.04, 2.65)
		Standard	108	110	98.2 (93.6, 99.8)		2576 (1788, 3712)	
	D365	Fractional	118	118	100.0 (96.9, 100.0)	0.87 (-3.02, 4.10)	2862 (2091, 3918)	1.26 (0.80, 1.99)
		Standard	115	114	99.1 (95.3, 100.0)		2269 (1624, 3171)	
Sanofi Pasteur	D10	Fractional	120	83	69.2 (60.1, 77.3)	-8.98 (-20.2, 1.90)	76.4 (53.0, 110.2)	0.65 (0.38, 1.10)
		Standard	119	93	78.2 (69.6, 85.2)		117.5 (80.4, 171.7)	
	D28	Fractional	113	113	100.0 (96.8, 100.0)	0.0 (-3.32, 3.29)	5545 (4106, 7488)	1.30 (0.83, 2.04)
		Standard	112	112	100.0 (96.8, 100.0)		4252 (3033, 5961)	
	D365	Fractional	116	114	98.3 (93.9, 99.8)	-1.72 (-5.14, 2.62)	5339 (3955, 7206)	1.38 (0.91, 2.09)
		Standard	117	117	100.0 (96.9, 100.0)		3876 (2886, 5205)	

Note: Difference: Fractional-Standard; Ratio: Fractional/Standard
Day 28 in per-protocol population (i.e. seronegative at baseline). Day 10 and D365 in the intent-to-treat population (i.e. any participant with at least one $PRNT_{50}$ result post-vaccination

References

1. WHO. Vaccines and vaccination against yellow fever. WHO Position Paper. Wkly Epidemiol Rec. 2013;27(88):269–84.

2. WHO. International Coordinating Group (ICG) on Vaccine Provision [Internet]. Online Q&A. 2016 [cited 2020 Mar 31]. Available from: http://www.who.int/csr/disease/icg/qa/en/

3. Monath TP, Woodall JP, Gubler DJ, Yuill TM, Mackenzie JS, Martins RM, et al. Yellow fever vaccine supply: a possible solution. Comment. Lancet. 2016;387(10028):1599–600.

4. WHO. Fractional dose yellow fever vaccine as a dose-sparing option for outbreak response. WHO Secretariat information paper. [Internet]. 2016. Available from: https://www.who.int/immunization/sage/meetings/2016/october/3_Fractional_dose_secretariat_report_full_version.pdf

5. WHO. Yellow fever vaccine: WHO position on the use of fractional doses - June 2017. Addendum to Vaccines and vaccination against yellow fever WHO Position Paper - June 2013. Wkly Epidemiol Rec. 2017;25(92):345–50.

6. WHO. Yellow fever mass vaccination campaign using fractional dose in Kinshasa, DRC. SAGE meeting report [Internet]. 2016. p. 5. Available from: https://www.who.int/immunization/sage/meetings/2016/october/4_Yellow_fever_mass_vaccination_campaign_using_fractional_dose_in_Kinshasa.pdf

7. WHO. Yellow fever in Africa and the Americas, 2018. Wkly Epidemiol Rec. 2019;(33):365–80.

8. Lopes de Souza O, de Almeida Guimaraes SSD, de Carvalho R. Studies on yellow fever vaccine III-dose response in volunteers. J Biol Stand. 1988;16:77–82.

9. Roukens AH, Vossen AC, Bredenbeek PJ, van Dissel JT, Visser LG. Intradermally Administered Yellow Fever Vaccine at Reduced Dose Induces a Protective Immune Response: A Randomized Controlled Non-Inferiority Trial. PLoS One. 2008;3(4).

10. Martins RM, Maia M de LS, Farias RHG, Camacho LAB, Freire MS, Galler R, et al. 17DD yellow fever vaccine: a double blind, randomized clinical trial of immunogenicity and safety on a dose-response study. Hum Vaccin Immunother. 2013;9(4):879–88.

11. Campi-Azevedo AC, de Almeida Estevam P, Coelho-Dos-Reis JG, Peruhype-Magalhães V, Villela-Rezende G, Quaresma PF, et al. Subdoses of 17DD yellow fever vaccine elicit equivalent virological/immunological kinetics timeline. BMC Infect Dis. 2014;14:391.

12. WHO. Annex 5. Recommendations to assure the quality, safety and efficacy of live attenuated yellow fever vaccines. In: WHO Expert Committee on Biological Standardization. Sixty-firs. Geneva; 2013. p. 241–314.

13. Ferguson M, Shin Jinho J, Knezevic I, Minor P, Barrett A. WHO Working Group on Technical Specifications for Manufacture and Evaluation of Yellow Fever Vaccines, Geneva, Switzerland, 13–14 May 2009. Vaccine. 2010;28(52):8236–45.

14. Casey RM, Harris JB, Ahuka-Mundeke S, Dixon MG, Kizito GM, Nsele PM, et al. Immunogenicity of Fractional-Dose Vaccine during a Yellow Fever Outbreak — Final Report. N Engl J Med. 2019 Aug 1;381(5):444–54.

15. Kimathi D, Juan A, Bejon P, Grais RF, Warimwe GM. Randomized, double-blinded, controlled non-inferiority trials evaluating the immunogenicity and safety of fractional doses of Yellow Fever vaccines in Kenya and Uganda. Wellcome Open Res. 2019;4:182.

16. Kwagonza L, Masiira B, Kyobe-Bosa H, Kadobera D, Atuheire EB, Lubwama B, et al. Outbreak of yellow fever in central and southwestern Uganda, February-may 2016. BMC

515   Infect Dis. 2018;18(1):1–9.

516   17.   Hombach J, Solomon T, Kurane I, Jacobson J, Wood D. Report on a WHO consultation on
517   immunological endpoints for evaluation of new Japanese encephalitis vaccines, WHO, Geneva,
518   2-3 September, 2004. Vaccine. 2005 Nov 1;23(45):5205–11.

519   18.   Roehrig JT, Hombach J, Barrett ADT. Guidelines for Plaque-Reduction Neutralization Testing
520   of Human Antibodies to Dengue Viruses. Viral Immunol. 2008;21(2):123–32.

521   19.   De Madrid AT, Porterfield JS. A simple micro-culture method for the study of group B
522   arboviruses. Bull World Health Organ [Internet]. 1969 [cited 2019 Oct 18];40(1):113–21.
523   Available from: http://www.ncbi.nlm.nih.gov/pubmed/4183812

524   20.   WHO. Detection and investigation of Serious Adverse Events following Yellow Fever
525   vaccination. Guidance from an informal consultation of experts, 18–19 November 2008,
526   Geneva, Switzerland [Internet]. 2010. Available from:
527   https://www.who.int/csr/resources/publications/HSE_GAR_ERI_2010_2/en/

528   21.   Food and Drug Administration. Guidance for Industry. Toxicity grading scale for healthy
529   volunteers enrolled in preventive vaccine clinical trials. U.S. Department of Health and Human
530   Services. Center for Biologics Evaluation and Research; 2007.

531   22.   WHO. A Global Strategy to Eliminate Yellow Fever Epidemics (EYE) 2017-2026 [Internet].
532   2017. Available from: www.who.int/csr/disease/yellowfev/eye-strategy/en/

533   23.   Wu JT, Peak CM, Leung GM, Lipsitch M. Fractional Dosing of Yellow Fever Vaccine to
534   Extend Supply: A Modeling Study. Lancet. 2016;388(10062):2904–11.

535   24.   Mason RA, Tauraso NM, Spertzel RO, Ginn RK. Yellow Fever Vaccine: Direct Challenge of
536   Monkeys Given Graded Doses of 17D Vaccine. Appl Microbiol. 1973;25(4):539.

537   25.   Menezes R De, Lourdes M De, Maia S, Maria S, Lima B De, Guimarães T, et al. Duration of
538   post-vaccination immunity to yellow fever in volunteers eight years after a dose-response
539   study. Vaccine [Internet]. 2018;36(28):4112–7. Available from:
540   https://doi.org/10.1016/j.vaccine.2018.05.041

541   26.   Roukens AHE, van Halem K, de Visser AW, Visser LG. Long-Term Protection After
542   Fractional-Dose Yellow Fever Vaccination: Follow-up Study of a Randomized, Controlled,
543   Noninferiority Trial. Ann Intern Med. 2018;169(11):761–5.

544   27.   Monath TP. Yellow fever: An update. Lancet Infect Dis. 2001;1(1):11–20.

545   28.   Muyanja E, Ssemaganda A, Ngauv P, Cubas R, Perrin H, Srinivasan D, et al. Immune
546   activation alters cellular and humoral responses to yellow fever 17D vaccine. J Clin Invest.
547   2014;124(7):3147–58.

548   29.   Vannice K, Wilder-Smith A, Hombach J. Fractional-Dose yellow fever vaccination —
549   advancing the evidence base. Vol. 379, New England Journal of Medicine. Massachussetts
550   Medical Society; 2018. p. 603–5.