The Effects of Coronavirus Victimization Distress and Coronavirus Racial Bias on Mental Health Among Black, Indigenous and Latinx Young Adults in the United States

Celia B. Fisher [1,2] [*], Xiangyu Tao [1], and Tiffany Yip [1]

[1]Department of Psychology, Fordham University, Bronx, New York, U.S.

[2]Center for Ethics Education, Fordham University, Bronx, New York, U.S.

* Corresponding Author: Fisher@fordham.edu

Brief Title: Effects of Coronavirus Victimization Distress and Racial Bias on Mental Health Among U.S. Ethnic Minority Adults

The Effects of Coronavirus Victimization Distress and Coronavirus Racial Bias on Mental Health Among Black, Indigenous and Latinx Young Adults in the United States

Abstract

Background. People of color in the U.S. have been disproportionately impacted by the COVID-19 pandemic in terms of rates of infection and morbidity. Explanations for these disparities include over-representation as essential workers and long-standing inequities in access to health services. Prior to the pandemic, racial discrimination has been associated with depression and general anxiety. However, the effect of discrimination and racial bias specific to the Coronavirus on mental health has not been examined. The aim of this study was to assess the effects of Coronavirus victimization distress and Coronavirus racial bias beliefs on the mental health of young adult people of color.

Method. An online survey administered to a national sample of 350 Black, Indigenous and Latinx adults (18 – 25 years) included Coronavirus health risks, prescription and financial security, measure of depression and anxiety and 2 new psychometrically validated measures for Coronavirus related victimization distress and racial bias.

Results. Employment, number of Coronavirus health risks, Coronavirus victimization distress and Coronavirus racial bias were positively correlated with each other and with depression and anxiety. By contrast, household income and perceived financial and prescription security were negatively correlated with Coronavirus victimization, Coronavirus racial bias and with the mental health indices. Structural equation modeling controlling for demographic variables indicated perceived Coronavirus racial bias mediated the effect of Coronavirus victimization distress on both mental health measures across all groups.

38 **Conclusions.** Results suggest the COVID-19 pandemic has created new pathways to mental

39 health disparities among young adults of color by reversing formerly protective factors such as

40 employment, and by exacerbating structural and societal inequities linked to race. Findings

41 highlight the necessity of creating mental health services tailored to the specific needs of

42 racial/ethnic minorities during the current and future health crises.

Introduction

The COVID-19 pandemic has amplified existing racial/ethnic disparities in the United

States. People of color have higher risks of COVID-19 infection and mortality. Black,

Indigenous and Latinx adults are at disproportionally higher risk. For example, as of July 28,

2020, Black, Indigenous, and Latinx people in the U.S. compared with other racial/ethnic groups

accounted for greater proportions of Coronavirus infections among young adults ages 18 – 29

years (17.7%, 1.3%, 30.28%, respectively) and an even greater proportion of deaths (34.6%,

4.5%, 38.66%, respectively) (1). Disparities faced by people of color during the pandemic are

rooted in long-standing inequities in access to health services and health outcomes (2-4). These

include lower levels of treatment satisfaction and trust, utilization rate, and service outcome

resulting in delay and lack of adherence to treatment uptake (3-5). Compared with other

racial/ethnic groups Black, Indigenous and Latinx persons in the US also have higher rates of

CDC identified medical conditions associated with risk of severe illness from COVID-19,

including obesity, diabetes, asthma, and HIV (6-14). In addition, Black, Indigenous, and Latinx

adults are more likely to be employed as lower skilled workers, and during the current crises, as

essential workers (15, 16), which increases risk of COVID-19 exposure. In 2018, approximately

20% of employed Black and Latinx men worked in service occupations, compared to 13% of

White and Asian men (16). Although 11.9% of all workers are Black, they account for 17% of

workers in Coronavirus frontline industries such as grocery and drug stores, public

transportation, trucking, and health care (15). In addition to these longstanding disparities in

healthcare, stay at home orders and reallocation of pharmaceuticals for emergency care have

created barriers to prescriptions for health conditions related to higher risk for Coronavirus,

placing people in lower economic strata and racially segregated communities at higher risk (17).

Mental health among people of color

People of color also suffer higher mental health risks compared to non-Hispanic whites

(5, 18, 19). The National Center for Health Statistics (20) reported that Indigenous populations

are 50% more likely to report hopelessness, worthlessness, and feelings of nervousness or

restlessness compared to non-Hispanic Whites and 80% more likely to report frequent sadness.

Black and Latinx women also report significantly greater depressive symptomatology than non-

Hispanic White females (18, 21). Although the prevalence rates of major depression and general

anxiety among Black and Latinx are often reported to be similar or lower than non-Hispanic

Whites, these findings have been attributed to the fact people of color in the US seek mental

health clinical care at rates well below their need and face discrimination during diagnosis and

treatment, misdiagnosis and clinician bias (2, 5, 22-24). Other studies suggest the long-lasting

effects of depression may be higher for African American and Caribbean Black men compared to

non-Hispanic Whites (56% versus 38.6%) (25).

Factors contributing to mental health problems among people of color

The health, economic and employment disparities described above for people of color

also contribute to increased internalizing disorders. Medical disparities currently identified as

increasing Coronavirus risk, have previously been shown to be associated with poor mental

health in studies on the general population and on people of color specifically. For example,

obesity and asthma have been associated with mood disorders and health-related quality of life

among Black and Latinx populations in the U.S. (26-30). Relatedly, inadequate access to health

care also contributes to depression and anxiety (31). Lower socioeconomic status (SES),

financial and food insecurity in general have been associated with higher levels of depression

and anxiety among Black and Latinx populations (32-39); although some studies found that

higher SES is associated with depression among Black people (40-42). Past research has

repeatedly reported on the protective effect of employment on mental health (43, 44). However,

during the pandemic, employed individuals, especially those in positions involving in-person

contact with others during working hours, are at higher risks of COVID-19 infection (45). Since

racial/ethnic minorities are more likely to be employed in health care or low-wage jobs that are

considered essential during the pandemic (31, 46), emerging reports indicate they are

experiencing higher levels of depression and anxiety (47). Finally, although in general gender

and sexual minority status among young adults is associated with increases in depression and

anxiety (48, 49), intersecting ethnic and sexual minority identities have been identified as an

increased risk factor for depression and suicidal ideation among Latinx and Black sexual

minority men and women (50-54) and across people of color identified as transgender females

(55, 56).

The relationship between racial/ethnic discrimination and systemic

racism on depression and anxiety among people of color

People of color experience unique stressors related to their marginalized social identity

that adversely contribute to depression and anxiety (57), including explicit racial/ethnic

discrimination and microaggressions (58-61). Intergenerational trauma created by American

slavery, colonialism, Jim Crow laws, and segregation, has been shown to effect both physical

and psychological health among Black people (62-64). Loss and violation of sacred lands and a

history of segregation of American Indian boarding schools and food programs is a significant

stressor for Indigenous persons (12, 65). For Latinx people, studies have found significant

association among documentation status (66, 67), fear of deportation (68), acculturation stress

(69-71), and mental health status. Although to date, the effect of racial bias associated

specifically with the Coronavirus on mental health has not been examined, the negative impact

of racial/ethnic discrimination on physical and mental health has been well documented (58, 72-

76). Based on the stress and coping framework (77), racial/ethnic discrimination is a social

stressor that affects physiological and psychological responses such as increased levels of heart

rate, cortisol secretions, blood pressure, emotional reactivity, and reliance on maladaptive

emotion-regulation strategies (72). These in turn lead to higher risks of both physical health

problems such as obesity and cardiovascular disease (78-80), and internalizing disorders such as

depression and anxiety (73).

Prior research on populations facing contagious disease discrimination

Research on victimization distress due to contagious diseases such as HIV, H1N1, and

the 2003 severe acute respiratory syndrome (SARS) has found that contagious disease

discrimination is associated with long-term negative mental health outcomes including

depression and anxiety (81-84). For example, in a study on SARS-related stigma in Hong Kong,

residents who lived in the first officially identified site of community outbreak reported they

were marginalized in their work, social, and home environments and experienced high levels of

psychological distress, which did not decrease during the post-SARS era (85, 86). Prior biases

against marginalized social groups also play a role in disease discrimination. For example, in

Malaysia, during the HINI Flu epidemic, farmers held misconceptions that societal "out-groups"

such as sexual minorities, the homeless and sex workers had higher risks of infections, which in

turn led to discrimination against these groups (87).

Individuals who have personally experienced a negative event may attribute the

victimization to various factors, such as personal reasons and discrimination (88). Following an

experience of Coronavirus related bullying or discrimination, people of color may attribute the

event to racial/ethnic discrimination based on beliefs that the Coronavirus has increased systemic

racism in the US. The attributional ambiguity perspective suggests that attributions to racial

discrimination rather than personal attributions may be self-protective, buffer the harmful effect

of negative events, and contribute to better mental health (89). However, recent studies have

suggested that there are still harmful effects on mental health with attributions to discrimination

(73). This suggests that perceived increases in systemic racism as a consequence of the

Coronavirus pandemic may not only be directly related to increases in depression and anxiety but

may mediate the influence of Coronavirus victimization on these mental health indices.

The current study

Although the devastating impact of COVID-19 on the health of people of color in the

U.S. has been well established, there is a paucity of research on how individual and societal

factors are affecting the mental health of these populations. The current study conducted an

anonymous online survey in April 2020 to examine the association among demographic

variables, Coronavirus victimization distress, and Coronavirus racial bias with depression and

anxiety among Black, Latinx and Indigenous young adults. The following hypotheses were

tested:

1. Employment status, sexual and gender minority identity, lower household income,

financial and prescription insecurity, and number of COVID-19 health risks will be associated

with higher levels of depression and anxiety among young adults of color.

2. Higher levels of victimization distress and perceived racial bias specifically related to

the Coronavirus will be directly associated with depression and anxiety among young adults of

color.

3. Coronavirus racial bias will mediate the association between Coronavirus victimization

distress and depression and anxiety among young adults of color. The conceptual model for this

mediating effect is provided in Table 1.

Figure 1. Conceptual model for the mediating effect of Coronavirus racial bias on the association

between Coronavirus victimization distress and depression and anxiety.

Note. Conceptual model of the mediating effect of Coronavirus racial bias on the association between

Coronavirus victimization distress and depression and anxiety. Structural equation modeling will test the

model with the following covariates held constant: race/ethnicity, age, gender, sexual orientation,

employment, financial security, prescription security and COVID-19 health risks.

Methods

Participants

Data were collected during April 2020 as part of a larger Internet-based national survey

on the biological, psychological and social impact of the Coronavirus pandemic among young

adults from diverse racial/ethnic groups. For the current study, eligible participants self-

identified as Black, Indigenous, or Latinx; were between 18 – 25 years of age; self-reported they

did not currently have the Coronavirus; lived in the United States for more than one year; and

could read English at an eighth-grade level.

Procedure and data validation

Recruitment was conducted by Qualtrics XM. Qualtrics XM is an aggregator of survey

panel websites and can recruit individuals who have signed up to take paid surveys across

multiple sites. Participants are paid in points which they can exchange for gift cards. For the

current study, Qualtrics XM sent emails or provided posts to young adults who had signed up to

take surveys through various survey websites and offered compensation worth $16.50 converted

into the survey panel's point system. The emails and posts briefly described the study and

provided a link to a screener on another Qualtrics site. The screener included questions on

participants' age, race/ethnicity, assigned sex at birth, gender, sexual orientation, the US state

and zip code in which they lived, length of time living in the U.S., geographic region, living

situation, employment and student status.

The target goal was to recruit a minimum of 100 participants from each of the 3

racial/ethnic groups and to over sample for individuals living in rural areas to obtain a

representative geographic sample. A Qualtrics system feature excluded participants who did not

meet eligibility criteria and prevented them from re-entering the screener. Manual data validation

protocols were established to exclude fraudulent or repeat participants; consistency between age

and date of birth; inconsistency between reported city in which the survey was taken and zip

code, and responses with identical IP addresses. Qualtrics also includes a speed check to exclude

participants who respond in less than half the time of the median survey response.

Approximately 57,000 people were sent emails or posts. A total of 418 Black, 223 Indigenous,

and 195 Latinx participants took the screener, with 469 meeting eligibility requirements. Of

those 350 (74.63%) completed the survey and passed the speed check. For the purposes of this

study, an additional 38 respondents were eliminated if they reported they had the Coronavirus

and 7 were eliminated for missing data on the Coronavirus Racial Bias Scale, resulting in a final

sample of 305 (42% Black, 28% Indigenous; 30% Latinx).

Following the screener, eligible individuals were immediately sent to a page providing

informed consent information. Participants who selected "I agree" at the bottom of the informed

consent page, were redirected to the main survey which consisted of 204 items which took on

average 14.02 minutes (SD = 33.84); some participants took an hour or more to complete the

survey resulting in the large standard deviation. Participants were able to quit the survey at any

time by closing the survey window and their data was not included in analysis. At the end of the

survey, online resources on health information and Coronavirus prevention were provided.

Qualtrics uses unique identification numbers for each participant so that identifiable information

and survey data are not stored on Qualtrics servers. The identity of participants and their contact

information were therefore unknown to the investigators. Participants received the agreed

amount of incentive within seven days after they completed the survey to allow for data quality

checks. The study was approved by the University Institutional Review Board.

Measures

Demographic information

Demographic information collected during screening included age, gender, race and

ethnicity, sexual orientation, employment (full or part-time; whether one was employed as an

"essential worker"), student status, and region (urban, suburban, exurban, rural). Additional

information included: household income; financial security; prescription security; and an 8 item

checklist on CDC COVID-19 health risks. Item choices are provided in Table 1

Table 1. Frequency and percentages for sociodemographic characteristics and COVID-19 health

risks for Black, Indigenous, and Latinx participants young adults ages 18 – 25 years.

	Black (N = 128)		Indigenous (N = 86)		Latinx (N = 91)		Total (N = 305)		X^2_{df}	p
	N	%	N	%	N	%	N	%		
Gender									X^2_4=2.02	.73
Cisgender Male	31	14.21	23	26.74	17	18.68	71	23.28		
Cisgender Female	67	52.34	46	53.49	53	58.24	166	54.43		
Gender Minority	30	23.44	17	19.77	21	23.08	68	22.30		

									X^2	p
Sexual Orientation									$X^2_4= .65$	.96
Heterosexual	91	71.09	59	68.6	61	67.03	211	69.18		
Sexual Minority	33	25.78	25	29.07	27	29.67	85	27.87		
Do not wish to answer	4	3.13	2	2.33	3	3.3	9	2.95		
Employment Status									$X^2_4=1.45$	.83
Not employed	55	42.97	34	39.53	32	35.16	121	39.67		
Essential workers	44	34.38	31	36.05	34	37.36	109	35.74		
Non-essential workers	29	22.67	21	24.42	25	27.47	75	24.59		
Student	73	57.03	45	52.33	56	61.54	174	57.05	$X^2_2=1.53$	.46
Financial Security									$X^2_4=7.61$	.11
I can't make ends meet	46	35.94	23	26.74	22	24.18	91	29.84		
I have just enough	49	38.28	40	46.51	51	56.04	140	45.90		
I am comfortable	33	25.78	23	26.74	18	19.78	74	24.26		
Household Annual Income									$X^2_6=1.83$	.93
Less than $30,999	58	45.31	42	48.84	39	42.86	139	45.57		
$31,000 – $79,999	38	29.69	25	29.07	31	34.07	94	30.82		
More than $80,000	11	8.59	9	10.47	8	8.79	28	9.18		

I do not want to answer/I do not know	21	16.41	10	11.63	13	14.29	44	14.43		
Education Level									X^2_8=13.89	.085
Partial high school or less	11	8.59	9	10.47	14	15.38	34	11.15		
High school graduate	59	46.09	29	33.72	23	25.27	111	36.39		
Some college	41	21.09	34	27.91	37	23.08	112	36.72		
Graduate degree	15	11.72	12	13.95	17	18.68	44	14.43		
Missing data	2	1.56	2	2.33	0	0	4	1.31		
Region									X^2_6=**22.60**	**.001**
Urban	35	27.34	25	29.07	30	32.97	90	29.51		
Suburban	43	33.59	35	40.7	42	46.15	120	39.34		
Exurban	4	3.13	9	10.47	9	9.89	22	7.21		
Rural	46	35.94	17	19.77	10	10.99	73	23.93		
Prescription Security									X^2_6=6.81	.57
No, because of costs or lack of insurance	19	14.84	9	10.47	16	17.58	44	14.43		
No, unable to reach a doctor to get a prescription I need	25	19.53	10	11.63	14	15.38	49	16.07		
No, have not been able to get to a pharmacy to pick up my prescription	8	6.25	7	8.14	4	4.4	19	6.23		

Yes, have been able to fill my prescription needs	26	20.31	20	23.26	24	26.37	70	22.95		
Have not had any prescription needs	50	39.06	40	46.51	33	36.26	123	40.33		
COVID-19 Health risks [a]										
Asthma or chronic lung disease	21	16.41	24	27.91	23	25.27	68	22.30	$X^2_1 = 4.59$	.10
Heart condition	16	12.50	12	13.95	9	9.89	37	12.13	$X^2_1 = .71$	.70
HIV	7	5.47	4	4.65	6	6.59	17	5.57	$X^2_1 = .32$	.85
Rheumatoid Arthritis	10	7.81	8	9.30	8	8.79	26	8.52	$X^2_1 = .16$	.94
Other immunocompromised disease	8	6.25	11	12.79	9	9.89	28	9.18	$X^2_1 = 2.72$	.26
Cancer treatment	6	4.69	7	8.14	5	5.49	18	5.90	$X^2_1 = 1.14$	.56
Diabetes	11	8.59	9	10.47	8	8.79	28	9.18	$X^2_1 = .24$	.89
Obesity	17	13.28	14	16.28	14	15.38	45	14.75	$X^2_1 = .41$	.81

223 [a] Mean number of COVID-19 Health risks, standard deviations and range for each group are provided in
224 Table 5.

Coronavirus Victimization Distress Scale (CVDS)

The Coronavirus Victimization Distress Scale (CVDS) is a 5-item scale adapted from

items in the Daily Heterosexist Experiences Questionnaire (90), the LGBT People of Color

Microaggressions Scale (91), and the Adolescent Discrimination Distress Index (ADDI) (75).

The items assess distress in response to different instances of victimization (e.g. "teased or

bullied", "physically threatened") resulting from other's believing the participant was a

Coronavirus risk. Each item was formatted as "I have been […] because someone thought I was

infected with the Coronavirus." Responses were on a 5-point Likert-type scale (1 = It never

happened, 2 = It happened but did not upset me; 3 = It happened and upset me a little; 4 = It

happened and upset me moderately; 5 = It happened and upset me quite a bit).

Coronavirus Racial Bias Scale (CRBS)

The Coronavirus Racial Bias Scale (CRBS) is a 9-item scale developed to assess

participants' beliefs about how the Coronavirus is negatively affecting societal attitudes toward

one's race/ethnicity. Sample items include "I believe the country has become more dangerous for

people in my racial/ethnic group because of fear of the Coronavirus"; "People of my

race/ethnicity are more likely to lose their job because of the Coronavirus"; Negative social

media posts against people of my race/ethnicity have increased because of the Coronavirus."

Participants indicated their Coronavirus racial bias beliefs on a 4-point Likert-type scale (1 –

Strongly disagree, 4 – Strongly agree).

Patient Health Questionnaire (PHQ-9)

The 9-item Patient Health Questionnaire (PHQ-9) was used to assess the frequency of

past-month experiences with depressive symptoms (92). Sample Items include "Trouble falling

asleep, staying asleep, or sleeping too much" and "Feeling down, depressed or hopeless".

Responses are scored on a 4-point scale, anchored by (0) Not at All and (3) Nearly Every Day.

Composite scale scores are computed by the average of item responses. In prior studies,

Cronbach's alpha for this scale ranged from .79 to .91 across adults identifying as Black,

Indigenous, and Latinx (93-95). The measurement invariance of the PHQ-9 across Black,

Indigenous, Latinx and Non-Hispanic White young adults has been established (96-98).

General Anxiety Disorder Screener (GAD-7)

The 7-item Generalized Anxiety Disorder Screener (GAD-7) (99) was used to assess

anxiety symptoms of participants during the past month. Sample items include "Being so restless

that it is hard to sit still" and "Feeling afraid as if something awful might happen". Participants

respond to a 4-point scale, anchored by (0) *not at all* and (3) *nearly every day*. Composite scale

scores are computed by the average of item responses. In prior studies, Cronbach's alpha for

GAD-7 ranged from .79 to .91 across adults identifying as Black, Indigenous, and Latinx (100-

104).

Data analysis plan

Descriptive statistics were calculated for all demographic variables. This was followed by

analysis of variance (ANOVA) and Chi-square tests to assess differences on these variables

among the 3 racial/ethnic groups. To determine the independence and structure of the

Coronavirus Victimization Distress Scale and the Coronavirus Racial Bias Scale, several

confirmatory factor analyses (CFA) were conducted followed by Cronbach alphas to assess

resulting scale reliability. Measurement invariance was then tested to assess whether the

constructs of the Coronavirus Victimization Distress Scale and Coronavirus Racial Bias Scale

were the same across the three racial/ethnic minority groups (105). Correlational analyses and

ANOVAs were then conducted to examine: (1) the associations among Coronavirus

victimization distress, Coronavirus racial bias, and demographic variables; and (2) differences on

depression and anxiety levels based on demographic variables, Coronavirus victimization

distress and Coronavirus racial bias.

Structural equation modeling (SEM) using the R-4.0.1. and *lavaan* package (106, 107) to

test the a priori conceptual model based on research hypotheses, which assessed the hypothesis

that Coronavirus racial bias exerted a mediating role of Coronavirus racial bias on the effect of

Coronavirus victimization distress on depression and anxiety. Two SEM analyses were

conducted, with the first analysis estimated the effect across all racial/ethnic groups, and the

second multigroup analysis estimated the effect for each racial/ethnic group separately. An

alternative model was then examined, which reversed the relationship between Coronavirus

racial bias and Coronavirus victimization distress, with the Coronavirus victimization distress

mediated the relationship between Coronavirus racial bias and depression or anxiety.

Goodness of fit indices included the comparative fit index (CFI), Tucker-Lewis Index

(TLI), and the root-mean-square error of approximation (RMSEA). A fit of > 0.95 for the CFI

and TLI and < 0.06 for RMSEA was considered to indicate adequate fit (108). To test the

indirect effects for statistical significance, the bias-corrected bootstrapping approach was

adopted as it is robust against the violation of normal distribution assumptions for both the

sampling distribution and indirect effect (109). One thousand re-samples were drawn to estimate

the standard errors of the indirect effects and their 95% confidence intervals.

Results

Descriptive data

The sample included 305 participants (Mean age = 20.84, SD = 2.19, range =18 – 25)

self-identified as, Black (N = 128; Mean age = 20.71, SD = 2.15), Indigenous (N = 86; Mean age

= 21.30, SD = 2.41) and Latinx (N = 91%; Mean age = 20.58, SD = 1.99). An ANOVA

indicated no significant age differences across groups ($F_{2, 203}$ = 2.80, p = .06). Demographic data,

percentages and Chi square tests of significance provided in Table 1 for each racial/ethnic group.

Across race/ethnicity the majority identified as cisgender females or cisgender males, reported

heterosexual sexual orientation, were full or part-time employed (60.33%; including 35.74%

essential workers); and approximately half were students. Nearly one third (29.8%) indicated

they felt financially insecure ("can't make ends meet"). Approximately 45.57% reporting an

annual household income below the national poverty rate for a family of 5 ($30,680) (110).

On average participants reported 1.52 COVID-19 pre-existing health risks (range = 0 – 8).

The mean and standard deviation (SD) for health risks for each group was: Black (Mean = .75,

SD = 1.34, range = 0 – 7); Indigenous (Mean = 1.03, SD = 1.85, range = 0 – 8); and Latinx

(Mean = .90, SD = 1.43, range = 0 – 6). The distributions are highly skewed (skewness > 2 for

each group), as more than half of Black (65.62%), Indigenous (56.98%), and Latinx (53.84%)

participants did not report COVID-19 health risks. As indicated in Table 1, across groups thirty-

seven percent reported difficulty filling prescriptions during the past month. Chi-square tests

yielded no significant differences among the three racial/ethnic groups for the above

demographic categories, except for region, which included a higher percentage of Black

participants in rural areas.

Psychometric validation of the Coronavirus Victimization Distress

Scale (CVDS) and Coronavirus Racial Bias Scale (CRBS)

As a prerequisite for the SEM analysis, three confirmatory factor analyses were

conducted to determine the independence and structure of the Coronavirus Victimization

Distress Scale (CVDS) and the Coronavirus Racial Bias Scale (CRBS). Model 1 included all

items on both scales. Table 2 provides factor loadings of items on each scale. In this model,

modification indices (MI = 39.54) indicated a strong correlated error between item 7 on the

CRBS ("Due to the Coronavirus I have been cyberbullied because of my race/ethnicity") and the

CVDS (111). For that reason, item 7 on the CRBS was removed from the scale. Model 2

included all five items on the CVDS and 8 items on the CRBS. Finally, to yield higher reliability

and communalities for the SEM (112), Model 3 was developed, with all five items on CVDS and

four parcels constructed from the 8-item CRBS, which were created by randomly pairing two

items together and calculating the mean.

Table 2. Factor loadings for Coronavirus Victimization Distress Scale and Coronavirus Racial Bias

Scale based on CFA Model 1.

	Coronavirus Victimization Distress	Coronavirus Racial Bias
I have been teased or bullied because someone thought I was infected with the Coronavirus	**.752**	.321
I have been physically threatened, hit or beaten up because someone thought I was infected with the Coronavirus	**.870**	.339
I have been treated rudely or unfairly because someone thought I was infected with the Coronavirus	**.825**	.312

I have been verbally taunted or called bad names in public because someone thought I was infected with the Coronavirus	**.787**	.303
I have been cyberbullied because someone thought I was infected with the Coronavirus	**.863**	.322
Due to the Coronavirus I have been cyberbullied because of my race/ethnicity	.570	.659 [a]
Since the Coronavirus I have seen a lot more cyberbullying of people of my race/ethnicity	.424	**.689**
Negative social media posts against people of my race/ethnicity have increased because of the Coronavirus	.377	**.744**
I worry about people thinking I have the Coronavirus simply because of my race/ethnicity.	.339	**.731**
People of my race/ethnicity are more likely to get the Coronavirus	.338	**.766**
Most social and mass media reports about the Coronavirus create bias against people of my racial/ethnic group.	.277	**.719**
People of my race/ethnicity will not receive Coronavirus healthcare as good the care received by other groups	.209	**.547**
I believe the country has become more dangerous for people in my racial/ethnic group because of fear of the Coronavirus.	.192	**.613**

People of my race/ethnicity are more likely to lose their job because of the Coronavirus.	.042	**.480**

Note. Extraction Method: Maximum Likelihood. Rotation Method: Oblimin with Kaiser Normalization.

[a] This item was removed from final scale based on analysis of confirmatory factor analysis.

Table 3 shows the $\chi2$ statistic and the global model fit indices of the three hypothesized

CFA models. Model 2 had a relatively good fit based on all three model fit indices including the

global fit index root mean square error of approximation (RMSEA), the Comparative Fix Index

(CFI), and the Tucker-Lewis Index (TLI). However, model 3 fit the data best among all tested

models, and met rule of thumb criteria for adequate fit (RMSEA < .06, CFI and TLI > .95) (108).

Both resultant scales had high demonstrated internal consistency (Cronbach's alpha = .91 for

CVDS, Cronbach's alpha = .86 for CRBS). The two scales were positively correlated ($r = .43$, p

< .001).

Table 3. Global model fit indices of CFA models for Coronavirus Victimization Distress and

Coronavirus Racial Bias items.

Model	$\chi2$	df	RMSEA	CFI	TLI
1	220.21	76	.079	.935	.922
2	139.66	64	.062	.962	.953
3	34.24	26	.032	.995	.993

Note. CFA = confirmatory factor analysis; RMSEA = root-mean-square error of approximation; CFI =

Comparative Fix Index; TLI = Tucker-Lewis Index.

Measurement invariance of the CVDS and the CRBS across Black, Latinx

and Indigenous

The measurement invariance of model 3 across Black, Indigenous, and Latinx participants was

examined (See Table 4 for detailed results). The sequence of tests of measurement invariance

included configural invariance, metric invariance, scalar invariance, and strict invariance. The

null hypothesis of the $\chi 2$ difference test was that the more restricted invariance model (e.g.,

metric invariance) fit the data equally as well as the less restricted invariance model (e.g.,

configural invariance). Thus, the $\chi 2$ difference test of the two nested invariance models in the

sequence should ideally be non-significant or there should be no material change in RMSEA

(105, 113). With the exception of the invariance model, the models were not statistically

different, and the RMSEA change of strict invariance from scalar invariance did not increase

noticeably (change $\leq .01$), supporting measurement invariance of the CVDS and CRBS across

the three racial/ethnic groups.

Table 4. Global model fit indices of measurement invariance tests for the CVDS and CRBS

Invariance Model	χ^2	df	χ^2 difference test			RMSEA
			Comparison	$\chi^2(df)$	p	
Configural	109.10	78				.063
Metric	124.31	92	Configural	15.21 (14)	.36	.059
Scalar	145.87	106	Metric	21.56 (14)	.088	.061
Strict	160.57	110	Scalar	14.70 (4)	.0054	.067

Racial/ethnic and demographic differences on the CVDS and CRBS

Mean scores on the CVDS and CRBS among the 3 racial/ethnic groups are illustrated in

Table 5. An ANOVA followed by Bonferroni post-hoc comparison ($p = .001$) indicated Black

and Latinx participants had significantly higher scores on the CRBS than Indigenous respondents,

but no significant racial/ethnic differences emerged for the CVDS. Gender minority persons

reported higher scores on the CVDS ($F(2,302) = 7.78$, $p = .001$) and CRBS ($F(2,302) = 4.39$, p

$= .013$) than cisgender females ($p=.01$, $p<.001$, respectively), but did not differ from males.

There were no significant differences between participants employed as essential workers and

those employed in other positions, so employment was combined as one category. Employed

participants compared to non-employed reported higher scores on the CVDS ($F(1,303) = 15.65$,

$p < .001$) and the CRBS ($F(1,303) = 4.77$, $p =.03$). There were no significant differences between

participants who did not have any prescription needs or were able to fill their prescriptions and as

a group these participants reported lower levels of CVDS ($F(1,303) = 70.33$, $p < .001$) and

CRBS ($F(1,303) = 17.15$, $p < .001$) than those reporting prescription difficulties. Pearson's

correlation tests indicated significant positive correlations between the number of COVID-19

health risks and scores on the CVDS ($r = .39$, $p < .001$) and CRBS ($r = 17$, $p < .01$). There were

no differences on CVDS and CRBS based on age, sexual orientation, financial security, or

annual household income.

Table 5. Means, standard deviations, and ranges for COVID-19 health risks, CVDS, CRBS, PHQ-9,

and GAD-7 for Black, Indigenous, and Latinx participants.

	Black (N = 128)	Indigenous (N = 86)	Latinx (N = 91)	Total (N = 305)		
	Mean (SD)	Mean (SD)	Mean (SD)	Mean (SD)	*F*	*p*

	Range	Range	Range	Range	(df1, df2)	
COVID-19 Health Risks	.75 (1.34) Range = 0 − 7	1.03 (1.85) Range = 0 − 8	.90 (1.43) Range = 0 − 6	.88 (1.52) Range = 0 − 8	.92 (2, 302)	.55
Coronavirus Victimization Distress Scale (CVDS)	1.67 (.96) Range = 1 − 5	1.60 (.98) Range = 1 − 5	1.65 (.90) Range = 1 − 3.6	1.64 (.94) Range = 1 − 5	.17 (2,302)	.84
Coronavirus Racial Bias Scale (CRBS)	2.35 (.74) Range = 1 − 4	1.97 (.74) Range = 1 − 4	2.26 (.67) Range = 1 − 4	2.21 (.74) Range = 1 − 4	7.11 (2, 203)	**.001**
Patient Health Questionnaire□9 (PHQ□9)	2.28 (.81) Range = 1 − 4	2.37 (.83) Range = 1 − 4	2.32 (.76) Range = 1 − 4	2.32 (.80) Range = 1 − 4	.30 (2, 203)	.74
Generalized Anxiety Disorder Screener (GAD-7)	2.31 (.87) Range = 1 − 4	2.47 (.84) Range = 1 − 4	2.46 (.79) Range = 1 − 4	2.40 (.84) Range = 1 − 4	1.24 (2, 203)	.29

Relationship of race/ethnicity, CVDS and CRBS and other demographic variables with mental health indices

Before examining the SEM model, Pearson's correlation analyses were conducted to

examine relationships among demographic variables and the CVDS and CRBS and the mental

health indices (See Table 6). Consistent with hypotheses 1, employment status, sexual

orientation, and number of COVID-19 health risks were significantly and positively associated

with scores on the PHQ-9 and GAD-7 and financial and prescription security were significantly

and negatively associated with the mental health indices. There were no racial/ethnic, age,

gender, or household income differences on either mental health indices.

Table 6. Correlation matrix of all study variables and potential covariates.

Variable	1	2	3	4	5	6	7	8	9	10	11	12	13
1. Coronavirus Victimization Distress Scale (CVDS)	1	.38***	.25***	.19***	.027	-.042	.11	.090	.22***	.39***	-.085	-.43***	.069
2. Coronavirus Racial Bias Scale (CRBS)		1	.30***	.25***	.17***	-.10	.086	.059	.12*	.17**	-.061	-.23***	-.021
3. Depression (PHQ-9)			1	.82***	-.031	-.11	.109	.24***	.16**	.25***	-.23***	-.19***	.046
4. Anxiety (GAD-7)				1	-.018	-.089	.057	.21***	.14*	.22***	-.20***	-.15*	.041
5. Race/ethnicity					1	-.13*	.063	.00	.022	-.046	-.033	-.069	.017
6. Age						1	-.090	-.048	.12*	-.042	-.071	.003	.034
7. Gender							1	.22***	-.022	.12*	-.041	-.021	-.15*
8. Sexual Orientation								1	.029	.056	-.098	-.040	-.010
9. Employment									1	.22***	.085	-.23***	.16**
10. COVID-19 Health Risks										1	-.13*	-.19***	-.001

											1	.14**	.19**
11. Financial Security											1	.14**	.19**
12. Prescription Security												1	.062
13. Household Income													1

390 Note. N = 305. CVDS and CRBS are both mean scores from the two scales. * $p < .05.$ ** $p < .01$ *** p

391 $< .001.$

Tests of the mediation hypothesis

As illustrated in Table 6, consistent with hypothesis 2, scores on the CVDS and CRBS

were significantly and positively associated with depression and anxiety measures. Structural

equation modeling (SEM) was conducted to test the third hypothesis that perceived Coronavirus

racial bias mediates the effect of Coronavirus victimization distress on depression and anxiety.

Gender, age, sexual orientation, financial security, COVID-19 health risks, prescription security,

and employment status were added as covariates based on significant associations between these

factors and studied variables. Household income was not included as a covariate since there were

14.43% missing data, and it was unrelated to mental health, CVDS or CRBS. To accommodate

the use of categorical exogenous variables, parameters were estimated using weighted least

squares with robust standard errors (WLSMV) (107). Missing data (N = 9) on sexual orientation

were handled by listwise deletion (107). Model fit statistics are summarized for two SEM

analyses in Table 7. The first analysis, which posited Coronavirus racial bias as the mediator of

the effect of Coronavirus victimization distress on depression and anxiety, with racial/ethnic

groups added as a covariate, showed adequate fit on all three fit indices including the RMSEA

(.034), 90% CI [.019 .047], CFI (.980), and TCL (.972). Based on the same model, a multigroup

analysis for three racial/ethnic minority groups was then conducted and adequate fit (RMSEA

= .048, 90% CI [.029 .063], CFI = .965, TLI = .954) was found.

Table 7. Global model fit indices of structural equation models.

	$\chi2$	*df*	RMSEA	CFI	TLI
Main Model	158.85	118	.034	.980	.972
Multigroup Analysis	389.31	320	.047	.967	.956
Alternative Model	226.89	119	.055	.947	.927

Note. CFA = confirmatory factor analysis; RMSEA = root-mean-square error of approximation; CFI =

Comparative Fix Index; TLI = Tucker-Lewis Index.

The first analysis (Fig 2) which included all participants and race/ethnicity as a covariate

generated the best model fit statistics with sufficient power. Table 8 illustrates the standardized

direct, indirect, and total effects of Coronavirus victimization distress on depression and anxiety

through Coronavirus racial bias. In this model, Coronavirus victimization distress had significant

indirect effects on depression and anxiety (ß = .102, p =.006, 95% CI [.023, 0.115], ß = .095, p

=.006, 95% CI [.024, .117], respectively); direct and total effects were non-significant. The

analysis indicates that Coronavirus racial bias beliefs fully accounted for the influence of

Coronavirus victimization distress on depression and anxiety.

Figure 2. Standardized results for structural equation model 1 with bootstrapping approach testing

the mediating effect of Coronavirus racial bias on the association between Coronavirus

victimization distress and depression and anxiety.

Note. Fig 2. Structural equation model 1 standardized results with bootstrapping approach testing the

mediating effect of Coronavirus racial bias on the association between Coronavirus victimization distress

and depression and anxiety. Covariates included race/ethnicity, age, gender, sexual orientation,

employment status, financial security, healthcare, and COVID-19 health risks

* p < .05, ** p < .01

Table 8. Total, direct and indirect standardized effects of Coronavirus victimization distress on

depression and anxiety based on the first SEM analysis

Paths		*B*	95% CI
Coronavirus Victimization Distress → Coronavirus Racial Bias → Depression	Total Effect	.142	[-.004, .164]
	Indirect Effect	.102	[.023, .115]*
	Direct Effect	.040	[-.087, .116]

Coronavirus Victimization Distress → Coronavirus Racial Bias → Anxiety	Total Effect	.104	[-.022, .153]
	Indirect Effect	.095	[.024, .117]*
	Direct Effect	.009	[-.102, .100]

* Statistically significant result based on 95% CI.

Test of Alternative Hypothesis

To further investigate support for the model above, a third SEM analysis considered an alternative model which posited Coronavirus victimization distress as the mediator of the effect of Coronavirus racial bias on depression and anxiety. We found an acceptable fit based on values of RMSEA and CFI, but the TLI (.927) did not support adequate fit (See Table 7). Moreover, the standardized direct, indirect, and total effects were not statistically significant.

Exploratory Analysis

The Black subsample had adequate power (=.80) to conduct the SEM; however, the Indigenous and Latinx groups needed another six and two participants, respectively, to achieve sufficient power (114). To inform future studies, we conducted exploratory analysis to examine the same model separately for the three racial/ethnic minority groups. Table 9 presents the standardized direct, indirect, and total effects of Coronavirus victimization distress on depression and anxiety through Coronavirus racial bias for each racial/ethnic minority group. Similar to the results of the main model, significant indirect effects on depression were found for both Indigenous (ß = .146, p =. 083, 95% CI [.011, .226]) and Black (ß = .090, p =.032, 95% CI

[.012, .120]) participants, yet indirect effects on anxiety were not significant. Unexpectedly, for

Latinx participants, Coronavirus victimization distress had significant direct (ß = .270, p =.034,

95% CI [.010, .321], ß = .250, p =.031, 95% CI [.009, .314], respectively) and total effects (ß

= .331, p =.002, 95% CI [.064, .327], ß = .334, p =.001, 95% CI [.074 .343], respectively) on

depression and anxiety, although the indirect effects were not significant.

Table 9. Total, direct and indirect standardized effects of Coronavirus victimization distress on

depression and anxiety for each racial/ethnic minority group based on the multigroup SEM

analysis

Paths		B	95% CI
Black			
Coronavirus Victimization Distress → Coronavirus Racial Bias → Depression	Total Effect	.162	[-.054, .278]
	Indirect Effect	.090	[.009, .110]*
	Direct Effect	.072	[-.121, .228]
Coronavirus Victimization Distress → Coronavirus Racial Bias → Anxiety	Total Effect	.083	[-.126, .196]
	Indirect Effect	.076	[-.003, .123]
	Direct Effect	-.038	[-.196, .157]

Indigenous			
Coronavirus Victimization Distress → Coronavirus Racial Bias → Depression	Total Effect	.033	[-.135, .133]
	Indirect Effect	.146	[.011, .231]*
	Direct Effect	-.112	[-.274, .055]
Coronavirus Victimization Distress → Coronavirus Racial Bias → Anxiety	Total Effect	.040	[-.129, .146]
	Indirect Effect	.125	[-.012, .203]
	Direct Effect	-.084	[-.243, .098]
Latinx			
Coronavirus Victimization Distress → Coronavirus Racial Bias → Depression	Total Effect	.331	[.062, .334]*
	Indirect Effect	.061	[-.047, .130]
	Direct Effect	.270	[.002, .334]*
Coronavirus Victimization Distress → Coronavirus Racial	Total Effect	.334	[.074, .362]*

Bias → Anxiety	Indirect Effect	.084	[-.031, .144]
	Direct Effect	.250	[.011, .324]*

458 * Statistically significant result based on 95% CI.

Discussion

People of color in the United States have been disproportionately impacted by the

COVID-19 pandemic in terms of rates of infection and morbidity. Explanations for these

disparities include over-representation as essential workers and long-standing inequities in

financial security and access to health services associated with racial/ethnic discrimination. Prior

to the pandemic, racial and ethnic discrimination has been associated with poor mental health.

Our data demonstrate that in addition to employment and health risks, victimization experiences

and perceived increases in systemic racial biases specifically associated with the COVID-19

pandemic are related to higher levels of depression and anxiety among Black, Indigenous and

Latinx young adults across geographic regions of the U.S.

Prior studies have indicated that employment is a protective factor against mental health

problems (43, 44). However, as indicated by the present study, the COVID-19 pandemic

represents a unique situation in which young adults of color who are employed face greater risk

for depression and anxiety. Our study suggests this is due not only to the infection risks

associated with work outside the home during this period, but to victimization by others based on

social perceptions that people of color are more likely to be infected with the virus. Past research

has also shown that sexual and gender minority status, medical conditions and financial

insecurity have direct impact on depression and anxiety among Black, Indigenous and Latinx

young adults (26, 27, 32-42, 48-56). Our results are consistent with past research indicating the

above factors are associated with internalizing disorders, and also demonstrates that during the

COVID-19 pandemic, prescription insecurity can contribute to mental health risk. Prior research

has shown an association between the pre-existing conditions now associated with increased

Coronavirus risk. Since the incidence of these pre-existing conditions are disproportionately high

among people of color, in particular obesity, diabetes, cardiovascular disease, and chronic illness

(6-14, 115-119), it is not surprising that our study demonstrated an association between the

number of Coronavirus health risks and mental health indices. Of note, is that 28% and 22% of

our participants identified as sexual or gender minorities, respectively. As both persons of color

and members of sexual and gender minority groups, the current COVID-19 crisis is likely

exacerbating mental health stressors due to lack of access to the medical care resulting from

multiple barriers such as medical mistrust, provider bias, and lack of insurance (2, 5, 22-24).

489 Coronavirus victimization distress and Coronavirus racial bias

This study provides evidence for the psychometric validity of two new scales: The

Coronavirus Victimization Distress Scale and the Coronavirus Racial Bias Scale. These

scales offered the opportunity to examine the relationship among Coronavirus specific

discrimination and perceived racial bias, demographic factors, and mental health. We

found that participants who were employed and reported prescription insecurity and more

COVID-19 health risks reported higher levels of Coronavirus victimization and

Coronavirus racial bias. These findings underscore the fact, that in addition to health and

economic disparities, Black, Indigenous and Latinx young adults are bearing a

disproportionate social burden during the current health crises.

The level of Coronavirus victimization distress and percentage of participants

reporting other risk factors did not differ across different racial/ethnic groups, with one

exception: Black and Latinx respondents reported higher levels of Coronavirus racial bias

than Indigenous young adults. This finding may be explained by news and social media

reports during April 2020 at the time this study was conducted. These reports identified

COVID-19 infections among Black and Latinx as the most vulnerable groups for

COVID-19 infection and death (6, 120), with outbreaks on tribal lands largely ignored by

the federal government, mainstream media and online tracking (121); although recent

studies suggest a relationship between household characteristics and infection rates on

reservations (122). Thus, at the time this survey was conducted, Indigenous participants

may have been less likely to have been exposed to public expressions of Coronavirus

related racial bias against their group.

Results of our main SEM analysis for the full sample indicate that Coronavirus

victimization distress was not directly associated with mental health, as the effect was fully

mediated by Coronavirus racial bias beliefs. This finding was strengthened by the results of the

alternative SEM model which did not find any mediation effect of Coronavirus victimization

distress on the relationship of Coronavirus racial bias to mental health indices. These findings

may be explained by prior work on attribution theory, suggesting that in certain instances

members of marginalized groups may attribute negative social experiences to racial

discrimination, which in turn has a harmful effect on mental health (73, 88). Thus, participants in

the present study may have attributed the bullying, threats, and unfair treatment described in the

Coronavirus Victimization Distress Scale items to perceived increases in racial biases created by

the pandemic.

Although the results of separate group SEM analysis must be viewed with caution given

the smaller sample size resulting in lower statistical power, they raise questions for future

research. Similar to the results of the main model, for Black and Indigenous participants,

Coronavirus racial bias continued to mediate the effects of Coronavirus victimization distress on

depression, although the effect was not significant for anxiety. By contrast, for Latinx

participants, Coronavirus victimization distress had significant direct effects on depression and

anxiety which was not mediated by perceived Coronavirus racial bias. One explanation for this

finding, which needs further investigation, is the statistically powerful effect that sexual minority

status had on mental health indices among the Latinx respondents. Thirty-three percent of Latinx

participants identified as sexual minorities in this sample. Thus, these young adults may have

been more likely to attribute Coronavirus victimization to social biases against sexual minorities

than to discrimination based on their ethnicity (123).

Limitations and future directions

To our knowledge, this is the first study to examine the relationship to mental health of

victimization distress and perceived racial bias directly related to the Coronavirus among Black,

Indigenous and Latinx young adults. However, due to the cross-sectional nature of the survey,

we are unable to interpret the results as definitive of mediation or causation. It will be important

for future studies to examine the longitudinal impact of Coronavirus victimization distress and

associated racial bias beliefs on mental health among young adults within these populations. In

addition, our data collection and recruitment methods yielded a geographically diverse national

sample of Black, Indigenous and Latinx young adults; however, this methodology does not allow

for absolute certainty that inclusion criteria were met and also limited participation those with

Internet or mobile phone access and who had previously registered for online surveys.

This study developed and psychometrically validated new measures of victimization and

racial bias associated with the current pandemic. This validation was limited to the sample

surveyed and future research will assess the extent to which it will be validated on other Black,

Indigenous and Latinx populations as well as other racial/ethnic groups in the U.S. For example,

although the infection and morbidity rate of Asian heritage groups in the U.S. has not met levels

seen among the populations included in this study, since April, reports of Coronavirus

victimization and racial bias against Asian groups has increased and is worthy of study (124).

Relatedly we found somewhat different patterns in the relationship of victimization distress and

racial bias to mental health indices among the 3 different groups. Studies with larger sample

sizes to achieve necessary power for each of these groups will help understand the extent to

which these differences will be replicated. Finally, our data also suggest, that for some

populations, minority stress associated with sexual orientation may have a greater influence on

interpretations of Coronavirus related victimization. It will be fruitful to examine if this is tied to

the legacy of HIV/AIDS discrimination among sexual minority young adults (125).

Conclusions

The mental health of people of color in the United States has been associated with

longstanding experiences of racial and ethnic discrimination and systemic bias. This study

highlights how the current COVID-19 pandemic has added to these mental health burdens

through increases in Coronavirus victimization and perceived racial bias experienced by Black,

Indigenous and Latinx young adults. These findings also demonstrate how an infectious disease

crisis can reverse the usual protective effects of employment on mental health, when working

people of color are more likely to be employed in settings that increase their exposure to

infection and subsequent social bias. In addition, this study underscores the deleterious effects on

mental health of Coronavirus health risks and associated prescription insecurity and the

continuing vulnerabilities associated with financial insecurity and sexual minority status. In

addition to ongoing efforts to increase multicultural mental health services, these findings

highlight the urgency of mental health services tailored to the specific needs of racial/ethnic

minorities during the current and future health crises.

573   References

1. CDC. Demographic Trends of COVID-19 cases and deaths in the US reported to CDC. 2020.

2. Nestor BA, Cheek SM, Liu RT. Ethnic and racial differences in mental health service

utilization for suicidal ideation and behavior in a nationally representative sample of adolescents.

Journal of affective disorders. 2016;202:197-202.

3. Manuel JI. Racial/Ethnic and Gender Disparities in Health Care Use and Access. Health

Services Research. 2018;53(3):1407-29.

4. Ben J, Cormack D, Harris R, Paradies Y. Racism and health service utilisation: A systematic

review and meta-analysis. PLOS ONE. 2017;12(12):e0189900.

5. Chen JA. Psychiatric Symptoms and Diagnoses Among U.S. College Students: A Comparison

by Race and Ethnicity. Psychiatric services (Washington, DC). 2019;70(6):442-9.

6. El Chaar M, King K, Galvez Lima A. Are black and Hispanic persons disproportionately

affected by COVID-19 because of higher obesity rates? Surgery for Obesity and Related

Diseases. 2020.

7. Cho P, Geiss LS, Burrows NR, Roberts DL, Bullock AK, Toedt ME. Diabetes-Related

Mortality Among American Indians and Alaska Natives, 1990–2009. American Journal of Public

Health. 2014;104(S3):S496-S503.

8. Gold DR, Wright R. Population disparities in asthma. Annu Rev Public Health. 2005;26:89-

113.

9. Peek ME, Cargill A, Huang ES. Diabetes health disparities. Medical Care Research and

Review. 2007;64(5_suppl):101S-56S.

10. Beavis AL, Gravitt PE, Rositch AF. Hysterectomy☐corrected cervical cancer mortality rates

reveal a larger racial disparity in the United States. Cancer. 2017;123(6):1044-50.

11.Bancks MP, Kershaw K, Carson AP, Gordon-Larsen P, Schreiner PJ, Carnethon MR.

Association of Modifiable Risk Factors in Young Adulthood With Racial Disparity in Incident

Type 2 Diabetes During Middle Adulthood. JAMA. 2017;318(24):2457.

12. Warne D, Lajimodiere D. American Indian health disparities: Psychosocial influences. Social

and Personality Psychology Compass. 2015;9(10):567-79.

13. Peng TY, Ehrlich SF, Crites Y, Kitzmiller JL, Kuzniewicz MW, Hedderson MM, et al.

Trends and racial and ethnic disparities in the prevalence of pregestational type 1 and type 2

diabetes in Northern California: 1996–2014. American Journal of Obstetrics and Gynecology.

2017;216(2):177.e1-.e8.

14. Flegal KM, Kruszon-Moran D, Carroll MD, Fryar CD, Ogden CL. Trends in Obesity Among

Adults in the United States, 2005 to 2014. JAMA. 2016;315(21):2284.

15.Rho HJ, Brown H, Fremstad S. A basic demographic profile of workers in frontline industries.

Center for Economic and Policy Research. 2020.

16. Bureau of Labor Statistics. Labor force characteristics by race and ethnicity, 2018. US

Bureau of Labor Statistics. 2019.

17. CDC. Groups at Higher Risk for Severe Illness 2020 [updated May 14, 2020. Available from:

https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/groups-at-higher-risk.html.

613 18. Longmire-Avital B. Young, Depressed, and Black: A Comparative Exploration of

614 Depressive Symptomatology Among Black and White Collegiate Women. Journal of college

615 student psychotherapy. 2018;32(1):53-72.

616 19. Baldwin JA, Eaves ER, Brown BG, Elwell K, Williamson HJ. The Behavioral Health of

617 American Indian/Alaska Native Populations: Risk and Resiliency. Springer International

618 Publishing; 2020. p. 205-30.

619 20. National Center for Health Statistics. Health, United States, 2015: with special feature on

620 racial and ethnic health disparities. Washington (DC): US Government Printing Office. 2016.

621 21. Shattell MM, Smith KM, Quinlan-Colwell A, Villalba JA. Factors Contributing to

622 Depression in Latinas of Mexican Origin Residing in the United States: Implications for Nurses.

623 2008;14(3):193-204.

624 22. Alegría M, Canino G, Shrout PE, Woo M, Duan N, Vila D, et al. Prevalence of mental illness

625 in immigrant and non-immigrant U.S. Latino groups. The American journal of psychiatry.

626 2008;165(3):359-69.

627 23. Vilsaint CL, NeMoyer A, Fillbrunn M, Sadikova E, Kessler RC, Sampson NA, et al.

628 Racial/ethnic differences in 12-month prevalence and persistence of mood, anxiety, and

629 substance use disorders: Variation by nativity and socioeconomic status. Comprehensive

630 psychiatry. 2019;89:52-60.

631 24. Shao Z, Richie WD, Bailey RK. Racial and Ethnic Disparity in Major Depressive Disorder.

632 Journal of Racial and Ethnic Health Disparities. 2016;3(4):692-705.

633 25. Williams DR, Gonzalez HM, Neighbors H, Nesse R, Abelson JM, Sweetman J, et al.

634 Prevalence and distribution of major depressive disorder in African Americans, Caribbean blacks,

and non-Hispanic whites: results from the National Survey of American Life. Archives of

general psychiatry. 2007;64(3):305-15.

26. Sherr L, Cluver L. World Health Day focus on HIV and depression-a comorbidity with

specific challenges. J Int AIDS Soc. 2017;20(1).

27. Schachter J, Martel J, Lin C-S, Chang C-J, Wu T-R, Lu C-C, et al. Effects of obesity on

depression: a role for inflammation and the gut microbiota. Brain, Behavior, and Immunity.

2018;69:1-8.

28. Assari S. Association between obesity and depression among American Blacks: role of

ethnicity and gender. Journal of Racial and Ethnic Health Disparities. 2014;1(1):36-44.

29. Pate CA, Zahran HS, Bailey CM. Impaired health-related quality of life and related risk

factors among US adults with asthma. Journal of Asthma. 2019;56(4):431-9.

30. Oh H, Stickley A, Singh F, Koyanagi A. Self-reported asthma diagnosis and mental health:

Findings from the Collaborative Psychiatric Epidemiology Surveys. Psychiatry research.

2019;271:721-5.

31. Schneider EC, Zaslavsky AM, Epstein AM. Racial disparities in the quality of care for

enrollees in Medicare managed care. Jama. 2002;287(10):1288-94.

32. Hoebel J, Maske UE, Zeeb H, Lampert T. Social inequalities and depressive symptoms in

adults: the role of objective and subjective socioeconomic status. PloS one. 2017;12(1):e0169764.

33. Arenas DJ, Thomas A, Wang J, DeLisser HM. A Systematic Review and Meta-analysis of

Depression, Anxiety, and Sleep Disorders in US Adults with Food Insecurity. J Gen Intern Med.

2019;34(12):2874-82.

34. Lemstra M, Neudorf C, D'Arcy C, Kunst A, Warren LM, Bennett NR. A Systematic Review

of Depressed Mood and Anxiety by SES in Youth Aged 10–15 Years. Canadian Journal of

Public Health. 2008;99(2):125-9.

35. Nagata JM, Palar K, Gooding HC, Garber AK, Whittle HJ, Bibbins-Domingo K, et al. Food

Insecurity Is Associated With Poorer Mental Health and Sleep Outcomes in Young Adults. J

Adolesc Health. 2019;65(6):805-11.

36. Weinstein N, Stone DN. Need depriving effects of financial insecurity: Implications for well-

being and financial behaviors. Journal of Experimental Psychology: General. 2018;147(10):1503.

37. Peplinski B, McClelland R, Szklo M. Associations between socioeconomic status markers

and depressive symptoms by race and gender: results from the Multi-Ethnic Study of

Atherosclerosis (MESA). Annals of Epidemiology. 2018;28(8):535-42.e1.

38. McCurley JL, Gutierrez AP, Bravin JI, Schneiderman N, Reina SA, Khambaty T, et al.

Association of social adversity with comorbid diabetes and depression symptoms in the Hispanic

Community Health Study/Study of Latinos Sociocultural Ancillary Study: a syndemic

framework. Annals of Behavioral Medicine. 2019;53(11):975-87.

39. Villegas E, Hannon B, Hammons A, Teran☐Garcia M, Wiley A. Is Food Security

Influencing Emotional Health? A Study with Hispanic Immigrant Mothers. The FASEB Journal.

2017;31:791.27-.27.

40. Walton QL, Boone C. Voices Unheard: An Intersectional Approach to Understanding

Depression among Middle-Class Black Women. Women & Therapy. 2019;42(3-4):301-19.

41. Assari S, Gibbons FX, Simons R. Depression among black youth; interaction of class and

place. Brain sciences. 2018;8(6):108.

42. Assari S. Social Determinants of Depression: The Intersections of Race, Gender, and

Socioeconomic Status. Brain Sciences. 2017;7(12):156.

43. McGee RE, Thompson NJ. Unemployment and depression among emerging adults in 12

states, Behavioral Risk Factor Surveillance System, 2010. Prev Chronic Dis. 2015;12:E38.

44. Paul KI, Moser K. Unemployment impairs mental health: Meta-analyses. Journal of

Vocational Behavior. 2009;74(3):264-82.

45. Bonanno GA, Brewin CR, Kaniasty K, Greca AML. Weighing the costs of disaster:

Consequences, risks, and resilience in individuals, families, and communities. Psychological

science in the public interest. 2010;11(1):1-49.

46. Solomon D, Maxwell C, Castro A. Systematic Inequality and Economic Opportunity.

Washington: Center for American Progress. 2019.

47. Mehdi M, Waseem M, Rehm MH, Aziz N, Anjum S, Javid MA. Depression and Anxiety in

Health Care Workers during COVID-19. Biomedica. 2020;36.

48. Krueger EA, Meyer IH, Upchurch DM. Sexual orientation group differences in perceived

stress and depressive symptoms among young adults in the United States. LGBT health.

2018;5(4):242-9.

49. Heterogeneity of transgender identity nonaffirmation microaggressions and their association

with depression symptoms and suicidality among transgender persons [press release]. US:

Educational Publishing Foundation2019.

50. Bostwick WB, Meyer I, Aranda F, Russell S, Hughes T, Birkett M, et al. Mental health and

suicidality among racially/ethnically diverse sexual minority youths. American journal of public

health. 2014;104(6):1129-36.

51. Vu M, Li J, Haardörfer R, Windle M, Berg CJ. Mental health and substance use among

women and men at the intersections of identities and experiences of discrimination: insights from

the intersectionality framework. BMC public health. 2019;19(1):108.

52. Rhodes SD, Martinez O, Song E-Y, Daniel J, Alonzo J, Eng E, et al. Depressive symptoms

among immigrant Latino sexual minorities. American Journal of Health Behavior.

2013;37(3):404-13.

53. Race/ethnicity and sexual orientation disparities in mental health, sexism, and social support

among women veterans [press release]. US: Educational Publishing Foundation2019.

54. Flanders CE, Shuler SA, Desnoyers SA, Vankim NA. Relationships Between Social Support,

Identity, Anxiety, and Depression Among Young Bisexual People of Color. Journal of

Bisexuality. 2019;19:2.

55. Jefferson K, Neilands TB, Sevelius J. Transgender women of color: discrimination and

depression symptoms. Ethn Inequal Health Soc Care. 2013;6(4):121-36.

56. Lefevor GT, Janis RA, Franklin A, Stone W-M. Distress and Therapeutic Outcomes Among

Transgender and Gender Nonconforming People of Color. The Counseling Psychologist.

2019;47(1):34-58.

57. Stein GL, Gonzalez LM, Huq N. Cultural stressors and the hopelessness model of depressive

symptoms in Latino adolescents. Journal of youth and adolescence. 2012;41(10):1339-49.

58. Benner AD, Wang Y, Shen Y, Boyle AE, Polk R, Cheng Y-P. Racial/ethnic discrimination

and well-being during adolescence: A meta-analytic review. American Psychologist.

2018;73(7):855.

59. Douglass S, Mirpuri S, English D, Yip T. "They were just making jokes": Ethnic/racial

teasing and discrimination among adolescents. Cultural diversity & ethnic minority psychology.

2016;22(1):69-82.

60. Lui PP. Racial Microaggression, Overt Discrimination, and Distress: (In)Direct Associations

With Psychological Adjustment. The Counseling Psychologist. 2020;48(4):551-82.

61. Forrest-Bank SS, Cuellar MJ. The Mediating Effects of Ethnic Identity on the Relationships

between Racial Microaggression and Psychological Well-Being. Social Work Research.

2018;42(1):44-56.

62. Barlow JN. Restoring Optimal Black Mental Health and Reversing Intergenerational Trauma

in an Era of Black Lives Matter. Biography. 2018;41(4):895-908.

63. Farisi AB, Habibi H, Mariska PN. Cultural Identity of Colonialism: Traumatic Effects of

Slavery and Racism. Cultura Interpreta. 2019;9(3):117-24.

64. Grills CN, Aird EG, Rowe D. Breathe, baby, breathe: Clearing the way for the emotional

emancipation of Black people. Cultural Studies? Critical Methodologies. 2016;16(3):333-43.

65. Gone JP, Hartmann WE, Pomerville A, Wendt DC, Klem SH, Burrage RL. The impact of

historical trauma on health outcomes for indigenous populations in the USA and Canada: A

systematic review. American Psychologist. 2019;74(1):20.

66. Documentation status and psychological distress among New York City community college

students [press release]. US: Educational Publishing Foundation2020.

67. Cobb CL, Meca A, Branscombe NR, Schwartz SJ, Xie D, Zea MC, et al. Perceived

discrimination and well-being among unauthorized Hispanic immigrants: The moderating role of

ethnic/racial group identity centrality. Cultural Diversity and Ethnic Minority Psychology.

2019;25(2):280-7.

68. Stafford AM, Bigatti SM, Draucker CB. Cultural stressors experienced by young Latinas

with depressive symptoms living in a tumultuous sociopolitical climate in the United States.

Archives of Psychiatric Nursing. 2019;33(5):36-42.

69. Cervantes RC, Cardoso JB, Goldbach JT. Examining differences in culturally based stress

among clinical and nonclinical Hispanic adolescents. Cultural Diversity and Ethnic Minority

Psychology. 2015;21(3):458-67.

70. Cariello AN, Perrin PB, Williams CD, Espinoza GA, Morlett-Paredes A, Moreno OA, et al.

Moderating influence of enculturation on the relations between minority stressors and physical

health via anxiety in Latinx immigrants. Cultural Diversity and Ethnic Minority Psychology.

2019.

71. Chu J, Maruyama B, Batchelder H, Goldblum P, Bongar B, Wickham RE. Cultural pathways

for suicidal ideation and behaviors. Cultural Diversity and Ethnic Minority Psychology.

2020;26(3):367-77.

72. Pascoe EA, Smart Richman L. Perceived discrimination and health: a meta-analytic review.

Psychological bulletin. 2009;135(4):531.

73. Schmitt MT, Branscombe NR, Postmes T, Garcia A. The consequences of perceived

discrimination for psychological well-being: a meta-analytic review. Psychological bulletin.

2014;140(4):921.

74. Priest N, Truong M, Chong S, Paradies Y, King T, Kavanagh A, et al. Experiences of racial

discrimination and cardiometabolic risk among Australian children. Brain, Behavior, and

Immunity. 2020.

75. Fisher CB, Wallace SA, Fenton RE. Discrimination distress during adolescence. Journal of

youth and adolescence. 2000;29(6):679-95.

76. Yip T. The effects of ethnic/racial discrimination and sleep quality on depressive symptoms

and self-esteem trajectories among diverse adolescents. Journal of youth and adolescence.

2015;44(2):419-30.

77. Clark R, Anderson NB, Clark VR, Williams DR. Racism as a stressor for African Americans:

A biopsychosocial model. American psychologist. 1999;54(10):805.

78. Nolen-Hoeksema S, Wisco BE, Lyubomirsky S. Rethinking rumination. Perspectives on

psychological science. 2008;3(5):400-24.

79. Adam EK, Heissel JA, Zeiders KH, Richeson JA, Ross EC, Ehrlich KB, et al. Developmental

histories of perceived racial discrimination and diurnal cortisol profiles in adulthood: A 20-year

prospective study. Psychoneuroendocrinology. 2015;62:279-91.

80. Brondolo E, Libby DJ, Denton EG, Thompson S, Beatty DL, Schwartz J, et al. Racism and

ambulatory blood pressure in a community sample. Psychosom Med. 2008;70(1):49-56.

81. Williams J, Gonzalez-Medina D. Infectious diseases and social stigma. Applied Innovations

and Technologies. 2011;4(1):58-70.

82. Person B, Sy F, Holton K, Govert B, Liang A, National Center for Inectious Diseases SCOT.

Fear and stigma: the epidemic within the SARS outbreak. Emerg Infect Dis. 2004;10(2):358-63.

83. Crockett KB, Kalichman SC, Kalichman MO, Cruess DG, Katner HP. Experiences of HIV-

related discrimination and consequences for internalised stigma, depression and alcohol use.

Psychology & health. 2019;34(7):796-810.

84. Baral SC, Karki DK, Newell JN. Causes of stigma and discrimination associated with

tuberculosis in Nepal: a qualitative study. BMC public health. 2007;7(1):1-10.

85. Lee S, Chan LY, Chau AM, Kwok KP, Kleinman A. The experience of SARS-related stigma

at Amoy Gardens. Social science & medicine. 2005;61(9):2038-46.

86. Siu JY-m. The SARS-associated stigma of SARS victims in the post-SARS era of Hong

Kong. Qualitative health research. 2008;18(6):729-38.

87. Goodwin R, Haque S, Neto F, Myers LB. Initial psychological responses to Influenza A,

H1N1 ("Swine flu"). BMC Infectious Diseases. 2009;9(1):166.

88. Major B, Quinton WJ, McCoy SK. Antecedents and consequences of attributions to

discrimination: Theoretical and empirical advances. 2002.

89. Major B, Kaiser CR, McCoy SK. It's not my fault: When and why attributions to prejudice

protect self-esteem. Personality and Social Psychology Bulletin. 2003;29(6):772-81.

90. Balsam KF, Beadnell B, Molina Y. The Daily Heterosexist Experiences Questionnaire:

Measuring minority stress among lesbian, gay, bisexual, and transgender adults. Measurement

and Evaluation in Counseling and Development. 2013;46(1):3-25.

91. Balsam KF, Molina Y, Beadnell B, Simoni J, Walters K. Measuring multiple minority stress:

the LGBT People of Color Microaggressions Scale. Cultural Diversity and Ethnic Minority

Psychology. 2011;17(2):163.

92. Spitzer RL. Validation and Utility of a Self-report Version of PRIME-MD: The PHQ

Primary Care Study. JAMA. 1999;282(18):1737.

93. Huang FY, Chung H, Kroenke K, Delucchi KL, Spitzer RL. Using the patient health

questionnaire-9 to measure depression among racially and ethnically diverse primary care

patients. Journal of general internal medicine. 2006;21(6):547-52.

94. Kroenke K, Spitzer RL, Williams JBW, Löwe B. The Patient Health Questionnaire Somatic,

Anxiety, and Depressive Symptom Scales: a systematic review. General Hospital Psychiatry.

2010;32(4):345-59.

95. Easton SD, Roh S, Kong J, Lee Y-S. Childhood Sexual Abuse and Depression among

American Indians in Adulthood. Health & social work. 2019;44(2):95-103.

96. Keum BT, Miller MJ, Inkelas KK. Testing the factor structure and measurement invariance

of the PHQ-9 across racially diverse US college students. Psychol Assess. 2018;30(8):1096.

97. Harry ML, Waring SC. The measurement invariance of the Patient Health Questionnaire-9

for American Indian adults. Journal of affective disorders. 2019;254:59-68.

98. Patel JS, Oh Y, Rand KL, Wu W, Cyders MA, Kroenke K, Stewart JC. Measurement

invariance of the patient health questionnaire-9 (PHQ-9) depression screener in US adults across

sex, race/ethnicity, and education level: NHANES 2005–2016. Depression and anxiety. 2019

Sep;36(9):813-23.

99. Spitzer RL, Kroenke K, Williams JB, Löwe B. A brief measure for assessing generalized

anxiety disorder: the GAD-7. Archives of internal medicine. 2006;166(10):1092-7.

100. Löwe B, Decker O, Müller S, Brähler E, Schellberg D, Herzog W, et al. Validation and

standardization of the Generalized Anxiety Disorder Screener (GAD-7) in the general population.

Medical care. 2008:266-74.

101. Dear BF, Titov N, Sunderland M, McMillan D, Anderson T, Lorian C, et al. Psychometric

comparison of the generalized anxiety disorder scale-7 and the Penn State Worry Questionnaire

for measuring response during treatment of generalised anxiety disorder. Cognitive behaviour

therapy. 2011;40(3):216-27.

102. Wurster HE, Sarche M, Trucksess C, Morse B, Biringen Z. Parents' adverse childhood

experiences and parent–child emotional availability in an American Indian community: Relations

with young children's social–emotional development. Development and Psychopathology.

2019:1-12.

103. Saunders HE. Adverse childhood experiences, stress, and emotional availability: an

American Indian context: Colorado State University. Libraries; 2016.

104. Parkerson HA, Thibodeau MA, Brandt CP, Zvolensky MJ, Asmundson GJG. Cultural-

based biases of the GAD-7. Journal of Anxiety Disorders. 2015;31:38-42.

105. Satorra A, Bentler PM. A scaled difference chi-square test statistic for moment structure

analysis. Psychometrika. 2001;66(4):507-14.

106. R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R

Foundation for Statistical Computing; 2020.

107. Rosseel Y, Oberski D, Byrnes J, Vanbrabant L, Savalei V, Merkle E, et al. Package 'lavaan'.

2017. p. 2017.

108. Hu Lt, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis:

Conventional criteria versus new alternatives. Structural equation modeling: a multidisciplinary

journal. 1999;6(1):1-55.

109. MacKinnon DP, Lockwood CM, Williams J. Confidence limits for the indirect effect:

Distribution of the product and resampling methods. Multivariate behavioral research.

2004;39(1):99-128.

110. HHS. HHS POVERTY GUIDELINES FOR 2020 2020 [Available from:

https://aspe.hhs.gov/poverty-guidelines.

111. MacCallum R. Specification searches in covariance structure modeling. 1986;100:107-20.

112. Little TD, Cunningham WA, Shahar G, Widaman KF. To parcel or not to parcel: Exploring

the question, weighing the merits. Structural equation modeling. 2002;9(2):151-73.

113. Cheung GW, Rensvold RB. Evaluating goodness-of-fit indexes for testing measurement

invariance. Structural equation modeling. 2002;9(2):233-55.

114. MacCallum RC, Browne MW, Sugawara HM. Power analysis and determination of sample

size for covariance structure modeling. Psychological methods. 1996;1(2):130.

115. Raifman MA, Raifman JR. Disparities in the Population at Risk of Severe Illness From

COVID-19 by Race/Ethnicity and Income. American journal of preventive medicine.

2020;59(1):137-9.

116. CDC. Summary Health Statistics: National Health Interview Survey: 2018. Table A-15a

2020 [cited 2020 August 10]. Available from:

https://ftp.cdc.gov/pub/Health_Statistics/NCHS/NHIS/SHS/2018_SHS_Table_A-15.pdf.

117. CDC. National diabetes statistics report, 2020. Atlanta, GA: Centers for Disease Control

and Prevention, US Department of Health and Human Services. 2020.

118. CDC. HIV surveillance report, 2018 (preliminary). Atlanta: CDC. 2019.

119. Twice M. Racial and Ethnic Disparities in Heart Disease 2019 [Available from:

https://www.cdc.gov/nchs/hus/spotlight/HeartDiseaseSpotlight_2019_0404.pdf.

120. Owen WF, Carmona R, Pomeroy C. Failing another national stress test on health disparities.

Jama. 2020.

121. Akee R. How COVID-19 is impacting indigenous peoples in the U.S. PBS News Hour.

2020.

122. Rodriguez-Lonebear D, Barceló NE, Akee R, Carroll SR. American Indian Reservations

and COVID-19: Correlates of Early Infection Rates in the Pandemic. Journal of Public Health

Management and Practice. 2020;26(4):371-7.

123. Cohen JM, Blasey C, Barr Taylor C, Weiss BJ, Newman MG. Anxiety and Related

Disorders and Concealment in Sexual Minority Young Adults. Behavior Therapy.

2016;47(1):91-101.

124. Ruiz NG, Horowitz JM, Tamir C. Many Black and Asian Americans Say They Have

Experienced Discrimination Amid the COVID-19 Outbreak 2020 [updated July 1, 2020; cited

2020 August 12]. Available from: https://www.pewsocialtrends.org/2020/07/01/many-black-and-

asian-americans-say-they-have-experienced-discrimination-amid-the-covid-19-outbreak/.

125. Fitzsimons T. LGBTQ History Month: The early days of America's AIDS crisis 2018

[updated Oct. 15, 2018; cited 2020 August 14]. Available from:

https://www.nbcnews.com/feature/nbc-out/lgbtq-history-month-early-days-america-s-aids-crisis-

n919701.