Full Title

Effects of an urban sanitation intervention on childhood enteric infection and diarrhea in Maputo,

Mozambique: a controlled before-and-after trial

Author list

Jackie Knee, PhD;[1,2] Trent Sumner, BS;[2] Zaida Adriano, BA;[3] Claire Anderson, BS;[2] Farran

Bush, BS;[4] Drew Capone, PhD;[2,6] Veronica Casmo, MSc;[5] David Holcomb, PhD;[6,7] Pete Kolsky,

PhD;[6] Amy Macdougall, PhD;[8] Evgeniya Molotkova, BS;[9] Judite Monteiro Braga, MSc;[5] Celina

Russo, BS;[2] Wolf Peter Schmidt, PhD;[1] Jill Stewart, PhD;[6] Winnie Zambrana, BS;[2] Valentina

Zuin, PhD;[10] Rassul Nalá, MPH;[5] Oliver Cumming, MSc;[1] Joe Brown, PhD[6,2]

[1] London School of Hygiene & Tropical Medicine, Faculty of Infectious Tropical Diseases,
Disease Control Department, Keppel Street, Bloomsbury, London WC1E 7HT, United Kingdom

[2] Georgia Institute of Technology, School of Civil and Environmental Engineering, 311 Ferst
Drive, Atlanta, GA, USA 30332, USA

[3] WE Consult ltd, Rua Tomás Ribeiro 177, RC, Postal code 3248, Maputo, Moçambique

[4] Georgia Institute of Technology, School of Chemical and Biomolecular Engineering, 311 Ferst
Drive, Atlanta, GA 30332, USA

[5] Instituto Nacional de Saúde, Estrada Nacional N°1, Bairro da Vila, Distrito de Marracuene,
Província de Maputo, Moçambique

[6] University of North Carolina at Chapel Hill, Gillings School of Global Public Health,
Department of Environmental Sciences and Engineering, 135 Dauer Drive, Chapel Hill, NC,
27599, USA

[7] University of North Carolina at Chapel Hill, Gillings School of Global Public Health,
Department of Epidemiology, 135 Dauer Drive, Chapel Hill, NC, 27599, USA

[8] London School of Hygiene & Tropical Medicine, Faculty of Epidemiology and Population
Health, Department of Medical Statistics, Keppel Street, Bloomsbury, London WC1E 7HT,
United Kingdom

[9] Georgia Institute of Technology, School of Biological Sciences, 310 Ferst Drive NW, Atlanta,
GA 30332, USA

[10] Yale-NUS College, Division of Social Science, 28 College Ave West, 138527, Singapore

Correspondence to:

Joe Brown (0000-0002-5200-4148)
University of North Carolina at Chapel Hill, Gillings School of Global Public Health,
Department of Environmental Sciences and Engineering, 135 Dauer Drive, Chapel Hill, NC,
27599, USA, Email: Joebrown@unc.edu Phone: +1 919 966 7892

Abstract

We conducted a controlled before-and-after trial to evaluate the impact of an onsite urban

sanitation intervention on the prevalence of enteric infection, soil transmitted helminth re-

infection, and diarrhea among children in Maputo, Mozambique. A non-governmental

organization replaced existing poor-quality latrines with pour-flush toilets with septic tanks

serving household clusters. We enrolled children aged 1-48 months at baseline and measured

outcomes before and 12 and 24 months after the intervention, with concurrent measurement

among children in a comparable control arm. Despite nearly exclusive use, we found no evidence

that intervention affected the prevalence of any measured outcome after 12 or 24 months of

exposure. Among children born into study sites after intervention, we observed a reduced

prevalence of *Trichuris* and *Shigella* infection relative to the same age group at baseline (<2

years old). Protection from birth may be important to reduce exposure to and infection with

enteric pathogens in this setting.

Introduction

Rapid urbanization has led to the expansion of informal settlements in many low- and middle-

income countries (LMICs). Such settlements often have very limited sanitation infrastructure

(UN-Habitat, 2016). Separation of human waste from human contact can prevent exposure to

enteric pathogens that cause infection, diarrhea (Liu et al., 2016), and potentially long-term

health effects such as environmental enteric dysfunction (EED) (Kosek et al., 2017), linear

growth deficits (Rogawski et al., 2018), impaired cognitive development (MAL-ED Network

Investigators, 2018), and reduced oral vaccine immunogenicity (Parker et al., 2018). Children

living in densely populated slum areas where fecal contamination is pervasive and sanitation

infrastructure is limited may be at an increased risk of adverse health effects due to frequent

exposure to enteric pathogens (Ezeh et al., 2017; Fink, Günther, & Hill, 2014).

Household-level sewerage has demonstrated health benefits (Barreto et al., 2010, 2007; Norman,

Pedley, & Takkouche, 2010) and remains an important long-term goal for many urban settings

despite limited evidence from controlled trials (Norman et al., 2010; Wolf et al., 2018). Such

systems may not be feasible short-term solutions due to cost, space, and logistical constraints,

challenges that have also impeded their evaluation via randomized trials (Norman et al., 2010).

Further, in densely populated areas, there may not be space for household-level sanitation of any

type. Shared sanitation is a subject of considerable debate but may represent the only near-term

sanitation option in some settings (Evans et al., 2017; Heijnen et al., 2014; Tidwell et al., 2020).

Yet, while shared, onsite systems may fill the growing need for safe sanitation in rapidly

expanding urban areas in LMICs, to date, there has been little evidence of their health impacts in

these settings. Recent large-scale, rigorous evaluations of onsite sanitation interventions and

combined water, sanitation, and hygiene interventions have demonstrated mixed effects on

health (Clasen et al., 2014; Humphrey et al., 2019; Luby et al., 2018; Null et al., 2018; Patil et

al., 2014; Pickering, Djebbari, Lopez, Coulibaly, & Alzua, 2015) but all were conducted in rural

areas with household-level interventions, and their findings may have limited generalizability to

urban areas. A recent meta-analysis estimated that non-sewered interventions reduced the risk of

self-reported diarrhea by 16% but did not estimate effects on objective health outcomes, such as

enteric infection (Brown & Cumming, 2019), and could not stratify estimates by rural versus

urban setting given the lack of evidence in urban areas (Wolf et al., 2018). To-date, no controlled

trials of urban onsite sanitation have been conducted despite over 740 million urban residents

relying on such technologies (Berendes, Sumner, & Brown, 2017).

The Maputo Sanitation (MapSan) trial was the first controlled trial to evaluate an onsite, shared

sanitation intervention in an urban setting and the first to use the prevalence of enteric infection,

as detected by molecular methods, as the primary study outcome (Brown et al., 2015). The study

was located in densely populated, low-income, informal neighborhoods of Maputo, Mozambique

where the sanitary conditions are poor and disease burden high (Knee et al., 2018). As of 2017,

only half of urban residents in Mozambique had access to at least basic sanitation infrastructure,

3% had access to sewerage, and 9% shared sanitation with multiple households, often in poor

neighborhoods where space and resources are limited (UNICEF/WHO, 2019). We investigated

whether an engineered, onsite, shared sanitation intervention could reduce enteric infection and

diarrhea in young children living in these low-income, densely populated neighborhoods in

Maputo, Mozambique.

Results

The MapSan trial was a controlled before-and-after trial designed to evaluate the impact of an

onsite sanitation intervention on child health after 12 and 24 months of follow-up. The

intervention consisted of pour-flush toilets to septic tanks with soakaway pits to discharge the

liquid portion of the waste. A non-governmental organization (NGO) delivered the intervention

to clusters of households known as compounds, replacing the existing poor-condition shared

facilities. Control compounds did not receive the intervention and continued to use their poor-

condition sanitation for the duration of the study. We assessed several measures of child health,

including enteric infection measured via stool-based molecular methods, soil-transmitted

helminth (STH) re-infection measured via Kato-Katz, and diarrhea measured via caregiver report

in both intervention and control children during three phases: baseline (pre-intervention), 12-

month follow-up, and 24-month follow-up. Children were eligible for baseline enrollment if they

were less than four years old (1-48 months old). At follow-up, children were eligible for

enrollment if they were less than four years old or if they would have been less than four years

old during baseline.

We enrolled 987 children in 495 compounds during the baseline phase (February 2015 –

February 2016) and collected stool samples (whole stool or diaper samples containing liquid

diarrhea) from 765 children (78%) (Figure 1). During the 12-month follow-up phase (March

2016 – April 2017), we enrolled or revisited 939 children in 438 compounds and collected 805

stool samples (86%). During the 24-month follow-up phase (April 2017 – August 2018), we

enrolled or revisited 1001 children in 408 compounds and collected stool samples from 922

(90%). To improve the success rate of stool sample collection during the 12- and 24-month

follow-up visits, we collected rectal swabs from children who did not provide a whole stool

sample after multiple collection attempts. The proportion of each type of sample (whole stool,

diaper sample, and rectal swab) was similar between arms at each phase (Supplemental Figure

1). Fewer than 5% of all samples were diapers and approximately 7% of 12-month samples and

25% of 24-month samples were rectal swabs (Supplemental Table 1). The NGO delivered

interventions to 15 control compounds after baseline and children in those compounds were

censored at the time of intervention receipt (Figure 1). Children living in control compounds that

independently upgraded their latrines were included in the main analyses. However, as inclusion

of these control children may have diluted the intervention effect, they were excluded from

sensitivity analyses designed to understand the impact of the intervention when compared with

controls with poor-condition sanitation throughout the study. Children in intervention and control

compounds were enrolled at similar rates during each phase (Supplemental Figure 2). Due to

migration out of the compound, we collected longitudinal data from 62% of children (59%

controls, 67% interventions) between baseline and 12-month and 51% of children (46% controls,

58% interventions) between baseline and 24-month.

At baseline enrollment, intervention compounds had more residents, households, and on-premise

water taps than controls, though the number of shared latrines was similar (Table 1). Animals

were observed in over half of all compounds. Intervention and control households had similar

wealth scores, though intervention households had more members and were more crowded while

control households more often had walls made of sturdy materials. All households used a

municipal water tap as their primary drinking water source with 78% reporting use of a tap on

the compound grounds. At baseline, latrines used by intervention households more often had

pedestals or slabs, drop-hole covers, and sturdy walls compared with controls. Consistent with

previous estimates in urban Maputo (Satterthwaite, Beard, Mitlin, & Du, 2019), open defecation

was rare in our study population with only one control household reporting open defecation at

baseline. Baseline characteristics of intervention and control children were similar: the average

age at enrollment was 23 months (SD = 13), 51% were female, and 32% were still breastfeeding

(Table 1). The age distributions of intervention and control children were similar at baseline and

both follow-up phases (Supplemental Figure 3).

We used the Luminex Gastrointestinal Pathogen Panel (GPP), a qualitative multiplex molecular

assay, to simultaneously test for 15 enteric pathogens in stool samples, including nine bacteria,

three protozoa, and three viruses. We detected ≥1 bacterial or protozoan enteric infection, our

pre-defined primary outcome, in 78% (591/753) of children with stools available at baseline. We

measured our pre-defined secondary outcome, ≥1 STH re-infection, using the Kato-Katz

microscope method and detected ≥1 STH in 45% (308/698) of stools at baseline. The

prevalences of pre-defined outcomes, individual pathogens, and pathogen types were similar

between the intervention and control arms at baseline (Table 2). The prevalence of most

bacterial, protozoan, and STH infections increased with age while the prevalence of enteric

viruses decreased with age (Supplemental Table 2 and Supplemental Figure 4).

The characteristics of children with repeated observations (including baseline) were similar to

characteristics of children measured at baseline only and to characteristics of children measured

at 12-month and/or 24-month only with the exception of age-related characteristics

(Supplemental Tables 3-6). Over half of the children enrolled after baseline were born into study

sites (336/622 [54%], Figure 1).

Our main analyses included observations from all eligible children enrolled at baseline (mean

sampling age 664 days, SD=393) and the 12-month (940 days, SD=498) and 24-month (1137

days, SD=603) follow-up visits (Table 2). We used a difference-in-difference (DID) analysis to

estimate the intervention effect and adjust for baseline differences between intervention and

control compounds. We present effect estimates from the DID analyses as prevalence ratios

(ratio of ratios). To assess the validity of the parallel trend assumption, a key assumption of DID

analyses, we ran "placebo tests" by replacing outcomes with variables unrelated to the

intervention, such as child age, respondent role, and presence of animals. Placebo tests showed

no effect of the intervention on these variables, suggesting the parallel trend assumption was

valid. We found no evidence that the intervention had an effect on the prevalence of any

bacterial or protozoan infection (adjusted PR 1.04, 95% CI [0.94 – 1.15]), or any STH re-

infection (1.11 [0.89 – 1.38]) 12 months after implementation (Table 2) despite household

respondents reporting almost exclusive use of the intervention latrine (97%, 404/417). The

prevalence of diarrhea remained fairly constant in both arms in all three phases with the

exception of the 12-month measure in the control arm which was lower, resulting in a larger

effect estimate with low precision (1.69 [0.89-3.21]).

The intervention had no meaningful effect at 12 months on the prevalence of infection with any

of the three pathogen types measured by the GPP (bacterial, protozoan, viral), pathogen

coinfection, or on any individual pathogen (Table 2). There was poor precision in the effect

estimates for infrequently detected pathogens, evident from their wide confidence intervals.

Therefore, some estimates suggestive of a large protective or detrimental effect (*Campylobacter*,

C. difficile, *E. coli* O157, STEC, Norovirus GI/GII, Adenovirus 40/41) may have arisen by

chance. While the National Deworming Campaign (NDC) provided albendazole to all compound

members following baseline, during 12-month visitation only 58% of caregivers (56% control,

60% intervention) confirmed that their child was dewormed during these visits. A sensitivity

analysis restricted to children confirmed to have been dewormed produced similar results to the

main analysis (Supplemental Table 7). By the 12-month visit, 19 control compounds (19/240

[8.0%]) had independently upgraded their facilities to pour-flush toilets. Results from sensitivity

analyses excluding children living in control compounds with independently upgraded facilities

were consistent with the main results (Supplemental Table 8).

There was no evidence that the intervention had an effect on the prevalence of any bacterial or

protozoan infection, any STH re-infection, or diarrhea after 24 months among all enrolled

children (Table 2). We also found limited evidence of effect on the prevalence of any pathogen

type or coinfection with ≥2 GPP pathogens 24 months after intervention. Results for several

individual outcomes were suggestive of a protective (STEC, *E. coli* O157, *Cryptosporidium*,

STH coinfection) or adverse (*Campylobacter*, *C. difficile*) effect, but evidence was weak as

estimates were accompanied by wide confidence intervals and chance discoveries were possible

given multiple comparisons. At the 24 month visits, caregivers confirmed baseline and/or 12-

month deworming more frequently for intervention children (339/502 [68%]) than for control

children (286/499 [57%]). Adjustment for deworming status or time since deworming had no

impact on effect estimates (Supplemental Table 7). Excluding children from control compounds

which independently upgraded their facilities by the 24-month visit (35/211 compounds, [17%])

did not impact the results (Supplemental Table 8).

Point estimates of effect and associated confidence intervals were largely similar in unadjusted

and adjusted models with few exceptions (e.g. ETEC at 24-month) (Table 2). Multivariable

models for GPP outcomes and STH outcomes were adjusted for covariates selected *a priori*

(child age, sex, caregiver education, and household wealth index). No other variables met our

inclusion criteria for multivariable models, which included being imbalanced between

intervention and control at baseline and meaningfully changing 12-month effect estimates (>10%

change in prevalence ratios) (Supplemental Table 9). While the relationship between age and

pathogen prevalence appeared to be non-linear for many pathogens (Supplemental Figure 4), the

inclusion of a higher order age term (age squared) did not meaningfully change effect estimates

in the main or sub-group analyses (Supplemental Table 10). Three measures of seasonality were

considered for inclusion in multivariable models to adjust for any difference in seasonal

distributions of data collection: (1) a binary variable defining the 'rainy' (November – April) and

'dry' seasons (May – October) in Maputo, (2) a measure of cumulative rainfall (mm) in the 30

days prior to data collection, and (3) sine and cosine terms representing dates of sample

collection. While there was some imbalance between arms in data collected during the wet and

dry seasons at baseline (Supplemental Table 9), no measure of seasonality meaningfully changed

effect estimates in the 12- and 24-month analyses and seasonality was excluded from final

multivariable models (Supplemental Table 9 and 11). For diarrhea, two variables in addition to

variables selected *a priori* met our inclusion criteria and were included in adjusted models:

presence of a latrine drop-hole cover at baseline and reported use of a water tap located within

the compound grounds at baseline (Supplemental Table 9). The magnitude of effect estimates

were larger and confidence intervals wider for diarrhea in adjusted versus unadjusted models in

the 12-month and 24-month analyses (Table 2). In addition to the main analyses which included

all enrolled children, we also performed two sub-group analyses. The first included children who

were born after the intervention was implemented (or after baseline in control compounds) and

present at the 12- and/or 24-month follow-up visit. This analysis allowed us to evaluate the

impact of the intervention on young children who were never exposed to the poor sanitation at

baseline. The second sub-group analysis included only children with repeated measures at

baseline and 12- and/or 24-month follow-up.

In sub-group analyses comparing children born into study compounds before the 24-month visit

with children of similar ages at baseline (<2 years old), there was suggestive evidence that the

intervention reduced the prevalence of infection with any STH by half (n=522; adjusted

prevalence ratio 0.51, [95% CI 0.27 - 0.95]), *Trichuris* by 76% (n=522; 0.24, [0.10 - 0.60]), and

Shigella by 51% (n=630; 0.49, [0.28 - 0.85]) (Table 3). These effects were attenuated in sub-

group analyses restricted to older children (>24 months) who were born before the intervention

was implemented and present at the 24-month phase (Supplemental Table 12). We did not

observe intervention effects among children born into the study by the 12-month visit, but the

sample size was small, resulting in high uncertainty in effect estimates (Supplemental Table 13).

Longitudinal sub-group analyses explored the effect of the intervention on children with repeated

measures at baseline and 12-month (for unadjusted analyses: n=870 data points [435 children

with repeat measures] for GPP outcomes, n=572 [286] for Kato-Katz outcomes, and n=1112

[556] for diarrhea) and at baseline and 24-month (n=716 (358), n=402 (201), n=834 (417)).

Effect estimates were consistent with results from the main analyses (Supplemental Tables 14-

15) but less precise due to the reduced sample numbers.

Discussion

We found no evidence that this urban, onsite shared sanitation intervention was protective against our pre-specified child health outcomes of enteric infection, STH re-infection, or diarrhea. We also found no strong evidence that the intervention affected prevalence of any individual pathogen, pathogen type, or coinfection with ≥2 enteric pathogens or STH. In exploratory sub-group analyses, we found suggestive evidence that the intervention reduced the prevalence of any STH, *Trichuris,* and *Shigella* infections among children born into the study by the 24-month follow-up visit. Studying children born into intervention sites after implementation allowed us to examine the effect of the intervention from birth through the first two years of life. These results suggest that the intervention delayed pathogen exposure and the accumulation of enteric infections during early childhood, but it needs to be treated with caution as this was an exploratory subgroup analysis.

The trial was neither designed nor powered to detect differences in sub-groups of children such as those born after the intervention was implemented, potentially limiting our ability to detect small effects in such analyses. Further, all exploratory sub-group analyses included multiple comparisons, increasing the likelihood of chance discoveries. However, the magnitude of the effect estimates for the outcomes of any STH, *Trichuris*, and *Shigella* observed among children born into the study by the 24-month visit, and the directional consistency of effect estimates among most other outcomes in this sub-group analysis, strengthens the plausibility of these findings.

There are several reasons we observed suggestive evidence of an effect for some outcomes among this sub-group of young children but not among older children or in the main analyses.

Children's exposures vary by age, particularly as they become mobile and begin independent

exploration of their environment. It is possible that the intervention reduced exposure via

pathways that are important for very young children but may represent just minor pathways of

exposure among older children (Kwong et al., 2020) Additionally, young children may

experience fewer exposures outside of the compound. Reductions in exposure and subsequent

infection early in life may delay or prevent the development of environmental enteric

dysfunction (EED), a subclinical condition that affects the structure and function of the gut and

may increase susceptibility to future infection (Keusch et al., 2014; Prendergast & Kelly, 2016).

Results from the EED sub-study of the WASH Benefits cluster randomized controlled trial

(cRCT) in Bangladesh suggest that the intervention delayed but did not prevent the onset of EED

(Lin et al., 2019). If this intervention similarly delayed the development of EED among children

born into intervention sites, they may have been less susceptible to infection than children of a

similar age at baseline. Finally, some pathogens, like *Giardia* and certain STH, can cause

persistent infections that can remain active for months or years if not treated (Else et al., 2020;

Rogawski et al., 2017). The intervention would have no effect on such infections, highlighting

the potentially important role of protection from birth.

Notably, both *Shigella* and *Trichuris* are primarily anthroponotic, and infection was strongly age-

dependent in this study population (Knee et al., 2018). These factors may help explain the

differing intervention effects observed both among pathogens and age groups. The intervention

was unlikely to limit exposure to animal feces, reducing the likelihood that it would impact

infection prevalence of zoonotic pathogens like *Campylobacter* or *Giardia*. The strong positive

associations between age and prevalence for *Shigella* and *Trichuris* suggest that exposure

increases with age. This supports the hypothesis that the intervention may have reduced the

overall frequency or intensity of exposure enough to impact *Shigella* and *Trichuris* infection

among young children but not older children.

Rapid urbanization is expanding informal settlements and out-pacing the expansion of sanitation

services in many cities, widening the gap in sanitation access between the urban rich and poor

(UNICEF/WHO, 2019). To our knowledge, MapSan was the first trial to estimate the health

impact of an urban, onsite shared sanitation intervention and the first to use enteric infection as

the primary trial outcome. Most of the urban sanitation literature published to date has evaluated

the expansion of sewerage, an important and ambitious goal that is out of reach for many cities in

the near-term (Norman et al., 2010). Access to sewerage is associated with a 30-60% reduction

of diarrheal disease depending on starting conditions, and an approximately 30% reduction in

enteric parasite detection, though most studies are observational and few controlled trials exist

(Barreto et al., 2010; Norman et al., 2010; Wolf et al., 2018).

Most studies of onsite sanitation interventions have occurred in rural areas. Despite good

evidence that onsite sanitation is associated with reductions in diarrheal disease (M. C. Freeman

et al., 2017; Wolf et al., 2018), several recent rural trials of basic sanitation and combined

WASH interventions with good uptake and use reported mixed effects on child health outcomes

including diarrhea, linear growth, and more recently, enteric infection (Ercumen et al., 2019;

Grembi et al., 2020; Humphrey et al., 2019; Lin et al., 2018; Luby et al., 2018; Null et al., 2018;

Pickering et al., 2019; Rogawski McQuade, Platts-Mills, et al., 2020).

The Sanitation, Hygiene, Infant Nutrition Efficacy (SHINE) trial in rural Zimbabwe found no

impact of a combined WASH intervention on diarrhea, growth, or the prevalence of a suite of

enteric pathogens among children aged <12 months old but did report a small reduction in the

number of parasitic pathogens detected.(Humphrey et al., 2019; Rogawski McQuade, Platts-

Mills, et al., 2020)

While the WASH Benefits Bangladesh cRCT reported no effect of any WASH intervention on

child growth, the sanitation, hygiene, and combined WASH study arms reduced the prevalence

of diarrheal disease from 5.7% to 3.5% (Luby et al., 2018), accompanied by absolute reductions

in *Giardia* prevalence of 6-9% among children aged 2-3 years in the same arms (Lin et al.,

2018). The sanitation arm also reduced the prevalence of *T. trichiura* among children 2-3 years

old (from 5.2% to 3.2%) but had no impact on *A. lumbricoides* or hookworm, the only other

parasites detected frequently enough to estimate effects in that study (Ercumen et al., 2019). In a

parallel analysis, only the water treatment and combined WASH interventions of the WASH

Benefits Kenya cRCT reduced the prevalence *A. lumbricoides*, suggesting that the reduction in

prevalence in the combined WASH arm may be attributable to the water treatment intervention

(Pickering et al., 2019). The sanitation-only arm had no impact on any parasite measured, though

T. trichiura was too infrequently detected to estimate effects (Pickering et al., 2019). An

evaluation of a comprehensive suite of 34 enteric pathogens reported reduced prevalence and

quantity of enteric viruses, but not bacteria or parasites, among children aged 14 months old in

the combined WASH arms in the Bangladesh trial (Grembi et al., 2020). Together with our

findings, these results suggest that sanitation and combined WASH interventions can reduce the

prevalence of enteric infection in some settings but that effects may vary by pathogen, child age,

intervention, and setting.

We previously published two baseline risk factor analyses to identify demographic,

environmental, and WASH-related predictors of infection and environmental fecal contamination

in our study setting prior to the intervention implementation (Holcomb et al., 2020; Knee et al.,

2018). Age was an important predictor of infection, though the direction of its effect varied by

pathogen type. Increasing age was associated with increased risk of bacterial and protozoan

infections and decreased risk of viral infections (Knee et al., 2018). Other socio-demographic

predictors of infection included breastfeeding, which was associated with a decreased risk of any

infection (driven by its strong association with protozoan infection), and female sex which was

associated with an increased risk of viral infection. Few sanitation-related or environmental

variables were associated with infection at baseline and the magnitude of associations were often

small. The presence of a latrine superstructure and drop-hole cover were associated with small

reductions in risk of bacterial or protozoan infection, often only in unadjusted analyses, but other

latrine features (e.g. presence of a cleanable slab) were not. The observation of feces or used

diapers around the compound grounds was associated with increased risk of bacterial and

protozoan infection but most other environmental and sanitary hazards were not (Knee et al.,

2018).

Fecal contamination was common among all environmental reservoirs tested (water, soil, food

preparation surfaces) at baseline. We detected one or more microbial markers of contamination

in over 95% of environmental samples (Holcomb et al., 2020). *E. coli* was the most frequently

detected and abundant marker of contamination among all sample types, and human-associated

markers were most frequently detected in soil (59%) and stored drinking water (17%) samples.

Measures of latrine quality that were associated with small reductions in infection risk (e.g. drop-

hole covers, latrine superstructures) were not associated with decreased odds of fecal

contamination in this setting. Overall, we found few consistent relationships between markers of

fecal contamination and environmental, WASH-related, and demographic characteristics at

baseline (Holcomb et al., 2020).

While these results suggest WASH-related and environmental risk factors may be poor

determinants of child health in this setting, the lack of heterogeneity in WASH conditions at

baseline, given the selection criterion that compounds must share sanitation in "poor condition,"

may have limited our ability to identify strong WASH-related predictors of infection or

environmental fecal contamination. Results from a forthcoming companion study suggests the

intervention had mixed effects on environmental fecal contamination. The intervention may have

reduced the concentration of *E. coli* by an order of magnitude in soil collected from latrine

entrances after 12 months, however, there was no effect on the prevalence or concentration of

indicators of fecal contamination in any other environmental compartment sampled at that time

(Holcomb et al., 2021). It is unlikely that the observed reductions in fecal contamination in soils

alone would be sufficient to impact health outcomes in this setting. Other studies that have

evaluated the impact of sanitation interventions on fecal contamination of the surrounding

environment have found limited evidence of effect (Clasen et al., 2014; Ercumen, Mertens, et al.,

2018; Ercumen, Pickering, et al., 2018; Fuhrmeister et al., 2020; Patil et al., 2014; Pickering et

al., 2015; Gloria D. Sclar et al., 2016; Steinbaum et al., 2019).

In this setting, where fecal contamination was pervasive and burden of infection high, even

considerable reductions in contamination and exposure may have been insufficient to realize

measurable health gains as the intervention did not address all potential transmission pathways

(Briscoe, 1984; Julian, 2016; Robb et al., 2017). For example, the intervention did not address

child feces disposal practices or handwashing behaviors and it is unlikely that the intervention

infrastructure would have changed these (Majorin, Torondel, Chan, & Clasen, 2019). Previous

studies of sanitation interventions have found no reduction in hand contamination (Ercumen,

Pickering, et al., 2018), which has been associated with increased incident diarrheal disease in

young children (Pickering et al., 2018). The intervention may not have reduced exposure via

consumption of contaminated food – particularly foods contaminated prior to arrival in the

compound – likely an important source of enteric pathogen transmission in some settings (Julian,

2016; Kwong et al., 2020). Children's exposure to animal feces has been documented in rural,

peri-urban, and urban settings and could be an important, unmitigated source of exposure to

enteric pathogens in both intervention and control arms where animals were frequently observed

(Delahoy et al., 2018; Kwong et al., 2020; Penakalapati et al., 2017). Observation of animals in

compounds was examined as a potential confounder but did not change effect estimates.

The intervention was delivered at the compound level, not the community level, and was not

designed to achieve any specified threshold of sanitation coverage in the study neighborhoods.

Previous studies have suggested that achieving a certain level of community sanitation coverage

may be necessary to reduce disease burdens (Barreto et al., 2007; Fuller & Eisenberg, 2016;

Fuller, Villamor, Cevallos, Trostle, & Eisenberg, 2016; Harris, Alzua, Osbert, & Pickering,

2017; Jung, Lou, & Cheng, 2017; Spears, Ghosh, & Cumming, 2013; Wolf et al., 2018). For

example, a study of a large-scale sewerage expansion in urban Brazil found that the intervention

reduced diarrheal disease by 22%, with neighborhood coverage level being the single most

important explanatory variable (Barreto et al., 2007). We did not measure neighborhood-level

sanitation coverage, but previous estimates show that while coverage is high and open defecation

is limited (1%), only 9% of sanitation systems are safely managed (Satterthwaite et al., 2019).

Further, in the Nhlamankulu district where many of our study sites are located, the majority of

households (56%) rely on pit latrines serving individual households, most of which are in poor

condition (Devamani, Norman, & Schmidt, 2014; Satterthwaite et al., 2019). Together with our

results, this suggests that both the extent and quality of community coverage are likely important

to reducing overall transmission. Sanitation coverage and quality may be especially important in

urban areas given the proximity of compounds and the opportunity for person-to-person contact,

neighborhood-level exposure, and for external sources of contamination (e.g. a neighbor's

flooded pit latrine) to influence compound-level exposures (Barreto et al., 2007). We did not

measure neighborhood-level exposures, which may be important for young children in slum

settings (Ezeh et al., 2017; Medgyesi et al., 2019), and their impact on our health outcomes is

unclear. In addition to neighborhood-level exposures, the transience of the study population

meant that trips to and from provinces outside of Maputo, where exposures were varied and

unmeasured, were common.

It is unlikely that our findings are due to poor intervention fidelity or use, a challenge

encountered in some trials of rural sanitation interventions (Clasen et al., 2014; Patil et al., 2014).

The use of the intervention required minimal behavior change as compound members switched

from using their existing latrine in poor condition, which was removed following construction of

the intervention latrine, to using the new hygienic latrine. The results of a forthcoming process

evaluation demonstrate that 96% of intervention latrines were well-maintained two or more years

after construction, suggesting continued use by compound members (Bick et al., 2021). Further,

only 3% of intervention compounds (8/270) had a secondary, non-intervention latrine in use after

two or more years, indicating that members of most intervention compounds exclusively used the

intervention latrines   (Bick et al., 2021). It is possible that development in the study

neighborhoods, including changes to sanitation facilities in control compounds, contributed to

the limited effect of the intervention. However, results from sensitivity analyses that excluded

control compounds with upgraded sanitation were consistent with results from the main analyses.

The two intervention designs we evaluated in this study – communal sanitation blocks and

shared latrines – utilized the same basic sanitation technology but differed in the number of

cabins and amenities available. While it is possible that this heterogeneity in design may have

modified the effect of the intervention, this study was not powered to test this. Moreover, all

intervention compounds were encouraged to independently upgrade their facilities by adding

features like electricity and handwashing stations, or by connecting existing handwashing

stations to the water supply, resulting in heterogeneity even within the two broad categories of

intervention type.

While the NDC dewormed every study compound annually during the study period, it is possible

that not all study participants received, or took, the medication and that the time between

deworming and subsequent measurement of STH re-infection varied among children.

Additionally, single-dose albendazole can have limited effectiveness against certain STH,

notably *Trichuris* (Moser, Schindler, & Keiser, 2017). Inadequate or ineffective deworming

could have limited our ability to detect an effect on STH outcomes. Sensitivity analyses

adjusting for caregiver-confirmed deworming and for estimated time between deworming and re-

infection measurement produced similar results to the main analysis.

There are several important limitations of this study. As the intervention was pre-planned and not

implemented by the study team, we could not randomize its allocation, increasing the risk of

confounding. We assessed potential confounding variables at baseline and used a DID analysis,

which accounts for baseline outcome measures, to limit the effect of unmeasured, residual

confounding. While we attempted to enroll intervention and control compounds with comparable

numbers of residents, the NGO which identified and implemented the intervention selected most

of the largest eligible compounds for intervention. This resulted in intervention compounds

having a slightly higher mean number of residents than control compounds (Table 1). Crowding

has been identified as a risk factor for pathogen transmission and poor health outcomes in other

studies, (Halpenny, Koski, Valdés, & Scott, 2012; Rahman, Wojtyniak, Mujibur Rahaman, &

Aziz, 1985; Rogawski McQuade, Shaheen, et al., 2020) though we found limited evidence of this

in our study population at baseline (Knee et al., 2018). Further, we assessed the number of

compound residents as a potential confounder but found that it did not meaningfully change the

DID estimates for our pre-defined outcomes (Supplemental Table 9). We consider our analysis to

be robust to small differences in study arms at baseline, however, we cannot exclude the

possibility of residual confounding due to such differences, a limitation of non-randomized

designs.

It was not possible to mask participants to their intervention status, and our measure of caregiver-

reported diarrhea could be subject to respondent and recall biases. To reduce the risk of

respondent bias, the MapSan field enumerator team and implementation team were different, and

respondents were not informed explicitly that the MapSan team was evaluating the health effect

of the intervention. To limit recall bias, we used a 7-day recall period (Arnold et al., 2013). Our

other pre-specified outcomes were objective measures of pathogen infection and not subject to

the same biases (Brown & Cumming, 2019).

Due to the greater than expected losses to follow-up in both study arms, we were not able to

follow all children enrolled at baseline through time as expected, but we still achieved our target

enrollment numbers due to migration and births into study compounds. We conducted the

originally planned longitudinal analysis as a sub-group analysis. It also served as a sensitivity

analysis to estimate the impact of migration on our effect estimates. Results from this sub-group

analysis were largely similar to results of the main analysis which treated measures as repeated

cross-sections, though the reduction in sample size led to wider confidence intervals

(Supplemental Tables 14-15). Measures of outcomes and covariates in children with and without

repeated measures were mostly similar, further limiting the likelihood that changes in the study

population biased our results.

While molecular detection of enteric pathogens in stool is evidence of pathogen exposure, it is

not necessarily evidence of active infection, making its clinical significance less clear (Brown &

Cumming, 2019). We assumed pathogen detection by the GPP indicated infection because the

assay's limits of detection exceeded the median infectious dose of most pathogens. While the

GPP detects many enteric pathogens recognized as important causes of childhood diarrhea in

LMICs, (Liu et al., 2016) it does not detect all enteric pathogens of importance. Further,

qualitative, cross-sectional analysis of stools does not provide information on the duration or

intensity of infection or pathogen carriage. Quantitative results, like those produced by multiplex

quantitative PCR panels, can be used to aid identification of etiologic agents of diarrhea,

especially in cases of coinfection, and to differentiate between low-level enteric pathogen

detection of unknown clinical relevance and higher concentration shedding which is more clearly

associated with disease (Liu et al., 2014, 2016; Platts-Mills, Liu, & Houpt, 2013). Some studies

have demonstrated overall good performance of the GPP but observed elevated false positive

detection rates for the *Salmonella* targets (Duong et al., 2016; Kellner et al., 2019). For this

reason, we removed *Salmonella* results from our pre-specified outcome definition. Results from

analyses including and excluding *Salmonella* were similar. In addition, some studies have

observed reduced sensitivity or specificity for some GPP targets compared with qPCR-based

methods, including norovirus, adenovirus, *Campylobacter*, *Yersinia enterocolitica*, ETEC, and

Salmonella, though inconsistencies between studies exist and are likely due to differences in

comparator assays or sample storage and processing (Chhabra et al., 2017; Deng et al., 2015;

Duong et al., 2016; Huang et al., 2016; Zhan et al., 2020; Zhuo et al., 2017). Further, the lack of

an adequate reference standard in most comparative studies complicates interpretation (K.

Freeman et al., 2017).

Our ability to detect an effect on our primary outcome, the prevalence of ≥1 bacterial or

protozoan infection, may have been limited by (1) the extended duration of shedding of some

pathogens following active infection; (2) the overall high burden of disease in our study

population, particularly among older children; and (3) residual confounding by age given the

strong observed relationship between age and infection status (particularly for protozoan

pathogens), all of which may have biased our results toward the null. Further, the intervention

may have impacted the concentration of pathogens shed (Grembi et al., 2020; Lin et al., 2019),

but our binary outcome was not sensitive to such differences The qualitative nature of the GPP

did not allow us to interrogate this question.

We analyzed a smaller number of stool samples for STH than for other enteric pathogens due to

requirements of the Kato-Katz method used for STH detection. The Kato-Katz method can only

be performed on whole, solid stool. Diarrheal samples and rectal swabs, the latter of which were

introduced during the 12-month follow-up phase, were not eligible for STH analysis by Kato-

Katz. Further, when limited stool material was collected, we prioritized the molecular analysis

used for the primary outcome. While the smaller sample size available for the STH analyses may

have reduced our ability to detect small effects, the proportions of whole stool, diarrheal diaper

samples, and rectal swabs were similar between arms at each phase (Supplemental Table 1). This

limited the potential impact that sample type could have on our results.

While the Kato-Katz method performs similarly to other microscope-based and molecular

methods for detection of moderate to high intensity infections, it may be less sensitive than

molecular methods in detecting low intensity infections (Benjamin-Chung et al., 2020; Cools et

al., 2019). A recent study has also suggested reduced specificity of the Kato-Katz method for

detection of low-intensity *A. lumbricoides* infections (Benjamin-Chung et al., 2020). In settings

where low-intensity infections are common, or where STH may be targeted for elimination,

methods with better diagnostic accuracy, like qPCR, may be considered.

We had limited ability to evaluate the impact of seasonality or weather-related trends on our

effect estimates due to drought conditions during the 2015/2016 rainy season. We adjusted

models for cumulative 30-day rainfall, a binary indicator of wet/dry season, and sine/cosine

terms of sample collection date (Stolwijk, Straatman, & Zielhuis, 1999) but excluded all

seasonality terms from final multivariable models because they did not meaningfully change

effect estimates.

Our results demonstrate that access to hygienic, shared onsite sanitation systems was not

sufficient to reduce enteric infection or diarrhea in children aged 6 years or younger (≤4 at

baseline) 12-24 months after implementation. Results from our sub-group analysis of children

born into intervention sites showed a substantial reduction in the prevalence of any STH,

Trichuris, and *Shigella* infection, suggesting that children may require protection from birth to

reduce or delay infection burdens. Our results do not suggest that shared sanitation is inadvisable

in this setting, as we did not compare against household-level sanitation improvements, nor do

they account for the many non-health related benefits associated with this intervention or

upgraded sanitation generally (Caruso et al., 2018; G.D. Sclar et al., 2018; Shiras et al., 2018).

The need for effective sanitation solutions may be most urgent in densely populated, low-

income, informal communities like our study setting where ubiquitous fecal contamination drives

high infection burdens. Disease transmission in these settings may be driven by multiple

interrelated pathways, complicated by frequent migration and the diversity of circulating

pathogens, and therefore difficult to interrupt. While decades of research have demonstrated

meaningful health gains following sanitation improvements, the results of this study, and other

rigorous trials of sanitation interventions, suggest that the relationship between sanitation and

health is complex, difficult to measure, and may not be generalizable across diverse settings and

populations.

Methods

Study design and intervention

MapSan was a controlled before-and-after trial, and details of the study design and analysis plan

have been published previously (Brown et al., 2015). We conducted the study in 17 densely

populated, low-income, informal neighborhoods in Maputo, Mozambique. The intervention was

delivered to compounds, typically groups of three to five households (though larger and smaller

compounds exist) often delineated by a wall or barrier, that shared sanitation and outdoor living

space. Shared compound sanitation facilities are not considered public facilities. We collected

data in an open cohort of children in intervention and control compounds at three time-points:

baseline (pre-intervention), 12 months post-intervention, and 24 months post-intervention.

The NGO Water and Sanitation for the Urban Poor selected intervention compounds and

designed and built 300 intervention facilities - pour-flush toilets discharging to septic tanks, the

liquid effluent of which flows to the soil through soakaway pits (Supplemental Figures 5a and

5b). There were two intervention designs with the same basic sanitation technology: communal

sanitation blocks (CSBs) and shared latrines (SLs) (Supplemental Figure 6a and 6b). The

primary difference between CSBs and SLs was size. CSBs (n=50) included multiple stalls with

toilets and served compounds of 21 or more people with one stall allocated per 20 residents.

CSBs also included rainwater harvesting systems, a municipal shared water connection, elevated

water tanks for storage of municipal water, a handwashing basin, a laundry facility, and a well-

drained area for bathing. Shared piped water connections were part of the municipal water

system and could be used for drinking in addition to other domestic purposes. Rainwater was

intended for cleaning and flushing but not drinking. Shared latrines (n=250) were single-stall

facilities serving fewer than 21 people. All septic tanks were sized to require emptying after

approximately two years.

Intervention compounds were located in 11 neighborhoods of the Nhlamankulu and KaMaxakeni

districts of Maputo (Supplemental Figure 7). The NGO selected intervention compounds using

the following criteria: (1) residents shared sanitation in poor condition as determined by an

engineer; (2) the compound was located in the pre-defined implementation neighborhoods; (3)

there were no fewer than 12 residents; (4) residents were willing to contribute financially to

construction costs; (5) sufficient space was available for construction of the new facility; (6) the

compound was accessible for transportation of construction materials and tank-emptying

activities; (7) the compound had access to a legal piped water supply; and (8) the groundwater

level was deep enough for construction of a septic tank. Intervention compounds were expected

to pay approximately 10-15% of the construction costs (~$64 for shared latrines and ~$97 for

CSBs) within one year of construction, with 25% of the total due upfront. Presence of a child

was not a selection criterion and therefore not all intervention sites were included in the study.

Opening of newly constructed intervention latrines occurred between February 2015 and

February 2016. The study team used criteria 1, 3, 4, and 7 to select control sites that had at least

one child younger than 48 months old in residence. We enrolled intervention and control

compounds concurrently to limit any differential effects of seasonality or other secular trends on

the outcomes (Supplemental Figure 2). Additionally, we attempted to enroll control compounds

with similar numbers of residents as intervention compounds. Willingness to pay for facilities

among controls was assessed using hypothetical versions of questions posed to interventions.

Control compounds were located within the 11 intervention neighborhoods and six adjacent but

similar neighborhoods due to the limited availability of eligible compounds remaining within

intervention neighborhoods (Supplemental Figure 7). Intervention selection criteria (5), (6), and

(8) were not used to select control sites as they were deemed to be related to intervention

construction and maintenance and unlikely to influence our outcomes. It was not possible to

blind participants or enumerators to intervention status.

Participants

We enrolled eligible children at three time points: baseline (0 months), 12 months post-intervention, and 24 months post-intervention. Children aged 1- 48 months old were eligible for baseline enrollment if we received written informed consent from a parent or guardian and if the head of the compound provided verbal assent for the compound to be included in the study. Children were eligible for enrollment at 12- and 24-month visits if they were aged 1-48 months or if they were eligible for enrollment at baseline but absent during that study visit. Children who moved into the compound fewer than six months before the 12-month or 24-month visit were not eligible for enrollment during that phase given their limited exposure to their new compound.

Ethics

The study protocol was approved by the Comité Nacional de Bioética para a Saúde (CNBS), Ministério da Saúde (333/CNBS/14), the Research Ethics Committee of the London School of Hygiene & Tropical Medicine (reference # 8345), and the Institutional Review Board of the Georgia Institute of Technology (protocol # H15160).

Procedures

Trained field enumerators completed consent procedures and surveys in the participant's preferred language (Portuguese or Changana) and collected biological specimens from enrolled children (Supplemental Methods). At baseline, we aimed to visit intervention compounds two weeks prior to the opening of the new latrines. We scheduled follow-up visits to be 12 months (±2 weeks) and 24 months (±2 weeks) from the date compound members began using their new latrines, with visits to control compounds made concurrently (±2 weeks).

We collected stool samples independently of reported symptomology. If we were unable to

collect a stool sample after multiple attempts, a registered nurse collected a rectal swab after

obtaining written consent for the procedure from a parent or guardian. Stool samples were kept

cold and delivered to the Laboratory of Molecular Parasitology at the Instituto Nacional de

Saúde (INS) within six hours of collection for analysis and storage at -80°C.

Samples were shipped frozen with temperatures monitors to the Georgia Institute of Technology

(Atlanta, USA) where we used the xTAG Gastrointestinal Pathogen Panel (Luminex Corp,

Austin, USA), a qualitative multiplex molecular assay, to detect 15 enteric pathogens in stool

samples: *Campylobacter jejuni/coli/lari*; *Clostridium difficile*, toxin A/B; enterotoxigenic

Escherichia coli (ETEC) LT/ST; Shiga-like toxin producing *E. coli* (STEC) stx1/stx2; *E. coli*

O157; *Salmonella*; *Shigella boydii/sonnei/flexneri/dysenteriae*; *Vibrio cholerae*; *Yersinia*

enterocolitica; *Giardia lamblia*; *Cryptosporidium parvum/hominis*; *Entamoeba histolytica;*

adenovirus 40/41; norovirus GI/GII; and rotavirus. The GPP has been rigorously tested and

extensively used for stool-based enteric pathogen detection (Chisenga et al., 2018; Claas,

Burnham, Mazzulli, Templeton, & Topin, 2013; Deng et al., 2015; Duong et al., 2016; Huang et

al., 2016; Kellner et al., 2019; Khare et al., 2014; Navidad, Griswold, Gradus, & Bhattacharyya,

2013; Patel, Navidad, & Bhattacharyya, 2014). We analyzed samples according to manufacturer

instructions with the addition of elution steps for the pretreatment of rectal swabs and diaper

material saturated with liquid stool (Supplemental Methods). Technicians at INS assessed stool

samples for the presence of soil-transmitted helminths (STH) using the single-slide Kato-Katz

microscope method (Vestergaard Frandsen, Lausanne, Switzerland).

Representatives of the National Deworming Campaign (NDC) at the Mozambican Ministério da

Saúde (MISAU) offered single-dose albendazole (400 mg, 200 mg for children aged six to 12

months) to all eligible members of intervention and control compounds following sample

collection activities of each phase. Eligibility was defined by the NDC and included compound

members older than six months who were not pregnant.

Outcomes

For the 12-month analysis, we pre-specified the primary outcome as infection with one or more

of the 12 bacterial or protozoan enteric pathogens detected by the GPP and secondary outcomes

as re-infection with one or more STH as detected by Kato-Katz (following albendazole treatment

at baseline), and seven-day period prevalence of caregiver-reported diarrhea. All three outcomes

were considered secondary outcomes in the 24-month analysis. We defined diarrhea as the

passage of three or more loose or liquid stools in a 24-hour period or any stool with blood

(Arnold et al., 2013; Baqui et al., 1991). We excluded viral enteric pathogens from the primary

outcome definition. The intervention may not have interrupted virus transmission due to their

low infectious doses, high concentration shed in feces and extended period of shedding,

environmental persistence, and capability for direct person-to-person transmission (Julian, 2016).

Following reported specificity issues with the *Salmonella* target of the GPP, we removed it from

our GPP-based outcome definitions (Duong et al., 2016; Kellner et al., 2019). In addition to the

pre-specified outcomes, we evaluated the effect of the intervention on specific pathogen types

(bacterial, protozoan, viral) and on individual pathogens. The results for other secondary

outcomes listed in the trial registration (growth and environmental enteric dysfunction) will be

published separately.

 Statistical analysis

Our sample size calculation has been described previously (Brown et al., 2015). We included all

enrolled children at each visit and analysed data as repeated cross-sectional observations. We

examined the effect of the intervention at the 12-month and 24-month phases separately. We

conducted two sets of exploratory sub-group analyses. The first assessed the effect of the

intervention on children with repeat observations at baseline and 12-months and at baseline and

24-months visits. These longitudinal analyses also served as sensitivity analyses of the impact of

participant migration on effect estimates. The second sub-group analysis compared children who

were born into study sites after the intervention (or after baseline in controls) but before the 12-

month or 24-month visit with children of a similar age group at baseline. For example, children

born after baseline but before the 24-month visit were compared with children aged two years

old or younger at baseline. These analyses allowed us to explore whether exposure to the

intervention from birth would reduce enteric pathogen infection during the first 1-2 years of life.

We used a DID approach to assess the impact of the intervention on all outcomes at the 12- and

24-month visits. We used generalized estimating equations (GEE) to fit Poisson regression

models with robust standard errors. Our GEE models accounted for clustering at the compound

level because it was the highest level of nested data and the level of the intervention allocation

(Bottomley, Kirby, Lindsay, & Alexander, 2016). We estimated the effect of the intervention as

the interaction of variables representing treatment status (intervention versus control) and phase

(pre- or post-intervention). Therefore, effect estimates from our DID analysis are presented as

ratio measures (ratio of prevalence ratios) instead of absolute differences. Multivariable models

were adjusted for covariates determined *a priori* as potentially predictive of our outcomes,

including child age and sex, caregiver's education, and household wealth. Given the important

and potentially non-linear relationship between age and pathogen prevalence (Supplemental

Figure 4), we also considered inclusion of a higher order age term (age squared) in our models

(Supplemental Table 10). Additional covariates (Supplemental Table 9) were considered for

inclusion in multivariable models if they were imbalanced between arms at baseline (>0.1

standardized difference in prevalence or mean) and resulted in a meaningful change in the DID

effect estimate (±10% change in 12-month DID prevalence ratio). We assessed the potential

impact of seasonality on our results in three ways: (1) inclusion of binary indicator of wet

(November – April) and dry (May – October) season in multivariable models, (2) inclusion of a

variable representing cumulative rainfall (mm) 30 days prior to sample or survey collection in

multivariable models, and (3) inclusion of sine and cosine functions of sample and survey dates

in multivariable models (Supplemental Tables 9 and 11). We used the same statistical approach

for sub-group analyses. All analyses were performed on complete case data, and a missing data

table is presented in the supplemental information (Supplemental Table 16). We performed all

statistical analyses with Stata version 16 (StataCorp, College Station, USA).

Registration

The trial was pre-registered at ClinicalTrials.gov (NCT02362932).

Acknowledgements

We gratefully acknowledge data collection services and other support provided by the WE

Consult team and in particular Wouter Rhebergen and Ellen de Bruijn, and the hard work of the

enumerators Isabel Maninha Chiquele, Sérgio Adriano Macumbe, Carolina Zavale, Maria Celina

Macuacua, Guilherme Zimba, and Anabela Mondlane. Olimpio Zavale coordinated logistics for

early field work. We would like to thank staff at the Instituto Nacional de Saúde, specifically

Josina Mate, Acacio Sabonete, and Jeronimo Langa, for their support throughout the project. The

study would not have been possible without our implementing partners at Water and Sanitation

for the Urban Poor (WSUP), in particular Guy Norman, Carla Costa, Vasco Parente, and

Jonathan Stokes. At the Georgia Institute of Technology, we gratefully acknowledge the project

and laboratory support provided by Aaron Bivins, Kevin Zhu, David Berendes, Fred Goddard,

Olivia Ginn, Sarah Lowry, Olivia Stehr, Haley Lewis, Jonathan Pennie, Derek Whaler, Mio

Unno, Catherine Reynolds, Joel Seibel, Diana Chumak, Felicitas Schneider, Katherine Brand,

Jiaxin Li, and Meredith Lockwood. We would also like to thank Ben Arnold, Jack Colford, Radu

Ban, Jay Graham, Tony Kolb, Eddy Perez, Jan Willem Rosenboom, Tom Slaymaker, Larry

Moulton, and Darren Saywell for technical and study design input. We would like to recognize

the early contributions of the late Dr. Jeroen Ensink to the MapSan trial protocol. As a colleague,

and as a friend, he is sorely missed. Finally, we would like to acknowledge and thank all of the

participants and their families who graciously welcomed us into their homes and were so

generous with their time.

Role of the Funding Source

This study was funded by the United States Agency for International Development

(www.usaid.gov) under Translating Research into Action (Cooperative Agreement No. GHS-A-

00-09-00015-00) and the Bill and Melinda Gates Foundation (www.gatesfoundation.org) grant

OPP1137224. The funders had no role in study design, data collection and analysis, decision to

publish, or preparation of the manuscript.

Contributors

JB as principal investigator and OC as co-principal investigator designed the trial and drafted the

study protocol with input from WPS, JK, DH, PK, JS, VZ, and RN. JK was the study manager,

co-led laboratory work and data analysis with TS, and drafted the manuscript. TS curated the

data, designed data collection tools and activities with JK, and produced figures. ZA led field

data collection. CA, FB, DC, VC, EM, JMB, CR, WZ helped with sample organization and

laboratory analysis. WPS designed the analytical approach and JK, DH, and AM helped refine it.

JB and OC secured funding for the trial. All authors contributed critically to the final version of

the manuscript.

Data sharing

De-identified participant data which underlie the results reported in this manuscript is publicly

available on the MapSan trial Open Science Forum website (https://osf.io/p5shk). The published

trial protocol can be accessed at: https://bmjopen.bmj.com/content/5/6/e008215 .

Competing interests

All authors have completed the ICMJE uniform disclosure form

at *www.icmje.org/coi_disclosure.pdf* and declare: no support from any organization for the

submitted work; no financial relationships with any organizations that might have an interest in

the submitted work in the previous three years; no other relationships or activities that could

appear to have influenced the submitted work.

Source data for Figures and Tables

Figure 1 source data 1 and source code 1

Table 1 source data 1 and source code 1

Table 2 source data 1 and source code 1

Table 3 source data 1 and source code 1

References

References

Arnold, B. F., Galiani, S., Ram, P. K., Hubbard, A. E., Briceño, B., Gertler, P. J., & Colford, J.

M. (2013). Optimal Recall Period for Caregiver-reported Illness in Risk Factor and

Intervention Studies: A Multicountry Study. *American Journal of Epidemiology*, *177*(4),

361–370. https://doi.org/10.1093/aje/kws281

Baqui, A. H., Black, R. E., Yunus, M., Hoque, A. R. A., Chowdhury, H. R., & Sack, R. B.

(1991). Methodological issues in diarrhoeal diseases epidemiology: Definition of diarrhoeal

episodes. *International Journal of Epidemiology*, *20*(4), 1057–1063.

https://doi.org/10.1093/ije/20.4.1057

Barreto, M. L., Genser, B., Strina, A., Teixeira, M. G., Assis, A. M. O., Rego, R. F., …

Cairncross, S. (2010). Impact of a citywide sanitation program in Northeast Brazil on

intestinal parasites infection in young children. *Environmental Health Perspectives*,

118(11), 1637–1642. https://doi.org/10.1289/ehp.1002058

Barreto, M. L., Genser, B., Strina, A., Teixeira, M. G., Marlucia, A., Assis, O., … Cairncross, S.

(2007). Effect of city-wide sanitation programme on reduction in rate of childhood

diarrhoea in northeast Brazil: assessment by two cohort studies. In *www.thelancet.com* (Vol.

370). Retrieved from www.thelancet.com

Benjamin-Chung, J., Pilotte, N., Ercumen, A., Grant, J. R., Maasch, J. R. M. A., Gonzalez, A.

M., … Colford, J. M. (2020). Comparison of multi-parallel qPCR and double-slide kato-

katz for detection of soil-transmitted helminth infection among children in rural

Bangladesh. *PLoS Neglected Tropical Diseases*, *14*(4), 1–23.

https://doi.org/10.1371/journal.pntd.0008087

Berendes, D. M., Sumner, T. A., & Brown, J. M. (2017). Safely Managed Sanitation for All

Means Fecal Sludge Management for at Least 1.8 Billion People in Low and Middle

Income Countries. *Environmental Science and Technology*, *51*(5), 3074–3083.

https://doi.org/10.1021/acs.est.6b06019

Bick, S., Buxton, H., Chase, R. P., Ross, I., Adriano, Z., Capone, D., … Dreibelbis, R. (2021).

Using path analysis to test Theory of Change: a quantitative process evaluation of the

MapSan trial. *BMC Public Health (Under Review)*. https://doi.org/10.21203/rs.3.rs-

234718/v1

Bottomley, C., Kirby, M. J., Lindsay, S. W., & Alexander, N. (2016). Can the buck always be

passed to the highest level of clustering? *BMC Medical Research Methodology*, *16*(1).

https://doi.org/10.1186/s12874-016-0127-1

Briscoe, J. (1984). Intervention studies and the definition of dominant transmission routes.

American Journal of Epidemiology, *120*(3), 449–456.

https://doi.org/10.1093/oxfordjournals.aje.a113909

Brown, J., & Cumming, O. (2019). Stool-Based Pathogen Detection Offers Advantages as an

Outcome Measure for Water, Sanitation, and Hygiene Trials. *Am. J. Trop. Med. Hyg*, *0*(0),

1–2. https://doi.org/10.4269/ajtmh.19-0639

Brown, J., Cumming, O., Bartram, J., Cairncross, S., Ensink, J., Holcomb, D., … Schmidt, W.-P.

(2015). A controlled, before-and-after trial of an urban sanitation intervention to reduce

enteric infections in children: research protocol for the Maputo Sanitation (MapSan) study,

Mozambique. *Bmj Open*, *5*(6), e008215. https://doi.org/10.1136/bmjopen-2015-008215

Caruso, B. A., Cooper, H. L. F., Haardörfer, R., Yount, K. M., Routray, P., Torondel, B., &

Clasen, T. (2018). The association between women's sanitation experiences and mental

health: A cross-sectional study in Rural, Odisha India. *SSM - Population Health*, *5*, 257–

266. https://doi.org/10.1016/j.ssmph.2018.06.005

Chhabra, P., Gregoricus, N., Weinberg, G. A., Halasa, N., Chappell, J., Hassan, F., … Vinjé, J.

(2017). Comparison of three multiplex gastrointestinal platforms for the detection of

gastroenteritis viruses. *Journal of Clinical Virology*, *95*, 66–71.

https://doi.org/10.1016/j.jcv.2017.08.012

Chisenga, C. C., Bosomprah, S., Makabilo Laban, N., Mwila- Kazimbaya, K., Mwaba, J.,

Simuyandi, M., & Chilengi, R. (2018). Aetiology of Diarrhoea in Children Under Five in

Zambia Detected Using Luminex xTAG Gastrointestinal Pathogen Panel. *Pediatric*

Infectious Diseases: Open Access, *03*(02), 1–6. https://doi.org/10.21767/2573-0282.100064

Claas, E. C., Burnham, C. A. D., Mazzulli, T., Templeton, K., & Topin, F. (2013). Performance

of the xTAG® gastrointestinal pathogen panel, a multiplex molecular assay for

simultaneous detection of bacterial, viral, and parasitic causes of infectious gastroenteritis.

Journal of Microbiology and Biotechnology, *23*(7), 1041–1045.

https://doi.org/10.4014/jmb.1212.12042

Clasen, T., Boisson, S., Routray, P., Torondel, B., Bell, M., Cumming, O., … Schmidt, W.-P.

(2014). Effectiveness of a rural sanitation programme on diarrhoea, soil-transmitted

helminth infection, and child malnutrition in Odisha, India: a cluster-randomised trial.

Lancet Global Health, *2*(11), E645–E653. https://doi.org/10.1016/s2214-109x(14)70307-9

Cools, P., Vlaminck, J., Albonico, M., Ame, S., Ayana, M., José Antonio, B. P., … Levecke, B.

(2019). Diagnostic performance of a single and duplicate Kato-Katz, Mini-FLOTAC,

FECPAKG2 and qPCR for the detection and quantification of soil-transmitted helminths in

three endemic countries. *PLOS Neglected Tropical Diseases*, *13*(8), e0007446.

https://doi.org/10.1371/journal.pntd.0007446

Delahoy, M. J., Wodnik, B., McAliley, L., Penakalapati, G., Swarthout, J., Freeman, M. C., &

Levy, K. (2018). Pathogens transmitted in animal feces in low- and middle-income

countries. *International Journal of Hygiene and Environmental Health*, *221*(4), 661–676.

https://doi.org/10.1016/j.ijheh.2018.03.005

Deng, J., Luo, X., Wang, R., Jiang, L., Ding, X., Hao, W., … Che, X. (2015). A comparison of

Luminex xTAG® Gastrointestinal Pathogen Panel (xTAG GPP) and routine tests for the

detection of enteropathogens circulating in Southern China. *Diagnostic Microbiology and*

Infectious Disease, *83*(3), 325–330. https://doi.org/10.1016/j.diagmicrobio.2015.07.024

Devamani, C., Norman, G., & Schmidt, W.-P. (2014). A Simple Microbiological Tool to

Evaluate the Effect of Environmental Health Interventions on Hand Contamination.

International Journal of Environmental Research and Public Health, *11*(11), 11846.

Retrieved from http://www.mdpi.com/1660-4601/11/11/11846

Duong, V. T., Phat, V. V., Tuyen, H. T., Dung, T. T. N., Trung, P. D., Minh, P. Van, … Baker,

S. (2016). Evaluation of luminex xTAG gastrointestinal pathogen panel assay for detection

of multiple diarrheal pathogens in fecal samples in Vietnam. *Journal of Clinical*

Microbiology, *54*(4), 1094–1100. https://doi.org/10.1128/JCM.03321-15

Else, K. J., Keiser, J., Holland, C. V., Grencis, R. K., Sattelle, D. B., Fujiwara, R. T., … Cooper,

P. J. (2020, December 1). Whipworm and roundworm infections. *Nature Reviews Disease*

Primers, Vol. 6, pp. 1–23. Nature Research. https://doi.org/10.1038/s41572-020-0171-3

Ercumen, A., Benjamin-Chung, J., Arnold, B. F., Lin, A., Hubbard, A. E., Stewart, C., … Luby,

S. P. (2019). Effects of water, sanitation, handwashing and nutritional interventions on soil-

transmitted helminth infections in young children: A cluster-randomized controlled trial in

rural Bangladesh. *PLOS Neglected Tropical Diseases*, *13*(5), e0007323.

https://doi.org/10.1371/journal.pntd.0007323

Ercumen, A., Mertens, A., Arnold, B. F., Benjamin-Chung, J., Hubbard, A. E., Ahmed, M. A.,

… Colford, J. M. (2018). Effects of single and combined water, sanitation and handwashing

interventions on fecal contamination in the domestic environment: a cluster-randomized

controlled trial in rural Bangladesh. *Environmental Science & Technology*, *22*,

acs.est.8b05153. https://doi.org/10.1021/acs.est.8b05153

Ercumen, A., Pickering, A. J., Kwong, L. H., Mertens, A., Arnold, B. F., Benjamin-Chung, J., …

Colford, J. M. (2018). Do Sanitation Improvements Reduce Fecal Contamination of Water,

Hands, Food, Soil, and Flies? Evidence from a Cluster-Randomized Controlled Trial in

Rural Bangladesh. *Environmental Science and Technology*, *52*(21), 12089–12097.

https://doi.org/10.1021/acs.est.8b02988

Evans, B., Hueso, A., Johnston, R., Norman, G., Pérez, E., Slaymaker, T., & Trémolet, S. (2017).

Limited services? The role of shared sanitation in the 2030 agenda for sustainable

development. *Journal of Water Sanitation and Hygiene for Development*, *7*(3), 349–351.

https://doi.org/10.2166/washdev.2017.023

Ezeh, A., Oyebode, O., Satterthwaite, D., Chen, Y. F., Ndugwa, R., Sartori, J., … Lilford, R. J.

(2017). The history, geography, and sociology of slums and the health problems of people

who live in slums. *The Lancet*, *389*(10068), 547–558. https://doi.org/10.1016/S0140-

6736(16)31650-6

Fink, G., Günther, I., & Hill, K. (2014). Slum Residence and Child Health in Developing

Countries. *Demography*, *51*, 1175–1197. https://doi.org/10.1

Freeman, K., Tsertsvadze, A., Taylor-Phillips, S., McCarthy, N., Mistry, H., Manuel, R., &

Mason, J. (2017). Agreement between gastrointestinal panel testing and standard

microbiology methods for detecting pathogens in suspected infectious gastroenteritis: Test

evaluation and meta-analysis in the absence of a reference standard. *PLOS ONE*, *12*(3),

e0173196. https://doi.org/10.1371/journal.pone.0173196

Freeman, M. C., Garn, J. V., Sclar, G. D., Boisson, S., Medlicott, K., Alexander, K. T., …

Clasen, T. F. (2017). The impact of sanitation on infectious disease and nutritional status: A

systematic review and meta-analysis. *International Journal of Hygiene and Environmental*

Health, *220*(6), 928–949. https://doi.org/10.1016/j.ijheh.2017.05.007

Fuhrmeister, E. R., Ercumen, A., Pickering, A. J., Jeanis, K. M., Crider, Y., Ahmed, M., …

Nelson, K. L. (2020). Effect of Sanitation Improvements on Pathogens and Microbial

Source Tracking Markers in the Rural Bangladeshi Household Environment. *Environmental*

877 *Science and Technology*, *54*(7), 4316–4326. https://doi.org/10.1021/acs.est.9b04835

878 Fuller, J. A., & Eisenberg, J. N. S. (2016). Herd Protection from Drinking Water, Sanitation, and

879  Hygiene Interventions. *The American Journal of Tropical Medicine and Hygiene*, *95*(5),

880  1201–1210. https://doi.org/10.4269/ajtmh.15-0677

881 Fuller, J. A., Villamor, E., Cevallos, W., Trostle, J., & Eisenberg, J. N. S. (2016). I get height

882  with a little help from my friends: herd protection from sanitation on child growth in rural

883  Ecuador. *International Journal of Epidemiology*. https://doi.org/10.1093/ije/dyv368

884 Grembi, J. A., Lin, A., Karim, M. A., Islam, M. O., Miah, R., Arnold, B. F., … Luby, S. P.

885  (2020). Effect of Water, Sanitation, Handwashing, and Nutrition Interventions on

886  Enteropathogens in Children 14 Months Old: A Cluster-Randomized Controlled Trial in

887  Rural Bangladesh. *The Journal of Infectious Diseases*. https://doi.org/10.1093/infdis/jiaa549

888 Halpenny, C. M., Koski, K. G., Valdés, V. E., & Scott, M. E. (2012). Prediction of Child Health

889  by Household Density and Asset-Based Indices in Impoverished Indigenous Villages in

890  Rural Panamá. *The American Journal of Tropical Medicine and Hygiene*, *86*(2), 280–291.

891  https://doi.org/10.4269/ajtmh.2012.11-0289

892 Harris, M., Alzua, M. L., Osbert, N., & Pickering, A. (2017). Community-Level Sanitation

893  Coverage More Strongly Associated with Child Growth and Household Drinking Water

894  Quality than Access to a Private Toilet in Rural Mali. *Environmental Science &*

895  *Technology*, *51*(12), 7219–7227. https://doi.org/10.1021/acs.est.7b00178

896 Heijnen, M., Cumming, O., Peletz, R., Chan, G. K.-S., Brown, J., Baker, K., & Clasen, T.

(2014). Shared Sanitation versus Individual Household Latrines: A Systematic Review of

Health Outcomes. *PLoS ONE*, *9*(4), e93300. https://doi.org/10.1371/journal.pone.0093300

Holcomb, D. A., Knee, J., Capone, D., Sumner, T., Adriano, Z., Nalá, R., … Stewart, J. R.

(2021). Bayesian analysis of source tracking markers to estimate the effects of an urban

sanitation intervention on human fecal contamination in Mozambique. *BioRxiv*,

2021.02.19.432000. https://doi.org/10.1101/2021.02.19.432000

Holcomb, D. A., Knee, J., Sumner, T., Adriano, Z., de Bruijn, E., Nalá, R., … Stewart, J. R.

(2020). Human fecal contamination of water, soil, and surfaces in households sharing poor-

quality sanitation facilities in Maputo, Mozambique. *International Journal of Hygiene and*

Environmental Health, *226*(February), 113496. https://doi.org/10.1016/j.ijheh.2020.113496

Huang, R. S. P., Johnson, C. L., Pritchard, L., Hepler, R., Ton, T. T., & Dunn, J. J. (2016).

Performance of the Verigene® enteric pathogens test, Biofire FilmArray^{TM} gastrointestinal

panel and Luminex xTAG® gastrointestinal pathogen panel for detection of common

enteric pathogens. *Diagnostic Microbiology and Infectious Disease*, *86*(4), 336–339.

https://doi.org/10.1016/j.diagmicrobio.2016.09.013

Humphrey, J. H., Mbuya, M. N. N., Ntozini, R., Moulton, L. H., Stoltzfus, R. J., Tavengwa, N.

V, … Makoni, T. (2019). Independent and combined effects of improved water, sanitation,

and hygiene, and improved complementary feeding, on child stunting and anaemia in rural

Zimbabwe: a cluster-randomised trial. *The Lancet. Global Health*, *7*(1), e132–e147.

https://doi.org/10.1016/S2214-109X(18)30374-7

Julian, T. R. (2016). Environmental transmission of diarrheal pathogens in low and middle

income countries. *Environmental Science: Processes and Impacts*, *18*(8), 944–955.

https://doi.org/10.1039/c6em00222f

Jung, Y. T., Lou, W., & Cheng, Y.-L. (2017). Exposure-response relationship of neighbourhood

sanitation and children's diarrhoea. *Tropical Medicine & International Health*, *22*(7), 857–

865. https://doi.org/10.1111/tmi.12886

Kellner, T., Parsons, B., Chui, L., Berenger, B. M., Xie, J., Burnham, C. A. D., … Freedman, S.

B. (2019). Comparative evaluation of enteric bacterial culture and a molecular multiplex

syndromic panel in children with acute gastroenteritis. *Journal of Clinical Microbiology*,

57(6). https://doi.org/10.1128/JCM.00205-19

Keusch, G. T., Denno, D. M., Black, R. E., Duggan, C., Guerrant, R. L., Lavery, J. V, …

Brewer, T. (2014). Environmental Enteric Dysfunction: Pathogenesis, Diagnosis, and

Clinical Consequences. *Clinical Infectious Diseases*, *59*, S207–S212.

https://doi.org/10.1093/cid/ciu485

Khare, R., Espy, M. J., Cebelinski, E., Boxrud, D., Sloan, L. M., Cunningham, S. A., …

Binnicker, M. J. (2014). Comparative evaluation of two commercial multiplex panels for

detection of gastrointestinal pathogens by use of clinical stool specimens. *Journal of*

Clinical Microbiology, *52*(10), 3667–3673. https://doi.org/10.1128/JCM.01637-14

Knee, J., Sumner, T., Adriano, Z., Berendes, D., de Bruijn, E., Schmidt, W.-P., … Brown, J.

(2018). Risk factors for childhood enteric infection in urban Maputo, Mozambique: A cross-

sectional study. *PLOS Neglected Tropical Diseases*, *12*(11), e0006956.

https://doi.org/10.1371/journal.pntd.0006956

Kosek, M. N., Ahmed, T., Bhutta, Z. Z. A., Caulfield, L., Guerrant, R. L., Houpt, E., … Trigoso,

D. R. (2017). Causal Pathways from Enteropathogens to Environmental Enteropathy:

Findings from the MAL-ED Birth Cohort Study. *EBioMedicine*, *18*, 109–117.

https://doi.org/http://dx.doi.org/10.1016/j.ebiom.2017.02.024

Kwong, L. H., Ercumen, A., Pickering, A. J., Arsenault, J. E., Islam, M., Parvez, S. M., … Luby,

S. P. (2020). Ingestion of Fecal Bacteria along Multiple Pathways by Young Children in

Rural Bangladesh Participating in a Cluster-Randomized Trial of Water, Sanitation, and

Hygiene Interventions (WASH Benefits). *Environmental Science & Technology*,

acs.est.0c02606. https://doi.org/10.1021/acs.est.0c02606

Lin, A., Ali, S., Arnold, B. F., Rahman, M. Z., Alauddin, M., Grembi, J., … Luby, S. P. (2019).

Effects of Water, Sanitation, Handwashing, and Nutritional Interventions on Environmental

Enteric Dysfunction in Young Children: A Cluster-randomized, Controlled Trial in Rural

Bangladesh. *Clinical Infectious Diseases*. https://doi.org/10.1093/cid/ciz291

Lin, A., Ercumen, A., Benjamin-Chung, J., Arnold, B. F., Das, S., Haque, R., … Luby, S. P.

(2018). Effects of Water, Sanitation, Handwashing, and Nutritional Interventions on Child

Enteric Protozoan Infections in Rural Bangladesh: A Cluster-Randomized Controlled Trial.

Clinical Infectious Diseases, *67*(10), 1515–1522. https://doi.org/10.1093/cid/ciy320

Liu, J., Kabir, F., Manneh, J., Lertsethtakarm, P., Begum, S., Gratz, J., … Houpt, E. R. (2014).

Development and assessment of molecular diagnostic tests for 15 enteropathogens causing

childhood diarrhoea: a multicentre study. *Lancet Infect Dis*, *14*(8), 716–724.

https://doi.org/10.1016/s1473-3099(14)70808-4

Liu, J., Platts-Mills, J. A., Juma, J., Kabir, F., Nkeze, J., Okoi, C., … Houpt, E. R. (2016). Use of

quantitative molecular diagnostic methods to identify causes of diarrhoea in children: a

reanalysis of the GEMS case-control study. *The Lancet*, *388*(10051), 1291–1301.

https://doi.org/https://doi.org/10.1016/S0140-6736(16)31529-X

Luby, S. P., Rahman, M., Arnold, B. F., Unicomb, L., Ashraf, S., Winch, P. J., … Colford, J. M.

(2018, January). Effects of water quality, sanitation, handwashing, and nutritional

interventions on diarrhoea and child growth in rural Bangladesh: A cluster randomised

controlled trial. *The Lancet Global Health*. https://doi.org/10.1016/S2214-109X(17)30490-4

Majorin, F., Torondel, B., Chan, G. K. S., & Clasen, T. (2019). Interventions to improve disposal

of child faeces for preventing diarrhoea and soil-transmitted helminth infection. *Cochrane*

Database of Systematic Reviews, *2019*(9).

https://doi.org/10.1002/14651858.CD011055.pub2

MAL-ED Network Investigators. (2018). Early childhood cognitive development is affected by

interactions among illness, diet, enteropathogens and the home environment: findings from

the MAL-ED birth cohort study. *BMJ Global Health*, *3*(4), e000752.

https://doi.org/10.1136/bmjgh-2018-000752

Medgyesi, D., Sewell, D., Senesac, R., Cumming, O., Mumma, J., & Baker, K. K. (2019). The

landscape of enteric pathogen exposure of young children in public domains of low-income,

urban Kenya: The influence of exposure pathway and spatial range of play on multi-

pathogen exposure risks. *PLOS Neglected Tropical Diseases*, *13*(3), e0007292.

https://doi.org/10.1371/journal.pntd.0007292

Moser, W., Schindler, C., & Keiser, J. (2017). Efficacy of recommended drugs against soil

transmitted helminths: systematic review and network meta-analysis. *BMJ*, *358*, 4307.

https://doi.org/10.1136/bmj.j4307

Navidad, J. F., Griswold, D. J., Gradus, M. S., & Bhattacharyya, S. (2013). Evaluation of

Luminex xTAG gastrointestinal pathogen analyte-specific reagents for high-throughput,

simultaneous detection of bacteria, viruses, and parasites of clinical and public health

importance. *J Clin Microbiol*, *51*(9), 3018–3024. https://doi.org/10.1128/jcm.00896-13

Norman, G., Pedley, S., & Takkouche, B. (2010). Effects of sewerage on diarrhoea and enteric

infections: a systematic review and meta-analysis. *The Lancet Infectious Diseases*, *10*(8),

536–544. https://doi.org/10.1016/S1473-3099(10)70123-7

Null, C., Stewart, C. P., Pickering, A. J., Dentz, H. N., Arnold, B. F., Arnold, C. D., … Colford,

J. M. (2018). Effects of water quality, sanitation, handwashing, and nutritional interventions

on diarrhoea and child growth in rural Kenya: a cluster-randomised controlled trial. *The*

Lancet Global Health, *6*(3), e316–e329. https://doi.org/10.1016/S2214-109X(18)30005-6

Parker, E. P., Ramani, S., Lopman, B. A., Church, J. A., Iturriza-Gómara, M., Prendergast, A. J.,

& Grassly, N. C. (2018). Causes of impaired oral vaccine efficacy in developing countries.

Future Microbiology, *13*(1), 97–118. https://doi.org/10.2217/fmb-2017-0128

Patel, A., Navidad, J., & Bhattacharyya, S. (2014). Site-specific clinical evaluation of the

Luminex xTAG gastrointestinal pathogen panel for detection of infectious gastroenteritis in

fecal specimens. *Journal of Clinical Microbiology*, *52*(8), 3068–3071.

https://doi.org/10.1128/JCM.01393-14

Patil, S. R., Arnold, B. F., Salvatore, A. L., Briceno, B., Ganguly, S., Colford Jr., J. M., &

Gertler, P. J. (2014). The Effect of India's Total Sanitation Campaign on Defecation

Behaviors and Child Health in Rural Madhya Pradesh: A Cluster Randomized Controlled

Trial. *Plos Medicine*, *11*(8). https://doi.org/10.1371/journal.pmed.1001709

Penakalapati, G., Swarthout, J., Delahoy, M. J., McAliley, L., Wodnik, B., Levy, K., & Freeman,

M. C. (2017). Exposure to Animal Feces and Human Health: A Systematic Review and

Proposed Research Priorities. *Environmental Science & Technology*, *51*(20), 11537–11552.

https://doi.org/10.1021/acs.est.7b02811

Pickering, A. J., Djebbari, H., Lopez, C., Coulibaly, M., & Alzua, M. L. (2015). Effect of a

community-led sanitation intervention on child diarrhoea and child growth in rural Mali: a

cluster-randomised controlled trial. *The Lancet Global Health*, *3*(11), e701–e711.

https://doi.org/10.1016/S2214-109X(15)00144-8

Pickering, A. J., Ercumen, A., Arnold, B. F., Kwong, L. H., Parvez, S. M., Alam, M., … Luby,

S. P. (2018). Fecal Indicator Bacteria along Multiple Environmental Transmission Pathways

(Water, Hands, Food, Soil, Flies) and Subsequent Child Diarrhea in Rural Bangladesh.

Environmental Science & Technology, *52*(14), 7928–7936.

https://doi.org/10.1021/acs.est.8b00928

Pickering, A. J., Njenga, S. M., Steinbaum, L., Swarthout, J., Lin, A., Arnold, B. F., … Null, C.

(2019). Effects of single and integrated water, sanitation, handwashing, and nutrition

interventions on child soil-transmitted helminth and Giardia infections: A cluster-

randomized controlled trial in rural Kenya. *PLOS Medicine*, *16*(6), e1002841.

https://doi.org/10.1371/journal.pmed.1002841

Platts-Mills, J. A., Liu, J., & Houpt, E. R. (2013). New concepts in diagnostics for infectious

diarrhea. *Mucosal Immunology*, *6*(5), 876–885. https://doi.org/10.1038/mi.2013.50

Prendergast, A. J., & Kelly, P. (2016). Interactions between intestinal pathogens, enteropathy and

malnutrition in developing countries. *Current Opinion in Infectious Diseases*, *29*(3), 229–

236. https://doi.org/10.1097/QCO.0000000000000261

Rahman, M., Wojtyniak, B., Mujibur Rahaman, M., & Aziz, K. M. S. (1985). Impact of

Environmental Sanitation and Crowding on Infant Mortality in Rural Bangladesh. *The*

Lancet, *326*(8445), 28–30. https://doi.org/10.1016/S0140-6736(85)90068-6

Robb, K., Null, C., Teunis, P., Yakubu, H., Armah, G., & Moe, C. L. (2017). Assessment of

Fecal Exposure Pathways in Low-Income Urban Neighborhoods in Accra, Ghana:

Rationale, Design, Methods, and Key Findings of the SaniPath Study. *Am. J. Trop. Med.*

Hyg, *97*(4), 1020–1032. https://doi.org/10.4269/ajtmh.16-0508

Rogawski, E. T., Bartelt, L. A., Platts-Mills, J. A., Seidman, J. C., Samie, A., Havt, A., … MAL-

ED Network Investigators, the. (2017). Determinants and Impact of Giardia Infection in the

First 2 Years of Life in the MAL-ED Birth Cohort. *Journal of the Pediatric Infectious*

Diseases Society Giardia Epidemiology and Impact • JPIDS, *2017*(6), 153–160.

https://doi.org/10.1093/jpids/piw082

Rogawski, E. T., Liu, J., Platts-Mills, J. A., Kabir, F., Lertsethtakarn, P., Siguas, M., … Nyathi,

E. (2018). Use of quantitative molecular diagnostic methods to investigate the effect of

enteropathogen infections on linear growth in children in low-resource settings: longitudinal

analysis of results from the MAL-ED cohort study. *The Lancet Global Health*, *6*(12),

e1319–e1328. https://doi.org/10.1016/S2214-109X(18)30351-6

Rogawski McQuade, E. T., Platts-Mills, J. A., Gratz, J., Zhang, J., Moulton, L. H., Mutasa, K.,

… Houpt, E. R. (2020). Impact of water quality, sanitation, handwashing, and nutritional

interventions on enteric infections in rural zimbabwe: The sanitation hygiene infant

nutrition efficacy (SHINE) trial. *Journal of Infectious Diseases*, *221*(8), 1379–1386.

https://doi.org/10.1093/infdis/jiz179

Rogawski McQuade, E. T., Shaheen, F., Kabir, F., Rizvi, A., Platts-Mills, J. A., Aziz, F., …

Iqbal, N. T. (2020). Epidemiology of Shigella infections and diarrhea in the first two years

of life using culture-independent diagnostics in 8 low-resource settings. *PLOS Neglected*

Tropical Diseases, *14*(8), e0008536. https://doi.org/10.1371/journal.pntd.0008536

Satterthwaite, D., Beard, V. A., Mitlin, D., & Du, J. (2019). *Untreated and Unsafe□: Solving the*

Urban Sanitation Crisis in the Global South Solving the Urban Sanitation Crisis in the

Global South Untreated and Unsafe□: Washington DC. Retrieved from

www.citiesforall.org

Sclar, G.D., Penakalapati, G., Caruso, B. A., Rehfuess, E. A., Garn, J. V., Alexander, K. T., …

Clasen, T. (2018). Exploring the relationship between sanitation and mental and social well-

being: A systematic review and qualitative synthesis. *Social Science & Medicine*, *217*, 121–

134. https://doi.org/10.1016/J.SOCSCIMED.2018.09.016

Sclar, Gloria D., Penakalapati, G., Amato, H. K., Garn, J. V., Alexander, K., Freeman, M. C., …

Clasen, T. (2016). Assessing the impact of sanitation on indicators of fecal exposure along

principal transmission pathways: A systematic review. *International Journal of Hygiene*

and Environmental Health, *219*(8), 709–723.

https://doi.org/https://doi.org/10.1016/j.ijheh.2016.09.021

Shiras, T., Cumming, O., Brown, J., Muneme, B., Nala, R., & Dreibelbis, R. (2018). Shared

latrines in Maputo, Mozambique: Exploring emotional well-being and psychosocial stress.

BMC International Health and Human Rights, *18*(1). https://doi.org/10.1186/s12914-018-

0169-z

Spears, D., Ghosh, A., & Cumming, O. (2013). Open Defecation and Childhood Stunting in

India: An Ecological Analysis of New Data from 112 Districts. *PLoS ONE*, *8*(9), e73784.

https://doi.org/10.1371/journal.pone.0073784

Steinbaum, L., Mboya, J., Mahoney, R., Njenga, S. M., Null, C., & Pickering, A. J. (2019).

Effect of a sanitation intervention on soil-transmitted helminth prevalence and concentration

in household soil: A cluster-randomized controlled trial and risk factor analysis. *PLOS*

Neglected Tropical Diseases, *13*(2), e0007180.

https://doi.org/10.1371/journal.pntd.0007180

Stolwijk, A. M., Straatman, H., & Zielhuis, G. A. (1999). Studying seasonality by using sine and

cosine functions in regression analysis. *Journal of Epidemiology and Community Health*,

53(4), 235–238. https://doi.org/10.1136/jech.53.4.235

Tidwell, J. B., Chipungu, J., Ross, I., Antwi-Agyei, P., Alam, M.-U., Tumwebaze, I. K., …

Simiyu, S. (2020). Where Shared Sanitation is the Only Immediate Option: A Research

Agenda for Shared Sanitation in Densely Populated Low-Income Urban Settings. *The*

American Journal of Tropical Medicine and Hygiene, *104*(2), 429–432.

https://doi.org/10.4269/ajtmh.20-0985

UN-Habitat. (2016). *World Cities Report: Urbanization and Development Emerging futures*.

Nairobi. Retrieved from http://wcr.unhabitat.org/wp-content/uploads/2017/03/Chapter1-

WCR-2016.pdf

UNICEF/WHO. (2019). *Progress on household drinking water, sanitation and hygiene 2000-*

2017. Special focus on inequalities. New York. Retrieved from https://washdata.org

Wolf, J., Hunter, P. R., Freeman, M. C., Cumming, O., Clasen, T., Bartram, J., … Prüss-Ustün,

A. (2018). Impact of drinking water, sanitation and handwashing with soap on childhood

diarrhoeal disease: updated meta-analysis and meta-regression. *Tropical Medicine &*

International Health, *23*(5), 508–525. https://doi.org/10.1111/tmi.13051

Zhan, Z., Guo, J., Xiao, Y., He, Z., Xia, X., Huang, Z., … Zhang, J. (2020). Comparison of

BioFire FilmArray gastrointestinal panel versus Luminex xTAG Gastrointestinal Pathogen

Panel (xTAG GPP) for diarrheal pathogen detection in China. *International Journal of*

Infectious Diseases, *99*, 414–420. https://doi.org/10.1016/j.ijid.2020.08.020

Zhuo, R., Cho, J., Qiu, Y. Y., Parsons, B. D., Lee, B. E., Chui, L., … Pang, X. (2017). High

genetic variability of norovirus leads to diagnostic test challenges. *Journal of Clinical*

Virology, *96*(May), 94–98. https://doi.org/10.1016/j.jcv.2017.10.003

1104

1105

1106   Figure 1: Trial profile. †Eligible for enrollment at baseline and/or 12-month but traveling at time of visit. ‡Children
1107   removed from 24-month analysis because their compound received an intervention after completion of the baseline
1108   phase. Source files available in Figure 1 – source data 1 and Figure 1 – source code 1.

 Table 1: Baseline characteristics of enrolled children, households, and compounds

	Control		Intervention	
	N	**n (%) or mean (SD)**	**N**	**n (%) or mean (SD)**
Child level variables				
Age at survey, days[†]	520	700 (405)	441	694 (403)
Sex, female	520	266 (51%)	444	227 (51%)
Child is breastfed with or without complementary feeding	526	169 (32%)	448	143 (32%)
Child is exclusively breastfed	526	49 (9.3%)	448	37 (8.3%)
Child feces reported to be disposed of in a latrine	526	148 (28%)	448	141 (31%)
Child wears diapers	526	342 (65%)	447	294 (66%)
Caregiver completed primary school	528	287 (54%)	451	239 (53%)
Child's mother is alive	513	503 (98%)	435	426 (98%)
Respondent is child's mother	519	368 (71%)	443	284 (64%)
Household level variables				
Household population	441	5.4 (2.4)	365	6.1 (3.0)
Household wealth score, 0 (poorer) - 1 (wealthier)[*]	440	0.45 (0.10)	365	0.44 (0.10)
Household crowding, >3 persons/room	440	54 (12%)	365	60 (17%)
Household floor is covered[‡]	440	426 (97%)	365	333 (91%)
Household wall made of bricks, concrete, or similar[‡]	440	304 (69%)	365	215 (59%)
Household drinking water source inside compound	435	324 (74%)	360	294 (82%)
Latrine used by household has a ceramic or masonry pedestal[‡]	432	153 (35%)	359	142 (40%)
Latrine used by household has a drop-hole cover[‡]	434	232 (53%)	359	224 (62%)
Compound level variables				
Number of compound members	287	14 (6.2)	208	19 (12)
Number of households	287	3.8 (2.1)	208	4.4 (3.7)
Number of water taps in compound	283	0.98 (0.95)	207	1.4 (1.6)
Number of latrines in compound	287	1.0 (0.20)	207	1.1 (0.57)

Number of people sharing a latrine	285	14 (6.2)	197	17 (8.9)
Number of households sharing a latrine	285	3.7 (1.8)	197	4.0 (2.8)
Latrine walls made of brick, concrete or similar[‡]	282	72 (26%)	204	67 (33%)
Compound population density, persons/square meter[▯]	281	0.071 (0.04)	205	0.087 (0.05)
Compound has electricity that normally functions	287	251 (87%)	208	189 (91%)
Compound is prone to flooding	287	184 (64%)	208	120 (58%)
Any animals observed in compound[‡]	287	170 (59%)	208	132 (63%)
Dog(s) observed[‡]	287	14 (4.9%)	208	14 (6.7%)
Chicken(s) or duck(s) observed[‡]	287	40 (14%)	208	30 (14%)
Cat(s) observed[‡]	287	149 (52%)	208	116 (56%)

1110 Data are n (%) or mean (standard deviation) and collected by questionnaire unless otherwise noted. [†] Age range 32-1819 days, IQR 339-1021 days. Age
1111 distributions available in Supplemental Figure 3. [*]Assessed using Simple Poverty Scorecard for Mozambique
1112 (http://www.simplepovertyscorecard.com/MOZ_2008_ENG.pdf), [‡]Data collected by direct observation. [▯]Calculated as # of people living in the compound
1113 divided by the area of the compound in square meters. Source files available in Table 1 – source data 1 and Table 1 – source code 1.

1114 **Table 2: Effect of the intervention on bacterial, protozoan, and STH infection and diarrhea at 12 and 24 months post-intervention.**

	Prevalence			12 month Prevalence ratio (95% CI), p-value *		24 month Prevalence ratio (95% CI), p-value ⬜	
	Baseline	**12-month**	**24-month**	**unadjusted**	**adjusted†**	**unadjusted**	**adjusted†**
Any bacterial or protozoan infection‡							
Control	313/392 (80%)	334/395 (85%)	403/459 (88%)	..	..	..	..
Intervention	278/361 (77%)	347/408 (85%)	392/462 (85%)	1.04 (0.94 - 1.15), p=0.41	1.04 (0.94 - 1.15), p=0.41	1.00 (0.91 - 1.10), p=1.0	0.99 (0.91 - 1.09), p=0.89
Any STH infection‡							
Control	170/360 (47%)	143/283 (51%)	142/253 (56%)	..	..	..	..
Intervention	138/329 (42%)	150/305 (49%)	136/292 (47%)	1.12 (0.89 - 1.40), p=0.33	1.11 (0.89 - 1.38), p=0.35	0.94 (0.75 - 1.17), p=0.59	0.95 (0.77 - 1.17), p=0.62
Diarrhea‡							
Control	67/526 (13%)	40/430 (9.3%)	53/390 (14%)	..	..	..	..
Intervention	59/448 (13%)	59/436 (14%)	53/410 (13%)	1.41 (0.80 - 2.48), p=0.24	1.69 (0.89 - 3.21), p=0.11	0.92 (0.55 - 1.54), p=0.76	0.84 (0.47 - 1.51), p=0.56
Any Bacteria							
Control	271/392 (69%)	285/395 (72%)	345/459 (75%)	..	..	..	..
Intervention	227/361 (63%)	292/408 (72%)	324/462 (70%)	1.09 (0.95 - 1.25), p=0.25	1.09 (0.95 - 1.26), p=0.20	1.03 (0.90 - 1.18), p=0.69	1.00 (0.87 - 1.15), p=0.96
Shigella							
Control	179/392 (46%)	204/395 (52%)	269/459 (59%)	..	..	..	..
Intervention	152/361 (42%)	218/408 (53%)	245/462 (53%)	1.13 (0.91 - 1.39), p=0.28	1.12 (0.92 - 1.38), p=0.27	0.98 (0.80 - 1.20), p=0.86	0.95 (0.79 - 1.16), p=0.64
ETEC							
Control	116/392 (30%)	142/395 (36%)	127/459 (28%)	..	..	..	..
Intervention	110/361 (30%)	143/408 (35%)	126/462 (27%)	0.93 (0.68 - 1.28), p=0.66	0.96 (0.69 - 1.33), p=0.81	0.95 (0.67 - 1.35), p=0.77	0.83 (0.57 - 1.19), p=0.31
Campylobacter							
Control	39/392 (9.9%)	32/395 (8.1%)	48/459 (10%)	..	..	..	..
Intervention	21/361 (5.8%)	35/408 (8.6%)	34/462 (7.4%)	1.78 (0.89 - 3.56), p=0.10	1.68 (0.82 - 3.45), p=0.16	1.20 (0.60 - 2.39), p=0.60	1.28 (0.62 - 2.62), 0.50
C. difficile							
Control	22/392 (5.6%)	13/395 (3.3%)	13/459 (2.8%)	..	..	..	..

Intervention	13/361 (3.6%)	17/408 (4.2%)	11/462 (2.4%)	1.95 (0.71 - 5.35), p=0.20	2.09 (0.77 - 5.64), p=0.15	1.32 (0.47 - 3.73), p=0.60	1.41 (0.46 - 4.30), p=0.54
E. coli O157							
Control	13/392 (3.3%)	19/395 (4.8%)	25/459 (5.5%)	..	..	..	..
Intervention	18/361 (5.0%)	14/408 (3.4%)	16/462 (3.5%)	0.48 (0.18 - 1.27), p=0.14	0.46 (0.18 - 1.21), p=0.12	0.43 (0.15 - 1.29), p=0.13	0.52 (0.17 - 1.59), p=0.25
STEC							
Control	3/392 (0.77%)	9/395 (2.3%)	17/459 (3.7%)	..	..	..	..
Intervention	10/361 (2.8%)	5/408 (1.2%)	15/462 (3.3%)	0.14 (0.03 - 0.67), p=0.014	0.15 (0.03 - 0.70), p=0.016	0.23 (0.05 - 1.03), p=0.055	0.24 (0.05 - 1.01), p=0.052
Any Protozoa							
Control	205/392 (52%)	236/395 (60%)	303/459 (66%)	..	..	..	..
Intervention	195/361 (54%)	259/408 (63%)	296/462 (64%)	1.04 (0.87 - 1.24), p=0.69	1.03 (0.86 - 1.22), p=0.76	0.93 (0.78 - 1.11), p=0.40	0.91 (0.76 - 1.09), p=0.29
Giardia							
Control	201/392 (51%)	230/395 (58%)	294/459 (64%)	..	..	..	..
Intervention	186/361 (52%)	251/408 (62%)	289/462 (63%)	1.06 (0.88 - 1.27), p=0.55	1.05 (0.88 - 1.25), p=0.58	0.96 (0.80 - 1.14), p=0.61	0.93 (0.78 - 1.11), p=0.44
Cryptosporidium							
Control	8/392 (2%)	8/395 (2%)	14/459 (3.0%)	..	..	..	..
Intervention	16/361 (4.4%)	15/408 (3.7%)	15/462 (3.3%)	0.89 (0.23 - 3.43), p=0.87	0.89 (0.24 - 3.31), p=0.86	0.46 (0.11 - 1.93), p=0.29	0.53 (0.13 - 2.14), p=0.37
Any virus							
Control	53/392 (14%)	52/395 (13%)	59/459 (13%)	..	..	..	..
Intervention	52/361 (14%)	45/408 (11%)	62/462 (13%)	0.77 (0.45 - 1.32), p=0.35	0.75 (0.44 - 1.27), p=0.29	0.96 (0.55 - 1.68), p=0.88	1.03 (0.57 - 1.86), p=0.92
Norovirus GI/GII							
Control	38/392 (9.7%)	44/395 (11%)	47/459 (10%)	..	..	..	..
Intervention	39/361 (11%)	37/408 (9.1%)	55/462 (12%)	0.71 (0.38 - 1.33), p=0.28	0.68 (0.36 - 1.27), p=0.23	1.00 (0.52 - 1.93), p=0.99	1.10 (0.55 - 2.18), p=0.79
Adenovirus 40/41							
Control	13/392 (3.3%)	9/395 (2.3%)	7/459 (1.5%)	..	..	..	..
Intervention	9/361 (2.5%)	9/408 (2.2%)	6/462 (1.3%)	1.34 (0.34 - 5.23), p=0.68	1.24 (0.32 - 4.83), p=0.76	1.18 (0.23 - 5.98), p=0.84	0.97 (0.18 - 5.19), p=0.97
Coinfection, ≥2 GPP							

pathogens							
Control	206/392 (53%)	237/395 (60%)	302/459 (66%)	..	..	..	..
Intervention	185/361 (51%)	257/408 (63%)	282/462 (61%)	1.08 (0.90 - 1.29), p=0.39	1.08 (0.91 - 1.29), p=0.37	0.95 (0.80 - 1.12), p=0.54	0.93 (0.79 - 1.10), p=0.41
Trichuris							
Control	139/360 (39%)	116/283 (41%)	124/253 (49%)	..	..	..	..
Intervention	117/329 (36%)	120/305 (39%)	117/292 (40%)	1.05 (0.82 - 1.35), p=0.68	1.01 (0.79 - 1.28), p=0.96	0.89 (0.69 - 1.16), p=0.40	0.86 (0.67 - 1.10), p=0.22
Ascaris							
Control	95/360 (26%)	82/283 (29%)	78/253 (31%)	..	..	..	..
Intervention	68/329 (21%)	87/305 (29%)	56/292 (19%)	1.26 (0.87 - 1.82), p=0.22	1.33 (0.92 - 1.93), p=0.13	0.80 (0.52 - 1.21), p=29	0.83 (0.54 - 1.27), p=0.39
Coinfection, ≥2 STH							
Control	64/360 (18%)	55/283 (19%)	60/253 (24%)	..	..	..	..
Intervention	47/329 (14%)	57/305 (19%)	37/292 (13%)	1.16 (0.76 - 1.77), p=0.50	1.17 (0.76 - 1.79), p=0.49	0.67 (0.40 - 1.13), p=0.13	0.63 (0.37 - 1.07), p=0.084

Prevalence results are presented as (n/N (%)). All effect estimates are presented as prevalence ratios (ratio of ratios) and estimated using
generalized estimating equations to fit Poisson regression models with robust standard errors. *Analysis includes all children measured at baseline
and 12-month visits. □Analysis includes all children measured at baseline and 24-month visits. ‡Outcome was pre-specified in trial registration.
All other outcomes are exploratory. †Pathogen outcomes adjusted for child age and sex, caregiver's education, and household wealth index.
Reported diarrhea was also adjusted for baseline presence of a drop-hole cover and reported use of a tap on compound grounds as primary
drinking water source. Sample sizes for adjusted analyses are slightly smaller than numbers presented in prevalence estimates due to missing
covariate data. *Y. enterocolitica, V. cholerae, E. histolytica*, and rotavirus were detected in <2% of samples in each arm at each phase. Descriptive
data for these pathogens are available in the Supplemental Table 2. Source files available in Table 2 – source data 1 and Table 2 – source code 1.

Table 3: Effect of intervention on bacterial, protozoan, and STH infection and reported diarrhea in children born into study sites post-intervention (post-baseline)
but by 24-month visit compared with children of a similar age at baseline (<2 years old).

	Prevalence (<2 years old)		Prevalence ratio (95% CI), p-value	
	Baseline	**24-month, Born-in**	**unadjusted**	**adjusted†**
Any bacterial or protozoan infection‡				
Control	158/228 (69%)	79/106 (75%)	..	..
Intervention	129/201 (64%)	71/107 (66%)	0.96 (0.77 - 1.21), p=0.74	0.99 (0.80 - 1.22), p=0.92
Any STH infection‡				
Control	67/205 (33%)	25/68 (37%)	..	..
Intervention	52/183 (28%)	13/75 (17%)	0.52 (0.26 - 1.05), p=0.069	0.51 (0.27 - 0.95), p=0.035
Diarrhea‡				
Control	46/283 (16%)	18/105 (17%)	..	..
Intervention	43/238 (18%)	22/100 (22%)	1.20 (0.57 -2.5), p=0.64	1.37 (0.47 - 4.03), p=0.57
Any Bacteria				
Control	142/228 (62%)	70/106 (66%)	..	..
Intervention	102/201 (51%)	51/107 (48%)	0.89 (0.66 - 1.20), p=0.44	0.90 (0.67 - 1.19), p=0.45
Shigella				
Control	67/228 (29%)	36/106 (34%)	..	..
Intervention	49/201 (24%)	15/107 (14%)	0.48 (0.28 - 0.83), p=0.009	0.49 (0.28 - 0.85), p=0.011
ETEC				
Control	70/228 (31%)	30/106 (28%)	..	..
Intervention	58/201 (29%)	24/107 (22%)	0.84 (0.46 - 1.52), p=0.56	0.85 (0.48 - 1.51), p=0.58
Campylobacter				
Control	27/228 (12%)	14/106 (13%)	..	..
Intervention	14/201 (7%)	13/107 (12%)	1.75 (0.63 - 4.87), p=0.29	1.75 (0.61 - 4.98), p=0.30
C. difficile				
Control	20/228 (8.8%)	7/106 (6.6%)	..	..
Intervention	13/201 (6.5%)	7/107 (6.5%)	1.33 (0.36 - 4.86), p=0.67	1.49 (0.41 - 5.44), p=0.55

E. coli O157				
Control	7/228 (3.1%)	3/106 (2.8%)	..	..
Intervention	9/201 (4.5%)	2/107 (1.9%)	0.45 (0.06 - 3.66), p=0.46	0.53 (0.07 - 4.24), p=0.55
STEC				
Control	1/228 (0.44%)	2/106 (1.9%)	..	..
Intervention	9/201 (4.5%)	1/107 (0.93%)	0.05 (0.00 - 1.13), p=0.059	0.05 (0.00 - 1.26), p=0.070
Any Protozoa				
Control	82/228 (36%)	47/106 (44%)	..	..
Intervention	74/201 (37%)	43/107 (40%)	0.84 (0.55 - 1.28), p=0.42	0.90 (0.62 - 1.30), p=0.58
Giardia				
Control	79/228 (35%)	44/106 (42%)	..	..
Intervention	68/201 (34%)	41/107 (38%)	0.90 (0.58 - 1.39), p=0.63	0.93 (0.64 - 1.36), p=0.70
Cryptosporidium				
Control	7/228 (3.1%)	5/106 (4.7%)	..	..
Intervention	12/201 (6%)	5/107 (4.7%)	0.45 (0.08 - 2.57), p=0.37	0.64 (0.12 - 3.51), p=0.61
Any virus				
Control	34/228 (15%)	18/106 (17%)	..	..
Intervention	36/201 (18%)	18/107 (17%)	0.83 (0.37 - 1.83), p=0.64	0.83 (0.37 - 1.87), p=0.66
Norovirus GI/GII				
Control	26/228 (11%)	12/106 (11%)	..	..
Intervention	26/201 (13%)	17/107 (16%)	1.24 (0.48 - 3.17), p=0.66	1.29 (0.49 - 3.41), p=0.61
Adenovirus 40/41				
Control	7/228 (3.1%)	4/106 (3.8%)	..	..
Intervention	7/201 (3.5%)	0/107 (0.0%)	..⬜	..⬜
Coinfection, ≥2 GPP pathogens				
Control	92/228 (40%)	52/106 (49%)	..	..
Intervention	74/201 (37%)	39/107 (36%)	0.82 (0.56 - 1.21), p=0.33	0.86 (0.59 - 1.24), p=0.41
Trichuris				

Control	48/205 (23%)	18/68 (26%)	..	..
Intervention	41/183 (22%)	5/75 (6.7%)	0.25 (0.09 - 0.68), p=0.006	0.24 (0.10 - 0.60), p=0.002
Ascaris				
Control	45/205 (22%)	16/68 (24%)	..	..
Intervention	29/183 (16%)	9/75 (12%)	0.70 (0.30 - 1.64), p=0.42	0.68 (0.30 - 1.54), p=0.36
Coinfection, ≥2 STH				
Control	26/205 (13%)	9/68 (13%)	..	..
Intervention	18/183 (9.8%)	1/75 (1.3%)	0.13 (0.02 - 1.08), p=0.059	0.12 (0.01 - 1.02), p=0.052

Analysis includes children <2 years old at baseline and children born into the study after baseline and <2 years old at the time of the 24-month
visit. Prevalence results are presented as (n/N (%)). All effect estimates are presented as prevalence ratios (ratio of ratios) and estimated using
generalized estimating equations to fit Poisson regression models with robust standard errors ‡Outcome was pre-specified in trial registration. All
other outcomes are exploratory. †Pathogen outcomes adjusted for child age and sex, caregiver's education, and household wealth index. Reported
diarrhea was also adjusted for baseline presence of a drop-hole cover and reported use of a tap on compound grounds as primary drinking water
source. Sample sizes for adjusted analyses are slightly smaller than numbers presented in prevalence estimates due to missing covariate data.
⬚Models would not converge due to sparse data. *Y. enterocolitica, V. cholerae, E. histolytica*, and rotavirus were detected in <2% of samples in
each arm at each phase and excluded. Descriptive data for these pathogens are available in the Supplemental Table 2. Source files available in
Table 3 – source data 1 and Table 3 – source code 1.

Supplemental Files

Supplemental Figure 1: Proportion of each type of sample collected during the baseline, 12-
month, and 24-month phases.

Supplemental Table 1: Number and proportion of sample types collected in each arm at each
phase.

Supplemental Figure 2: Enrollment and stool sample collection profile.

Supplemental Figure 3: Distribution of age (years) of enrolled children at each phase.

Supplemental Table 2: Age stratified baseline prevalence of health outcomes.

Supplemental Figure 4: Prevalence of pathogens by age at baseline, 12-month, and 24-month
phases.

Supplemental Table 3: Baseline enrollment characteristics of children with and without repeated
measures at the 12-month phase

Supplemental Table 4: Baseline enrollment characteristics of children with and without repeated
measures at the 24-month phase

Supplemental Table 5: Balance of characteristics measured at 12-month visits between children
with repeat observations at baseline and 12-month and children with observations at the 12-
month phase only.

Supplemental Table 6: Balance of characteristics measured at 24-month visits between children
with repeat observations at baseline and 24-month and children with observations at the 24-
month phase only.

Supplemental Table 7: Sensitivity analysis assessing the impact of reported deworming on STH
effect estimates 12 and 24 months after the intervention.

Supplemental Table 8: Sensitivity analysis assessing impact of independent upgrading of control
sanitation facilities on effect estimates.

Supplemental Table 9: Confounding assessment for primary outcome and both secondary
outcomes (any STH, diarrhea) at 12-month.

Supplemental Table 10: Effect estimates (prevalence ratios) for main analyses and all sub-group
analyses adjusted for *a priori* covariates and age-squared

Supplemental Table 11: Comparison of effect estimates (prevalence ratios) at 12- and 24 month
adjusted for a priori covariates only and for *a priori* covariates and seasonality.

Types of Collected Samples

Elevated water storage tank (500L)

Door (entry)

Laundry basin

Septic tank

Pour-flush toilet (pedestal design)

Study Sites

■ Intervention Sites
◆ Control Sites

250 0   250 500 750 1000 m