

1 Title Page for Submission to Paediatrics and International Child Health

2

3 Title: Where will they grow from here? Association between height at 24 months and adult
4 attained height in a population with a high prevalence of linear growth restriction.

5

6 Running Head: Where will they grow from here?

7

8 Author list

9 Julia Schwarz¹, Pablo Peñataro Yori^{2,3}, William K. Pan⁴, Maribel Paredes Olortegui^{2,3}, Robert
10 Klapheke⁵, Margaret N. Kosek^{2,3*}

11

12 ¹Icahn School of Medicine at Mount Sinai, 1 Gustave L. Levy Pl, New York, NY.

13 ²Asociación Benéfica Proyectos de Informática, Salud, Medicina, y Agricultura (A.B. PRISMA),
14 Iquitos, Peru.

15 ³Department of International Health, Johns Hopkins Bloomberg School of Public Health,
16 Baltimore, Maryland.

17 ⁴Duke Global Health Institute, Duke University, Durham, NC.

* Corresponding author info:

Margaret N. Kosek

Department of Infectious Diseases and International Health
University of Virginia School of Medicine

PO Box 801340
Charlottesville, VA 22908

mkosek@virginia.edu

18 ⁵ Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, MD, USA.

19 Keywords: Stunting; length-for-age; Peru; linear growth

20 Abstract Word Count: 180

21 Text Word Count: 4306

22 Figures: 2

23 Tables: 5

24

25 **Abstract**

26

27 This was a longitudinal observational cohort study to investigate the association of
28 stunting at 24 months of age and attained adult height. A cohort of 104 Peruvian males (n=47)
29 and females (n=57) was studied from 24 months to after puberty in a rural community in the
30 Peruvian Amazon located 15 km southeast from the urban center of Iquitos. Anthropometric
31 measures were made at 24 months and after puberty. Height for Age Z (HAZ) Scores were
32 calculated and compared to assess the association of HAZ score at 24 months and adult height.
33 67.7% of males and 54.1% of females who were stunted at 24 m (HAZ<-2.0) recovered (HAZ>-
34 2.0) after puberty, whereas 6.25% of males and 6.06% of females who were not stunted at 24 m
35 (HAZ>-2.0) became stunted (HAZ<-2.0) after puberty. The first 1000 days are not the only
36 critical window for growth promotion. Global and national initiatives to reduce stunting should
37 not exclusively focus on the first two years of life, but also work to promote the nutritional
38 initiatives to promote optimal nutrition and growth throughout childhood and adolescence.

39

40 Keywords: Child growth; adolescent growth; nutrition; stunting; Peru; Iquitos

41 **Introduction**

42 Linear growth is a key measure of child health. A link between stunting and both
43 physical and cognitive damage has been demonstrated in the last several decades with increasing
44 validation and stunting is a commonly accepted surrogate for loss of human potential at the
45 population level.^{1, 2, 3, 4, 5} In previous studies, 1 cm gained in the cohort's average height was
46 associated with a 6% increase in income per capita.⁶

47 Stunting is now considered a major global health priority and is the focus of several
48 high-profile initiatives, including Scaling Up Nutrition, the Zero Hunger Challenge and the
49 Nutrition for Growth Summit.⁷ It is estimated that in 2010, 26.7% (95% CI 24.8,28.7) of children
50 or 171 million (167 million in developing countries) were stunted, meaning they had a height-
51 for-age- Z-Score (HAZ) <-2.⁸ For the last decade, growth faltering research has focused on the
52 first 1000 days of life as the window of opportunity for growth promotion.⁹ These studies led to
53 the creation of programs that specifically promote nutrition during the first 24 months. However,
54 recent longitudinal cohort studies have suggested that recovery from growth stunting is possible
55 after the first two years of life.^{10, 11, 12}

56 This study investigates the veracity of the belief that stunting in the first two years
57 dictates height throughout adulthood. Human growth velocity is highly age dependent and has
58 classically been described as occurring in the infancy phase, which begins in midterm gestation
59 and ends at 3-4 years with a pattern of rapid deceleration, the childhood phase of slowly
60 decelerating linear growth velocity, and the pubertal phase, which is characterized by an increase
61 in growth velocity followed by a rapid and steep decline after puberty growth when the
62 epiphyseal plate closes and linear growth ends.^{13, 14} The biology of the childhood and pubertal
63 phases is not accounted for in the many public health programs designed with primary outcomes

64 based on measurements taken at 2 and 5 years. The evidence presented here offers an
65 uncommon perspective in that it follows a cohort from early childhood to puberty and
66 concentrates on HAZ scores at the individual level. Here we analyzed the attained height of
67 children at 12-16 years old who were and were not stunted at 24 months to evaluate the
68 predictive capacity of early childhood measures on final attained adult stature.

69 **Methods**

70 Study setting

71 Santa Clara is a rural community in the Peruvian Amazon located 15 km southeast from the
72 urban center of Iquitos. Stunting in this community is high, whereas acute undernutrition is
73 relatively uncommon. Previous studies offer additional details of this community.¹⁵

74

75 Study design

76 Between October 1, 2002 and April 15, 2006, the team conducted a 43-month long prospective,
77 community-based diarrheal disease surveillance in 442 children <72 months of age in Santa
78 Clara. The cohort and study design have been previously described.^{16, 17} The study cohort was
79 chosen from a community census that generated a list of children <72 months of age. Enrollment
80 was limited to one child per household. Selection included every third child on the census list
81 sorted according to date of birth. Ninety-six percent of families invited to participate gave
82 informed consent for enrollment in the study protocol. During the months they were enrolled in
83 the study subject's height was measured monthly by using a marked platform with a sliding
84 footboard. After 72 months of age, the child was removed from surveillance and the next live
85 birth in that community was enrolled.

86 Follow-up measurements were conducted between September and December 2016. Contact was
87 attempted with all participants in the original cohort. Community health workers returned to the
88 households they had visited monthly during the first stage of the cohort study to see if
89 participants would consent to further measurements. Retention is detailed in Figure 1. Subjects
90 had their height and weight measured and age of puberty was assessed by maternal report of the
91 onset of menses (for female subjects) or the age of voice change (for male subjects). Menarche is
92 an established indicator as the final stage of female puberty¹⁸ and age of voice change has been
93 shown to be representative of physical maturity in men.^{19, 20} The age of pubertal onset is an
94 important indicator of adult height, as the release of gonadal steroids (estrogen, testosterone)
95 causes closure of the epiphyseal growth plates and the end of statural growth. Therefore, we
96 included only those in the cohort that had gone through puberty as a marker of having passed
97 through the late adolescent growth spurt and be at, or very close to, terminal adult height.

98

99 Statistical Analysis

100 Data was analyzed to look at the association of height at 24 months on height, weight and
101 BMI post-puberty as well as on the age of puberty onset.

102 The anthropometrical measurements closest to 24 months (ranging from 23-30 months)
103 was compared with the WHO Reference Data 2006 to calculate the HAZ score for each child at
104 24 months. Participants were then categorized as stunted ($HAZ < -2$) or not stunted ($HAZ > -2$) at
105 24 months based on the HAZ calculated for the date closest to 24 months. For those retained at
106 follow-up, HAZ and BMI post puberty were calculated. The participants were again categorized
107 into “HAZ < -2 post-puberty” and “HAZ > -2 post puberty” based on the anthropometrical
108 measurements after maternal report of puberty.

109 Based on the change in HAZ score from 24 months to post puberty, participants were
110 divided into four categories. Children who were stunted at 24 months were categorized as
111 “recovered children” if their post-puberty HAZ was greater than negative two, or as “stunted
112 throughout” if their post-puberty HAZ was less than negative two. Among “recovered children,”
113 a subcategory of “fully recovered” was created for those with a post puberty HAZ greater than
114 negative one. Children who were not stunted at 24 months were categorized either as “late
115 stunting” if their post puberty HAZ was less than negative two , or “non-stunted throughout” if
116 their post-puberty HAZ was greater than negative two. The percentage of male, female and total
117 cohort members in each category was calculated. A chi-squared test was used to compare the
118 percentage of male and female participants in each category. In addition, mean attained height
119 among male and female participants was compared across each of the four categories. A one-
120 way ANOVA was conducted to compare the average age of puberty between each category
121 followed by post-hoc comparison using a Tukey test.

122 Selection bias due to loss-to-follow up was evaluated by comparing sex, HAZ score, and
123 stunting at 24 months among children lost to follow-up and those retained. In addition, using
124 data from 2002-2006, when the study began, average years of maternal education and average
125 monthly income was compared among children lost to follow-up and those retained. Data was
126 missing for 5 members in the group who were measured for adult-follow and 13 members who
127 were lost-to follow up. Two sample t-tests were used to compare means and chi-squared tests
128 compared percentages.

129 Post-puberty HAZ, weight and BMI were compared across three levels of HAZ at 24
130 months: HAZ less than negative two, HAZ between negative one and negative two, and HAZ
131 greater than -1. A one -way ANOVA was conducted to compare the association of HAZ at 24

132 months on HAZ, BMI, and weight at puberty. This was followed by post-hoc comparison using a
133 Tukey test.

134 Odds ratios were calculated for the overall likelihood a child with a HAZ less than
135 negative two at 24 months will be stunted post puberty compared with a child with a HAZ
136 greater than negative two at 24 months.

137 The average age of puberty onset was calculated and analyzed for each category. Because
138 puberty onset was reported in years, when puberty onset was studied in months it would have
139 been falsely clustered in the first month of the year. To give a more accurate representation, a
140 random normally distributed covariate with a mean of 6 and standard deviation of 4 was added to
141 the reported age of puberty onset in months.

142

143 Ethical Review

144 The study protocol was reviewed and approved by the Institutional Review Board of the Johns
145 Hopkins Bloomberg School of Public Health (Baltimore, MD) and the Ethics Committee of
146 Asociacion Benefica PRISMA (Lima, Peru).

147

148 **Results**

149

150 Subject Characteristics

151 A description of the participants that were lost to follow-up and those that were retained
152 is shown in Table 1. 45 percent of those retained for follow-up were male. The average HAZ was
153 -1.99 and 53 percent were stunted at 24 months. The average years of maternal education was

154 8.04 years and the average per capita monthly income was \$31.52. The average age at puberty
155 was 12.64 among males and 11.79 among females in the group retained for follow-up. There was
156 no significant difference in the gender breakdown, HAZ, percentage stunted at 24 months, years
157 of maternal education or per capita monthly income in the children lost to follow-up.

158

159 Association of HAZ at 24 Months on HAZ, Weight, and BMI at Puberty

160 The mean HAZ, weight and BMI at puberty are presented in Table 2 categorized by
161 participants who had a HAZ less than -2, HAZ between -2 and -1, and HAZ greater than -1 at 24
162 months.

163 When analyzing the association of HAZ at 24 months on HAZ at puberty there was a
164 significant difference between groups as determined by one-way ANOVA ($F(2,101) = 26.944$, p
165 $< .001$). A Tukey post hoc test revealed that the HAZ score at puberty was significantly lower if
166 the child had a HAZ less than -2 at 24 months (-1.75 ± 0.76 , $p < .001$) and if the child has a HAZ
167 between -2 and -1 at 24 months (-0.96 ± 0.70 , $p = .011$) compared to those who HAZ greater than -
168 1 at 24 months (-0.24 ± 0.82). The HAZ score at puberty was also significantly lower if the child
169 had a HAZ less than -2 at 24 months (-1.75 ± 0.76 , $p < .001$) compared to those who had a HAZ
170 between -2 and -1 at 24 months (-0.96 ± 0.70).

171 The association of HAZ at 24 months on weight at puberty also showed a significant
172 difference between groups as determined by one-way ANOVA ($F(2,101) = 7.127$, $p = .001$). A
173 Tukey post hoc test revealed that the weight at puberty was significantly lower if the child had a
174 HAZ less than -2 at 24 months (50.0 ± 8.6 , $p = .002$) and if the child had a HAZ between -2 and -1
175 at 24 months (49.6 ± 6.4 , $p = .002$) compared to those who HAZ greater than -1 at 24 months ($-$
176 59.1 ± 11.1). There was no significant difference in weight at puberty between children who had a

177 HAZ less than -2 at 24 months and those who had a HAZ between -2 and -1 at 24 months
178 (p=.975).

179 The association of HAZ at 24 months on BMI at puberty also showed a significant
180 difference between groups as determined by one-way ANOVA ($F(2,101) = 3.608, p = .031$). A
181 Tukey post hoc test revealed that the BMI at puberty was significantly lower if the child had a
182 HAZ between -2 and -1 at 24 months ($20.55 \pm 2.6, p = .034$) compared to those who HAZ greater
183 than -1 at 24 months (23.1 ± 3.7). There was no significant difference in BMI at puberty between
184 children who had a HAZ less than -2 at 24 months and those who has a HAZ between -1 and -2
185 at 24 months ($p = .167$). Similarly, there was no significant difference in BMI at puberty between
186 children who had a HAZ less than -2 at 24 months and those who has a HAZ greater than -1 at
187 24 months ($p = .347$). In the three groups the mean BMI ranged from 20.55 to 23.1.

188

189 Comparing Age Specific Prevalence of Stunting and Recovery by Gender

190 The overall prevalence of stunting was significantly different between boys and girls with
191 rates of 66% and 42% ($p = 0.015$), respectively at 24 months (Table 3). The frequency of recovery
192 in boys and girls recovered from stunting at 24 months to post-puberty was similar with boys at
193 68% and girls at 54%. Although there were different rates of stunting at 24 months the
194 prevalence of stunting post-puberty was 23% for both sexes. Recovery from HAZ < -2 at 24
195 months to HAZ > -1 post-puberty was also similar among boys (23%) and girls (8%) at 24
196 months.

197 The average age of puberty, as delineated by menarche in girls and voice change in boys
198 is

199 shown by categorization of children who recovered from stunting at 24 months, those who
200 remained stunted, those that became stunted in late childhood, and those that were neither
201 stunted at 24 months or at their measurement following puberty in Table 5. When analyzing the
202 association of recovering from stunting on age at puberty in males there was a significant
203 difference between groups as determined by one-way ANOVA ($F(3,43) = 3.539, p = .022$). A
204 Tukey post hoc test revealed males who recovered from stunting had a significantly lower age of
205 puberty ($12.28 \pm 0.90, p = .007$) compared to those who were stunted throughout (13.50 ± 1.18).
206 There was no significant difference in age of puberty between those who were stunted and those
207 who were never stunted ($p = .079$) or between those who recovered from stunting and those who
208 were never stunted ($p = .621$). When analyzing the association of recovering from stunting on age
209 at puberty in females there was no significant difference between groups as determined by one-
210 way ANOVA ($F(3,53) = .139, p = .936$).

211

212 Rates of Recovery

213 In this cohort, it was found that a child who had a HAZ < -2 at 24 months was 9.5 times
214 (CI 2.6-34.3, p-value 0.006) more likely to be stunted post-puberty compared to a non-stunted
215 child at 24 months (Table 4). Although if a child is stunted at 24 months it is more likely that
216 they will be stunted post puberty, the rates of recovery in this cohort were unexpectedly high.
217 Figure 2 displays a matched comparison of attained length at 24 months and attained height after
218 puberty demonstrates that in both boys and girls there is a trend towards recovery, with even
219 those with severe stunting ($HAZ \leq -3$) normalizing by the time they reached their adult stature.
220 On the other hand, almost no children who were not stunted at 24 months became stunted later in
221 childhood.

222 Among the birth cohort in Santa Clara, there was recovery from stunting among both
223 girls and boys, with 62% (n=34) of children who had a HAZ <-2 at 24 months (n=55) achieving
224 a HAZ >-2 post-puberty (Table 3). More specifically, 16% (n=9) of children who were stunted at
225 24 months (n=55) had a HAZ>-1 post-puberty.

226 Despite high rates of recovery from stunting, the mean attained height among cohort
227 members is well below the expected attained height as reported by the WHO. Among the male
228 cohort members, the mean anthropometrical measurements post puberty were a height of was
229 156.38 cm (SD 6.34), HAZ score of -1.34 (SD 1.01), weight of 52.15 kg (SD 9.79) and BMI of
230 21.26 (SD 3.27). Among the female cohort members the mean anthropometrical measurements
231 post puberty were a height of 151.69 (SD 5.74), HAZ score of -1.24 (SD 0.84), weight of 50.00
232 (SD 7.78) and BMI of 21.73 (SD 3.16). This result indicated that the average man in Iquitos is
233 11.31 cm shorter and the average woman in Iquitos is 9.20 cm shorter than mean attained adult
234 standard as indicated by WHO 2006 norms.

235

236

237

238 Discussion:

239 Our results indicate significantly higher rate of stunting in boys than girls at 24 months. After
240 puberty, boys still had slightly higher but not statistically significant rate of stunting. Although
241 being stunted at 24 months of age was a risk factor for having a constrained adult height, the
242 majority of children who were stunted at 24 months exhibited growth recovery, including
243 children with HAZ of ≤ -3 at 24 months. If children were not stunted by 24 months of age, they
244 were unlikely to become stunted in later childhood. Our interpretation of this data is that the
245 potential for linear growth recovery in childhood does exist. The use of the HAZ at 24 months is
246 a useful one to target populations of children who would may benefit from intensified dietary
247 and health interventions to maximize statural growth in late childhood. On the individual level,
248 parents of children who are stunted at 24 months should be encouraged to optimize current diet
249 to promote growth in later childhood, which may allow for normal adult stature to be attained.

250 A weakness in the present study was the loss to follow-up, which was 48%. Nearly all this
251 loss was due to migration outside the study area. Of the group that was not able to be assessed
252 for the ascertainment of puberty, 48% had a known migratory destination, which was most likely
253 to be Lima. When examined, there was no significant difference in the HAZ at 24 months and
254 eventual migratory status. Even if located the change in environment would have limited our
255 ability to interpret the data, loss to follow-up is difficult to avoid with a gap in time to follow-up
256 of 12-14 years.

257 This study is novel in showing that higher rates of stunting among boys than girls in early
258 childhood dissipate after puberty. This brings up the question of when the crucial catch-up
259 growth occurs in boys. It has been suggested that late childhood presents another critical window
260 when interventions can be made to increase height growth.²¹ A calcium carbonate supplement

261 given for 13 months to British boys age 16-18 years was associated with a modest increase of 7
262 mm compared to the control group.²² Additionally, a 12 month prepubertal calcium carbonate
263 supplementation given to Gambian boys age 8-12 years increased heights two centimeters at age
264 15.5 in comparison to the control group. Although the calcium group had a greater stature in
265 adolescence, subsequently, the calcium group stopped growing earlier and was 3.5 at a mean age
266 of 23.5 y. This population commonly has late onset puberty and low calcium intake.²³ This
267 seems to suggest that interventions in late adolescence may be more effective in males, but as
268 with most nutritional supplementation studies, the absolute amount of the total attained height
269 deficit corrected by the intervention is quite limited. Clearly though, more work needs to be done
270 to study this window for catch-up growth. Tumilowicz et al. advocates specifically for future
271 nutrition research that explores standard indicators to measure adolescent dietary quality to
272 determine causal effects between dietary intake and growth.²⁴

273 Among boys and girls, this nonintervention-study showed high rates of recovery with 62% of
274 children in the birth cohort who had a HAZ<-2 at 24 months having a HAZ>-2 post-puberty, and
275 16% of children stunted at 24 months fully recovering with HAZ>-1 post-puberty. These rates of
276 recovery suggest that catch-up growth after 24 months is possible and common among children
277 who stay in resource constrained environments with a high prevalence of acquired linear growth
278 deficits. Our study adds to a large amount of literature that suggests there is a natural high rate of
279 recovery from stunting after the first thousand days. Walker et al. found 87% of participants who
280 received a nutritional intervention from 24-48 months were no longer stunted at follow-up at 7 to
281 8 years. This rate of recovery, while promising, did not differ from the control, a group of
282 children who were stunted at 24 months and not given any intervention.²⁵ Satarayana et al.
283 found that Indian girls stunted in childhood gained significantly more height in the years before

284 finishing puberty than girls who were not stunted in childhood, even with no nutritional
285 intervention.²⁶ The Young Lives study in Peru-which includes participants from urban and rural
286 areas and represents coastal, highland, and jungle regions-found that one-third of children were
287 stunted around 1 year of age had recovered (HAZ>-2) at age 5, again with no nutritional
288 intervention.²⁷

289 The adolescent growth spurt has been posited as a critical period of growth beyond the first
290 1000 days.²¹ This may explain why our study, in which we did follow-up measurements after
291 puberty, showed much higher rates of recovery from stunting than other studies. The Young
292 Lives study in Peru, which includes participants from urban and rural areas and represents
293 coastal, highland, and jungle regions, found that one-third of children were stunted around 1 year
294 of age had recovered (HAZ>-2) at age 5.²⁷ Sterling et al. found among 27 participants that were
295 stunted at 12 months only 11 recovered by adolescence (40%).²⁸ In contrast, we found that of 55
296 participants who were stunted at 24 months, 34 had recovered post puberty (62%) despite higher
297 rates of stunting in early childhood as would be expected given the location of the study site in
298 one of the poorest regions of Peru. The major difference between these studies and ours is that
299 follow-up in our study was done only after participants went through puberty. This suggests that
300 there is significant room for recovery from stunting through adolescence, even in conditions
301 where stunting is common and severe.

302 Puberty is a time of significant growth among girls and boys and it is key to look at the
303 onset of the end of puberty when looking at catch-up growth. Our study found the mean age of
304 menarche to be 11.8 (median 12) and the mean age of voice change to be 12.6 (median 12),
305 whereas the median age at menarche was found to be 12.8 years and the median
306 for voice change 15.1 years among adolescents in Germany.²⁹ The only group that had a

307 significantly later onset of puberty was boys who were stunted both at 24 months and post
308 puberty. The early onset of puberty in this population is another signal of increased nutrition, as
309 in many other populations with high percentages of early childhood stunting, menarche and
310 voice change are significantly delayed.^{30, 31, 32} The early onset of puberty may be one reason that
311 although significant recovery was made from stunting, the average height attained by the cohort
312 was still more than one standard deviation below the WHO standards. Studies following
313 malnourished children adopted to more developed countries have found that early puberty can
314 negate the gains made in childhood.³³ This is of particular attention as obesity is an ever-growing
315 problem even in developing countries. Within this cohort all categories of children had a healthy mean
316 BMI. Many think over-nutrition can lead to early puberty, meaning that nutritional interventions must be
317 carefully timed and carried out both to avoid obesity and extend the pubertal growth spurt.³⁴

318 One of the most damaging aspects of growth stunting is its association with delayed
319 cognitive development in childhood, reduced height and lower earning potential as an adult.⁶
320 There has been controversy over whether or not recovery in stature from growth stunting
321 corresponds to recovery from the cognitive effects of stunting. The majority of the literature,
322 however, now seems to agree that those who recover from stunting have similar cognitive ability
323 to those who were never stunted.^{27, 35, 36, 37}

324 In addition to the risk of delayed cognitive development, stunted maternal height causes a
325 multigenerational burden. An analysis of 109 Demographic and Health Surveys in 54 countries
326 conducted between 1991 and 2008 found that a 1-cm increase in maternal height was associated
327 with a decreased risk in underweight (RR, 0.968; 95% CI, 0.968–0.969), stunting (RR, 0.968;
328 95% CI, 0.967–0.968), and wasting (RR, 0.994; 95% CI, 0.993–0.995).³⁸ Maternal stunting
329 increases the risk of giving birth to children who are small-for-gestational age (SGA). This is of

330 particular concern because multiple studies show that birth size affects development ³⁹.
331 Additionally, Black et al. estimated that one-fifth of all stunting originates in the fetal period,
332 resulting from children born SGA, which perpetuates the cycle of stunting that occurs in many
333 LMICs. Nutritional interventions focused on adolescent women can reduce risks of
334 complications in pregnancy and improve the fetal growth and development for their future
335 children. Providing iron-rich school meals has been shown to reduce iron-deficiency anemia,
336 which increases risk maternal mortality, low birthweight, and perinatal mortality.³⁹

337 Multiple micronutrient interventions are another strategy that has been successful in
338 improving height and weight gains among both male and female children. These studies, which
339 ranged in length, provided the daily-recommended allowance of iron, zinc, vitamin A, folic acid,
340 and B vitamins to children ranging from 6 months to 11 years in Latin America, Africa, and
341 Asia.⁴⁰ While effective, multiple micronutrient are difficult to carry out on a large scale because
342 of high costs. Nutritional-sensitive interventions, or programs that address the underlying causes
343 of malnutrition and improve the effects of nutrition-specific interventions, may be a more
344 effective long-term strategy to improve nutrition throughout a lifecycle. Biofortification of crops
345 is one promising strategy to help provide a daily dose of micronutrients and prevent
346 deficiencies.⁴¹ Further research is necessary to evaluate the impact of biofortification on
347 stunting. Continued promotion of schooling, for boys and girls, is also an important strategy for
348 the nutritional health of the next generation. Maternal schooling level has been positively
349 associated with child nutritional status.⁴²

350

351 This study adds to the growing literature that recovery from stunting is possible and common
352 and that the predictive capacity of growth at two years, while an important risk indicator for poor

353 outcomes, is not as tightly associated with adult height as is commonly held. More research
354 needs to be done to identify what promotes catch-up linear growth in late childhood. Some
355 studies have suggested that the major predictor of adult stature is the severity of stunting in
356 infancy, though this study did not find that.²⁷ The first 2 years of life are crucial for growth and
357 development, and efforts should continue to make sure that every child receives proper nutrition
358 and care during this critical time. However, a child who is stunted at 24 months should not be
359 viewed as stunted- in stature or cognition-for life. Throughout childhood and adolescence
360 nutritional programs have value to allow stunted children to achieve full genetic height potential
361 by the time they become adults, and to reduce stunting and associated morbidities in future
362 generations.

363

364 Financial Support

365 This work was supported by NIH grant K01-TW05717, the Bill and Melinda Gates Foundation
366 OPP1066146, and from the Sherrilyn and Ken Fischer Center for Environmental Infectious
367 Diseases at the Johns Hopkins School of Medicine to MK. The Parker Huang Undergraduate
368 Travel Fellowship from Yale University, which funded JS while working on this project and the
369 Leitman Award from the Johns Hopkins Center for Global Health (RK).

370

371 Disclosures

372 None

373

374

375

376

377 List of Authors Information

378 Julia Schwarz, Icahn School of Medicine 1 Gustave L. Levy Pl, New York, NY 10029;

379 Julia.Schwarz@icahn.mssm.edu

380

381 Pablo Peñataro Yori, Johns Hopkins Bloomberg School of Public 615 N. Wolfe Street Room

382 E5532 Baltimore, Maryland 21205; pyori@jhu.edu

383

384 William K. Pan, Duke University 227 Trent Hall, 310 Trent Drive, Durham, NC 27708;

385 william.pan@duke.edu

386

387 Maribel Paredes Olortegui, Asociación Benéfica Prisma Calle Ramirez Hurtado, 622 Iquitos –

388 Perú; mparedeso@prisma.org.pe

389

390 Robert Klapheke, UC San Diego School of Medicine 9500 Gilman Drive La Joalla, CA 92093;

391 rklaphe1@jhmi.edu

392

393 Margaret N. Kosek, Johns Hopkins Bloomberg School of Public Health 615 N Wolfe St

394 Baltimore, MD 21205; mkosek@jhu.edu

395

396 References:

397

- 398 1. Grantham-McGregor S, Cheung YB, Cueto S, Glewwe P, Richter L, Strupp B,
399 International Child Development Steering G, 2007. Developmental potential in the
400 first 5 years for children in developing countries. *Lancet* 369: 60-70.
- 401 2. Walker SP, Wachs TD, Gardner JM, Lozoff B, Wasserman GA, Pollitt E, Carter JA,
402 International Child Development Steering G, 2007. Child development: risk factors
403 for adverse outcomes in developing countries. *Lancet* 369: 145-57.
- 404 3. Woldehanna T, Behrman JR, Araya MW, 2017. The effect of early childhood stunting
405 on children's cognitive achievements: Evidence from young lives Ethiopia. *Ethiop J*
406 *Health Dev* 31: 75-84.
- 407 4. Hoddinott J, Behrman JR, Maluccio JA, Melgar P, Quisumbing AR, Ramirez-Zea M,
408 Stein AD, Yount KM, Martorell R, 2013. Adult consequences of growth failure in
409 early childhood. *Am J Clin Nutr* 98: 1170-8.
- 410 5. Victora CG, Adair L, Fall C, Hallal PC, Martorell R, Richter L, Sachdev HS, Maternal,
411 Child Undernutrition Study G, 2008. Maternal and child undernutrition:
412 consequences for adult health and human capital. *Lancet* 371: 340-57.
- 413 6. Akachi Y, Canning D, 2015. Inferring the economic standard of living and health
414 from cohort height: Evidence from modern populations in developing countries.
415 *Econ Hum Biol* 19: 114-28.
- 416 7. de Onis M, Branca F, 2016. Childhood stunting: a global perspective. *Matern Child*
417 *Nutr* 12 Suppl 1: 12-26.
- 418 8. de Onis M, Blossner M, Borghi E, 2012. Prevalence and trends of stunting among
419 pre-school children, 1990-2020. *Public Health Nutr* 15: 142-8.
- 420 9. Victora CG, de Onis M, Hallal PC, Blossner M, Shrimpton R, 2010. Worldwide timing
421 of growth faltering: revisiting implications for interventions. *Pediatrics* 125: e473-
422 80.
- 423 10. Svehors P, Rahman A, Ekstrom EC, Khan AI, Lindstrom E, Persson LA, Ekholm Selling
424 K, 2016. Stunted at 10 Years. Linear Growth Trajectories and Stunting from Birth to
425 Pre-Adolescence in a Rural Bangladeshi Cohort. *PLoS One* 11: e0149700.
- 426 11. Lundeen EA, Behrman JR, Crookston BT, Dearden KA, Engle P, Georgiadis A, Penny
427 ME, Stein AD, Young Lives D, Consequences of Child Growth Project T, 2014. Growth
428 faltering and recovery in children aged 1-8 years in four low- and middle-income
429 countries: Young Lives. *Public Health Nutr* 17: 2131-7.
- 430 12. Coly AN, Milet J, Diallo A, Ndiaye T, Benefice E, Simondon F, Wade S, Simondon KB,
431 2006. Preschool stunting, adolescent migration, catch-up growth, and adult height in
432 young senegalese men and women of rural origin. *J Nutr* 136: 2412-20.
- 433 13. Tanner JM, Davies PS, 1985. Clinical longitudinal standards for height and height
434 velocity for North American children. *J Pediatr* 107: 317-29.
- 435 14. Tanner JM, Whitehouse RH, Takaishi M, 1966. Standards from birth to maturity for
436 height, weight, height velocity, and weight velocity: British children, 1965. II. *Arch*
437 *Dis Child* 41: 613-35.

- 438 15. Yori PP, Lee G, Olortegui MP, Chavez CB, Flores JT, Vasquez AO, Burga R, Pinedo SR,
439 Asayag CR, Black RE, Caulfield LE, Kosek M, 2014. Santa Clara de Nanay: the MAL-ED
440 cohort in Peru. *Clin Infect Dis* 59 Suppl 4: S310-6.
- 441 16. Kosek M, Yori PP, Pan WK, Olortegui MP, Gilman RH, Perez J, Chavez CB, Sanchez
442 GM, Burga R, Hall E, 2008. Epidemiology of highly endemic multiply antibiotic-
443 resistant shigellosis in children in the Peruvian Amazon. *Pediatrics* 122: e541-9.
- 444 17. Lee G, Yori P, Olortegui MP, Pan W, Caulfield L, Gilman RH, Sanders JW, Delgado HS,
445 Kosek M, 2012. Comparative effects of vivax malaria, fever and diarrhoea on child
446 growth. *Int J Epidemiol* 41: 531-9.
- 447 18. Kaprio J, Rimpela A, Winter T, Viken RJ, Rimpela M, Rose RJ, 1995. Common genetic
448 influences on BMI and age at menarche. *Hum Biol* 67: 739-53.
- 449 19. E dLR, 2000. Age at Puberty of Girls and Boys in France: Measurements from a
450 Survey on Adolescent
451 Sexuality. *Population* 12: 51-79.
- 452 20. Status WECOP, 1995. Physical Status: the Use and Interpretation of Anthropometry.
453 Geneva: WHO.
- 454 21. Prentice AM, Ward KA, Goldberg GR, Jarjou LM, Moore SE, Fulford AJ, Prentice A,
455 2013. Critical windows for nutritional interventions against stunting. *Am J Clin Nutr*
456 97: 911-8.
- 457 22. Prentice A, Ginty F, Stear SJ, Jones SC, Laskey MA, Cole TJ, 2005. Calcium
458 supplementation increases stature and bone mineral mass of 16- to 18-year-old
459 boys. *J Clin Endocrinol Metab* 90: 3153-61.
- 460 23. Prentice A, Dibba B, Sawo Y, Cole TJ, 2012. The effect of prepubertal calcium
461 carbonate supplementation on the age of peak height velocity in Gambian
462 adolescents. *Am J Clin Nutr* 96: 1042-50.
- 463 24. Tumilowicz A, Beal T, Neufeld LM, Frongillo EA, 2019. Perspective: Challenges in Use
464 of Adolescent Anthropometry for Understanding the Burden of Malnutrition. *Adv*
465 *Nutr* 10: 563-575.
- 466 25. Walker SP, Grantham-McGregor SM, Himes JH, Powell CA, Chang SM, 1996. Early
467 childhood supplementation does not benefit the long-term growth of stunted
468 children in Jamaica. *J Nutr* 126: 3017-24.
- 469 26. Satyanarayana K, Nadamuni Naidu A, Swaminathan MC, Narasinga Rao BS, 1981.
470 Effect of nutritional deprivation in early childhood on later growth--a community
471 study without intervention. *Am J Clin Nutr* 34: 1636-7.
- 472 27. Crookston BT, Penny ME, Alder SC, Dickerson TT, Merrill RM, Stanford JB, Porucznik
473 CA, Dearden KA, 2010. Children who recover from early stunting and children who
474 are not stunted demonstrate similar levels of cognition. *J Nutr* 140: 1996-2001.
- 475 28. Sterling R, Miranda JJ, Gilman RH, Cabrera L, Sterling CR, Bern C, Checkley W, 2012.
476 Early anthropometric indices predict short stature and overweight status in a
477 cohort of Peruvians in early adolescence. *Am J Phys Anthropol* 148: 451-61.
- 478 29. Kahl H, Schaffrath Rosario A, Schlaud M, 2007. [Sexual maturation of children and
479 adolescents in Germany. Results of the German Health Interview and Examination
480 Survey for Children and Adolescents (KiGGS)]. *Bundesgesundheitsblatt*
481 *Gesundheitsforschung Gesundheitsschutz* 50: 677-85.

- 482 30. Leenstra T, Petersen LT, Kariuki SK, Oloo AJ, Kager PA, ter Kuile FO, 2005.
483 Prevalence and severity of malnutrition and age at menarche; cross-sectional
484 studies in adolescent schoolgirls in western Kenya. *Eur J Clin Nutr* 59: 41-8.
- 485 31. Belachew T, Hadley C, Lindstrom D, Getachew Y, Duchateau L, Kolsteren P, 2011.
486 Food insecurity and age at menarche among adolescent girls in Jimma Zone
487 Southwest Ethiopia: a longitudinal study. *Reprod Biol Endocrinol* 9: 125.
- 488 32. Golden MH, 1994. Is complete catch-up possible for stunted malnourished children?
489 *Eur J Clin Nutr* 48 Suppl 1: S58-70; discussion S71.
- 490 33. Teilmann G, Pedersen CB, Skakkebaek NE, Jensen TK, 2006. Increased risk of
491 precocious puberty in internationally adopted children in Denmark. *Pediatrics* 118:
492 e391-9.
- 493 34. Stein AD, Wang M, Martorell R, Norris SA, Adair LS, Bas I, Sachdev HS, Bhargava SK,
494 Fall CH, Gigante DP, Victora CG, Cohorts G, 2010. Growth patterns in early childhood
495 and final attained stature: data from five birth cohorts from low- and middle-income
496 countries. *Am J Hum Biol* 22: 353-9.
- 497 35. Berkman DS, Lescano AG, Gilman RH, Lopez SL, Black MM, 2002. Effects of stunting,
498 diarrhoeal disease, and parasitic infection during infancy on cognition in late
499 childhood: a follow-up study. *Lancet* 359: 564-71.
- 500 36. Mendez MA, Adair LS, 1999. Severity and timing of stunting in the first two years of
501 life affect performance on cognitive tests in late childhood. *J Nutr* 129: 1555-62.
- 502 37. Crookston BT, Schott W, Cueto S, Dearden KA, Engle P, Georgiadis A, Lundeen EA,
503 Penny ME, Stein AD, Behrman JR, 2013. Postinfancy growth, schooling, and cognitive
504 achievement: Young Lives. *Am J Clin Nutr* 98: 1555-63.
- 505 38. Ozaltin E, Hill K, Subramanian SV, 2010. Association of maternal stature with
506 offspring mortality, underweight, and stunting in low- to middle-income countries.
507 *JAMA* 303: 1507-16.
- 508 39. Black RE, Victora CG, Walker SP, Bhutta ZA, Christian P, de Onis M, Ezzati M,
509 Grantham-McGregor S, Katz J, Martorell R, Uauy R, Maternal, Child Nutrition Study G,
510 2013. Maternal and child undernutrition and overweight in low-income and middle-
511 income countries. *Lancet* 382: 427-51.
- 512 40. Ramakrishnan U, Aburto N, McCabe G, Martorell R, 2004. Multimicronutrient
513 interventions but not vitamin a or iron interventions alone improve child growth:
514 results of 3 meta-analyses. *J Nutr* 134: 2592-602.
- 515 41. Bhutta ZA, Das JK, Rizvi A, Gaffey MF, Walker N, Horton S, Webb P, Lartey A, Black
516 RE, Lancet Nutrition Interventions Review G, Maternal, Child Nutrition Study G,
517 2013. Evidence-based interventions for improvement of maternal and child
518 nutrition: what can be done and at what cost? *Lancet* 382: 452-77.
- 519 42. Semba RD, de Pee S, Sun K, Sari M, Akhter N, Bloem MW, 2008. Effect of parental
520 formal education on risk of child stunting in Indonesia and Bangladesh: a cross-
521 sectional study. *Lancet* 371: 322-8.
- 522

523

524 Tables:

525

526 Table 1: Comparison of participants with follow-up and those lost to follow up by gender,
 527 average HAZ scores, percent stunted at 24 months, average maternal education in years and per
 528 capita monthly income.

	Adult follow-up	Lost to follow-up	p-value
	n=104	n=95	
Males (%)	45	52	0.32
Average HAZ Score	-1.99	-2.17	0.15
Stunted at 24 months (%)	53	60	0.32
	n=99	n=82	
Average Maternal Education in Years	8.04	8.57	0.23
Average Monthly Income	n=102 31.52	n=89 31.45	0.98
Average Age at Puberty	Males n=47 12.64	Females n=57 11.79	

529 HAZ = height-for-age Z-Scores.

530

531 Table 2: The mean HAZ at puberty, mean weight at puberty and mean BMI at puberty is shown
532 by categorization of children who had a HAZ<-2, those who had a HAZ between -2 and -1, and

	Mean HAZ post puberty	SD	Mean Weight post Puberty (kg)	SD	Mean BMI post Puberty (kg/m²)	SD
HAZ<-2 at 24 months (n=55)	-1.75	0.76	50.0	8.6	21.77	3.3
HAZ -2 to -1 at 24 months (n=36)	-0.96	0.70	49.6	6.4	20.55	2.6
HAZ >-1 at 24 months (n=13)	-0.24	0.82	59.1	11.1	23.1	3.7

533 those with a HAZ > -1 at 24 months.

534

535

536

537 BMI=Body Mass Index. Puberty was assessed by maternal report of the onset of menses (for
538 female subjects) or the age of voice change (for male subjects).

539

540 Table 3: Prevalence of stunting and recovery from 24 months to post-puberty in males and
 541 females shown by categorization of children who recovered from stunting at 24 months, fully
 542 recovered from stunting at 24 months, those who remained stunted, those that became stunted in
 543 late childhood, and those that were neither stunted at 24 months or at their measurement
 544 following puberty.
 545

	Male (%)	Male (n)	Female (%)	Female (n)	P-Value	Total(%)	Total (n)
HAZ<-2 at 24 months	66%	31	42%	24	.015	53%	55
Recovered children (HAZ <-2.00 @24 months and >-2.00 post-puberty)	68%	21	54%	13	.293	62%	34
Fully Recovered children (HAZ <-2.00 @24 months and >- 1.00 post-puberty)	23%	7	8%	2	.140	16%	5
Stunted throughout (HAZ <-2.00 @24 month and HAZ <-2.00 post-puberty)	32%	10	46%	11	.293	38%	21
HAZ>-2 at 24 months	34%	16	58%	33	.015	47%	49
Late stunting (HAZ >-2.00 @24 months and HAZ<-2.00 post-puberty)	6%	1	6%	2	1.00	6%	3
Healthy Throughout (HAZ>-2.00 @ 24 months and Post-Puberty)	94%	15	94%	31	1.00	94%	46
HAZ<-2 Post-Puberty	23%	11	23%	13	1.00	23%	24

546
547

548 Table 4: Unadjusted odds ratios of stunted and non-stunted children who will be stunted post-
549 puberty (PP)

	OR	CI	n	P-Value
Odds Male HAZ<-2 @ 24 m Stunted PP Compared to Non-Stunted Male @ 24 m	9.5	2.6, 34.3	47	0.006
Odds Female HAZ<-2 @ 24 m Stunted PP Compared to Non-Stunted Female @ 24 m	13.1	2.5, 67.6	57	0.002
Odds Child HAZ<-2 @ 24 m Stunted PP Compared to Non-Stunted Child @ 24 m	7.1	.82, 61.9	104	0.074

550

551

552 Table 5: The average age of puberty, as delineated by menarche in girls and voice change in boys
 553 is shown by categorization of children who recovered from stunting at 24 months, those who
 554 remained stunted, those that became stunted in late childhood, and those that were neither
 555 stunted at 24 months or at their measurement following puberty.
 556


	Male Average Age of Puberty (Years)	SD	(n)	Female Average Age of Puberty (Years)	SD	(n)	Average Time Elapsed between Puberty and Final Measurement (Months)	SD	(n)
Recovered children HAZ <- 2.00 @24 months and >-2.00 post- puberty	12.28	0.90	21	11.85	1.07	14	20.03	18.21	35
Stunted throughout HAZ <-2.00 @24 month and HAZ <2.00 post- puberty	13.50	1.18	10	11.64	1.12	11	26.97	18.67	21
Late stunting HAZ >-2.00 @24 months and HAZ<-2.00 post- puberty	12	NA	1	11.50	2.12	2	38.37	20.51	3
Non-Stunted Throughout HAZ>-2.00 @ 24 months and post- puberty	12.60	0.99	15	11.84	1.13	30	22.90	14.87	45
Overall Cohort	12.64	1.08	47	11.79	1.10	57	23.20	17.11	104

557
 558

559 Figures:

560

561 Figure 1: Flow chart showing identification of study population.



562

563

564

565

566

567

568

569

570

571

572

573

574

575

576


577

578

579

580

581 Figure 2: A matched comparison of attained length at 24 months and attained height after
582 puberty. X- axis shows HAZ score at 24 months and Y axis shows the corresponding HAZ score
583 after puberty. Each point represents a single participant. The graph is broken into four quadrants
584 to delineate children who at puberty had recovered from stunting at 24 months, those that were
585 neither stunted at 24 months or at their measurement post-puberty, those that became stunted in
586 late childhood, and those who remained stunted.
587



588
589