Genomic surveillance revealed prevalence of unique SARS-CoV-2 variants

bearing mutation in the RdRp gene among Nevada patients

Paul D. Hartley[1,4*], Richard L. Tillett[7*], Yanji Xu[4,5], David P. AuCoin[3,4], Joel R. Sevinsky[6],

Andrew Gorzalski[2,4], Mark Pandori[2,4], Erin Buttery[8], Holly Hansen[8], Michael A. Picker[8],

Cyprian C. Rossetto[3,4*#], Subhash C. Verma[3,4*#]

[1] Nevada Genomics Center

[2] Nevada State Public Health Laboratory

[3] Department of Microbiology & Immunology,

University of Nevada, Reno School of Medicine

[4] University of Nevada, Reno

[5] Nevada Center for Bioinformatics

[6] Theiagen Consulting, LLC

[7] Nevada Institute of Personalized Medicine, University of Nevada, Las Vegas,

[8] Southern Nevada Public Health Laboratory of the Southern Nevada Health District,
Las Vegas
18

* These authors contributed equally to this work and are listed in alphabetical order

[#] To whom correspondence should be addressed:

Cyprian C. Rossetto crossetto@med.unr.edu, Subhash C. Verma

scverma@med.unr.edu

Running title: Unique RdRp variant of SARS-CoV-2 in Nevada patients

Keywords: SARS-CoV-2, COVID-19, genome enrichment, nsp12, RdRp, orf1b 314

ABSTRACT:

Background:

SARS-CoV-2 is the etiological agent of coronavirus disease 2019 (COVID-19), which has

a wide range of disease manifestations, including relatively mild respiratory illness to

severe disease requiring hospitalization. Patients with signs of COVID-19 (fever,

coughing, shortness of breath) were tested for the virus by targeting specific SARS-COV-

2 genomic loci via RT-PCR using RNA extracted from nasopharyngeal/nasal swabs.

However, this testing does not yield genetic information required for identifying viral

evolution or variants.

Methods:

We sequenced 200 specimens from positively identified cases in Nevada through our

robust protocol for sequencing SARS-CoV-2 genomes directly from patients

nasopharyngeal/nasal swabs. This protocol enabled identification of specific nucleotide

variants including those for D614G and clade defining mutations. Additionally, these

sequences were used to determine the phylogenetic relationships of SARS-CoV-2

genomes of public health importance occurring in the state of Nevada.

Results:

Our study reports the finding of a novel variant in the nsp12 (RdRp-RNA dependent RNA

Polymerase) protein at residue 323 (314aa of orf1b) to Phenylalanine (F) from Proline

(P), present in the original isolate of SARS-CoV-2 (Wuhan-1). This 323F variant is found

at a very high frequency in Northern Nevada suggesting for a founder effect. Structural

modeling of the RdRP with P323F mutation did not show any significant difference in

protein conformation but the phenotypic effect is unknown and an area of active investigation.

Conclusions:

Our results highlight the introduction and spread of specific SARS-CoV-2 variants at very high frequency within a distinct geographic location that is important for clinical and public health perspectives in understanding the evolution and transmission of SARS-CoV-2.

BACKGROUND:

Severe Acute Respiratory Syndrome coronavirus 2 (SARS-CoV-2), the cause of coronavirus disease 2019 (COVID-19), was first identified and reported in December 2019 in Wuhan, Hubei province, China [1-3]. RNA sequencing and phylogenetic analysis of a specimens taken during the initial outbreak in Wuhan determined that the virus is most closely related (89.1% nucleotide similarity) to a group of SARS-like coronaviruses (genus Betacoribavirus, subgenus Sarbecovirus) which had previously been identify in bats in China [2, 4]. Coronaviruses have a recent history as emerging infections, first SARS-CoV in 2002-2003, and Middle East respiratory syndrome coronavirus (MERS-CoV) in 2012, both zoonotic infections that cause severe respiratory illness in humans [2, 5-8] [2,5-8]. Unlike SARS-CoV and MERS-CoV which displayed limited global spread, SARS-CoV-2 has spread around the world within a few months. There are specific characteristics of SARS-CoV-2 which have facilitated the transmission, including infections that result in asymptomatic or mild disease, allowing for under-characterized transmission.

SARS-CoV-2 is an enveloped, positive single-stranded RNA virus. Detection of

SARS-CoV-2 in patients has primarily occurred using RT-qPCR to detect viral RNA from

respiratory specimens (primarily nasal, nasopharyngeal swabs). While RT-PCR results

can be quantified through determination of a cycle threshold (Ct) value for each sample

it does not yield sequence data leading to the description of genomic variants. To further

a study of such variants, and to better understand the epidemiology of the virus in the

state of Nevada, we developed a workflow that allowed us to sequence SARS-CoV-2

genomic RNA from patient swabs containing a broad range of viral loads. Of the

sequences of SARS-CoV-2 currently submitted to common database, several were

obtained after the virus had been passed in Vero cells [9, 10] and others came directly

from patient specimens [11, 12]. Certain data has suggested a potential for lab acquired

mutations following passage in cell culture [9, 13]. Specifically, a report of SARS-CoV-2

passage in Vero cells which resulted in a spontaneous 9 amino acid deletion within the

spike (S) protein that overlaps with furin cleavage site [13]. The loss of this site is

suggested to increase the viral entry into Vero cells [14]. For both research and

epidemiological purposes, sequencing of SARS-CoV-2 directly from patient specimens

not only reduces the possibility of lab-adapted mutations following passage in cell culture

but also reduces the time that would be spent growing the virus from the patient

specimens and subsequently also reduces handling larger amounts of infectious virus.

Additionally, one of the goals in developing an optimized SARS-CoV-2 NGS protocol was

to be able to generate adequate depth of coverage of the viral genome while minimizing

the sequencing of non-viral RNA which would allow for more specimens to be multiplexed

together during sequencing.

Our workflow employs a combination of RNA amplification, conversion into

Illumina-compatible sequencing libraries and enrichment of SARS-CoV-2 library

molecules prior to sequencing. Using this novel methodology, we sequenced SARS-

CoV-2 from a total of 200 patient specimens collected over a 3-month period originating

from Nevada. Of the 200 selected, 173 were sequenced with enough quality to be used

for determining SARS-CoV-2 nucleotide variants to perform further phylogenetic analysis

and study the viral epidemiology within the state of Nevada. Analysis of the data suggests

a specific epidemiological course for the local epidemic within Northern Nevada. This was

characterized by an initial observation of variants closely resembling isolates originating

directly from China or Europe. Subsequent to government-mandated period of restrictions

on business and social activity, we observed that a viral isolate not seen elsewhere in the

world emerged within Northern Nevada cases (nucleotide 14,407 and 14,408). This

isolate contains an amino acid change in residue P323L/F of the RdRp (nsp12).

Furthermore, we found that sampled viral isolates in Southern Nevada, unlike those in

Northern Nevada, closely resembled the makeup of the United States in general.

METHODS:

SARS-CoV-2 specimen and library preparation. Nasal and Nasopharyngeal swab

specimens were received at the Nevada State Public Health Lab (NSPHL) or Southern

Nevada Public Health Lab (SNPHL) and RNA extraction was completed using either a

QIAamp Viral RNA Mini Kit (QIAGEN) or Mag-Bind Viral DNA/RNA kit (Omega Biotek).

Specimens were tested for the presence of coronaviral RNA using FDA-approved kits

that employed RT-PCR to detect SARS-COV-2 RNA.

A set of 200 coronavirus positive specimens were selected for genome sequencing. Specimens were treated with DNase I (QIAGEN) for 30 minutes at room temperature and concentrated using RNeasy Minelute spin columns (QIAGEN) based on the manufacturer supplied protocol. These concentrated samples were converted into Illumina-compatible sequencing libraries with a QIAseq FX Single Cell RNA Library kit (QIAGEN). RNA samples were annealed to a 1:12.5 dilution of QIAseq FastSelect -HMR probes (QIAGEN) to reduce subsequent amplification of human ribosomal RNA. After treatment to remove trace DNA from the samples, a reverse transcription reaction was carried out using random hexamers. The synthesized DNA was ligated to one another, followed by isothermal linear amplification. Amplified DNA (1 μg) was enzymatically sheared to an average insert size of 300 bp, and Illumina-compatible dual-indexed sequencing adapters were ligated to the ends. Next, about 300 ng of adapter-ligated sample was amplified with 6 cycles of PCR with KAPA HiFi HotStart polymerase (Roche Sequencing Solutions). Enrichment of library molecules containing SARS-CoV-2 sequence was conducted with a myBaits kit and coronavirus-specific biotinylated probes (Arbor Biosciences). Each enrichment used 500 ng of PCR-amplified DNA, was carried out based on manufacturer instructions at a hybridization temperature of 65° for 16 hours, and was completed with 8-16 cycles of PCR using KAPA HiFi HotStart polymerase. Samples were sequenced using an Illumina Next-seq mid-output (2 x 75). The generated FASTQ files from the sequencing reaction were analyzed as described below. The data files are available at GISAID, NCBI under the https://www.ncbi.nlm.nih.gov/bioproject/657893

Computational analysis. Sequence pair libraries were trimmed using Trimmomatic,

version 0.39 and adapter-clipping setting "2:30:10:2:keepBothReads" [15]. Read pairs

were aligned against the Wuhan reference genome (NC_045512.2) by Bowtie 2, version

2.3.5, local alignment [16]. PCR optical duplicates were removed via Picard

MarkDuplicates [17].

Variants were called using Freebayes, version 1.0.2, with ploidy set to 1, minimum

allele frequency 0.75, and minimum depth of 4 [18]. No variants were called in the first

200 bp and final 63 bp of the COVID-19 genome. High-quality variant sites were selected

where site "QUAL > 20" using *vcffilter*, VCFlib version 1.0.0_rc2 [19]. Individual genomes

were reconstructed by their filter-passing variants using *bcftools consensus* and only

where aligned coverage depth ≥ 4; bases with coverage below four are reported as

unknown (Ns) [20].

A set of 3,644 complete, high-coverage SARS-CoV-2 genomes reported in the

July 15, 2020 Nextstrain.org global subsample and metadata were obtained from GISAID

and combined with our own samples to determine their phylogenetic placement [21, 22].

Four of the global samples were set aside after screening for unexpected FASTA

characters. The combined sets of global and Nevadan samples were aligned together,

with metadata, by the *augur* phylodynamic pipelines of the *ncov* build of the *nextstrain*

command-line tool, version 2.0.0.post1 [23].

Cumulative frequency of D614G, clades (19A, 19B, 20A, 20B, 20C), and P323L/F

were calculated at each time point based on the total number of specimens up to the

indicated date. Plots and pie charts were generated using GraphPad Prism (version 8).

162 **nsp12 protein modeling.** Sequence of nsp12 (RdRp) protein for SARS-CoV-2

163 (YP_009725307.0) was retrieved from NCBI protein database and 3D model was

164 structured based on a previously published report (PDB ID: 6XEZ [24]. In addition to

165 nsp12 (chain A), the model also contains nsp7 (chain C), nsp8 (chain B and D), nsp13

166 (chain E and F), ligands (ZN, Mg) and RNA template and product strands. Mutational

167 changes to residue 323 within nsp12 were performed using PyMol Molecular Graphics

168 System (version 2.0, Schrödinger LLC). The original proline (P) was mutated to either

169 leucine (L) or phenylalanine (F) as indicated, these residues along with residues

170 containing side chains within 5 Å of P323L/F are shown as sticks. The rotamers for each

171 P323L/F were assessed and those with least rotational strain and steric hindrance were

172 used to generate the final image. To determine any NCBI deposited sequences which

173 contain the P323F variant, standard protein BLAST from the BLASTp suite was used to

174 find nsp12 protein sequences which contained FSTVFP**F**TSFGP (P323F is bold and

175 underline) from full length SARS-CoV-2 genomes. The P323F amino acid changes were

176 confirmed with the NCBI deposited nucleotide sequences.

178 **RESULTS:**

179 **RNA-seq workflow and assembly of SARS-CoV-2 genomes.**

180  A total of 200 SARS-CoV-2 positive specimens collected in Nevada from March

181 6th to June 5th were randomly selected to have their viral genomes sequenced for variant

182 analysis and subsequent epidemiological studies (Fig 1A and Materials & Methods). Of

183 the sequenced specimens, 173 had >90% coverage and sufficient depth to accurately

184 call those genomic positions with variants (Fig. 1B). An alignment of SARS-CoV-2

genomes from these specimens is presented as supplemental data (Supplemental Fig.

S1) showing 173 specimens with over 90% coverage. These 173 specimens represented

133 patient specimens from Northern Nevada (including Washoe County of which Reno

is the major city, the Carson-Tahoe area, and other northern, rural counties), 40 patient

specimens from Southern Nevada (Clark county which encompasses Las Vegas).

Nucleotide similarity and variants were determined and used to measure the phylogenetic

relationships (Supplemental Fig. S2). The combined nucleotide diversity across the entire

SARS-CoV-2 genome for the Nevada specimens is shown in figure 1D, along with the

genomic areas that were assessed for change in frequency corresponding to amino acids

D614G, P323L/F and nucleotide 379.

During the sequencing analysis we also examined the correlation between Ct

values from the diagnostic RT-PCR and percentage coverage of the viral genome to

determine the performance and robustness of our sequencing method in relation to

available viral RNA in a specimen of a given Ct (Fig. 1C). In general, a Ct value less than

31 resulted in at least 90% coverage to the SARS-CoV-2 genome. Importantly, our in-

house developed method for viral genome enrichment and sequencing directly from the

patient's specimens (nasal and nasopharyngeal swabs) was robust and yielded

sequences covering over 90% of the genome even in samples having very high Ct (~40)

of viral genome detection. This is highly significant and shows the power of our workflow

in sequencing of SARS-CoV-2 genome from a spectrum of samples including the ones

having inadequate amounts of specimen (due to the variability in collection) or lower viral

loads in nasal secretions. Consequently, our sequencing protocol avoids any molecular

epidemiological bias, which may get acquired through cell culture-based amplification

especially in those specimens with high Ct (low viral load) as our method eliminates the

need of virus culture.

Prevalence of amino acid variant D614G of SARS-CoV-2 spike protein in specimens

collected in Nevada. Earlier studies have revealed the emergence, spread and potential

importance of an alteration, D614G (genomic change at 23403A>G), of the spike protein

[25]. This missense mutation has become a clade-distinguishing locus that differentiates

viral isolates originating in Asia from those that have emerged from Europe. A total of 173

cases were analyzed to determine the number and relative proportion of the specimens

which carried the D614G spike protein variant in Nevada. The cumulative frequency for

D614 and G614 were plotted from March 6th to June 5th (Fig. 2A). Specimens from the

beginning of March represent the earliest known cases in Nevada, and of the 14

specimens sequenced during this time period (March 6-March 15) D614 was the

predominant variant. This shifted from March to June with an increasing frequency of the

G614 allele. The trend for specimens originating from either Northern Nevada (N-NV) and

Southern Nevada (S-NV) both show a higher frequency of G614 (Fig. 2B). We used a

subsampling of sequence data from Nextstrain.org to assess the frequency of D614G in

the United States and globally during the same time period (March 6th to June 5th) (Fig.

2B and 2C). The global trend by continent of D614G is also similar, with G614 at a higher

frequency, the one noted exception is in Asia, where D614 and G614 continue to exist in

equal proportions (Fig. 2C).

Frequency of SARS-CoV-2 clades in Nevada. Worldwide, there are currently 5 main

clades (19A, 19B, 20A, 20B, 20C) of SARS-CoV-2 differentiated based on specific

nucleotide profiles in the Year-letter scheme of https://clades.nextstrain.org. Clade 19A

and 19B are defined by C8782T and T28144C respectively. 20A is a derivative of 19A

and contains mutations C3037T, C14408T and A23403G (resulting in D614G). 20B is

defined by mutations G28881A, G28882A and G28883C, and 20C contains C1059T and

G25563T [26].

To assess the introduction and spread of the clades in Nevada the cumulative

frequency for the clades were plotted from March 6th to June 5th (Fig. 3A). The earliest

sequenced specimens from Nevada were collected in the beginning of March (March 6-

March 15) and are predominantly from clades 19A and 19B. Additional sequenced

specimens collected from March to June revealed a shift to a higher frequency of 20C

(Fig. 3A). We performed phylogenetic reconstruction of the Nevada specimens and

differentiated the clades on the circular dendrogram by color (Fig. 3B). There were

discordant trends in the dominant clade for specimens originating from either Northern

Nevada (N-NV) and Southern Nevada (S-NV) (Fig. 3C and 3D).   Specimens from

Northern Nevada (Washoe County, Carson-Tahoe, and other counties) showed a

prevalence of 20C, while the Southern Nevada specimens from Clark county had a larger

proportion of 20A (Fig. 3C and Supplemental Fig. S3a). We used a subsampling of

Nextstrain.org data to assess the frequency of clades in the United States and globally

by continent during the same time period (March 6th to June 5th). The dominant clade in

the United States was 20C, similar to the frequency seen in the total Nevada samples (N-

NV and S-NV) (Fig. 3A and 3C).   The global clade distributions were variable in areas

outside of Asia while clades 19A and 19B are noted to be more prevalent in Asia. (Fig.

3D).

Prevalence of amino acid variant P323L/F of SARS-CoV-2 nsp12 (RdRp) in Nevada.

Analysis of sequencing data revealed a novel observation for our specimens at bases 14,407 and 14,408 which results in a change at residue 323 in nsp12 (RdRp). For the Wuhan isolate at 14,407 and 14,408 there is CC for proline (P), the variants have CT for leucine (P323L) and TT for phenylalanine (P323F). To assess the introduction and spread of P323L/F in Nevada, the cumulative frequency of P323, L323 and F323 were plotted from March 6th to June 5th (Fig. 3A). Nevada specimens from the beginning of March (March 6-March 15) show P323 was the predominant variant. As additional specimens were collected and sequenced from March to June there was a shift to a higher frequency of L323 and F323 (Fig. 4A). We performed phylogenetic reconstruction of the Nevada specimens and noted the P323L/F variants on the circular dendrogram with the indicated colors (Fig. 4B). Interestingly, analysis of the Northern Nevada and Southern Nevada specimen showed very different dominant variants (Fig. 4B). In Northern Nevada the F323 was more prevalent, and in Southern Nevada L323 was more prevalent. We used a subsampling of Nextstrain.org data to assess the frequency of P323L/F in the United States and globally during the same time period (March 6th to June 5th). P323 is the predominant variant in Asia, while L323 is more prevalent in other areas of the world and F323 is only appreciably noted in North America (Fig. 4D and 4E).

DISCUSSION:

We have developed a novel method which combines specific depletion and enrichment strategies that results in efficient SARS-CoV-2 RNA-seq with high genome coverage and depth. An advantage of this protocol is that it generates sequence data

directly from swab specimens without the need to passage the virus in cell culture thereby

reducing the handling of infectious material and induction of culture-acquired mutations.

Another obstacle in sequencing directly from swab specimens is that most FDA-approved

commercially available RNA extraction kits are specifically optimized to recover low

amounts of total nucleic acids, include carrier polyA RNA that could be convertible into

sequence able molecules, as has been observed previously with RNA-seq of Lassa- or

Ebola-positive clinical specimens [27].

The workflow incorporates amplification of low-abundance RNA into micrograms

of DNA, followed by conversion from a fraction of the DNA into Illumina-compatible

sequencing libraries and enrichment of these libraries for SARS-CoV-2 sequences. In

addition, during the reverse transcription step a reagent was incorporated to reduce the

subsequent amplification of host ribosomal RNA. This approach is robust in that it

converts low amounts of RNA into microgram quantities of DNA representative of all the

RNA species (aside of rRNA) present in the specimen. This DNA can be stored

indefinitely to be interrogated by multiple techniques at a later date. Additionally, RNA

amplification is likely less sensitive to low viral abundance compared to RT-PCR. Finally,

the use of probes to enrich for coronavirus-specific sequencing library molecules is less

sensitive to variants compared to tiling PCR amplicon approaches [28-32].

The data herein implicate that early in the pandemic, before the "stay-at-home"

order on April 1st, there were multiple introductions of SARS-CoV-2 into the state of

Nevada. From April 1st to the beginning of June Nevada experienced a period of semi-

isolation, as the casinos and most hotels shut down, tourism and travel to the state

essentially stopped. Because of the stay-at-home order and social distancing measures

put in place there was less mobility of people within and between states. It is possible that

these measures, compounded by potential inherent transmission variability of some viral

isolates, influenced the change in the frequency of D614G, clades and P323L/F that we

noted during this time period within Nevada. In addition, we also found 379C>A had a

high prevalence in our study specimens compared to the subsampling of sequences from

the United States and international (supplemental Fig. S4). This is a synonymous

mutation in nsp1, hence the biological relevance of this nucleotide variant remains to be

elucidated.

Of the 14,885 complete SARS-CoV-2 genomes currently available (as of August

14, 2020) in NCBI there are only 6 genomes that have the P323F variant (accession

number: MT706208, LR860619, MT345877, MT627429, MT810889, MT811171). In this

study 62 of the 133 specimens from Northern Nevada contain P323F. That is 46% of

specimens from Northern Nevada contain P323F compared to 0.04% of NCBI deposited

SARS-CoV-2 isolates. We performed structural modeling of the P323L/F variation and

did not find any significant change to the nsp12 confirmation with either P323L or P323F

(Supplemental Fig. S5), therefore this variant is most likely a neutral mutation and does

not confer either a fitness advantage or disadvantage to transmission or pathogenicity of

SARS-CoV-2.

We find that the overall trend for D614G in Nevada during this time period was

similar to what was observed in other states and internationally, with the exception of

within Asia where the D614 allele had originated. We noted that there were differences

between Northern Nevada and Southern Nevada. In Northern Nevada clade 20C and

F323 were more frequent, while during this same time period in Southern Nevada clade

20A and L323 were more prevalent. These data indicate that there were distinct genomic

profiles of the SARS-CoV-2 viruses that were circulating in these populations during the

initial months of the pandemic while stay-at-home order were in place to help prevent

transmission of the virus.

CONCLUSIONS:

SARS-COV-2 is currently a global pandemic on a scale that has not been experienced

since the 1918 influenza pandemic. Viral genomes accumulate nucleotide variations while

passing in the human population and these mutations may confer phenotypic differences

including immune response and anti-viral drug resistance. We have developed a protocol

to enrich SARS-CoV-2 RNA directly from patient nasal/nasopharyngeal specimens that

does not rely on amplicon based sequencing strategies nor the need of passing the virus

into tissue culture and reducing the possible introduction of lab-adapted mutations. These

specimens were subjected to next generation sequencing (NGS) to determine the

phylogenetic relationship of SARS-CoV-2 sequences within Nevada during the early

months (March to June) of the pandemic. Our study identified a very rare mutation in the

RdRp protein (P323F) found to occur at a very high frequency among patients of Nevada.

This may be important for clinical and public health perspectives in understanding the

transmission and evolution of SARS-CoV-2 as it circulates in humans.

DATA AVAILABILITY:

All sequences are available at bio project:

https://www.ncbi.nlm.nih.gov/bioproject/657893. All reported data are deposited and

available at GISAID with the following accession numbers: EPI_ISL_515409,

EPI_ISL_515407, EPI_ISL_515408, EPI_ISL_515375, EPI_ISL_515376,

EPI_ISL_515373, EPI_ISL_515374, EPI_ISL_515371, EPI_ISL_515372,

EPI_ISL_515370, EPI_ISL_515416, EPI_ISL_515417, EPI_ISL_515414,

EPI_ISL_515415, EPI_ISL_515379, EPI_ISL_515412, EPI_ISL_515413,

EPI_ISL_515377, EPI_ISL_515410, EPI_ISL_515378, EPI_ISL_515411,

EPI_ISL_515364, EPI_ISL_515365, EPI_ISL_515362, EPI_ISL_515363,

EPI_ISL_515360, EPI_ISL_515361, EPI_ISL_515405, EPI_ISL_515406,

EPI_ISL_515403, EPI_ISL_515404, EPI_ISL_515368, EPI_ISL_515401,

EPI_ISL_515369, EPI_ISL_515402, EPI_ISL_515366, EPI_ISL_515367,

EPI_ISL_515400, EPI_ISL_515353, EPI_ISL_515354, EPI_ISL_515351,

EPI_ISL_515352, EPI_ISL_515350, EPI_ISL_515359, EPI_ISL_515357,

EPI_ISL_515358, EPI_ISL_515355, EPI_ISL_515356, EPI_ISL_515342,

EPI_ISL_515343, EPI_ISL_515340, EPI_ISL_515461, EPI_ISL_515341,

EPI_ISL_515462, EPI_ISL_515460, EPI_ISL_515348, EPI_ISL_515349,

EPI_ISL_515346, EPI_ISL_515347, EPI_ISL_515344, EPI_ISL_515345,

EPI_ISL_515331, EPI_ISL_515298, EPI_ISL_515452, EPI_ISL_515332,

EPI_ISL_515299, EPI_ISL_515453, EPI_ISL_515296, EPI_ISL_515450,

EPI_ISL_515330, EPI_ISL_515297, EPI_ISL_515451, EPI_ISL_515294,

EPI_ISL_515295, EPI_ISL_515293, EPI_ISL_515339, EPI_ISL_515337,

EPI_ISL_515458, EPI_ISL_515338, EPI_ISL_515459, EPI_ISL_515335,

EPI_ISL_515456, EPI_ISL_515336, EPI_ISL_515457, EPI_ISL_515333,

EPI_ISL_515454, EPI_ISL_515334, EPI_ISL_515455, EPI_ISL_515319,

EPI_ISL_515320, EPI_ISL_515441, EPI_ISL_515321, EPI_ISL_515442,

EPI_ISL_515440, EPI_ISL_515328, EPI_ISL_515449, EPI_ISL_515329,

EPI_ISL_515326, EPI_ISL_515447, EPI_ISL_515327, EPI_ISL_515448,

EPI_ISL_515324, EPI_ISL_515445, EPI_ISL_515325, EPI_ISL_515446,

EPI_ISL_515322, EPI_ISL_515443, EPI_ISL_515323, EPI_ISL_515444,

EPI_ISL_515308, EPI_ISL_515429, EPI_ISL_515309, EPI_ISL_515390,

EPI_ISL_515397, EPI_ISL_515430, EPI_ISL_515310, EPI_ISL_515398,

EPI_ISL_515431, EPI_ISL_515395, EPI_ISL_515396, EPI_ISL_515393,

EPI_ISL_515394, EPI_ISL_515391, EPI_ISL_515392, EPI_ISL_515317,

EPI_ISL_515438, EPI_ISL_515318, EPI_ISL_515439, EPI_ISL_515315,

EPI_ISL_515436, EPI_ISL_515316, EPI_ISL_515437, EPI_ISL_515313,

EPI_ISL_515434, EPI_ISL_515314, EPI_ISL_515435, EPI_ISL_515311,

EPI_ISL_515399, EPI_ISL_515432, EPI_ISL_515312, EPI_ISL_515433,

EPI_ISL_515418, EPI_ISL_515419, EPI_ISL_515386, EPI_ISL_515387,

EPI_ISL_515420, EPI_ISL_515384, EPI_ISL_515385, EPI_ISL_515382,

EPI_ISL_515383, EPI_ISL_515380, EPI_ISL_515381, EPI_ISL_515306,

EPI_ISL_515427, EPI_ISL_515307, EPI_ISL_515428, EPI_ISL_515304,

EPI_ISL_515425, EPI_ISL_515305, EPI_ISL_515426, EPI_ISL_515302,

EPI_ISL_515423, EPI_ISL_515303, EPI_ISL_515424, EPI_ISL_515300,

EPI_ISL_515388, EPI_ISL_515421, EPI_ISL_515301, EPI_ISL_515389,

EPI_ISL_515422

ACKNOWLEDGEMENTS:

We thank the staff at the Nevada State Public Health Lab (NSPHL). This work was

supported by NSPHL, University of Nevada, Reno Vice President for Research and

Innovation (VPRI), Department of Microbiology & Immunology, UNR School of Medicine,

Nevada IDeA Network of Biomedical Research Excellence (INBRE) from the National

Institute of General Medical Sciences (GM 103440 and GM 104944) from the National

Institutes of Health (NIH). The authors wish to acknowledge the support of Research &

Innovation and the Office of Information Technology at the University of Nevada, Reno

for computing time on the Pronghorn High-Performance Computing Cluster.

AUTHOR CONTRIBUTION:

PDH: conceptualization, formal analysis, methodology, writing – original draft preparation, writing

– review and editing

RLT: conceptualization, formal analysis, methodology, writing – original draft preparation, writing

– review and editing

XY: formal analysis

DPA: conceptualization, funding

JRS: formal analysis, review and editing

AG: methodology

EB: specimen procurement and diagnostic testing

HH: specimen procurement and diagnostic testing

MP: specimen procurement and diagnostic testing

MP: conceptualization, specimen procurement, diagnostic testing, formal analysis, project

administration, writing – review and editing

CCR: conceptualization, formal analysis, methodology, project administration, funding, writing –

original draft preparation, writing – review and editing

SCV: conceptualization, formal analysis, methodology, project administration, funding, writing –

review and editing

DECLARATIONS:

Ethics Approval:

Deidentified human specimens (nasal and nasopharyngeal swabs) were used for the

extraction of viral RNA all the experiments were done in accordance with guidelines of

the University of Nevada, Reno. The University of Nevada, Reno Institutional Review

Board (IRB) reviewed this project and determined this study to be EXEMPT FROM IRB

REVIEW according to federal regulations and University policy.

The Environmental and Biological Safety committee of the University of Nevada, Reno,

approved methods and techniques used in this study.

Competing Interests:

The authors declare that they have no competing interests with the contents of this article.

REFERENCES:

1.   Coronaviridae Study Group of the International Committee on Taxonomy of V: **The**

species Severe acute respiratory syndrome-related coronavirus: classifying

2019-nCoV and naming it SARS-CoV-2. *Nat Microbiol* 2020, **5**(4):536-544.

2.   Petrosillo N, Viceconte G, Ergonul O, Ippolito G, Petersen E: **COVID-19, SARS**

and MERS: are they closely related? *Clin Microbiol Infect* 2020, **26**(6):729-734.

3.   Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, Si HR, Zhu Y, Li B, Huang

CL *et al*: **A pneumonia outbreak associated with a new coronavirus of**

probable bat origin. *Nature* 2020, **579**(7798):270-273.

4.   Hu D, Zhu C, Ai L, He T, Wang Y, Ye F, Yang L, Ding C, Zhu X, Lv R *et al*: **Genomic**

characterization and infectivity of a novel SARS-like coronavirus in Chinese

bats. *Emerg Microbes Infect* 2018, **7**(1):154.

5.   de Wit E, van Doremalen N, Falzarano D, Munster VJ: **SARS and MERS: recent**

insights into emerging coronaviruses. *Nat Rev Microbiol* 2016, **14**(8):523-534.

6.   Peiris JS, Lai ST, Poon LL, Guan Y, Yam LY, Lim W, Nicholls J, Yee WK, Yan

WW, Cheung MT *et al*: **Coronavirus as a possible cause of severe acute**

respiratory syndrome. *Lancet* 2003, **361**(9366):1319-1325.

7.   Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus AD, Fouchier RA: **Isolation**

of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J*

Med 2012, **367**(19):1814-1820.

8.   Zhong NS, Zheng BJ, Li YM, Poon, Xie ZH, Chan KH, Li PH, Tan SY, Chang Q,

Xie JP *et al*: **Epidemiology and cause of severe acute respiratory syndrome**

(SARS) in Guangdong, People's Republic of China, in February, 2003. *Lancet*

2003, **362**(9393):1353-1358.

9.   Kim D, Lee JY, Yang JS, Kim JW, Kim VN, Chang H: **The Architecture of SARS-**

CoV-2 Transcriptome. *Cell* 2020, **181**(4):914-921 e910.

10. Licastro D, Rajasekharan S, Dal Monego S, Segat L, D'Agaro P, Marcello A:

Isolation and Full-Length Genome Characterization of SARS-CoV-2 from

COVID-19 Cases in Northern Italy. *J Virol* 2020, **94**(11).

11.   Holland LA, Kaelin EA, Maqsood R, Estifanos B, Wu LI, Varsani A, Halden RU,

462   Hogue BG, Scotch M, Lim ES: **An 81-Nucleotide Deletion in SARS-CoV-2**

463   **ORF7a Identified from Sentinel Surveillance in Arizona (January to March**

464   **2020)**. *J Virol* 2020, **94**(14).

465   12.   Wu F, Zhao S, Yu B, Chen YM, Wang W, Song ZG, Hu Y, Tao ZW, Tian JH, Pei

466   YY *et al*: **A new coronavirus associated with human respiratory disease in**

467   **China**. *Nature* 2020, **579**(7798):265-269.

468   13.   Davidson AD, Williamson MK, Lewis S, Shoemark D, Carroll MW, Heesom KJ,

469   Zambon M, Ellis J, Lewis PA, Hiscox JA *et al*: **Characterisation of the**

470   **transcriptome and proteome of SARS-CoV-2 reveals a cell passage induced**

471   **in-frame deletion of the furin-like cleavage site from the spike glycoprotein**.

472   *Genome Med* 2020, **12**(1):68.

473   14.   Walls AC, Park YJ, Tortorici MA, Wall A, McGuire AT, Veesler D: **Structure,**

474   **Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein**. *Cell* 2020,

475   **181**(2):281-292 e286.

476   15.   Bolger AM, Lohse M, Usadel B: **Trimmomatic: a flexible trimmer for Illumina**

477   **sequence data**. *Bioinformatics* 2014, **30**(15):2114-2120.

478   16.   Langmead B, Salzberg SL: **Fast gapped-read alignment with Bowtie 2**. *Nat*

479   *Methods* 2012, **9**(4):357-359.

480   17.   Picard-Toolkit.: **[cited 17 Aug 2020]**. *In: Github [Internet]* 2019, **Available:**

481   **http://broadinstitute.github.io/picard**.

482   18.   Garrison E, Marth G: **Haplotype-based variant detection from short-read**

483   **sequencing**. *arXiv [q-bioGN]* 2012, **Available: http://arxiv.org/abs/1207.3907**.

19. Garrison E: **Vcflib: A C++ library for parsing and manipulating VCF files**.

Available: https://githubcom/vcflib/vcflib 2019.

20. Li H: **A statistical framework for SNP calling, mutation discovery, association**

mapping and population genetical parameter estimation from sequencing

data. *Bioinformatics* 2011, **27**(21):2987-2993.

21. Elbe S, Buckland-Merrett G: **Data, disease and diplomacy: GISAID's innovative**

contribution to global health. *Glob Chall* 2017, **1**(1):33-46.

22. Nextstrain.: **Genomic epidemiology of novel coronavirus - Global**

subsampling. *In: Nextstrainorg [Internet] 15 Jul 2020 [cited 15 Jul 2020]* 2020,

Available: https://nextstrain.org/ncov/global/2020-07-

15?d=tree&l=clock&legend=closed.

23. Hadfield J, Megill C, Bell SM, Huddleston J, Potter B, Callender C, Sagulenko P,

Bedford T, Neher RA: **Nextstrain: real-time tracking of pathogen evolution**.

Bioinformatics 2018, **34**(23):4121-4123.

24. Chen J, Malone B, Llewellyn E, Grasso M, Shelton PMM, Olinares PDB, Maruthi

K, Eng ET, Vatandaslar H, Chait BT *et al*: **Structural Basis for Helicase-**

Polymerase Coupling in the SARS-CoV-2 Replication-Transcription

Complex. *Cell* 2020.

25. Korber B, Fischer WM, Gnanakaran S, Yoon H, Theiler J, Abfalterer W, Hengartner

N, Giorgi EE, Bhattacharya T, Foley B *et al*: **Tracking Changes in SARS-CoV-2**

Spike: Evidence that D614G Increases Infectivity of the COVID-19 Virus. *Cell*

2020, **182**(4):812-827 e819.

26. Nextclade: **[cited 17 Aug 2020]**. *Available: https://cladesnextstrainorg* 2020.

27. Matranga CB, Andersen KG, Winnicki S, Busby M, Gladden AD, Tewhey R,

Stremlau M, Berlin A, Gire SK, England E *et al*: **Enhanced methods for unbiased**

deep sequencing of Lassa and Ebola RNA viruses from clinical and

biological samples. *Genome Biol* 2014, **15**(11):519.

28. Briese T, Kapoor A, Mishra N, Jain K, Kumar A, Jabado OJ, Lipkin WI: **Virome**

Capture Sequencing Enables Sensitive Viral Diagnosis and Comprehensive

Virome Analysis. *mBio* 2015, **6**(5):e01491-01415.

29. O'Flaherty BM, Li Y, Tao Y, Paden CR, Queen K, Zhang J, Dinwiddie DL, Gross

SM, Schroth GP, Tong S: **Comprehensive viral enrichment enables sensitive**

respiratory virus genomic identification and analysis by next generation

sequencing. *Genome Res* 2018, **28**(6):869-877.

30. Paden CR, Tao Y, Queen K, Zhang J, Li Y, Uehara A, Tong S: **Rapid, Sensitive,**

Full-Genome Sequencing of Severe Acute Respiratory Syndrome

Coronavirus 2. *Emerg Infect Dis* 2020, **26**(10).

31. Paskey AC, Frey KG, Schroth G, Gross S, Hamilton T, Bishop-Lilly KA:

Enrichment post-library preparation enhances the sensitivity of high-

throughput sequencing-based detection and characterization of viruses

from complex samples. *BMC Genomics* 2019, **20**(1):155.

32. Xiao M, Liu X, Ji J, Li M, Li J, Yang L, Sun W, Ren P, Yang G, Zhao J *et al*: **Multiple**

approaches for massively parallel sequencing of SARS-CoV-2 genomes

directly from clinical samples. *Genome Med* 2020, **12**(1):57.

 FIGURE LEGENDS:

Figure 1. Workflow of SARS-CoV-2 genome sequencing and analysis from

nasopharyngeal patient specimens in Nevada. (A) RNA was extracted from Nasal or

Nasopharygeal (NP) swabs taken from patients in Nevada and first used to determine the

presence of SARS-CoV-2 genomes by RT-qPCR. Next generation sequencing (NGS)

libraries were prepared from positive specimens, this included steps for ribosomal RNA

depletion and SARS-CoV-2 enrichment. Subsequent libraries were pooled and used for

whole genome sequencing at the Nevada Genomics Center on the Illumina NextSeq 500

instrument. FASTQ files were aligned to the reference genome, and analyzed to

determine nucleotide variation and phylogenetic relationship. (B) A total of 200 specimens

were sequenced, of which 174 had over 99% coverage of the SARS-CoV-2 genome.

This included 133 patient specimens from Northern Nevada, 40 from Southern Nevada

and 1 specimen that was re-sequenced. (C) Correlation between RT-qPCR Ct value and

the percentage of coverage in the whole genome sequencing after trimming and

alignment. (D) Nucleotide variants across the SARS-CoV-2 genome in the 173 specimens

from Nevada from March 6th to June 5th.

Figure 2. Distribution of D614G in Nevada and comparison with the United States

and global proportion. (A) Cumulative frequency of D614G in 173 patient specimens

from Nevada from March 6th to June 5th, 2020 (D614 is indicated by teal, G614 is

indicated by yellow). Pie charts depict the cumulative proportion up to the indicated time

point (March 15, April 1, May 1, June 5). The total number of specimens included at each

time point is specified below each pie chart. Effective dates of emergency orders and

regulatory responses to SARS-CoV-2 spread in Nevada are indicated on the frequency

graph time axis. (B) Proportion of D614G in the United States from March 6th to June 5th,

specimens from Nevada are divided in the geographic area that they originated from,

Northern Nevada (N-NV) includes 133 specimens from Washoe county, Carson-Tahoe,

and other northern counties, and Southern Nevada (S-NV) includes 40 specimens from

Clark county. (C) Global proportion of D614G in the shown regions during the same time

period from a subsampling of sequences deposited in Nextstrain.org. The size of the pie

chart corresponds to the relative specimen number for each region.

Figure 3. Distribution of SARS-CoV-2 clades in Nevada. (A) Cumulative frequency of

SARS-CoV-2 clades in 173 patient specimens from Nevada during March 6th to June

5th. The five clades are colored 19A (blue), 19B (teal), 20A (green), 20B (yellow) and 20C

(orange). Pie charts depict the cumulative proportion up to the indicated time point (March

15, April 1, May 1, June 5). The total number of specimens included at each time point is

specified below each pie chart. Dates of emergency orders and regulations meant to slow

the spread of SARS-CoV-2 in Nevada are indicated on the time scale of the frequency

graph. (B) Circular dendrogram depicting clades from Nevada specimens. (C) Pie chart

of the clades from northern Nevada (N-NV), southern Nevada (S-NV) and the United

States. (D) Pie charts show the proportion of clades from global regions during the same

time period from a subsampling of sequences deposited in Nextstrain.org. The size of the

pie chart corresponds to the relative specimen number for each region.

Figure 4. Distribution of P323L/F (nsp12, RdRp) in Nevada. (A) Cumulative frequency

of P323L/F (nsp12, RdRp) in 173 patient specimens from Nevada during March 6th to

June 5th. The amino acid at position 323 is indicated by teal for proline (P), yellow for

leucine (L) and blue for phenylalanine (F). Pie charts depict the cumulative proportion up

to the indicated time point (March 15, April 1, May 1, June 5). The total number of

specimens included at each time point is specified below each pie chart. Dates of

emergency orders and regulations meant to slow the spread of SARS-CoV-2 in Nevada

are indicated on the time scale of the frequency graph. (B) Circular dendrogram

representing the distribution of amino acid change at residue 323 of nsp12 from a global

subsampling of sequences deposited in Nextstrain.org from March 6th to June 5th, the

larger dots indicate specimens from Nevada. (C) Pie chart indicating the ratio of P/L/F in

Northern NV and Southern NV specimens from this study. (D) Proportion of P323L/F from

a subsampling of sequences deposited in Nextstrain.org for the United States and (E)

global regions from March 6th to June 5th. The size of the pie chart corresponds to the

relative specimen number for each region.

598   **Figure 1:**

A.

B. **200** Total samples
   26 <90% coverage
   174 >90% coverage
   133 patient samples from Northern Nevada (N-NV)
          (Washoe County, Carson-Tahoe area and Other)
   40 patient samples from Southern Nevada (S-NV)
          (Clark county)
   1 Repeat sequencing of specimen

C.

D.

Figure 2:

Figure 3:

Figure 4:

1 **SUPPLEMENTARY INFORMATION**
2
3 **Figure S1:** Alignment of SARS-CoV-2 sequences from NV patient's specimens.
4
5

Figure S2. Dendrogram of Nevada specimens in context of other sequenced

specimens. (a) Nucleotide

mutation clock from

Nextstrain.org with the

SARS-CoV-2 genome

isolated from Washington

(USA/WA 1/2020) on

January 24th and first

specimen from Nevada

(A0004) on March 5th

indicated in red. (b) Circular

dendrogram of Nevada

specimens from March 6th

to June 5th positioned

within a subsample of

global sequences from

Nextstrain.org during the

same time period, the

larger dots indicate

specimens from Nevada.

The five clades are colored

19A (blue), 19B (teal), 20A (green), 20B (yellow) and 20C (orange).

Fig. S1a

Fig. S1b

Figure S3. Global distribution of clades from March 6th to June 5th. Pie charts depict

the proportion of the clades within (a) four main areas in Nevada, these include 91

specimens from

Washoe county

(upper left), 23

specimens from

Carson-Tahoe

(middle left), 40

specimens from

Clark county

(bottom right),

and 19 from rural

Nevada

(middle). The

five clades are

colored 19A

(blue), 19B

(teal), 20A

(green), 20B

(yellow) and 20C (orange). (b) Pie chart of the clades in each indicated country from

March 6th to June 5th generated from a subsampling of sequences deposited in

Nextstrain.org. The size of the pie chart corresponds to the relative specimen number for

each region.

Figure S4. Distribution of nucleotide variant 379C>A. (a) Green line at the far-left end of the genome denoted nucleotide position 379 of nsp1. (b) Circular dendrogram of global subsample of sequences from Nextstrain.org with NV specimens indicated by larger dots. (c) Pie chart indicating the proportion of sequences with either the cytidine (C) or adenosine (A) at position 379 from the Nevada specimens. Subsample of sequences from Nextstrain.org were used to generate the proportion of 379C>A in (d) the indicated states within the U.S. and (e) internationally. The size of the pie chart corresponds to the relative specimen number for each region.

Figure S5. Structure of SARS-CoV-2 nsp12 (RdRp) P323L/F. (a) Diagram depicting

ORF1b genomic location and encoded proteins. Below the linear protein schematic of nsp12 specific nucleotide variants at position 14,407 and 14,408 and the resulting amino acid changes are indicated. (b) SARS-CoV-2 replicase complex

modeled from 6XEZ template. This model includes nsp12, nsp7, nsp8, nsp13, ligands

(ZN, Mg), the template and product strand of RNA. P323L/F within the interface domain

of nsp12 is located at the left side of the complex. (c) Model nsp12 showing P323L/F

within the interface domain. Residue 323 is shown with either P, L or F and amino

acids with side chains within 5 Å of residue 323 are depicted as sticks in the cartoon

model.