

Assessment of the risk of SARS-CoV-2 reinfection in an intense re-exposure setting

Laith J. Abu Raddad PhD,^{1,2,3*} Hiam Chemaitelly MSc,^{1,2} Houssein H. Ayoub PhD,⁴ Zaina Al Kanaani PhD,⁵ Abdullatif Al Khal MD,⁵ Einas Al Kuwari MD,⁵ Adeel A. Butt MD,⁵ Peter Coyle MD,⁵ Andrew Jeremijenko MD,⁵ Anvar Hassan Kaleeckal MSc,⁵ Ali Nizar Latif MD,⁵ Hanan F. Abdul Rahim PhD,⁶ Mohamed G. Al Kuwari MD,⁷ Hamad Eid Al Romaihi MD,⁸ Sheikh Mohammad Al Thani MD,⁸ and Roberto Bertollini MD⁸

¹*Infectious Disease Epidemiology Group, Weill Cornell Medicine-Qatar, Cornell University, Doha, Qatar*

²*World Health Organization Collaborating Centre for Disease Epidemiology Analytics on HIV/AIDS, Sexually Transmitted Infections, and Viral Hepatitis, Weill Cornell Medicine-Qatar, Cornell University, Qatar Foundation – Education City, Doha, Qatar*

³*Department of Population Health Sciences, Weill Cornell Medicine, Cornell University, New York, New York, USA*

⁴*Department of Mathematics, Statistics, and Physics, Qatar University, Doha, Qatar*

⁵*Hamad Medical Corporation, Doha, Qatar*

⁶*College of Health Sciences, QU Health, Qatar University, Doha, Qatar*

⁷*Primary Health Care Corporation, Doha, Qatar*

⁸*Ministry of Public Health, Doha, Qatar*

* Address reprints requests or correspondence to Professor Laith J. Abu-Raddad, Infectious Disease Epidemiology Group, World Health Organization Collaborating Centre for Disease Epidemiology Analytics on HIV/AIDS, Sexually Transmitted Infections, and Viral Hepatitis, Weill Cornell Medicine - Qatar, Qatar Foundation - Education City, P.O. Box 24144, Doha, Qatar. Telephone: +(974) 4492-8321. Fax: +(974) 4492-8333. E-mail: lja2002@qatar-med.cornell.edu.

ABSTRACT

Background: Reinfection with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has been a subject of debate. We aimed to assess the risk and incidence rate of documented SARS-CoV-2 reinfection in a large cohort of laboratory-confirmed cases in Qatar.

Methods: All SARS-CoV-2 laboratory-confirmed cases with at least one PCR positive swab that is ≥ 45 days after a first positive swab were individually investigated for evidence of reinfection, and classified as showing *strong*, *good*, *some*, or *weak/no* evidence for reinfection. Risk and incidence rate of reinfection were estimated.

Results: Out of 133,266 laboratory-confirmed SARS-CoV-2 cases, 243 persons (0.18%) had at least one subsequent positive swab ≥ 45 days after the first positive swab. Of these, 54 cases (22.2%) had strong or good evidence for reinfection. Median time between first and reinfection swab was 64.5 days (range: 45-129). Twenty-three of the 54 cases (42.6%) were diagnosed at a health facility suggesting presence of symptoms, while 31 (57.4%) were identified incidentally through random testing campaigns/surveys or contact tracing. Only one person was hospitalized at or following time of reinfection swab, but still had relatively mild infection. No deaths were recorded. Risk of reinfection was estimated at 0.04% (95% CI: 0.03-0.05%) and incidence rate of reinfection was estimated at 1.09 (95% CI: 0.84-1.42) per 10,000 person-weeks.

Conclusions: SARS-CoV-2 reinfection appears to be a rare phenomenon suggestive of a strong protective immunity against reinfection that lasts for at least a few months post primary infection.

Keywords: SARS-CoV-2; epidemiology; COVID-19; incidence; infection; reinfection; immunity

INTRODUCTION

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has been spreading around the globe causing severe disruptions to social and economic activities [1-3]. Qatar, a peninsula in the Arabian Gulf region with a diverse population of 2.8 million [4, 5], has experienced a large epidemic with one of the highest laboratory-confirmed rates of infection at >45,000 infections per million population [6, 7]. Antibody testing and mathematical modeling indicated that about half of the population of Qatar has already been infected [6].

The intensity of the epidemic in this country with a high risk of re-exposure to the infection, as well as the availability of a centralized data-capture system of all laboratory-confirmed infections, provided an opportunity to epidemiologically assess the presence and incidence of reinfections, a debated feature of SARS-CoV-2 epidemiology whose elucidation is critical to inform global response, timing and intensity of future cycles, and impact and durability of potential vaccines [8-11].

Our primary aim was to assess the risk and incidence rate of documented reinfection in a cohort of 133,266 SARS-CoV-2 laboratory-confirmed infected persons. Since the relevant underlying question is whether risk of reinfection is appreciable or not, we implemented a conservative approach for assessing documented reinfections, that is prone to overestimate rather than underestimate risk of reinfection. However, we also conducted sensitivity analyses implementing more stringent criteria for assessing reinfection.

METHODS

Sources of data

We analyzed the centralized and standardized national SARS-CoV-2 testing and hospitalization database compiled at Hamad Medical Corporation (HMC), the main public healthcare provider in Qatar and the nationally-designated provider for Coronavirus Disease 2019 (COVID-19) healthcare needs. The database covers all SARS-CoV-2 cases in Qatar and encompasses data on all polymerase chain reaction (PCR) testing conducted from February 28-August 12, 2020, including testing of suspected SARS-CoV-2 cases and traced contacts as well as infection surveillance testing. The database further includes data on hospital admission of COVID-19 patients and the World Health Organization (WHO) severity classification for each infection, which is assessed through individual chart reviews by trained medical personnel [12]. More recently, data on serological testing for antibody on residual blood specimens collected for routine clinical care from outpatient and inpatient attendees at HMC were also incorporated [6].

Laboratory methods

All PCR testing was conducted at HMC Central Laboratory or at Sidra Medicine Laboratory, following standardized protocols. Nasopharyngeal and/or oropharyngeal swabs (Huachenyang Technology, China) were collected and placed in Universal Transport Medium (UTM). Aliquots of UTM were: extracted on the QIA Symphony platform (QIAGEN, USA) and tested with real-time reverse-transcription PCR (RT-qPCR) using the TaqPath™ COVID-19 Combo Kit (Thermo Fisher Scientific, USA) on a ABI 7500 FAST (ThermoFisher, USA); extracted using a custom protocol [13] on a Hamilton Microlab STAR (Hamilton, USA) and tested using the AccuPower SARS-CoV-2 Real-Time RT-PCR Kit (Bioneer, Korea) on a ABI 7500 FAST; or loaded directly to a Roche cobas® 6800 system and assayed with the cobas® SARS-CoV-2 Test (Roche, Switzerland). The first assay targets the S, N, and ORF1ab regions of the virus; the

second targets the virus' RdRp and E-gene regions; and the third targets the ORF1ab and E-gene regions.

Serological testing was performed using the Roche Elecsys[®] Anti-SARS-CoV-2 (Roche, Switzerland), an electrochemiluminescence immunoassay that uses a recombinant protein representing the nucleocapsid (N) antigen for the determination of antibodies against SARS-CoV-2. Qualitative anti-SARS-CoV-2 results were generated following the manufacturer's instructions (reactive: cutoff index ≥ 1.0 vs. non-reactive: cutoff index < 1.0).

Inclusion criteria

All SARS-CoV-2 laboratory-confirmed cases with at least one PCR positive swab that is ≥ 45 days after a first positive swab were considered as *suspected cases* of reinfection. The 45-day cutoff was informed by data from observational cohorts of SARS-CoV-2 infected persons [14, 15], and was set to account for the duration of prolonged PCR positivity of several weeks in these patients. The cutoff determination was further informed by the distribution of the time difference between the first positive swab and subsequent positive swabs among SARS-CoV-2 cases with multiple swabs (Figure 1). The tail of this distribution indicates that a cutoff of 45 days (at the 99th percentile) provides an appropriate mark for defining the end of prolonged PCR positivity: a subsequent positive swab within 45 days of the first positive swab is likely to reflect prolonged PCR positivity (due to non-viable virus fragments) rather than reinfection, and thus should not be included in analysis.

Suspected reinfection case classification

Suspected cases of reinfection, that is cases fitting above indicated inclusion criteria, were classified as showing either *strong* evidence, *good* evidence, *some* evidence, or *weak* (or *no*)

evidence for reinfection (Box 1). The classification was based on holistic quantitative and qualitative criteria applied to each investigated case of reinfection. The criteria included the pattern and magnitude of the change in PCR cycle threshold (Ct) value across repeated swabs, the time interval between subsequent swabs, PCR testing site (such as outpatients at primary care, hospital emergency, or inpatient hospitalization), purpose of PCR testing (such as appearance of symptoms, contact tracing, or survey/testing campaign), age, history of COVID-19-related hospital admission, and case severity per WHO classification [12].

Overall, swabs with Ct <30 (suggestive of active recent infection) at least 45 days after the first positive swab were considered as showing *strong evidence for reinfection*. Swabs with Ct \geq 30 at least 45 days after the first positive swab were considered as showing *good evidence for reinfection* if PCR positivity was associated with *contextual evidence* supporting the status of “reinfection” including appearance of symptoms (often as proxied by being diagnosed at a health facility), if the infection was diagnosed through contact tracing (indicating recent exposure to an infected person), if the change in Ct value from the last swab was to a lower Ct value (indicating increasing viral load), and/or if the repeated swabbing did not follow a regular pattern and time interval between repeated swabs was not short (to exclude cases under clinical management that are indicative of poor control of first infection).

Shorter durations bordering the cutoff of 45 days with Ct values \geq 30 and with no contextual evidence supporting the status of “reinfection” were indicative of *some evidence for reinfection*, but not strong nor good evidence for reinfection, as they are more likely to reflect the long tail of the prolonged PCR positivity distribution (Figure 1) [14, 15]. Age \geq 70 years, repeated swabs on hospitalized patients, and severe or critical WHO disease classifications were considered as contextual factors indicative of poor infection control of the first infection rather than reinfection.

Cases that had such contextual factors (and implicitly did not fit the criteria of strong, good, or some evidence for reinfection) were considered to have *weak (or no) evidence for reinfection*.

Of note that hospitalized COVID-19 cases often had multiple subsequent swabs administered to them as part of clinical care, and repeated swabbing was standard earlier in the epidemic, as the criteria for discharge from an isolation facility required at least two subsequent PCR negative swabs. This was changed later on to a time-based criteria per updated WHO recommendation [16].

Reinfection risk and rate

Documented reinfection *risk* was assessed by quantifying the proportion of cases with *strong or good evidence for reinfection* out of all laboratory-confirmed SARS-CoV-2 cases. *Incidence rate* of documented reinfection was calculated by dividing the number of cases with strong or good evidence for reinfection by the number of person-weeks contributed by all laboratory-confirmed cases who had their first positive swab ≥ 45 days before day of analysis. The follow-up person-time was calculated starting from 45 days after the first positive swab and up to the reinfection swab, all-cause death, or end-of-study censoring.

Sensitivity analyses

Since we implemented a conservative approach prone to overestimate risk of documented reinfection, several sensitivity analyses were conducted implementing more *stringent criteria* for assessing reinfection: 1) exclusion of cases where the Ct value for the first and/or subsequent positive swab was unknown or with a value ≥ 35 (to exclude potential PCR false-positive cases), 2) changing the ≥ 45 -day cutoff to a ≥ 60 -day cutoff to further exclude potential cases of long-term prolonged PCR positivity, and (*most stringent*) 3) setting definition of active recent

infection at Ct cutoff value of <25 (instead of <30) and excluding any suspected reinfection case with Ct value >25.

Ethical approval

Studies were approved by HMC and Weill Cornell Medicine-Qatar Institutional Review Boards.

RESULTS

Main analysis

Figure 2 illustrates the selection process of SARS-CoV-2 eligible cases and summarizes the results of their reinfection status' evaluation. Out of 133,266 laboratory-confirmed SARS-CoV-2 cases, 117,458 had only one single positive swab and thus were excluded from further analysis.

Of the remaining 15,808 cases with multiple swabs, only 243 persons had *at least* one subsequent positive swab that is ≥ 45 days from the first positive swab, and thus qualified for inclusion in analysis.

Individual investigation of each of the 299 positive swabs collected ≥ 45 days after the first positive swab for these 243 cases yielded 54 cases with *strong* or *good* evidence for reinfection.

Of these, 35 had *strong* evidence for reinfection, as the Ct value was <30 for the reinfection swab. The remaining 19 were considered to have *good* evidence for reinfection, as the reinfection swab Ct value was ≥ 30 . An additional 26 cases showed *some* evidence for reinfection, while evidence was *weak* for the remaining 163 cases.

Table 1 shows the characteristics of the 54 cases classified as showing strong or good evidence for reinfection. Almost all cases were males, but this reflects the focus of the epidemic in craft and manual workers [6]. The median age was 33 years (range: 16-57) and the median time between the *first* swab and the *reinfection* swab was 64.5 days (range: 45-129). The median Ct

value was 28 (range: 14-37); it was 22 (range 14-29) for the 35 swabs classified with strong evidence ($Ct < 30$) and 32 (range: 30-37) for the remaining swabs ($Ct \geq 30$).

Twenty-three cases (42.6%) were diagnosed at a health facility, suggesting presence of symptoms while 31 (57.4%) were identified incidentally either through random testing campaigns/surveys ($n=15$; 27.8%) or contact tracing ($n=16$; 29.6%), suggesting minimal symptoms if any.

Nine cases were hospitalized at any time, all but *one* occurred following the first infection episode and mostly for isolation purposes. Only one case had sufficient symptoms to warrant clinical assessment (during primary infection), but was classified with “mild” severity per WHO classification. No deaths were recorded. Of note that the vast majority of infections in Qatar occurred in young and healthy men and had limited severity [6].

Antibody test results were available for 48 out of the 243 assessed individuals (Table 2), of whom 30 (62.5%) had detectable antibodies. Of the 13 with strong evidence for reinfection *and* available antibody results, seven (53.9%) were sero-negative. Meanwhile, both individuals with good evidence for reinfection, three of the four individuals with some evidence for reinfection, and 19 of the 29 individuals with weak evidence for reinfection, were seropositive.

Risk of documented reinfection was estimated at 0.04% (95% CI: 0.03-0.05%)—that is a total of 54 reinfections in the cohort of 133,266 laboratory-confirmed SARS-CoV-2 infected persons. Incidence rate of reinfection was estimated at 1.09 (95% CI: 0.84-1.42) per 10,000 person-weeks—that is a total of 54 reinfection events in a follow-up person-time of 495,208.7 person-weeks.

Sensitivity analysis

The sensitivity analysis excluding all cases where Ct value for the first and/or subsequent positive swab was unknown or with a value ≥ 35 (to exclude potential PCR false-positive cases) reduced the number of cases with strong evidence to 21 and with good evidence to 5. Risk and incidence rate of reinfection were estimated at 0.02% (95% CI: 0.01-0.03%) and 0.52 (95% CI: 0.36-0.77) per 10,000 person-weeks, respectively.

The sensitivity analysis changing the ≥ 45 -day cutoff to a ≥ 60 -day cutoff to further exclude potential cases of long-term prolonged PCR positivity reduced number of cases with strong evidence to 25 and with good evidence to 7. Risk and incidence rate of reinfection were estimated at 0.02% (95% CI: 0.02-0.03%) and 1.06 (95% CI: 0.75-1.50) per 10,000 person-weeks, respectively.

The most stringent sensitivity analysis excluding any suspected reinfection with Ct value > 25 reduced number of cases to 19 (all with strong evidence). Risk and incidence rate of reinfection were estimated at 0.01% (95% CI: 0.01-0.02%) and 0.38 (95% CI: 0.24-0.60) per 10,000 person-weeks, respectively.

DISCUSSION

Results indicate supportive (albeit not conclusive) evidence for presence of reinfections in the SARS-CoV-2 epidemic of Qatar, but the risk for documented reinfection was rare at only 4 reinfections per 10,000 infected persons. This finding is striking as the epidemic in Qatar has been intense with half of the population estimated to have been infected [6]. Considering the strength of the force of infection, estimated at a *daily* probability of infection exceeding 1% at the epidemic peak around May 20 [6], it is all but certain that a significant proportion of the population has been repeatedly exposed to the infection, but such re-exposures hardly led to any documentable reinfections.

Indeed, of all identified reinfections, nearly two-thirds (57%) were discovered accidentally, either through random testing campaigns/surveys or through contact tracing. None were severe, critical, or fatal, all reinfections were asymptomatic or with minimal or mild symptoms. These findings suggest that most infected persons do develop immunity against reinfection that lasts for at least few months, and that reinfections (if they occur) are well tolerated and no more symptomatic than primary infections. Further follow up of this cohort of infected persons over time may allow elucidation of any potential effects of waning of immunity.

Other lines of evidence for this cohort also support this conclusion. Among 2,559 PCR positive persons where an antibody test outcome was available [6], and where the first positive PCR test was conducted >3 weeks before the serology test to accommodate for the delay in development of antibodies following onset of infection [14, 15], 91.7% were antibody positive [6]. The high antibody positivity was also stable for over three months [6], as described elsewhere [9]. The epidemic curve in Qatar was further characterized by rapid growth followed by rapid decline [6], at a time when levels of social and physical distancing restrictions were fairly stable. This points to susceptibles-infected-recovered “SIR” epidemic dynamics with most infections eliciting immunity against reinfection.

This assessment has limitations. We assessed risk of only *documented* reinfections, but other reinfections could have occurred but went undocumented, perhaps because of minimal or no symptoms. It is possible that with the primed immune system following the primary infection, reinfections are milder and shorter and thus less likely to be documented [10]. While risk of reinfection appears limited, reinfected persons could possibly transmit the virus as the Ct value was quite low, indicating high viral load, in some of the reinfections. Antibody testing outcomes

were available for only a number of cases, limiting use and inferences of the link between antibody status and risk of reinfection.

We found evidence for reinfections using epidemiological criteria, but this evidence cannot be conclusive unless it is confirmed with sequencing analysis of the paired viral samples. It is possible that some of the identified reinfections may not have been true reinfections. The possibility of a false positive swab cannot be excluded. It is also possible that some of the cases may just be poorly-controlled primary infections, long-term virus carriers, or persons shedding non-viable viral fragments for a prolonged time. The sensitivity analyses accounting for such limitations suggested that we may have overestimated the already low risk of reinfection.

In conclusion, SARS-CoV-2 reinfection appears to be a rare phenomenon. This suggests that immunity develops after the primary infection and lasts for at least a few months, and that immunity protects against reinfection.

Box 1. Classification of suspected cases of SARS-CoV-2 reinfection based on strength of supporting evidence.

Suspected cases of SARS-CoV-2 reinfection: all laboratory-confirmed cases with at least one polymerase chain reaction (PCR) positive swab that is ≥ 45 days after a first positive swab.

Strong evidence for reinfection: individuals having positive swabs with PCR cycle threshold (Ct) value < 30 at least 45 days after the first positive swab. No contextual evidence supporting poor control of first infection such as age ≥ 70 years, repeated swabs on hospitalized patients, and severe or critical World Health Organization disease classifications.

Good evidence for reinfection: individuals having positive swabs with PCR Ct value ≥ 30 at least 45 days after the first positive swab, but where PCR positivity was associated with contextual evidence supporting the status of reinfection:

- Appearance of symptoms (often as proxied by being diagnosed at a health facility)
- Infection diagnosis through contact tracing (indicating recent exposure to an infected person)
- Lower Ct value compared to last positive swab (indicating increasing viral load)
- Irregular and spaced-out pattern for repeated swabbing (to exclude cases under clinical management that are indicative of poor control of first infection).

No contextual evidence supporting poor control of first infection such as age ≥ 70 years, repeated swabs on hospitalized patients, and severe or critical World Health Organization disease classifications.

Some evidence for reinfection: individuals having positive swabs with PCR Ct value ≥ 30 at least 45 days after the first positive swab, but typically bordering the cutoff of 45 days. PCR positivity was **not** associated with evidence supporting the status of reinfection (listed above).

Weak evidence for reinfection: individuals having swabs with PCR Ct value ≥ 30 at least 45 days after the first positive swab, but typically bordering the cutoff of 45 days. PCR positivity was associated with contextual evidence indicative of poor infection control of the first infection rather than reinfection (such as age ≥ 70 years, repeated swabs on hospitalized patients, and severe or critical World Health Organization disease classifications).

Figure 1. Distribution of the time difference between the first swab and subsequent swabs among all laboratory-confirmed SARS-CoV-2 cases with more than one positive swab. The cutoff of 45 days was at the 99th percentile, and thus provides an appropriate mark for defining the end of the prolonged polymerase chain reaction (PCR) positivity.




Figure 2. Flow chart describing the selection process of SARS-CoV-2 eligible cases and summarizing the results of their reinfection status' evaluation.




Table 1. Characteristics of individuals classified as showing strong or good evidence for reinfection.

ID#	Socio-demographic			PCR testing				Hospitalization			Ab testing
	Sex	Age group*	Nationality	Sample type	PCR swab time†	Positive swab type	Average Ct value	Case severity‡	Hospital admission§	LOS (days)	Ab status¶
Strong evidence for reinfection											
1	Male	50-54	Pakistani	Survey**	Day 0	First pos swab	Unk	Not assessed††	Not hospitalized	0	Unk
	Male	50-54	Pakistani	Survey**	Day 75	Reinf swab	14	Not assessed††	Not hospitalized	0	Negative
2	Male	30-34	Qatari	Health facility	Day 0	First pos swab	36	Not assessed††	Not hospitalized	0	Unk
	Male	30-34	Qatari	Health facility	Day 55	Reinf swab	16	Not assessed††	Not hospitalized	0	Unk
3	Male	30-34	Iranian	Contact tracing	Day 0	First pos swab	Unk	Not assessed††	1 day in April	1	Unk
	Male	30-34	Iranian	Survey**	Day 85	Reinf swab	17	Not assessed††	Not hospitalized	0	Negative
4	Male	25-29	Nepalese	Health facility	Day 0	First pos swab	33	Not assessed††	Not hospitalized	0	Unk
	Male	25-29	Nepalese	Health facility	Day 76	Reinf swab	17	Not assessed††	Not hospitalized	0	Unk
5	Male	35-39	Qatari	Contact tracing	Day 0	First pos swab	Unk	Not assessed††	Not hospitalized	0	Unk
	Male	35-39	Qatari	Health facility	Day 20	Subs pos swab	24	Not assessed††	Not hospitalized	0	Unk
	Male	35-39	Qatari	Health facility	Day 129	Reinf swab	17	Not assessed††	Not hospitalized	0	Unk
6	Female	50-54	Qatari	Contact tracing	Day 0	First pos swab	34	Not assessed††	Not hospitalized	0	Unk
	Female	50-54	Qatari	Health facility	Day 53	Reinf swab	17	Not assessed††	Not hospitalized	0	Negative
7	Female	20-24	Qatari	Health facility	Day 0	First pos swab	35	Not assessed††	Not hospitalized	0	Unk
	Female	20-24	Qatari	Health facility	Day 84	Reinf swab	17	Not assessed††	Not hospitalized	0	Unk
8	Male	30-34	Yemeni	Health facility	Day 0	First pos swab	36	Not assessed††	Not hospitalized	0	Unk
	Male	30-34	Yemeni	Contact tracing	Day 60	Reinf swab	18	Not assessed††	Not hospitalized	0	Unk
9	Male	20-24	Palestinian	Contact tracing	Day 0	First pos swab	Unk	Not assessed††	Not hospitalized	0	Unk
	Male	20-24	Palestinian	Health facility	Day 97	Reinf swab	18	Not assessed††	Not hospitalized	0	Unk
10	Female	20-24	Qatari	Health facility	Day 0	First pos swab	Unk	Not assessed††	Not hospitalized	0	Unk
	Female	20-24	Qatari	Contact tracing	Day 91	Reinf swab	19	Not assessed††	Not hospitalized	0	Unk
11	Male	35-39	Qatari	Contact tracing	Day 0	First pos swab	Unk	Not assessed††	Not hospitalized	0	Unk
	Male	35-39	Qatari	Contact tracing	Day 86	Reinf swab	19	Not assessed††	Not hospitalized	0	Positive
12	Female	45-49	Qatari	Health facility	Day 0	First pos swab	30	Not assessed††	Not hospitalized	0	Unk
	Female	45-49	Qatari	Health facility	Day 74	Reinf swab	20	Not assessed††	Not hospitalized	0	Negative
13	Male	30-34	Palestinian	Contact tracing	Day 0	First pos swab	Unk	Not assessed††	1 day in April	1	Unk
	Male	30-34	Palestinian	Survey**	Day 57	Reinf swab	20	Not assessed††	Not hospitalized	0	Unk
14	Female	40-44	Qatari	Contact tracing	Day 0	First pos swab	24	Not assessed††	Not hospitalized	0	Unk
	Female	40-44	Qatari	Health facility	Day 103	Reinf swab	21	Not assessed††	Not hospitalized	0	Positive
15	Male	50-54	Qatari	Contact tracing	Day 0	First pos swab	34	Not assessed††	Not hospitalized	0	Unk
	Male	50-54	Qatari	Contact tracing	Day 57	Reinf swab	21	Not assessed††	Not hospitalized	0	Negative
16	Male	25-29	Nepalese	Health facility	Day 0	First pos swab	Unk	Not assessed††	5 days in March†	5	Unk
	Male	25-29	Nepalese	Contact tracing	Day 73	Reinf swab	21	Not assessed††	Not hospitalized	0	Unk
17	Male	20-24	Qatari	Health facility	Day 0	First pos swab	32	Not assessed††	Not hospitalized	0	Unk
	Male	20-24	Qatari	Health facility	Day 89	Reinf swab	22	Not assessed††	1 day in August	1	Unk
18	Female	20-24	Qatari	Health facility	Day 0	First pos swab	33	Not assessed††	Not hospitalized	0	Unk
	Female	20-24	Qatari	Health facility	Day 76	Reinf swab	22	Not assessed††	Not hospitalized	0	Negative

19	Male	40-44	Kenyan	Health facility	Day 0	First pos swab	23	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	40-44	Kenyan	Health facility	Day 65	Reinf swab	23	Not assessed ^{††}	Not hospitalized	0	Positive
20	Female	40-49	Palestinian	Health facility	Day 0	First pos swab	36	Not assessed ^{††}	Not hospitalized	0	Unk
	Female	40-49	Palestinian	Health facility	Day 88	Reinf swab	25	Not assessed ^{††}	Not hospitalized	0	Negative
21	Male	30-34	Indian	Survey ^{**}	Day 0	First pos swab	32	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	30-34	Indian	Survey ^{**}	Day 76	Reinf swab	25	Not assessed ^{††}	Not hospitalized	0	Positive
22	Male	20-24	Yemeni	Health facility	Day 0	First pos swab	28	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	20-24	Yemeni	Health facility	Day 66	Reinf swab	26	Not assessed ^{††}	Not hospitalized	0	Unk
23	Male	20-24	Qatari	Health facility	Day 0	First pos swab	31	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	20-24	Qatari	Contact tracing	Day 56	Reinf swab	27	Not assessed ^{††}	Not hospitalized	0	Unk
24	Male	35-39	Sudanese	Health facility	Day 0	First pos swab	29	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	35-39	Sudanese	Health facility	Day 56	Reinf swab	27	Not assessed ^{††}	Not hospitalized	0	Unk
25	Male	40-44	Nepalese	Health facility	Day 0	First pos swab	21	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	40-44	Nepalese	Contact tracing	Day 63	Reinf swab	28	Not assessed ^{††}	Not hospitalized	0	Unk
26	Male	40-44	Nepalese	Survey ^{**}	Day 0	First pos swab	17	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	40-44	Nepalese	Health facility	Day 14	Subs pos swab	32	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	40-44	Nepalese	Health facility	Day 22	Subs pos swab	28	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	40-44	Nepalese	Survey ^{**}	Day 51	Reinf swab	28	Not assessed ^{††}	Not hospitalized	0	Unk
27	Male	25-29	Indian	Health facility	Day 0	First pos swab	36	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	25-29	Indian	Health facility	Day 46	Reinf swab	28	Not assessed ^{††}	Not hospitalized	0	Unk
28	Male	40-44	Nepalese	Health facility	Day 0	First pos swab	Unk	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	40-44	Nepalese	Contact tracing	Day 89	Reinf swab	28	Not assessed ^{††}	Not hospitalized	0	Positive
29	Male	30-34	Pakistani	Survey ^{**}	Day 0	First pos swab	21	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	30-34	Pakistani	Survey ^{**}	Day 49	Reinf swab	28	Not assessed ^{††}	Not hospitalized	0	Unk
30	Male	15-19	Qatari	Health facility	Day 0	First pos swab	20	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	15-19	Qatari	Health facility	Day 64	Reinf swab	28	Not assessed ^{††}	Not hospitalized	0	Unk
31	Male	35-39	Nepalese	Health facility	Day 0	First pos swab	32	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	35-39	Nepalese	Survey ^{**}	Day 58	Reinf swab	29	Not assessed ^{††}	Not hospitalized	0	Unk
32	Male	35-39	Indonesian	Survey ^{**}	Day 0	First pos swab	17	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	35-39	Indonesian	Survey ^{**}	Day 49	Reinf swab	29	Not assessed ^{††}	Not hospitalized	0	Unk
33	Male	40-44	Egyptian	Health facility	Day 0	First pos swab	17	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	40-44	Egyptian	Health facility	Day 71	Reinf swab	29	Not assessed ^{††}	Not hospitalized	0	Unk
34	Male	40-44	Filipino	Health facility	Day 0	First pos swab	33	Not assessed ^{††}	1 day in May	1	Unk
	Male	40-44	Filipino	Contact tracing	Day 78	Reinf swab	29	Not assessed ^{††}	Not hospitalized	0	Unk
35	Male	35-39	Bangladeshi	Health facility	Day 0	First pos swab	25	Not assessed ^{††}	2 days in May-June	2	Unk
	Male	35-39	Bangladeshi	Health facility	Day 64	Reinf swab	29	Not assessed ^{††}	Not hospitalized	0	Positive
Good evidence for reinfection											
36	Male	30-34	Pakistani	Health facility	Day 0	First pos swab	22	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	30-34	Pakistani	Contact tracing	Day 57	Reinf swab	30	Not assessed ^{††}	Not hospitalized	0	Unk
37	Male	40-44	Sudanese	Health facility	Day 0	First pos swab	35	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	40-44	Sudanese	Health facility	Day 68	Reinf swab	30	Not assessed ^{††}	Not hospitalized	0	Unk
38	Male	25-29	Yemeni	Health facility	Day 0	First pos swab	35	Not assessed ^{††}	Not hospitalized	0	Unk

	Male	25-29	Yemeni	Contact tracing	Day 63	Reinf swab	30	Not assessed ^{††}	Not hospitalized	0	Unk
39	Male	20-24	Somali	Survey ^{**}	Day 0	First pos swab	Unk	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	20-24	Somali	Survey ^{**}	Day 4	Subs pos swab	Unk	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	20-24	Somali	Survey ^{**}	Day 14	Subs pos swab	36	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	20-24	Somali	Survey ^{**}	Day 55	Reinf swab	30	Not assessed ^{††}	Not hospitalized	0	Unk
40	Male	55-59	Bangladeshi	Health facility	Day 0	First pos swab	Unk	Mild	7 days in March [‡]	7	Unk
	Male	55-59	Bangladeshi	Health facility	Day 85	Reinf swab	31	Not assessed ^{††}	Not hospitalized	0	Positive
41	Male	30-34	Nepalese	Survey ^{**}	Day 0	First pos swab	26	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	30-34	Nepalese	Health facility	Day 14	Subs pos swab	36	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	30-34	Nepalese	Survey ^{**}	Day 50	Reinf swab	31	Not assessed ^{††}	Not hospitalized	0	Unk
42	Male	15-19	Pakistani	Health facility	Day 0	First pos swab	33	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	15-19	Pakistani	Health facility	Day 46	Reinf swab	32	Not assessed ^{††}	Not hospitalized	0	Unk
43	Male	15-19	Jordanian	Health facility	Day 0	First pos swab	Unk	Not assessed ^{††}	8 days in March [‡]	8	Unk
	Male	15-19	Jordanian	Contact tracing	Day 78	Reinf swab	32	Not assessed ^{††}	Not hospitalized	0	Unk
44	Male	25-29	Nepalese	Survey ^{**}	Day 0	First pos swab	30	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	25-29	Nepalese	Survey ^{**}	Day 20	Subs pos swab	34	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	25-29	Nepalese	Survey ^{**}	Day 55	Reinf swab	32	Not assessed ^{††}	Not hospitalized	0	Unk
45	Male	25-29	Nepalese	Contact tracing	Day 0	First pos swab	29	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	25-29	Nepalese	Contact tracing	Day 58	Reinf swab	33	Not assessed ^{††}	Not hospitalized	0	Unk
46	Male	20-24	Nepalese	Health facility	Day 0	First pos swab	20	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	20-24	Nepalese	Contact tracing	Day 58	Reinf swab	34	Not assessed ^{††}	Not hospitalized	0	Unk
47	Male	20-24	Nepalese	Survey ^{**}	Day 0	First pos swab	36	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	20-24	Nepalese	Survey ^{**}	Day 52	Reinf swab	34	Not assessed ^{††}	Not hospitalized	0	Unk
48	Male	35-39	Nepalese	Health facility	Day 0	First pos swab	20	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	35-39	Nepalese	Health facility	Day 69	Reinf swab	35	Not assessed ^{††}	Not hospitalized	0	Unk
49	Male	40-44	Nepalese	Survey ^{**}	Day 0	First pos swab	Unk	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	40-44	Nepalese	Health facility	Day 59	Reinf swab	35	Not assessed ^{††}	Not hospitalized	0	Unk
50	Male	50-54	Indian	Contact tracing	Day 0	First pos swab	Unk	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	50-54	Indian	Contact tracing	Day 52	Reinf swab	35	Not assessed ^{††}	Not hospitalized	0	Unk
51	Male	45-49	Indian	Health facility	Day 0	First pos swab	37	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	45-49	Indian	Survey ^{**}	Day 59	Reinf swab	36	Not assessed ^{††}	Not hospitalized	0	Positive
52	Male	30-34	Sudanese	Contact tracing	Day 0	First pos swab	Unk	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	30-34	Sudanese	Survey ^{**}	Day 78	Reinf swab	37	Not assessed ^{††}	Not hospitalized	0	Unk
53	Male	30-34	Indian	Contact tracing	Day 0	First pos swab	36	Not assessed ^{††}	Not hospitalized	0	Unk
	Male	30-34	Indian	Contact tracing	Day 45	Reinf swab	Unk	Not assessed ^{††}	Not hospitalized	0	Unk
54	Male	20-24	Sudanese	Health facility	Day 0	First pos swab	27	Not assessed ^{††}	17 days in April [‡]	17	Unk
	Male	20-24	Sudanese	Health facility	Day 15	Subs pos swab	36	Not assessed ^{††}	Inpatient	--	Unk
	Male	20-24	Sudanese	Survey ^{**}	Day 74	Reinf swab	Unk	Not assessed ^{††}	Not hospitalized	0	Unk

Ab, antibody; LOS, length of stay; Reinf, reinfection; PCR, polymerase chain reaction; Pos, positive; Subs, subsequent; Unk, unknown.

*Age was included as a range for anonymity.

[†]Precise swab dates not included for anonymity.

[‡]Severity classification per WHO guidelines was conducted only on a subset of all cases where it was deemed relevant. Asymptomatic cases or cases with minimal symptoms were not formally assessed for severity.

[§]It has been common to use hospitalization as a form of isolation especially early in the epidemic.

[¶]Precise antibody test dates not included for anonymity.

^{**}The category “survey” refers to surveillance testing campaigns conducted in workplaces and residential areas.

^{††}Not assessed because of no or minimal symptoms to warrant clinical assessment.

Table 2. Distribution of cases according to reinfection and antibody statuses (for individuals with complete information for these statuses).

Reinfection status	Seronegative	Seropositive	Total
	N (%)	N (%)	N (%)
Strong evidence for reinfection	7 (53.9)	6 (46.2)	13 (100.0)
Good evidence for reinfection	0 (0.0)	2 (100.0)	2 (100.0)
Some evidence for reinfection	1 (25.0)	3 (75.0)	4 (100.0)
Weak evidence for reinfection	10 (34.5)	19 (65.5)	29 (100.0)
Total N (%)	18 (37.5)	30 (62.5)	48 (100.0)

Author contributions

LJA conceived and co-designed the study and led the statistical analyses. HC co-designed the study, performed the data analyses, and wrote the first draft of the article. HHA contributed to analysis of data. All authors contributed to data collection and acquisition, database development, discussion and interpretation of the results, and to the writing of the manuscript. All authors have read and approved the final manuscript.

Competing interests

We declare no competing interests.

Acknowledgement

We would like to thank Her Excellency Dr. Hanan Al Kuwari, the Minister of Public Health, for her vision, guidance, leadership, and support. We also would like to thank Dr. Saad Al Kaabi, Chair of the System Wide Incident Command and Control (SWICC) Committee for the COVID-19 national healthcare response, for his leadership, analytical insights, and for his instrumental role in enacting the data information systems that made these studies possible. We further extend our appreciation to the SWICC Committee and the Scientific Reference and Research Taskforce (SRRT) members for their informative input, scientific technical advice, and enriching discussions. We also would like to acknowledge the dedicated efforts of the Clinical Coding Team and the COVID-19 Mortality Review Team, both at Hamad Medical Corporation, and the Surveillance Team at the Ministry of Public Health.

Funding

The authors are grateful for support provided by the Ministry of Public Health, Hamad Medical Corporation, and the Biomedical Research Program and the Biostatistics, Epidemiology, and

Biomathematics Research Core, both at Weill Cornell Medicine-Qatar. The statements made herein are solely the responsibility of the authors.

References

1. World Health Organization (WHO), *WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020*. Available from: <https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020>. Accessed on March 14, 2020. 2020.
2. De Walque D., et al., *How two tests can help contain COVID-19 and revive the economy*. Available from: <http://documents.worldbank.org/curated/en/766471586360658318/pdf/How-Two-Tests-Can-Help-Contain-COVID-19-and-Revive-the-Economy.pdf>. Accessed on April 16, 2020. Research & Policy Briefs, World Bank Malaysia Hub., 2020.
3. Kaplan J., Frias L., and McFall-Johnsen M., *A third of the global population is on coronavirus lockdown*. Available from: <https://www.businessinsider.com.au/countries-on-lockdown-coronavirus-italy-2020-3> Accessed on: April 25, 2020. Business Insider Australia, 2020.
4. Planning and Statistics Authority- State of Qatar, *Qatar Monthly Statistics*. Available from: <https://www.psa.gov.qa/en/pages/default.aspx>. Accessed on: may 26,2020. 2020.
5. Planning and Statistics Authority-State of Qatar, *The Simplified Census of Population, Housing & Establishments*. Available from: https://www.psa.gov.qa/en/statistics/Statistical%20Releases/Population/Population/2018/Population_social_1_2018_AE.pdf Accessed on: April 2, 2020. 2019.
6. Abu-Raddad, L.J., et al., *Characterizing the Qatar advanced-phase SARS-CoV-2 epidemic*. medRxiv, 2020: p. 2020.07.16.20155317v2.
7. Al Kuwari, H.M., et al., *Characterization of the SARS-CoV-2 outbreak in the State of Qatar, February 28-April 18, 2020*. medRxiv, 2020: p. 2020.07.15.20154211.
8. Seow, J., et al., *Longitudinal evaluation and decline of antibody responses in SARS-CoV-2 infection*. medRxiv, 2020: p. 2020.07.09.20148429.
9. Wajnberg, A., et al., *SARS-CoV-2 infection induces robust, neutralizing antibody responses that are stable for at least three months*. medRxiv, 2020: p. 2020.07.14.20151126.
10. Kellam, P. and W. Barclay, *The dynamics of humoral immune responses following SARS-CoV-2 infection and the potential for reinfection*. J Gen Virol, 2020.
11. Makhoul, M., et al., *Epidemiological impact of SARS-CoV-2 vaccination: mathematical modeling analyses*. medRxiv, 2020: p. 2020.04.19.20070805.
12. World Health Organization, *Clinical management of COVID-19*. Available from: <https://www.who.int/publications-detail/clinical-management-of-covid-19>. Accessed on: May 31st 2020. 2020.
13. Kalikiri, M.K.R., et al., *High-throughput extraction of SARS-CoV-2 RNA from nasopharyngeal swabs using solid-phase reverse immobilization beads*. medRxiv, 2020: p. 2020.04.08.20055731.
14. Sethuraman, N., S.S. Jeremiah, and A. Ryo, *Interpreting Diagnostic Tests for SARS-CoV-2*. JAMA, 2020.
15. Wajnberg, A., et al., *Humoral immune response and prolonged PCR positivity in a cohort of 1343 SARS-CoV-2 patients in the New York City region*. medRxiv, 2020: p. 2020.04.30.20085613.
16. World Health Organization, *Criteria for releasing COVID-19 patients from isolation*. Available from: <https://www.who.int/news-room/commentaries/detail/criteria-for-releasing-covid-19-patients-from-isolation>. Accessed on July 01, 2020. 2020.