1

Functional signatures in non-small-cell lung cancer: a systematic review and meta-analysis of sex-based differences in transcriptomic studies

Irene Pérez-Díez[1,2], Marta R. Hidalgo[1], Pablo Malmierca-Merlo[1,3], Zoraida Andreu[1], Rosa Farràs[4], María de la Iglesia-Vayá[2], Mariano Provencio[5], Atocha Romero[5] and Francisco García-García[1,6]*

[1]Bioinformatics and Biostatistics Unit, Principe Felipe Research Center (CIPF). Valencia, Spain.
[2]Biomedical Imaging Unit FISABIO-CIPF. Fundación para el Fomento de la Investigación Sanitario y Biomédica de la Comunidad Valenciana. Valencia, Spain.
[3]Atos Research Innovation (ARI), Spain.
[4]Department of Oncogenic Signalling, Principe Felipe Research Center (CIPF). Valencia, Spain.
[5]Medical Oncology Department, Hospital Universitario Puerta de Hierro-Majadahonda, Madrid, Spain.
[6]Spanish National Bioinformatics Institute, ELIXIR-Spain (INB, ELIXIR-ES).
* Correspondence: fgarcia@cipf.es

Abstract: While studies have established the existence of differences in the epidemiological and clinical patterns of lung adenocarcinoma between male and female patients, we know relatively little regarding the molecular mechanisms underlying such sex-based differences. In this study, we explore said differences through a meta-analysis of transcriptomic data. We performed a meta-analysis of the functional profiling of nine public datasets that included 1,366 samples from Gene Expression Omnibus and The Cancer Genome Atlas databases. Meta-analysis results from data merged, normalized, and corrected for batch effect show an enrichment for Gene Ontology terms and Kyoto Encyclopedia of Genes and Genomes pathways related to the immune response, nucleic acid metabolism, and purinergic signaling. We discovered the overrepresentation of terms associated with the immune response, particularly with the acute inflammatory response, and purinergic signaling in female lung adenocarcinoma patients, which could influence reported clinical differences. Further evaluations of the identified differential biological processes and pathways could lead to the discovery of new biomarkers and therapeutic targets. Our findings also emphasize the relevance of sex-specific analyses in biomedicine, which represents a crucial aspect influencing biological variability in disease.

Keywords: NSCLC; biomarkers; functional profiling; meta-analysis; sex characteristics

1. Introduction

Lung cancer is the most frequently diagnosed cancer and the leading cause of cancer-related death worldwide, representing 18.4% of all cancer deaths [1]. Exposure to tobacco, domestic biomass fuels, asbestos, and radon represent the most relevant lung cancer risk factors [1–3]; however, as has become evident from studies of other cancer types, sex-based differences (sexual dimorphisms) may also have significant relevance in lung cancer [1,4,5].

Lung cancer exhibits sex-based disparities in clinical characteristics and outcomes, with better survival observed in women [3,6,7]. While lung cancer incidence worldwide is higher in men, there exists an increasing trend in women that cannot be solely explained by tobacco consumption [1,2]. Furthermore, studies have reported sex-dependent differences in estrogen receptors and their impact on lung cancer [8–10]; however, conflicting results have attributed lung cancer susceptibility in women to genetic variants, hormonal factors, and molecular abnormalities, and oncogenic viruses [3,11–13].

Adenocarcinoma represents the most frequent non-small cell lung cancer (NSCLC) subtype in
both sexes [14], with a higher predominance in women compared to men (41% of cases in women
versus 34% in men) [3,7,13]. Interestingly, Wheatley-Price et al. demonstrated a more pronounced
survival rate difference between male and female lung adenocarcinoma patients when compared to
other tumor types [7]. The molecular causes underlying such sex-biased patterns remain still largely
unknown as limited efforts have been made for lung adenocarcinoma, with few studies considering
this differential perspective [15–18]. These limitations can be partially addressed through meta-
analysis, a robust methodology that combines information from related but independent studies to
derive results with increased statistical power and precision [19,20]. As current treatment strategies
do not cure most lung cancer patients, and aggressive diagnostic techniques (e.g., via biopsies and
bronchoscopies) often induce discomfort in patients, meta-analyses that improve our
understanding of sex-specific molecular mechanisms in lung adenocarcinoma may facilitate the
discovery of non-invasive prognostic and diagnostic biomarkers.

We performed a meta-analysis based on functional profiles of transcriptomic studies to explore
the molecular mechanisms underlying sex-based differences in early-stage lung adenocarcinoma.
We carried out exhaustive review and selection steps to guarantee the homogeneity of the selected
studies and the subsequent comparison and integration of the data in the meta-analysis with an
appreciation of this strategy's specific limitations. After the systematic review, we retrieved and
analyzed nine studies from Gene Expression Omnibus (GEO) [21] and The Cancer Genome Atlas
(TCGA) [22], and then combined the results in a random-effects meta-analysis. This approach
allowed the identification of functional alterations caused by lung adenocarcinoma in both male
and female patients. In this study, we identified immune responses, purinergic signaling, and lipid
metabolism as the main biological processes that display differences between male and female lung
adenocarcinoma patients, with the acute immune response increased in female patients. Overall,
our findings provide evidence that sex-based differences influence cancer biology and may impact
response to treatment. Furthermore, underlying sex-based differences may contribute to the
discovery of sex-specific prognostic and diagnostic biomarkers and the improvement of
personalized therapies.
80
81   **2. Results**
82
83   We organized our findings into three sections: the first describes the studies evaluated and
84   selected in the systematic review; the second section reports on the results of the bioinformatic
85   analysis of each of these selected studies as follows: i) exploratory analysis, ii) differential
86   expression, and iii) functional characterization; while the third section presents the results of the
87   differential functional profiling by sex.
88
89   *2.1. Study Search and Selection*
90
The systematic review identified 207 non-duplicated studies, of which 48.8% included both
male and female patients (**Figure S1**). We applied inclusion and exclusion criteria (see **Figure 1**) to
select a set of homogeneous and comparable studies to ensure the reliability of the subsequent
analyses. To ensure study homogeneity (and in the hope of contributing to the early diagnosis of
disease), we focused on those studies of early-stage disease and selected nine transcriptomic studies
for further analysis (**Table 1**). The selected studies represented a population of 1,366 early-stage
samples (369 controls and 997 cases), of which 44% were from men and 56% from women (**Figure
2**), with a median age of 65.54 years old. **Table 1**, **Figure 2**, and **Supplementary Table S1** contain
further information regarding the selected studies and the clinicopathological characteristics of the
study population.
101
102
103

104
105   **Table 1.** Studies selected after the systematic review.
106

Study	Platform	Publication
GSE10072	Affymetrix Human Genome U133A Array	[23]
GSE19188	Affymetrix Human Genome U133 Plus 2.0 Array	[24]
GSE31210	Affymetrix Human Genome U133 Plus 2.0 Array	[25,26]
GSE32863	Illumina HumanWG-6 v3.0 Expression BeadChip	[27]
GSE63459	Illumina HumanRef-8 v3.0 Expression BeadChip	[28]
GSE75037	Illumina HumanWG-6 v3.0 Expression BeadChip	[29]
GSE81089	Illumina HiSeq 2500	[30]
GSE87340	Illumina HiSeq 2000	[31]
TCGA	Illumina HiSeq 2000	[22]

107
108   *2.2. Individual Analysis*
109
As the normalized data derives from different platforms, we performed exploratory and
processing steps for the data set to ensure the comparability and integration of subsequent analyses.
The exploratory analysis found a lack of abnormal behavior except for three samples in the
principal component analysis (PCA) and unsupervised clustering; therefore, we excluded the
GSM47570 and GSM47578 samples in study GSE19188, and the GMS773784 sample in study
GSE31210 from further analysis.
The differential expression results for each study demonstrated a large number of differentially
expressed genes when comparing female lung adenocarcinoma patients to female control patients
and male adenocarcinoma patients to male control patients (**Supplementary Table S2**). However,
the evaluation of sex-based differences in lung adenocarcinoma patients provided a small number
of significantly affected genes (see **Supplementary Table S3**), with no intersecting genes.
We performed an individual functional enrichment analysis of Gene Ontology (GO) terms and
Kyoto Encyclopedia of Genes and Genomes (KEGG) to identify the possible implications of these
sex-specific differentially-expressed genes in pathways relevant to lung adenocarcinoma. The
identified pathways revealed a diversity of significant results among datasets, which we have
summarized in **Table 2**. When analyzing intersections, UpSet plots (**Figure 3,** equivalent to a Venn
diagram) illustrate the degree of intersection between studies, demonstrating that most significant
results are exclusive of each study. This data highlights the need for integrated strategies such as
meta-analysis to increase the statistical power of any findings.
129

Figure 1. Flow of information through the different phases of the systematic review, following PRISMA Statement guidelines [32].

2.3. Meta-analysis

We performed a functional meta-analysis for each of the 8,672 GO functions and KEGG pathways, including every term found in at least two studies. Results with a false discovery rate (FDR) of < 0.05 included 106 GO biological processes (BP), 3 GO molecular functions (MF), and 20 KEGG pathways, which were associated with 21 wider functional groups. We rejected potential bias on the significant results after the inspection of funnel plots; furthermore, sensitivity analyses failed to indicate alterations in the results due to the inclusion of any study. The results for the 129 significant GO terms and KEGG pathways are further detailed in **Supplementary Table S4**, including FDR, the log odds ratio (LOR), and its 95% confidence interval (CI), and the standard error (SE) of the LOR.

2.3.1. Upregulated Functions

We discovered that 43.88% of detected functions related to the immune response (**Supplementary Table S4** and **Figure 4**), which all displayed upregulation in female lung adenocarcinoma patients. This finding agrees with studies that report more robust innate and adaptive immune responses in women who, for example, present with more efficient antigen-presenting cells (APCs) than males [33,34]. The results provided evidence for the positive regulation of an acute inflammatory response, with CD8+ alpha-beta T cell differentiation and activation, B cell proliferation and activation, and an increase of interleukin (IL) biosynthesis, including IL-2, 6, 8, 10, and 17. Several immune-related signaling pathways also displayed differences between female and

male lung adenocarcinoma patients. We discovered the upregulation of the MyD88-independent
toll-like receptor signaling pathway, NIK/NF-kappa β signaling pathway, FC-epsilon receptor
signaling pathway, B-cell receptor signaling pathway, Toll-like receptor signaling pathway, NOD-
like receptor signaling pathway, and RIG-I-like receptor signaling pathways in female lung
adenocarcinoma patients. Overall, these findings suggest the relevance of sex-based differences in
immune responses to lung cancer, which may represent a significant contributor to the marked
differences observed during disease progression in male and female patients. Furthermore, such
findings may help to define novel therapeutic targets and may have important implications for
immunotherapy.

We also uncovered evident sex-based differences in cell metabolism. Studies have shown
the significant upregulation of lipid metabolism (ceramide and sphingolipid biosynthetic processes)
in females and the higher utilization of carbohydrates by males [35,36]. Furthermore, lipid
metabolism and signaling are widely accepted as major players in cancer biology [37]. "Metabolism-
Nucleic acids metabolism and signaling" was the second most abundant functional group
upregulated in female lung adenocarcinoma patients, comprising 8.63% of the altered functions.
These GO terms and KEGG pathways are mainly related to purinergic signaling through G protein-
coupled receptors and cytidine metabolism. Other functional groups upregulated in female lung
adenocarcinoma patients include cell migration and homeostasis. Overall, further explorations of
sex differences in metabolic pathways may provide new perspectives for treatment approaches
with sex-specific effects.

176
177   2.3.2. Downregulated Functions

178
23.26% of the significant functions exhibited lower activity in female compared to male lung
adenocarcinoma patients. Downregulated functional groups include those related to cell cycle
progression, cell junctions, DNA repair and telomere protection, mitochondrial processes, neural
development, post-translational changes, post-transcriptional changes, protein degradation, and
transcription regulation (**Supplementary Table S4** and **Figure 4**). Taken together, the
downregulated pathways in female patients suggest the existence of lower levels of oxidative stress
compared to male patients [38], which may contribute to the existence of a less permissive
tumorigenic environment.

187
188   *2.4. Metafun-NSCLC Web Tool*

189
The Metafun-NSCLC web tool (https://bioinfo.cipf.es/metafun-nsclc) contains
information related to the nine studies and 1,329 samples involved in this study. For each study,
this resource includes fold-changes of genes and LOR of functions and pathways that users can
explore to identify profiles of interest.
We carried out a total of 8,672 meta-analyses. For each of the 129 significant functions and
pathways, Metafun-NSCLC depicts the global activation level for all studies and each study's
specific contribution, using statistical indicators (LOR, CI, and p-value) and graphical
representations by function as forest and funnel plots. This open resource hopes to contribute to
data sharing between researchers, the elaboration of innovative studies, and the discovery of new
findings.
Here, we also highlight the importance of including/reporting sex-related data in the
results of clinical studies given their general importance in tumor risk, treatment response, and
outcomes in lung adenocarcinoma and other cancers/disorders. The integration of sex-based
differences in this manner has the potential to significantly impact the cancer biology field.
204

205
206   **Figure 2.** Number of samples per study, divided by sex and experimental group (ADC – lung
207   adenocarcinoma samples).

208
209   **Table 2.** Summary of functional enrichment analysis results by Gene Ontology functions (BP:
210   Biological Process, MF: Molecular Functions, CC: Cellular Component), and KEGG pathways. "Up"
211   terms are overrepresented in female lung adenocarcinoma patients, while "Down" terms are
212   overrepresented in male lung adenocarcinoma patients.
213

	Significant GO BP terms			Significant GO MF terms			Significant GO CC terms			Significant KEGG pathways		
	Up	Down	Total	Up	Down	Total	Up	Down	Total	Up	Down	Total
GSE10072	26	153	179	29	8	37	16	40	56	1	5	6
GSE19188	7	12	19	0	3	3	11	20	31	1	4	5
GSE31210	21	2	23	0	0	0	4	0	4	8	2	10
GSE32863	428	51	479	28	5	33	40	41	81	27	6	33
GSE63459	0	26	26	0	3	3	8	27	35	1	4	5
GSE75037	245	35	280	14	4	18	14	21	35	22	5	27
GSE81089	2	1	3	7	1	8	7	4	11	1	1	2
GSE87340	178	62	240	3	0	3	48	13	61	26	1	27
TCGA	294	228	522	28	30	58	21	70	91	30	17	47

Figure 3. The intersection of significant functions between studies. UpSet plots for **(a)** GO biological process, **(b)** GO molecular functions, **(c)** GO cellular components, and **(d)** KEGG pathways. UpSet plots detailing the number of common elements among GO terms in our functional enrichment analysis. Horizontal bars indicate the number of significant elements in each study. The vertical bars indicate the common elements in the sets, indicated with dots under each bar. The single points represent the number of unique elements in each group.

Figure 4. Summary dot plot of GO BP meta-analysis results. Only those significant terms with a LOR over 0.4 are shown.

225   **3. Discussion**
226
227   Despite the profuse evidence for the influence of sex on rates and patterns of metastasis, the
expression of prognostic biomarkers, and therapeutic responses in several cancer types [39,40], sex-
based differences have not been consistently considered when studying cancer, designing therapies,
or constructing clinical trials. Cases of NSCLC, including adenocarcinoma, exhibit differences in
incidence, prevalence, and severity in female and male patients [1,3,7,41]. Elucidating the molecular
basis for this sex-based differential impact will have clinical relevance, as this information can
guide/improve both diagnosis and treatment.
234   Biomedical research generally underrepresents female patients, with sex-based differences
rarely considered [42,43]. Our systematic review of transcriptomic studies revealed that only 48.8%
of lung adenocarcinoma-related datasets considered both sexes, a figure similar (49%) to that
reported by Woitowich et al. [43]. Sex-based differences impact disease biomarkers, drug responses,
and treatment [42], and, therefore, sex must represent a critical component of experimental design.
Added to this problem, we faced a lack of standardization among studies and detailed clinical
information (i.e., mutations, smoking status, stages) when searching for suitable datasets. The
consideration of Findable, Accessible, Interoperable, and Reusable (FAIR) data principles [44], a
requisite for quality science, would ensure that generated data can be of further use throughout the
scientific community.
To the best of our knowledge, only four studies have attempted to address the functional alterations
caused by lung adenocarcinoma in both male and female patients – those by Shi et al. [16], which
considered female patients, Araujo et al. [15], Yuan et al. [17], and Li et al. [18]. Shi et al. [16]
integrated samples from two datasets for a differential expression analysis followed by a functional
enrichment analysis, whereas Araujo et al. [15] independently processed six datasets and jointly
analyzed their results. Yuan et al. [17] compared male and female patients with various cancer
types (including lung adenocarcinoma) using the TCGA dataset, but did not include control
samples in the statistical comparison. Li et al. [18] evaluated the differences in lung adenocarcinoma
by focusing only on metabolic pathways. By including both male and female patients and controls
in our gene expression comparison, in contrast with the analysis performed by Shi et al. [16] and
Yuan et al. [17], we have effectively unveiled sex-based differences occurring in lung
adenocarcinoma. Of note, 88% of the results reported by Yuan et al. [17] relate to the sex
chromosome, but not necessarily due to cancer. Shi et al. [16] described the consequences of cancer
development in female patients, but the authors failed to compare said effects with male patients.
Although Araujo et al. [15] do not describe the statistical comparisons performed (and do not
perform a statistical integration such as meta-analysis), the authors describe the results obtained for
each dataset. Our study addressed sex-based differences in male and female lung adenocarcinoma
patients through meta-analysis to address previous limitations and improve on those approaches
employed by others. Despite certain supposed limitations to our approach (the presence of studies
with different sample sizes and types of platforms), meta-analyses can integrate selected studies by
eliminating inconsistency in individual studies, thereby increasing the statistical power and
highlighting robust disease-associated functions. We performed a meta-analysis using a random-
effects model on Gene Set Enrichment Analysis (GSEA) results independently obtained from each
study to evaluate the functions differentially altered between male and female lung
adenocarcinoma patients. To make results comparable and reduce biases in the type of analysis
used, we applied the same bioinformatics strategy from normalized expression matrices to GSEA

results. This robust methodology integrates groups of data and provides results with higher
statistical power and precision [19,20] and reveals findings that cannot be obtained through the
intersection or addition of results in individual studies. Nevertheless, we selected only samples
from early-stage lung adenocarcinoma patients to reduce variability and included smoking status in
the differential expression linear model. While the inclusion of the mutational status of relevant
genes to lung adenocarcinoma (e.g., EGFR) could have revealed important insight, the lack of this
information in the majority of the studies and the resultant limited sample size hampered this aim.
We would support the inclusion of this type of data as a requirement for the publication of new
data to repositories such as the GEO.

The immune system plays a crucial role in the development of cancer [45], and several studies
have reported sex-based differences in immune responses (Reviewed by Klein and Flanagan [33]).
Tumor cells evade the immune system using different strategies, including the modulation of
antigen-presentation and the suppression of regulatory T cells. Therefore, sex differences in APCs
and their downstream effector cells, among other components, may contribute to the sexual
disparity observed in various aspects of cancer development and may significantly impact
antitumor immunity and immunotherapy. Adult females generally present more robust innate and
adaptive immune responses than males, as evidenced by increased phagocytic activity of
neutrophils and macrophages, more efficient APCs, and differences in lymphocyte subsets (B cells,
CD4+T cells, CD8+T cells) and cytokine production. Accordingly, our results demonstrate an
enrichment of immune response-related terms in female lung adenocarcinoma patients, which
agrees with the findings of Araujo et al. [15]. The analyzed functions suggest the positive regulation
of CD8+ alpha-beta T cell activation and differentiation in female lung adenocarcinoma patients,
which play an essential role in antitumor immunity [46,47]. Furthermore, Ye et al. [47] discovered a
more abundant population of effector memory CD8+ T cells in female lung adenocarcinoma
patients, which agrees with our results. A previous study described CD8+ lymphocyte levels as a
prognostic biomarker in NSCLC [48], and specifically in lung adenocarcinoma [49], with a
correlation between higher levels of CD8+ lymphocytes with higher survival rates and lower
disease recurrence. Elevated levels of active CD8+ T cells in female lung adenocarcinoma patients
could form part of the molecular mechanism underlying higher survival rates when compared to
male lung adenocarcinoma patients. Activation of the Notch signaling pathway decreases CD8+ T
lymphocyte activity in lung adenocarcinoma [50]; therefore, the downregulation of the Notch
signaling pathway discovered in female lung adenocarcinoma patients could explain higher CD8+
T activity when compared to male lung adenocarcinoma patients.

Concerning the immune response, we also detected differences that supported the increased
production of IL-2, which is known to stimulate T cell proliferation and the production of effector T
cells, thereby amplifying the lymphocytic response [51]. Higher levels of IL-2 could also relate to
increased activity of CD8+ T cells in female lung adenocarcinoma patients. Increased levels of IL-10
are also supported in female lung adenocarcinoma patients and, although IL-10 has anti-
inflammatory and anti-immune activities [52,53], studies have suggested a dual role in cancer. In
advanced lung adenocarcinoma, high expression of IL-10 receptor 1 correlates with worse
prognosis [52], while IL-10 expression by T-regulatory cells inhibits apoptosis through Programmed
death-ligand 1 inhibition [53]. Nevertheless, IL-10 correlates with better prognosis when expressed
by CD8+ T cells in early-stage NSCLC [54], and it seems to activate the antitumor control of CD8+ T
cells [55]. IL-2 and IL-10 could display increased activity in early-stage female patients, alongside a
higher population of active CD8+ T cells than males, conferring women a survival advantage.

We also detected the positive regulation of IL-6 biosynthesis in female lung adenocarcinoma
patients, with increased IL-6 levels correlating with worse prognosis in NSCLC patients in previous
studies [56,57]. Network analysis in non-smoking female lung adenocarcinoma patients described
IL-6 as one of the pathology's central nodes [16], and these findings agree with our results, which
provide evidence of the critical role of IL-6 in tumor progression in female lung adenocarcinoma

patients. IL-8 and IL-17 exhibit increased production and biosynthesis in female lung
adenocarcinoma patients, with said interleukins known to influence tumor growth and metastasis
and correlate with worse prognosis [58–60].

Although altered immune responses can positively and negatively influence tumor
progression, our findings have detected GO terms that point to an elevated acute immune response
in female compared to male lung adenocarcinoma patients. Of note, sex-based immunological
differences in lung adenocarcinoma might have an impact on immunotherapy response. Different
studies have addressed the role of sex in immunotherapy [5,47,61,62] and established improved
survival for female NSCLC patients. The discovered molecular pathways differentially activated
between male and female lung adenocarcinoma patients may underlie phenotypic differences
regarding immunotherapy response.

Sex-based differences in metabolism occur under physiological conditions and in the
presence of cancer. Here, we have detected an upregulation of purinergic signaling and nucleic acid
metabolism in female lung adenocarcinoma patients, a finding not described in previous lung
cancer studies. An NSLC-based study described an antitumor effect of the P2X4 receptor [63],
which also exhibits sexual dimorphism in murine brain microglia [64]. Other P2 and A2 receptors
play a role in NSCLC [63], but evidence of sex-based differences in receptor expression in human
NSCLC patients has yet to be reported. Purinergic signaling and the role of purinergic receptors
may also have relevance to innate and adaptative responses in different inflammatory and
neurodegenerative diseases and several cancer types, including pancreatic ductal carcinoma
(PDAC), hepatocellular, hepatobiliary carcinoma cells, and breast cancer [65–69]. Of note, studies
have linked the upregulation of purinergic signaling with poor prognosis in PDAC [66]. We also
discovered significant differences in lipid metabolism, with the positive regulation of ceramide and
sphingolipid biosynthetic processes upregulated in female lung adenocarcinoma patients. The
presence of lipids can promote tumorigenesis, while higher adipose tissue levels are associated with
poorer outcomes in several cancers [37]. Thus, exploring the differential roles of purinergic
signaling and lipid metabolism between male and female lung adenocarcinoma patients may
represent an interesting proposition to improve sex-specific risk-stratification of patients,
prevention, diagnosis, and treatment.

DNA damage and repair-related genes also presented sex-based differences in lung
adenocarcinoma patients. In general, males present with a higher level of DNA damage, and
females present with a lower DNA repair capacity [70], and, in agreement, we detected DNA repair
and telomere protection as a downregulated functional group in female lung adenocarcinoma
patients (with both mechanisms involved in tumor growth prevention [71]). Furthermore, we
discovered the upregulation of DNA repair and an increase of DNA unwinding in male lung
adenocarcinoma patients. DNA unwinding has emerged as a new target in cancer therapy with a
primary focus on helicase inhibitors [72]. Besides, regarding DNA repair,the evaluation of poly
(ADP-ribose) polymerase inhibitors in NSCLC cell lines has suggested potential therapeutic activity
[73,74], and there may be value in exploring both treatment approaches in NSCLC patients,
especially male patients.

With these results in mind, we propose future studies focused on DNA repair and
lipid/purinergic metabolism in female lung adenocarcinoma patients and the immune response in
male lung adenocarcinoma patients in the hope of developing enhanced therapeutic strategies.

Our study has characterized functional differences between the sexes in lung adenocarcinoma,
shedding light on the functional basis behind this pathology in male and female patients. While our
meta-analysis confirmed the conclusions of other studies, we also report previously undescribed
alterations in biological processes that may broaden this field of study. Further knowledge
regarding how those factors related to the functional mechanisms described above differentially
impact male and female lung adenocarcinoma patients may improve our understanding of the
disease and improve treatment and diagnosis through biomarker identification.

370
371
372
373
374   **4. Materials and Methods**
375
376   Bioinformatics and statistical analysis employed R software v.3.5.3 [75]. **Supplementary Table**
377   **S5** details R packages and versions.
378
379   *4.1. Study Search and Selection*
380
Publicly available datasets were collected from GEO [21], ArrayExpress [76], and TCGA
[22]. A systematic search of studies published in the period 2004-2018 was conducted in 2019
following the preferred reporting items for systematic reviews and meta-analyses (PRISMA)
guidelines [32]. Two researchers involved in the study carried out the literature search, and the
consistency of the review and selection procedures used was evaluated and confirmed. Several
keywords were employed in the search, including lung adenocarcinoma (ADC), non-small-cell lung
carcinoma (NSCLC), Homo sapiens, and excluding cell lines.
Eleven variables were considered for each study, including the clinical characteristics of the
patients (e.g., sex and smoking habit) and experimental design (e.g., sample size and sample
extraction source). The final inclusion criteria were:
391
392   Sex, disease stage, and smoking habit variables registered
393   RNA extracted directly from human lung biopsies
394   Both normal and lung adenocarcinoma samples available
395   Patients had not undergone treatment before biopsy
396   Sample size of > 3 for case and control groups in both sexes
397
398   Finally, normalized gene expression data of six array NSCLC datasets (GSE10072, GSE19188,
399   GSE31210, GSE32863, GSE63459, and GSE75037) and counts matrix of three RNA-seq NSCLC
400   datasets (GSE81089, GSE87340, and TCGA-LUAD) were retrieved.
401
402   *4.2. Individual Transcriptomics Analysis*
403
Individual transcriptomics analysis consisted of three steps: pre-processing, differential
expression analysis, and functional enrichment analysis (**Figure 5a**).
Data pre-processing included the standardization of the nomenclature of the clinical variables
included in each study, normalization of RNA-seq counts matrix, and exploratory analysis. RNA-
seq counts were pre-processed with the *edgeR* [77] R package using the trimmed mean of m-values
(TMM) method [78]. We assessed the normalization methods performed by the original authors for
each dataset, and $\log_2$ transformed the matrices when necessary. Annotation from probe set to
Entrez identifiers from the National Center for Biotechnology Information [79] database and gene
symbol was carried out with the *biomaRt* [80] R package. When dealing with duplicated probe-to-
Entrez mappings, the median of their expression values was calculated. The exploratory analysis
included unsupervised clustering and PCA to detect patterns of expression between samples and
genes and the presence of batch effects in each study (**Figure 5b**).
Differential expression analyses were performed using the *limma* [81] R package. To detect
differentially expressed genes in male and female lung adenocarcinoma patients, the following
contrast was applied:
419
                    (ADC.W - Control.W) - (ADC.M - Control.M)

where ADC.W, Control.W, ADC.M and Control.M correspond to lung adenocarcinoma affected
women, control women, lung adenocarcinoma affected men, and control men, respectively. Paired
samples design was implemented, and tobacco consumption was included as a batch effect on the
limma linear model to reduce its impact on data. P-values were calculated and corrected for FDR
[82]. This comparison allows the detection of genes and functions altered by the disease and that
have higher or lower activity in women when compared to men. Significant functions and genes
were considered when FDR < 0.05.
Functional enrichment analyses were performed using the Gene Set Enrichment Analysis
(GSEA) implemented in the *mdgsa* [83] R package. P-values were, again, corrected for FDR. For
functional annotation, two functional databases were used: the KEGG Pathway database [84] and
GO [85]. GO terms were analyzed and propagated independently for each GO ontology: BPs, MFs,
and cellular components (CC). Those annotations excessively specific or generic were filtered out,
keeping functions with blocks of annotations between 10 and 500. Intersections within groups were
analyzed with UpSet plots [86] (**Figure 5c**).
434

Figure 5. Workflow and analysis design. **(a)** Summary of the analysis design followed in this work,
(b) PCA plot and HC analysis as an example of exploratory analysis (to explore possible batch
effects and to assure expected data behavior) performed at the pre-processing stage to assess for the
integrity of the data, **(c)** example of UpSet plot as an intersection analysis for functional enrichment
analysis results, and **(d)** examples of forest and funnel plots to assess meta-analysis results.

4.3. Functional Meta-analysis

Functional GSEA results were integrated into a functional meta-analysis [87] implemented
with *mdgsa* and *metafor* [88] R packages. Meta-analysis was applied under the DerSimonian & Laird
random-effects model [89], taking into account individual study heterogeneity. This model
considers the variability of individual studies by increasing the weights of studies with less
variability when computing meta-analysis results. Thus, the most robust functions between studies
are highlighted.
A total of 6,467 GO BP terms, 785 GO CC terms, 1207 GO MF terms, and 213 KEGG pathways
were evaluated. P-values, FDR corrected p-values, LOR, and 95% CIs of the LOR were calculated
for each evaluated function. Functions and pathways with FDR < 0.05 were considered significant,
and both funnel and forest plots were computed for each (**Figure 5d**). These representations were
checked to assess for possible biased results, where LOR represents the effect size of a function, and
the SE of the LOR serves as a study precision measure [90]. Sensitivity analysis (leave-one-out
cross-validation [88]) was conducted for each significant function to verify possible alterations in
the results due to the inclusion of any study.

4.4. Metafun-NSCLC Web Tool

All data and results generated in the different steps of the meta-analysis are available in the
Metafun-NSCLC web tool (https://bioinfo.cipf.es/metafun-nsclc), which is freely accessible to any
user and allows the confirmation of the results described in this manuscript and the exploration of
other results of interest.
The front-end was developed using the Angular Framework. All graphics used in this web
resource have been implemented with Plot.ly except for the exploratory analysis cluster plot, which
was generated with ggplot2 [91].
This easy-to-use resource is divided into five sections: 1) Summary of analysis results in
each phase. Then, for each of the studies, the detailed results of the 2) exploratory analysis, 3)
differential expression, and 4) functional profiling. The user can interact with the web tool through
graphics and tables and search for specific information for a gene or function. Finally, section 5)
provides several indicators for the significant functions identified in the meta-analysis that inform
whether they are more active in men or women.

5. Conclusions

Sex-based molecular differences may influence the incidence and outcome of lung
adenocarcinoma and, therefore, may have important clinical implications. We identified immune
responses, purinergic signaling, and lipid-related processes as the main biological processes altered
between male and female lung adenocarcinoma patients by a meta-analysis of transcriptomic
datasets. Said processes exhibit increased activity in female lung adenocarcinoma patients, whereas
other processes (such as DNA repair) are more active in male lung adenocarcinoma. Although
further studies are required to verify and fully explore these findings, our results provide new clues
to understand the molecular mechanisms of sex-based differences in lung adenocarcinoma patients
and new perspectives regarding the identification of biomarkers and therapeutic targets.

Data Availability*:* The data used for the analyses described in this work is publicly available at GEO (https://
www.ncbi.nlm.nih.gov/geo/) and the Genomic Data Commons Data Portal (TCGA,
https://portal.gdc.cancer.gov/). The accession numbers of the GEO datasets downloaded are: GSE10072,
GSE19188, GSE31210, GSE32863, GSE63459, GSE75037, GSE81089 and GSE87340. Only the samples associated
with the project TCGA-LUAD were downloaded from the TCGA.

Computer Code and Software: The code developed for the analyses described in this work is publicly available at GitLab (https://gitlab.com/ubb-cipf/metafunr). All software and versions used are detailed in **Supplementary Table S5**.

Supplementary Materials: The following materials are available online at www.mdpi.com/xxx/s1, **Figure S1**: Distribution of sex information among the reviewed studies, **Table S1**: Distribution of clinicopathological characteristics of each study population, **Table S2**: Summary of differential expression analysis results, **Table S3**: Significant genes differentially expressed between male and female lung adenocarcinoma patients, **Table S4**: All significant GO terms and KEGG pathways in the functional meta-analysis, **Table S5**: Software and versions used in this work.

Author Contributions: Conceptualization, F.G.G.; methodology, M.R.H., and F.G.G.; software, P.M.M., I.P.D., M.R.H., and F.G.G.; validation, P.M.M., A.R., M.B., R.F., M.R.H., and F.G.G.; formal analysis, I.P.D., M.R.H., and F.G.G; investigation, I.P.D., M.R.H., and F.G.G.; data curation, I.P.D.; writing—original draft preparation, I.P.D., M.R.H., and F.G.G.; writing—review and editing, I.P.D., M.R.H., A.R., M.B., Z.A., R.F., and F.G.G.; visualization, I.P.D., M.R.H., and F.G.G.; supervision, M.R.H., and F.G.G.; funding acquisition, F.G.G.; project administration, F.G.G. All authors have read and agreed to the published version of the manuscript.

Funding: This work was supported by Fondo de Investigación Sanitaria (ISCIII PI15-00209), GV/2020/186 and ISCIII PT17/0009/0015 FEDER.

Acknowledgments: The authors thank the Principe Felipe Research Center (CIPF) for providing access to the cluster, which is co-funded by European Regional Development Funds (FEDER) in Valencian Community 2014-2020. The results published here are in whole or part based upon data generated by the TCGA Research Network: https://www.cancer.gov/tcga.

Conflicts of Interest: The authors declare no conflicts of interest.

References

1. Bray, F.; Ferlay, J.; Soerjomataram, I.; Siegel, R.L.; Torre, L.A.; Jemal, A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA. Cancer J. Clin.* **2018**, *68*, 394–424, doi:10.3322/caac.21492.

2. Jemal, A.; Miller, K.D.; Ma, J.; Siegel, R.L.; Fedewa, S.A.; Islami, F.; Devesa, S.S.; Thun, M.J. Higher Lung Cancer Incidence in Young Women Than Young Men in the United States. *N. Engl. J. Med.* **2018**, *378*, 1999–2009, doi:10.1056/NEJMoa1715907.

3. Barta, J.A.; Powell, C.A.; Wisnivesky, J.P. Global Epidemiology of Lung Cancer. *Ann. Glob. Health* **2019**, *85*, 8, doi:10.5334/aogh.2419.

4. Siegel, R.L.; Miller, K.D.; Jemal, A. Cancer statistics, 2020. *CA. Cancer J. Clin.* **2020**, *70*, 7–30, doi:10.3322/caac.21590.

5. Conforti, F.; Pala, L.; Bagnardi, V.; De Pas, T.; Martinetti, M.; Viale, G.; Gelber, R.D.; Goldhirsch, A. Cancer immunotherapy efficacy and patients' sex: a systematic review and meta-analysis. *Lancet Oncol.* **2018**, *19*, 737–746, doi:10.1016/S1470-2045(18)30261-4.

6. Barquín, M.; Calvo, V.; García-García, F.; Nuñez, B.; Sánchez-Herrero, E.; Serna-Blasco, R.; Auglytė, M.; Carcereny, E.; Rodriguez-Abreu, D.; López Castro, R.; et al. Sex is a strong prognostic factor in stage IV non-small-cell lung cancer patients and should be considered in survival rate estimation. *Cancer Epidemiol.* **2020**, *67*, 101737, doi:10.1016/j.canep.2020.101737.

7. Wheatley-Price, P.; Blackhall, F.; Lee, S.-M.; Ma, C.; Ashcroft, L.; Jitlal, M.; Qian, W.; Hackshaw, A.; Rudd, R.; Booton, R.; et al. The influence of sex and histology on outcomes in non-small-cell lung cancer: a pooled analysis of five randomized trials. *Ann. Oncol.* **2010**, *21*, 2023–2028, doi:10.1093/annonc/mdq067.

8. Fasco, M.J.; Hurteau, G.J.; Spivack, S.D. Gender-dependent expression of alpha and beta estrogen receptors in human nontumor and tumor lung tissue. *Mol. Cell. Endocrinol.* **2002**, *188*, 125–140, doi:10.1016/S0303-7207(01)00750-X.

9.   Stabile, L.P.; Davis, A.L.G.; Gubish, C.T.; Hopkins, T.M.; Luketich, J.D.; Christie, N.;
Finkelstein, S.; Siegfried, J.M. Human Non-Small Cell Lung Tumors and Cells Derived from
Normal Lung Express Both Estrogen Receptor α and β and Show Biological Responses to
Estrogen. *Cancer Res.* **2002**, *62*, 2141–2150.
10.   Kawai, H.; Ishii, A.; Washiya, K.; Konno, T.; Kon, H.; Yamaya, C.; Ono, I.; Minamiya, Y.;
Ogawa, J. Estrogen Receptor α and β are Prognostic Factors in Non–Small Cell Lung Cancer.
Clin. Cancer Res. **2005**, *11*, 5084–5089, doi:10.1158/1078-0432.CCR-05-0200.
11.   Kligerman, S.; White, C. Epidemiology of Lung Cancer in Women: Risk Factors, Survival, and
Screening. *Am. J. Roentgenol.* **2011**, *196*, 287–295, doi:10.2214/AJR.10.5412.
12.   Bain, C.; Feskanich, D.; Speizer, F.E.; Thun, M.; Hertzmark, E.; Rosner, B.A.; Colditz, G.A.
Lung Cancer Rates in Men and Women With Comparable Histories of Smoking. *JNCI J. Natl.*
Cancer Inst. **2004**, *96*, 826–834, doi:10.1093/jnci/djh143.
13.   North, C.M.; Christiani, D.C. Women and Lung Cancer: What is New? *Semin. Thorac.*
Cardiovasc. Surg. **2013**, *25*, 87–94, doi:10.1053/j.semtcvs.2013.05.002.
14.   Travis, W.D.; Brambilla, E.; Noguchi, M.; Nicholson, A.G.; Geisinger, K.R.; Yatabe, Y.; Beer,
D.G.; Powell, C.A.; Riely, G.J.; Van Schil, P.E.; et al. International Association for the Study of
Lung Cancer/American Thoracic Society/European Respiratory Society International
Multidisciplinary Classification of Lung Adenocarcinoma. *J. Thorac. Oncol.* **2011**, *6*, 244–285,
doi:10.1097/JTO.0b013e318206a221.
15.   Araujo, J.M.; Prado, A.; Cardenas, N.K.; Zaharia, M.; Dyer, R.; Doimi, F.; Bravo, L.; Pinillos, L.;
Morante, Z.; Aguilar, A.; et al. Repeated observation of immune gene sets enrichment in
women with non-small cell lung cancer. *Oncotarget* **2016**, *7*, 20282–20292,
doi:10.18632/oncotarget.7943.
16.   Shi, K.; Li, N.; Yang, M.; Li, W. Identification of Key Genes and Pathways in Female Lung
Cancer Patients Who Never Smoked by a Bioinformatics Analysis. *J. Cancer* **2019**, *10*, 51–60,
doi:10.7150/jca.26908.
17.   Yuan, Y.; Liu, L.; Chen, H.; Wang, Y.; Xu, Y.; Mao, H.; Li, J.; Mills, G.B.; Shu, Y.; Li, L.; et al.
Comprehensive Characterization of Molecular Differences in Cancer between Male and
Female Patients. *Cancer Cell* **2016**, *29*, 711–722, doi:10.1016/j.ccell.2016.04.001.
18.   Li, Y.; He, C.-L.; Li, W.-X.; Zhang, R.-X.; Duan, Y. Transcriptome analysis reveals gender-
specific differences in overall metabolic response of male and female patients in lung
adenocarcinoma. *PLOS ONE* **2020**, *15*, e0230796, doi:10.1371/journal.pone.0230796.
19.   Normand, S.-L.T. Meta-analysis: formulating, evaluating, combining, and reporting. *Stat.*
Med. **1999**, *18*, 321–359, doi:10.1002/(SICI)1097-0258(19990215)18:3<321::AID-SIM28>3.0.CO;2-
P.
20.   Higgins, J.; Thomas, J. *Cochrane Handbook for Systematic Reviews of Interventions,* 2nd ed;
Chichester:*Wiley-Blackwell, 2019.*
21.   Edgar, R.; Domrachev, M.; Lash, A.E. Gene Expression Omnibus: NCBI gene expression and
hybridization array data repository. *Nucleic Acids Res.* **2002**, *30*, 207–210,
doi:10.1093/nar/30.1.207.
22.   Weinstein, J.N.; Collisson, E.A.; Mills, G.B.; Shaw, K.R.M.; Ozenberger, B.A.; Ellrott, K.;
Shmulevich, I.; Sander, C.; Stuart, J.M. The Cancer Genome Atlas Pan-Cancer analysis project.
Nat. Genet. **2013**, *45*, 1113–1120, doi:10.1038/ng.2764.
23.   Landi, M.T.; Dracheva, T.; Rotunno, M.; Figueroa, J.D.; Liu, H.; Dasgupta, A.; Mann, F.E.;
Fukuoka, J.; Hames, M.; Bergen, A.W.; et al. Gene Expression Signature of Cigarette Smoking
and Its Role in Lung Adenocarcinoma Development and Survival. *PLOS ONE* **2008**, *3*, e1651,
doi:10.1371/journal.pone.0001651.
24.   Hou, J.; Aerts, J.; den Hamer, B.; van IJcken, W.; den Bakker, M.; Riegman, P.; van der Leest,
C.; van der Spek, P.; Foekens, J.A.; Hoogsteden, H.C.; et al. Gene Expression-Based
Classification of Non-Small Cell Lung Carcinomas and Survival Prediction. *PLOS ONE* **2010**,
5, e10312, doi:10.1371/journal.pone.0010312.

25.   Okayama, H.; Kohno, T.; Ishii, Y.; Shimada, Y.; Shiraishi, K.; Iwakawa, R.; Furuta, K.; Tsuta,
K.; Shibata, T.; Yamamoto, S.; et al. Identification of Genes Upregulated in ALK-Positive and
EGFR/KRAS/ALK-Negative Lung Adenocarcinomas. *Cancer Res.* **2012**, *72*, 100,
doi:10.1158/0008-5472.CAN-11-1403.
26.   Yamauchi, M.; Yamaguchi, R.; Nakata, A.; Kohno, T.; Nagasaki, M.; Shimamura, T.; Imoto, S.;
Saito, A.; Ueno, K.; Hatanaka, Y.; et al. Epidermal Growth Factor Receptor Tyrosine Kinase
Defines Critical Prognostic Genes of Stage I Lung Adenocarcinoma. *PLOS ONE* **2012**, *7*,
e43923, doi:10.1371/journal.pone.0043923.
27.   Selamat, S.A.; Chung, B.S.; Girard, L.; Zhang, W.; Zhang, Y.; Campan, M.; Siegmund, K.D.;
Koss, M.N.; Hagen, J.A.; Lam, W.L.; et al. Genome-scale analysis of DNA methylation in lung
adenocarcinoma and integration with mRNA expression. *Genome Res.* **2012**, *22*, 1197–1211,
doi:10.1101/gr.132662.111.
28.   Robles, A.I.; Arai, E.; Mathé, E.A.; Okayama, H.; Schetter, A.J.; Brown, D.; Petersen, D.;
Bowman, E.D.; Noro, R.; Welsh, J.A.; et al. An Integrated Prognostic Classifier for Stage I Lung
Adenocarcinoma Based on mRNA, microRNA, and DNA Methylation Biomarkers. *J. Thorac.
Oncol.* **2015**, *10*, 1037–1048, doi:10.1097/JTO.0000000000000560.
29.   Girard, L.; Rodriguez-Canales, J.; Behrens, C.; Thompson, D.M.; Botros, I.W.; Tang, H.; Xie, Y.;
Rekhtman, N.; Travis, W.D.; Wistuba, I.I.; et al. An Expression Signature as an Aid to the
Histologic Classification of Non–Small Cell Lung Cancer. *Clin. Cancer Res.* **2016**, *22*, 4880,
doi:10.1158/1078-0432.CCR-15-2900.
30.   Mezheyeuski, A.; Bergsland, C.H.; Backman, M.; Djureinovic, D.; Sjöblom, T.; Bruun, J.; Micke,
P. Multispectral imaging for quantitative and compartment-specific immune infiltrates reveals
distinct immune profiles that classify lung cancer patients. *J. Pathol.* **2018**, *244*, 421–431,
doi:10.1002/path.5026.
31.   Sun, Z.; Wang, L.; Eckloff, B.W.; Deng, B.; Wang, Y.; Wampfler, J.A.; Jang, J.; Wieben, E.D.; Jen,
J.; You, M.; et al. Conserved recurrent gene mutations correlate with pathway deregulation
and clinical outcomes of lung adenocarcinoma in never-smokers. *BMC Med. Genomics* **2014**, *7*,
486, doi:10.1186/1755-8794-7-32.
32.   Moher, D.; Liberati, A.; Tetzlaff, J.; Altman, D.G.; Group, T.P. Preferred Reporting Items for
Systematic Reviews and Meta-Analyses: The PRISMA Statement. *PLOS Med.* **2009**, *6*,
e1000097, doi:10.1371/journal.pmed.1000097.
33.   Klein, S.L.; Flanagan, K.L. Sex differences in immune responses. *Nat. Rev. Immunol.* **2016**, *16*,
626–638, doi:10.1038/nri.2016.90.
34.   Cook, M.B.; Dawsey, S.M.; Freedman, N.D.; Inskip, P.D.; Wichner, S.M.; Quraishi, S.M.;
Devesa, S.S.; McGlynn, K.A. Sex Disparities in Cancer Incidence by Period and Age. *Cancer
Epidemiol. Prev. Biomark.* **2009**, *18*, 1174–1182, doi:10.1158/1055-9965.EPI-08-1118.
35.   Lukey, M.J.; Katt, W.P.; Cerione, R.A. Targeting amino acid metabolism for cancer therapy.
Drug Discov. Today **2017**, *22*, 796–804, doi:10.1016/j.drudis.2016.12.003.
36.   Poff, A.; Koutnik, A.P.; Egan, K.M.; Sahebjam, S.; D'Agostino, D.; Kumar, N.B. Targeting the
Warburg effect for cancer treatment: Ketogenic diets for management of glioma. *Semin. Cancer
Biol.* **2019**, *56*, 135–148, doi:10.1016/j.semcancer.2017.12.011.
37.   Butler, L.M.; Perone, Y.; Dehairs, J.; Lupien, L.E.; de Laat, V.; Talebi, A.; Loda, M.; Kinlaw,
W.B.; Swinnen, J.V. Lipids and cancer: Emerging roles in pathogenesis, diagnosis and
therapeutic intervention. *Adv. Drug Deliv. Rev.* **2020**, doi:10.1016/j.addr.2020.07.013.
38.   Gaignard, P.; Savouroux, S.; Liere, P.; Pianos, A.; Thérond, P.; Schumacher, M.; Slama, A.;
Guennoun, R. Effect of Sex Differences on Brain Mitochondrial Function and Its Suppression
by Ovariectomy and in Aged Mice. *Endocrinology* **2015**, *156*, 2893–2904, doi:10.1210/en.2014-
1913.
39.   Mervic, L. Time Course and Pattern of Metastasis of Cutaneous Melanoma Differ between
Men and Women. *PLOS ONE* **2012**, *7*, e32955, doi:10.1371/journal.pone.0032955.
40.   Pal, S.K.; Hurria, A. Impact of Age, Sex, and Comorbidity on Cancer Therapy and Disease

Progression. *J. Clin. Oncol.* **2010**, *28*, 4086–4093, doi:10.1200/JCO.2009.27.0579.

41. Sun, S.; Schiller, J.H.; Gazdar, A.F. Lung cancer in never smokers — a different disease. *Nat.*
Rev. Cancer **2007**, *7*, 778–790, doi:10.1038/nrc2190.

42. Kim, A.M.; Tingen, C.M.; Woodruff, T.K. Sex bias in trials and treatment must end. *Nature*
2010, *465*, 688–689, doi:10.1038/465688a.

43. Woitowich, N.C.; Beery, A.; Woodruff, T. A 10-year follow-up study of sex inclusion in the
biological sciences. *eLife* **2020**, *9*, e56344, doi:10.7554/eLife.56344.

44. Wilkinson, M.D.; Dumontier, M.; Aalbersberg, Ij.J.; Appleton, G.; Axton, M.; Baak, A.;
Blomberg, N.; Boiten, J.-W.; da Silva Santos, L.B.; Bourne, P.E.; et al. The FAIR Guiding
Principles for scientific data management and stewardship. *Sci. Data* **2016**, *3*, 160018,
doi:10.1038/sdata.2016.18.

45. Vinay, D.S.; Ryan, E.P.; Pawelec, G.; Talib, W.H.; Stagg, J.; Elkord, E.; Lichtor, T.; Decker,
W.K.; Whelan, R.L.; Kumara, H.M.C.S.; et al. Immune evasion in cancer: Mechanistic basis
and therapeutic strategies. *Semin. Cancer Biol.* **2015**, *35 Suppl*, S185–S198,
doi:10.1016/j.semcancer.2015.03.004.

46. Altorki, N.K.; Markowitz, G.J.; Gao, D.; Port, J.L.; Saxena, A.; Stiles, B.; McGraw, T.; Mittal, V.
The lung microenvironment: an important regulator of tumour growth and metastasis. *Nat.*
Rev. Cancer **2019**, *19*, 9–31, doi:10.1038/s41568-018-0081-9.

47. Ye, Y.; Jing, Y.; Li, L.; Mills, G.B.; Diao, L.; Liu, H.; Han, L. Sex-associated molecular
differences for cancer immunotherapy. *Nat. Commun.* **2020**, *11*, 1779, doi:10.1038/s41467-020-
15679-x.

48. Zeng, D.-Q.; Yu, Y.-F.; Ou, Q.-Y.; Li, X.-Y.; Zhong, R.-Z.; Xie, C.-M.; Hu, Q.-G. Prognostic and
predictive value of tumor-infiltrating lymphocytes for clinical therapeutic research in patients
with non-small cell lung cancer. *Oncotarget* **2016**, *7*.

49. Ye, S.-L.; Li, X.-Y.; Zhao, K.; Feng, T. High expression of CD8 predicts favorable prognosis in
patients with lung adenocarcinoma: A cohort study. *Medicine (Baltimore)* **2017**, *96*.

50. Li, S.; Wang, Z.; Li, X.-J. Notch signaling pathway suppresses CD8+ T cells activity in patients
with lung adenocarcinoma. *Int. Immunopharmacol.* **2018**, *63*, 129–136,
doi:10.1016/j.intimp.2018.07.033.

51. Abbas, A.K.; Trotta, E.; R. Simeonov, D.; Marson, A.; Bluestone, J.A. Revisiting IL-2: Biology
and therapeutic prospects. *Sci. Immunol.* **2018**, *3*, eaat1482, doi:10.1126/sciimmunol.aat1482.

52. Gao, Y.; Lu, J.; Zeng, C.; Yang, J.; Huang, B.; Zhang, N.; Li, L.; Fu, X. IL-10 suppresses IFN-γ-
mediated signaling in lung adenocarcinoma. *Clin. Exp. Med.* **2020**, *20*, 449–459,
doi:10.1007/s10238-020-00626-3.

53. Vahl, J.M.; Friedrich, J.; Mittler, S.; Trump, S.; Heim, L.; Kachler, K.; Balabko, L.; Fuhrich, N.;
Geppert, C.-I.; Trufa, D.I.; et al. Interleukin-10-regulated tumour tolerance in non-small cell
lung cancer. *Br. J. Cancer* **2017**, *117*, 1644–1655, doi:10.1038/bjc.2017.336.

54. Miotto, D.; Cascio, N.L.; Stendardo, M.; Querzoli, P.; Pedriali, M.; De Rosa, E.; Fabbri, L.M.;
Mapp, C.E.; Boschetto, P. CD8+ T cells expressing IL-10 are associated with a favourable
prognosis in lung cancer. *Lung Cancer* **2010**, *69*, 355–360, doi:10.1016/j.lungcan.2009.12.012.

55. Emmerich, J.; Mumm, J.B.; Chan, I.H.; LaFace, D.; Truong, H.; McClanahan, T.; Gorman, D.M.;
Oft, M. IL-10 Directly Activates and Expands Tumor-Resident CD8+ T Cells without De Novo
Infiltration from Secondary Lymphoid Organs. *Cancer Res.* **2012**, *72*, 3570, doi:10.1158/0008-
5472.CAN-12-0721.

56. Shang, G.-S.; Liu, L.; Qin, Y.-W. IL-6 and TNF-α promote metastasis of lung cancer by
inducing epithelial-mesenchymal transition. *Oncol. Lett.* **2017**, *13*, 4657–4660,
doi:10.3892/ol.2017.6048.

57. Silva, E.M.; Mariano, V.S.; Pastrez, P.R.A.; Pinto, M.C.; Castro, A.G.; Syrjanen, K.J.; Longatto-
Filho, A. High systemic IL-6 is associated with worse prognosis in patients with non-small cell
lung cancer. *PLOS ONE* **2017**, *12*, e0181125, doi:10.1371/journal.pone.0181125.

58. Lewis, A.M.; Varghese, S.; Xu, H.; Alexander, H.R. Interleukin-1 and cancer progression: the

emerging role of interleukin-1 receptor antagonist as a novel therapeutic agent in cancer
treatment. *J. Transl. Med.* **2006**, *4*, 48, doi:10.1186/1479-5876-4-48.

59. Gottschlich, A.; Endres, S.; Kobold, S. Can we use interleukin-1β blockade for lung cancer
treatment? *Transl. Lung Cancer Res.* **2018**, *7*, S160–S164.

60. Wu, F.; Xu, J.; Huang, Q.; Han, J.; Duan, L.; Fan, J.; Lv, Z.; Guo, M.; Hu, G.; Chen, L.; et al. The
Role of Interleukin-17 in Lung Cancer. *Mediators Inflamm.* **2016**, *2016*, 8494079,
doi:10.1155/2016/8494079.

61. Wang, C.; Qiao, W.; Jiang, Y.; Zhu, M.; Shao, J.; Ren, P.; Liu, D.; Li, W. Effect of sex on the
efficacy of patients receiving immune checkpoint inhibitors in advanced non-small cell lung
cancer. *Cancer Med.* **2019**, *8*, 4023–4031, doi:10.1002/cam4.2280.

62. Conforti, F.; Pala, L.; Bagnardi, V.; Viale, G.; De Pas, T.; Pagan, E.; Pennacchioli, E.;
Cocorocchio, E.; Ferrucci, P.F.; De Marinis, F.; et al. Sex-Based Heterogeneity in Response to
Lung Cancer Immunotherapy: A Systematic Review and Meta-Analysis. *JNCI J. Natl. Cancer*
Inst. **2019**, *111*, 772–781, doi:10.1093/jnci/djz094.

63. Vijayan, D.; Smyth, M.J.; Teng, M.W.L. Purinergic Receptors: Novel Targets for Cancer
Immunotherapy. In *Oncoimmunology: A Practical Guide for Cancer Immunotherapy*; Zitvogel, L.,
Kroemer, G., Eds.; Springer International Publishing: Cham, 2018; pp. 115–141 ISBN 978-3-
319-62431-0.

64. Crain, J.M.; Nikodemova, M.; Watters, J.J. Expression of P2 nucleotide receptors varies with
age and sex in murine brain microglia. *J. Neuroinflammation* **2009**, *6*, 24, doi:10.1186/1742-2094-
6-24.

65. Savio, L.E.B.; de Andrade Mello, P.; da Silva, C.G.; Coutinho-Silva, R. The P2X7 Receptor in
Inflammatory Diseases: Angel or Demon? *Front. Pharmacol.* **2018**, *9*, 52,
doi:10.3389/fphar.2018.00052.

66. Hu, L.-P.; Zhang, X.-X.; Jiang, S.-H.; Tao, L.-Y.; Li, Q.; Zhu, L.-L.; Yang, M.-W.; Huo, Y.-M.;
Jiang, Y.-S.; Tian, G.-A.; et al. Targeting Purinergic Receptor P2Y2 Prevents the Growth of
Pancreatic Ductal Adenocarcinoma by Inhibiting Cancer Cell Glycolysis. *Clin. Cancer Res. Off.*
J. Am. Assoc. Cancer Res. **2019**, *25*, 1318–1330, doi:10.1158/1078-0432.CCR-18-2297.

67. Tak, E.; Jun, D.Y.; Kim, S.-H.; Park, G.-C.; Lee, J.; Hwang, S.; Song, G.-W.; Lee, S.-G.
Upregulation of P2Y2 nucleotide receptor in human hepatocellular carcinoma cells. *J. Int.*
Med. Res. **2016**, *44*, 1234–1247, doi:10.1177/0300060516662135.

68. Asif, A.; Khalid, M.; Manzoor, S.; Ahmad, H.; Rehman, A.U. Role of purinergic receptors in
hepatobiliary carcinoma in Pakistani population: an approach towards proinflammatory role
of P2X4 and P2X7 receptors. *Purinergic Signal.* **2019**, *15*, 367–374, doi:10.1007/s11302-019-
09675-0.

69. Xu, M.; Almasi, S.; Yang, Y.; Yan, C.; Sterea, A.M.; Rizvi Syeda, A.K.; Shen, B.; Richard Derek,
C.; Huang, P.; Gujar, S.; et al. The lysosomal TRPML1 channel regulates triple negative breast
cancer development by promoting mTORC1 and purinergic signaling pathways. *Cell Calcium*
2019, *79*, 80–88, doi:10.1016/j.ceca.2019.02.010.

70. Zhu, Y.; Shao, X.; Wang, X.; Liu, L.; Liang, H. Sex disparities in cancer. *Cancer Lett.* **2019**, *466*,
35–38, doi:10.1016/j.canlet.2019.08.017.

71. Shay, J.W.; Wright, W.E. Senescence and immortalization: role of telomeres and telomerase.
Carcinogenesis **2005**, *26*, 867–874, doi:10.1093/carcin/bgh296.

72. Datta, A.; Brosh, R.M. New Insights Into DNA Helicases as Druggable Targets for Cancer
Therapy. *Front. Mol. Biosci.* **2018**, *5*, 59, doi:10.3389/fmolb.2018.00059.

73. Luo, J.; Dai, X.; Hu, H.; Chen, J.; Zhao, L.; Yang, C.; Sun, J.; Zhang, L.; Wang, Q.; Xu, S.; et al.
Fluzoparib increases radiation sensitivity of non-small cell lung cancer (NSCLC) cells without
BRCA1/2 mutation, a novel PARP1 inhibitor undergoing clinical trials. *J. Cancer Res. Clin.*
Oncol. **2020**, *146*, 721–737, doi:10.1007/s00432-019-03097-6.

74. Abbotts, R.; Topper, M.J.; Biondi, C.; Fontaine, D.; Goswami, R.; Stojanovic, L.; Choi, E.Y.;
McLaughlin, L.; Kogan, A.A.; Xia, L.; et al. DNA methyltransferase inhibitors induce a

BRCAness phenotype that sensitizes NSCLC to PARP inhibitor and ionizing radiation. *Proc.*
Natl. Acad. Sci. **2019**, *116*, 22609, doi:10.1073/pnas.1903765116.

75.   R Core Team *R: A Language and Environment for Statistical Computing*; R Foundation for
Statistical Computing: Vienna, Austria, 2014;

76.   Athar, A.; Füllgrabe, A.; George, N.; Iqbal, H.; Huerta, L.; Ali, A.; Snow, C.; Fonseca, N.A.;
Petryszak, R.; Papatheodorou, I.; et al. ArrayExpress update - from bulk to single-cell
expression data. *Nucleic Acids Res.* **2019**, *47*, D711–D715, doi:10.1093/nar/gky964.

77.   Robinson, M.D.; McCarthy, D.J.; Smyth, G.K. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. *Bioinformatics* **2010**, *26*, 139–140,
doi:10.1093/bioinformatics/btp616.

78.   Robinson, M.D.; Oshlack, A. A scaling normalization method for differential expression
analysis of RNA-seq data. *Genome Biol.* **2010**, *11*, R25, doi:10.1186/gb-2010-11-3-r25.

79.   Maglott, D.; Ostell, J.; Pruitt, K.D.; Tatusova, T. Entrez Gene: gene-centered information at
NCBI. *Nucleic Acids Res.* **2005**, *33*, D54–D58, doi:10.1093/nar/gki031.

80.   Durinck, S.; Moreau, Y.; Kasprzyk, A.; Davis, S.; De Moor, B.; Brazma, A.; Huber, W. BioMart
and Bioconductor: a powerful link between biological databases and microarray data analysis.
Bioinformatics **2005**, *21*, 3439–3440, doi:10.1093/bioinformatics/bti525.

81.   Ritchie, M.E.; Phipson, B.; Wu, D.; Hu, Y.; Law, C.W.; Shi, W.; Smyth, G.K. limma powers
differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids*
Res. **2015**, *43*, e47–e47, doi:10.1093/nar/gkv007.

82.   Benjamini, Y.; Hochberg, Y. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B Methodol.* **1995**, *57*, 289–300.

83.   Montaner, D.; Dopazo, J. Multidimensional Gene Set Analysis of Genomic Data. *PLOS ONE*
2010, *5*, e10348, doi:10.1371/journal.pone.0010348.

84.   Kanehisa, M.; Goto, S. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res.*
2000, *28*, 27–30, doi:10.1093/nar/28.1.27.

85.   Ashburner, M.; Ball, C.A.; Blake, J.A.; Botstein, D.; Butler, H.; Cherry, J.M.; Davis, A.P.;
Dolinski, K.; Dwight, S.S.; Eppig, J.T.; et al. Gene Ontology: tool for the unification of biology.
Nat. Genet. **2000**, *25*, 25–29, doi:10.1038/75556.

86.   Lex, A.; Gehlenborg, N.; Strobelt, H.; Vuillemot, R.; Pfister, H. UpSet: Visualization of
Intersecting Sets. *IEEE Trans. Vis. Comput. Graph.* **2014**, *20*, 1983–1992,
doi:10.1109/TVCG.2014.2346248.

87.   García-García, F. Methods of functional enrichment analysis in genomic studies, Universitat
de València: València, 2016.

88.   Viechtbauer, W. Conducting Meta-Analyses in R with the metafor Package. *J. Stat. Softw. Vol 1*
Issue 3 2010 **2010**.

89.   DerSimonian, R.; Laird, N. Meta-analysis in clinical trials. *Control. Clin. Trials* **1986**, *7*, 177–188,
doi:10.1016/0197-2456(86)90046-2.

90.   Sterne, J.A.C.; Egger, M. Funnel plots for detecting bias in meta-analysis: Guidelines on choice
of axis. *J. Clin. Epidemiol.* **2001**, *54*, 1046–1055, doi:10.1016/S0895-4356(01)00377-8.

91.   Wickham, H. *ggplot2: Elegant Graphics for Data Analysis*; Springer-Verlag New York, 2016;
ISBN 978-3-319-24277-4.