Inferring person-to-person networks of pathogen transmission:

is routine surveillance data up to the task?

John H. Huber[1*], Michelle S. Hsiang[2,3,4], Nomcebo Dlamini[5], Maxwell Murphy[6], Sibonakaliso

Vilakati[5], Nomcebo Nhlabathi[5], Anita Lerch[1], Rasmus Nielsen[7], Nyasatu Ntshalintshali[8],

Bryan Greenhouse[6,9], T. Alex Perkins[1*]

[1]Department of Biological Sciences, University of Notre Dame, Notre Dame, Indiana, United

States of America

[2]Department of Pediatrics, University of Texas Southwestern Medical Center, Dallas, Texas,

United States of America

[3]Malaria Elimination Initiative, Global Health Group, University of California, San Francisco,

California, United States of America

[4]Department of Pediatrics, University of California, San Francisco, California, United States of

America

[5]National Malaria Elimination Programme, Ministry of Health, Manzini, Eswatini

[6]Department of Medicine, University of California, San Francisco, California, United States of

America

[7]Department of Integrative Biology and Statistics, University of California, Berkeley, California,

United States of America

[8]Clinton Health Access Initiative, Eswatini Country Office, Mbabane, Eswatini

[9]Chan Zuckerberg Biohub, San Francisco, California, United States of America

*Corresponding Authors

Email: jhuber3@nd.edu (JHH); taperkins@nd.edu (TAP)

Short Title: Inferring person-to-person transmission networks

Abstract

Inference of person-to-person transmission networks using routinely collected surveillance data is being used increasingly to estimate spatiotemporal patterns of pathogen transmission. Several data types can be used to inform transmission network inferences, yet the sensitivity of those inferences to different data types is not routinely evaluated. We evaluated the influence of different combinations of spatial, temporal, and travel-history data on transmission network inferences for *Plasmodium falciparum*, the pathogen responsible for most human malaria. After developing a new inference framework and applying it to simulated data, we found that these data types have limited utility for inferring transmission networks and, in some combinations, tend to overestimate transmission. Only when outbreaks were highly focal in time or when travel histories were highly accurate was the inference algorithm able to accurately estimate the reproduction number under control, R_c, a key metric of transmission. Applying this approach to surveillance data from Eswatini indicated that inferences of R_c and spatiotemporal patterns therein are sensitive to the choice of data types and assumptions about the accuracy of travel-history data. Taken together, these results suggest that transmission network inferences made with routinely collected surveillance data should be interpreted with caution. As we have done here, future studies inferring transmission networks should apply their algorithm to data simulated under alternative assumptions to assess the robustness of their inferences.

Author Summary

Person-to-person networks of pathogen transmission can reveal epidemiological patterns that are not obvious based on routinely collected surveillance data alone. Several types of data can be used to inform estimates of these networks, including the timing of when cases present, the locations where those individuals reside, and whether those individuals recently traveled to an area with higher transmission. Little is known about whether transmission network inferences may depend on the types of data used to inform those inferences. To address this question, we developed an algorithm to estimate transmission networks of *Plasmodium falciparum* malaria. We validated this algorithm on simulated data and then applied it to surveillance data from Eswatini. Our results demonstrate that transmission network inferences are sensitive to which data types are used to inform them and to assumptions about how to interpret those data, such as whether data about recent travel are accurate. These results suggest that future work should carefully consider the epidemiological setting, which data types are used to inform transmission network inferences, and how those inferences are interpreted.

Introduction

Concomitant with improved epidemiological surveillance, there is growing interest to leverage the collected data to infer transmission networks for a wide range of pathogens and to use those inferences to inform public health efforts. Past studies have incorporated temporal data [1] and spatial data [2–5] to estimate pairwise probabilities of transmission between individual cases and to use those estimates to infer time-varying and spatially varying reproduction numbers, respectively. More recently, methods have been developed to incorporate this type of detailed, individual-level epidemiological data [6–8] to infer transmission networks for infectious diseases of humans, including severe acute respiratory syndrome [9] and tuberculosis [10], and of animals, such as rabies [11] and foot-and-mouth disease [12].

In addition to the diseases for which these methods have been applied to date, there is a growing need to apply similar methods to malaria in near-elimination settings. As incidence of malaria declines within a country, transmission becomes more heterogeneous in space and time [13]. Focal areas of high transmission, known as "hotspots," pose a serious risk of fueling resurgence if left untargeted, potentially reversing decades of progress towards elimination [14]. To this end, granular estimates of when and where transmission occurs are needed, as spatially aggregated estimates may obscure important heterogeneities of practical relevance to control efforts [15]. In addition to characterizing details of local transmission, measurement of progress towards malaria elimination hinges on correct classification of cases as imported or locally acquired [16,17], which is a byproduct of estimating transmission networks.

Previous work on malaria has made progress on the use of individual-level epidemiological data to infer transmission networks of *Plasmodium falciparum*, the parasite primarily responsible for human malaria in many regions of the world. Churcher *et al.* [18] used

temporal data to estimate the proportion of imported cases needed to confidently estimate the

reproduction number under control, R_c, below one and thereby provide evidence of controlled,

non-endemic malaria transmission. Reiner *et al.* [6] then built upon this work by incorporating

spatial data and inferring an individual-level transmission network of *P. falciparum* in Eswatini.

More recently, Routledge *et al.* [19,20] used related approaches to infer transmission networks

and R_c of *P. vivax* in El Salvador and China.

As the adoption of these methods increases, both for malaria and for other infectious

diseases, care should be taken to assess how the epidemiological setting and the inclusion or

exclusion of certain data types might affect transmission network inferences, as well as resultant

inferences about epidemiological quantities including R_c and spatiotemporal variation therein. To

date, transmission network inferences have relied on various types of epidemiological data,

ranging from only the use of timing of symptom onset [1] to more detailed spatiotemporal data

[2–6,20]. Additionally, studies that have incorporated travel-history information into

transmission network inferences have considered these data to be perfectly accurate, assuming

that all cases that reported travel were imported [6,18–20]. However, travel history may be an

imperfect indicator of importation owing to errors in recall [17] and the fact that travel to an area

of ongoing transmission alone does not guarantee that an individual was infected there [17,21].

Thus, how different choices about data types and different assumptions about possible errors in

travel-history data affect transmission network inferences remain largely unexplored [22].

Here, we present a Bayesian method for inferring transmission networks based on

temporal, spatial, and travel-history data for individual malaria cases. We use it to characterize

the sensitivity of transmission network inferences to the inclusion of different data types and to

different assumptions about the accuracy of travel histories. Our method builds upon previous

work by leveraging individual-level epidemiological data to obtain posterior estimates of

transmission networks and model parameters in a way that is flexible enough to accommodate

different assumptions about errors in travel histories. After establishing a proof-of-concept of our

inference method on simple test cases, we applied our method to real-world surveillance data

from Eswatini and additional simulated data sets to understand how the inclusion or exclusion of

different data types and different assumptions about travel-history error affect our ability to infer

transmission networks and estimate transmission metrics, such as R_c.

Results

To establish proof-of-concept, we first applied our inference method on three simple test cases

and evaluated how well our inferences recovered the true transmission networks. We then

applied our method to surveillance data collected in Eswatini during 2013-2017 and assessed

how our epidemiological conclusions changed with the inclusion or exclusion of different data

types and different assumptions about travel histories. These inference settings used: (1) spatial

and temporal data while estimating the accuracy of the travel history (default setting); (2) spatial

and temporal data while believing the travel history; (3) spatial and temporal data alone; (4)

temporal data while estimating the accuracy of the travel history; and (5) temporal data while

believing the travel history. To validate the inferences based on data from Eswatini, we

simulated data generated using posterior parameter estimates obtained from the data from

Eswatini and evaluated the ability of our inference method to recover the true transmission

networks along with the underlying parameters on those simulated data. Finally, we performed a

simulation sweep across different epidemiological settings to determine the range of conditions

under which our inference method yielded reliable estimates of transmission. A full description

of the analyses and additional results can be found in the S1 File.

Validation using a simple test case

We first validated our approach on three small simulated networks of twenty nodes. Although

these networks varied in their proportion of imported cases, the local transmission chains were

arranged optimally both temporally and spatially with perfect travel histories and complete

observation of cases, providing idealized test cases to validate our inference algorithm. We

measured the performance of our inference algorithm in terms of its ability to reconstruct

different features of the transmission network and correctly estimate R_c. As the proportion of

imported cases decreased from 90% to 5%, we found that the ability of the algorithm to correctly

classify cases as imported or locally acquired improved (Fig 1). For example, when we used the

default inference setting, classification accuracy improved from 85.7% (95% Uncertainty

Interval: 85.7 – 85.7%) to 100% (100 – 100%). As classification accuracy improved, our

estimates of R_c also improved, with all five inference settings yielding accurate estimates when

imported cases comprised only 5% of total cases (Fig 1C). Similarly, the ability to identify the

correct parent of each locally acquired case and assign it to the correct outbreak depended on the

extent of local transmission in the network. When 90% of the cases were imported, performance

was variable across inference settings (Fig 1A). Believing the travel history, regardless of

whether spatial data was included, enabled us to perfectly reconstruct the transmission network,

because the simulated travel-history data were accurate. However, as the proportion of imported

cases decreased, the benefit of believing the travel history diminished, and using all data types

resulted in the greatest accuracy. For example, in the most extreme case in which only 5% of

cases were imported, the accuracy of identifying the true parent ranged from 84.2% (68.4 –

94.7%) under the default inference settings to 63.2% (47.4 – 78.9%) using temporal data and

estimating the accuracy of the travel history (Fig 1C). In terms of identifying the outbreak to

which a case belongs, the algorithm was accurate under all inference settings, since there was

only one outbreak (Fig 1C).

Fig 1. Inference accuracies for three simulated transmission networks with (A) 90%, (B)

50%, and (C) 5% of cases as imported. Case classification refers to proportion of cases that are

correctly classified as imported vs. locally acquired. Transmission linkage denotes the proportion

of locally acquired cases for which the true parent is correctly identified, Outbreak is the

proportion of locally acquired cases for which the inferred parent belongs to the correct outbreak,

and R_c is the estimated reproduction number under control. Square points signify the median

posterior value, and bars are the 95% uncertainty intervals. The gray line indicates the true value

of R_c.

Application to Eswatini surveillance data

We next applied our method to surveillance data collected in Eswatini during 2013-2017. Under

the default inference setting, we estimated the diffusion coefficient D, which quantifies the

spatial spread of transmission, to be 4.42 km^2 day^{-1} (2.92 – 6.18 km^2 day^{-1}) (Fig 2A). This

corresponded to a median inferred transmission distance of 13.0 km (0.0160 – 65.9 km), a

median inferred serial interval of 47 days (-33 – 150 days) (Fig 3A & 3B), and median estimates

of τ_s, the probability that an imported case reported travel, of 0.61 (0.44 – 0.78) and τ_l, the

probability that a locally acquired case reported travel, of 0.57 (0.53 – 0.61) (Fig 2B & 2C). That

the 95% credible interval for τ_s contained 0.50 indicated that our inference algorithm found

limited use of travel-history data in discriminating between imported and locally acquired cases,

because that implies that imported cases have equal probabilities of reporting or not reporting

travel. The algorithm estimated the proportion of imported cases to be 0.052, corresponding to R_c

= 0.95. Mapping risk of importation and local transmission across Eswatini under the default

inference setting, we estimated consistently low risk of importation throughout the country and

transmission hotspots in the northeastern part of Eswatini, close to the border with Mozambique

(Fig 4A & 4B).

Fig 2. Marginal posterior distributions of parameters from Eswatini surveillance data.

Histograms represent the marginal posterior distribution of each parameter, color-coded by the

inference settings used. D is the diffusion coefficient, τ_s is the probability that an imported case

reports travel, and τ_l is the probability that a locally acquired case reports travel. Gray shapes

represent the prior distributions placed on each parameter. Inference settings in which a given

parameter was not estimated are indicated by NA.

Fig 3. Spatial and temporal scales of transmission in Eswatini. Density plots of the spatial

(km) and temporal (days) scales of transmission are reported and color-coded for each inference

setting. Dashed lines indicate the corresponding null distribution, generated from all random

196   pairs of cases in the Eswatini surveillance data set. The grey shape is the serial interval

197   distribution used in the likelihood.

Fig 4. Spatial distribution of importation and transmission risk in Eswatini. Maps of the

proportion of cases that are imported and the reproduction number under control (R_c) were

generated for each inference setting using a generalized additive model with a Gaussian process

basis function setting using the mgcv package in R [23,24]. In each plot, darker colors indicate

greater importation or transmission risk.

Parameter estimates and transmission network inferences differed under other inference

settings. When we believed the travel history, we estimated a larger median transmission

distance (Fig 3D). We attribute this increase in the spatial scale of transmission to clusters of

cases with positive travel histories located near metropolitan areas. By forcing those cases to be

imported, the algorithm tended to infer transmission across longer distances to explain the

origins of the remainder of cases that did not report travel and were thereby inferred to be locally

acquired. With respect to time, all five inference settings produced consistent serial interval

estimates, though the inclusion of spatial data allowed for a wider range of transmission linkages

in time (Fig 3A, 3C, & 3E). Finally, in the absence of spatial data, the model estimated higher

predictive power of travel histories in identifying imported cases (τ_s: 0.83, [0.60, 0.95]), though

the travel history was consistently found to be uninformative for identifying locally acquired

cases (τ_l: 0.57, [0.53, 0.60]) (Fig 2K & 2L).

Classification of cases as imported or locally acquired, key information for control

programs, was sensitive to the choice of inference setting. The proportion of cases classified as

imported was most sensitive to different assumptions about the accuracy of the travel histories

(Fig 4, left column). Believing the travel history yielded high estimates of importation in western

Eswatini (Fig 4C & 4I), whereas estimating or ignoring the travel history yielded low, relatively

homogeneous estimates of importation risk (Fig 4A, 4E, & 4G). For instance, using temporal

data and estimating the accuracy of the travel history produced probabilities of importation that

ranged 0.0043 – 0.0050, suggesting that nearly all cases resulted from local transmission (Fig

4G). Estimates of the spatial distribution of R_c depended most on the choice of which data types

we included (Fig 4, right column). Notably, inclusion of spatial and temporal data produced a

consistent spatial distribution of relative transmission risk, with transmission hotspots in

northeastern Eswatini (Fig 4B, 4D, & 4F). However, believing the travel history reduced the

magnitude of transmission that we inferred from a median R_c of 0.95 (Fig 4B) under default

settings to 0.41 (Fig 4D). Omitting spatial data changed the spatial distribution of transmission.

Estimating the accuracy of the travel history yielded high transmission estimates (median R_c:

1.00) in eastern Eswatini (Fig 4H), whereas believing the travel history inferred hotspots of

transmission (median R_c: 0.42) in southern Eswatini (Fig 4J).

Validation of inferences from Eswatini

Reconciling the different inferences under different inference settings in Figs. 2-4 was

challenging because the true, underlying network and parameters were unknown. Using median

posterior estimates from the Eswatini data under each inference setting, we simulated data and

assessed the ability of the inference method to recover the underlying parameters and

transmission networks (Table 1). We found that the model was able to estimate τ_s and τ_l

reasonably well, depending on the inference setting (Fig 5). One exception was that the

algorithm slightly overestimated τ_s under the default inference setting. We attribute this to the

low proportion (0.052) of imported infections in the simulated data set and the strong prior

placed on this parameter. This tendency was not observed under the inference setting where

spatial data was excluded, because the true value of τ_s closely matched the mean of the prior

distribution in that case (see S2 File for further discussion).

Table 1. Characteristics of simulated data generated using the branching process model. A

description of the simulated data used in the inference exercises are reported for each of the five

inference settings. The total number of nodes in the network, the number of distinct outbreaks,

the proportion of cases that are imported, and the underlying parameters are provided.

Inference Setting			Network Size	# of Outbreaks	Prop. Imported	D	τ_s	τ_l
Space	Time	Travel						
Yes	Yes	Estimate	775	43	0.052	4.42	0.61	0.57
Yes	Yes	Believe	775	492	0.59	3.70	1	0
Yes	Yes	No	775	36	0.043	4.96	NA	NA
No	Yes	Estimate	775	1	0.0013	NA	0.83	0.57
No	Yes	Believe	775	489	0.58	NA	1	0

Fig 5. Marginal posterior distributions for parameters inferred from simulated data. The

marginal posterior distributions are reported for each inference setting from its respective

simulated data set. Each line denotes the true value of the parameter, and the grey shapes

represent the prior distributions of the parameters. Inference settings in which a given parameter

was not estimated are indicated by NA.

With the exception of believing the travel history, the model consistently overestimated

the diffusion coefficient (Fig 5A, 5D, & 5G). We attribute the challenge of correctly estimating

the diffusion coefficient to an inability to correctly estimate the underlying transmission network,

the extent of local transmission in the network, and a numerical insensitivity in the overall

likelihood to changes in D. When we conditioned the likelihood of D on the true transmission

network when R_c was high, the true values of D fell close to the range of maximum-likelihood

estimates, suggesting that this parameter could be estimated correctly if the true network was

identified (S1 Fig). The likelihood around the true value was very flat, however, making it easy

for D to be estimated incorrectly. When R_c was low, we underestimated the diffusion coefficient,

because the likelihood of imported cases increases as D decreases.

The overall accuracy of classifying cases as imported or locally acquired was close to one

(Fig 6). Though seemingly promising, these high accuracies masked a tendency to overclassify

cases as locally acquired, because many more cases were simulated to be locally acquired than

imported. For example, under the default inference setting, the accuracy of correctly classifying

imported cases was 0.023 (0.023 – 0.067). Similarly, the accuracies of identifying the parent of

each transmission linkage were poor, despite simulating under the assumptions of the model,

with accuracies ranging from 0.042 (0.022 – 0.065) when using temporal data and believing the

travel history to 0.31 (0.28 – 0.34) when incorporating spatial and temporal data and estimating

the accuracy of the travel history (Fig 6, circle points). This suggests that, as the number of cases

increases within a fixed space-time window, the information content of routinely collected

epidemiological data decreases and the method becomes incapable of correctly estimating the

transmission network. Nevertheless, under some settings, the method was able to capture higher-

order summaries of the network, such as case classification and R_c (Fig 6, square and diamond

points).

Fig 6. Inference accuracies for validation exercises. Accuracy metrics are reported for each

inference setting applied to its respective simulated data set. Case Classification, represented by

squares, refers to the proportion of cases that are correctly classified as imported or locally

acquired. Transmission Linkage, denoted by circles, is the proportion of locally acquired cases

for which the true parent is correctly identified. Outbreak, represented by triangles, is the

proportion of locally acquired cases for which the inferred parent belongs to the correct outbreak.

Bars denote the 95% credible intervals, and the grey line is the true R_c value of the network.

Simulation Sweep

Validation of our inference algorithm revealed that its performance varied across simulated data

sets. When applied to a series of simple test cases in which the transmission networks were small

and in an optimal spatiotemporal arrangement, the inference method was able to reconstruct the

transmission network and correctly estimate R_c (Fig 1). When applied to larger transmission

networks in which outbreaks overlapped in space and time, performance of the inference method

was poor (Fig 6). This indicated that the performance of our inference algorithm depends on the

epidemiological setting to which it is applied. To address this observation, we generated 2,000

simulated data sets in which we varied the proportion of imported cases, the spatiotemporal

window over which imported cases were distributed, the diffusion coefficient, and the accuracies

of the travel history (i.e., τ_s and τ_l) (S2 Table). We then applied our inference algorithm under

three different inference settings and quantified the accuracy of reconstructing each transmission

network. The three inference settings used: (1) spatial and temporal data while estimating the

accuracy of the travel history (default setting); (2) spatial and temporal data while believing the

travel history; and (3) spatial and temporal data alone (S1 Table).

We observed that the accuracy of reconstructing transmission networks depended upon

both the inference setting used and the epidemiological features of the simulated data. When we

used spatial and temporal data and estimated the accuracy of the travel history or excluded it, the

accuracy of reconstructing transmission networks depended on the relative proportion and

temporal distribution of imported cases (S8 and S9 Figs). As the temporal window over which

imported cases are distributed increased, the accuracy of identifying the true parent and the true

outbreak of each locally acquired case increased. With an increasing temporal window,

outbreaks within the transmission network became relatively more focal in time, which made the

likelihoods of alternative transmission linkages more readily distinguishable. More accurate

estimates of R_c under these inference settings similarly depended on the temporal window over

which imported cases were distributed (Fig 7A, 7C). When the mean temporal interval between

imported infections was greater than two times the mean length of the serial interval (i.e.,

approximately 100 days), our estimates of R_c improved, though we generally overestimated it.

Furthermore, as the proportion of imported cases increased and R_c decreased, the accuracy of

identifying the correct outbreak of each locally acquired case decreased (S8 and S9 Figs). This

pattern reflected the relationship between R_c and the size of individual outbreaks. As R_c

decreased, the size of individual outbreaks decreased, and, consequently, the probability that the

inferred parent of a locally acquired case belonged to the same outbreak decreased.

Fig 7. Comparison of R_c estimates across inference settings. The inference algorithm was

applied to 2,000 simulated data sets. The estimated R_c is compared to the true R_c for each of the

inference settings: (A) spatial and temporal data while estimating the accuracy of the travel

history; (B) spatial and temporal data while believing the travel history; and (C) spatial and

temporal data alone. Each point represents a simulated data set. The darker, accented points are

simulated data sets with epidemiological features that improved performance. In (A) and (C), the

darker, accented points were simulated data sets where the mean temporal interval between

imported infections was greater than two times the mean serial interval. In (B), the darker,

accented points were simulated data sets where the proportion of cases reporting travel was

within 0.05 of the proportion of imported cases.

By contrast, when we believed the travel history, the accuracy of reconstructing

transmission networks depended most strongly on the accuracies of the travel history. As the

probability of reporting travel increased, the accuracy of classifying imported cases increased,

and the accuracy of classifying locally acquired cases decreased (S10 Fig). Under this inference

setting, our estimate of R_c depended only on the proportion of cases that reported travel. When

the proportion of cases that reported travel matched the proportion of cases that were imported,

we correctly estimated R_c (Fig 7B).

Discussion

Our results show that, in most settings, routinely collected surveillance data offer limited value

for reconstructing individual-level transmission networks of *P. falciparum* malaria and informing

estimates of the reproduction number under control, R_c. Using simulated data similar to the

Eswatini surveillance data that we analyzed, we observed that our inference algorithm correctly

identified transmission linkages less than 35% of the time. We attribute this inaccuracy primarily

to the inherently limited information content of spatiotemporal data on *P. falciparum*. Its

characteristically long serial interval [25] means that an appreciable number of cases presenting

within a short timeframe are difficult to link to each other based on their timing, even in a

relatively facile test case in which the generative process assumed in the likelihood function

matched that used to simulate the data.

Under most simulated scenarios and assumptions about the accuracy of travel-history

data, we overestimated the number of locally acquired cases, leading to overestimates of R_c.

Crucially, our simulation sweep demonstrated that routinely collected surveillance data was most

informative of individual-level transmission networks and R_c when local outbreaks were highly

focal in time. Otherwise, while we were able to reconstruct the true transmission network with

modest accuracy, we tended to misclassify truly imported cases as locally acquired, thereby

overestimating R_c. Taken together, these results suggest limited use of routinely collected

surveillance data for informing fine-scale estimates of *P. falciparum* transmission. At broader

spatial scales, however, routinely collected surveillance data may still have practical value,

because the spatial distribution of cases can reveal epidemiological risk factors relevant for

targeted control interventions [26,27].

Although we were able to reach some general conclusions about our inference algorithm,

our inferences were highly sensitive to which data types we included and which assumptions we

made about the accuracy of travel-history data. Applying our algorithm to surveillance data from

Eswatini, we observed that inferred patterns of transmission depended on which data types we

included. With the inclusion of spatial data, we captured a spatial pattern of transmission

consistent with another analysis from Eswatini [28] with data from a different time period.

Assumptions about the travel history appeared to have a strong influence on the overall

magnitude of transmission that we inferred, due to the direct relationship between R_c and the

proportion of imported cases [16]. As a result, believing the travel history, and thereby treating it

as perfectly accurate as in previous approaches [6,18–20], could bias R_c estimates if there are

errors in travel-history data. A study comparing community travel surveys to mobile-phone data

in Kenya found that travel histories considerably underestimated the volume of travel, suggesting

high rates of false negatives in community travel surveys [29]. Believing the travel history may,

then, underestimate the number of imported cases and overestimate R_c.

The method that we used only considered a single spatial model to infer transmission

linkages and assumed complete observation of cases, both of which are factors that could have

affected our inferences based on the Eswatini surveillance data. The diffusion model that we

used to represent spatial dispersion of parasites assumed that movement is isotropic in space and

did not consider landscape features, such as heterogeneity in human population densities and

environmental factors that may affect mosquito ecology. A study analyzing self-reported

movement patterns in Mali, Burkina Faso, Zambia, and Tanzania found that gravity and radiation

models of spatial dispersion fit the data well, though the appropriateness of each model depended

on the type of traveler, the travel distance, and the population size of the destination considered

[30]. Regarding the representation of *P. falciparum* infections in our data set from Eswatini,

there are asymptomatic and mild infections that are unlikely to have been recorded in the

surveillance system yet may comprise a substantial proportion of malaria infections within

Eswatini [13]. Accordingly, it is possible that our assumption of complete observation of cases

could have biased R_c estimates, likely downward due to the fact that missing cases will tend to

make offspring numbers appear smaller than they actually are [31,32]. Even so, we expect that

our conclusions about the sensitivity of transmission network inferences to the choice of data

types and assumptions about travel-history data are robust to these limitations of our study. This

reinforces our conclusion of the need for caution in attempting to reconstruct person-to-person

transmission networks from routine surveillance data.

Given that some of the limitations of our approach may be inherent to the information

content of these data types in this system, one potential avenue for improving inferences of fine-

scale patterns of *P. falciparum* transmission could involve the integration of additional data

streams. For example, mobile-phone data [33] and high-resolution friction surfaces [34] could

more realistically characterize mobility patterns, whereas travel-history information that details

the dates, duration, and location of each trip that has been used in programmatic contexts [26]

could more accurately identify importation events. Additionally, the inclusion of pathogen

genetic data, which has the potential to provide a more direct signal of parasite movement, could

complement traditional epidemiological data [35]. There is also scope for further methodological

development, such as relaxing our assumption of complete observation of infections and

incorporating an underlying mechanistic model of transmission (as in Lau *et al*. [8]; Guzzetta *et*

al. [36]). Incorporating an underlying mechanistic model would relax our uninformative prior

assumption on all possible transmission networks, ruling out transmission networks that are

epidemiologically implausible and allowing us to account for spatial differences in transmission

potential due to different epidemiological factors. To this end, we envision that leveraging the

strengths of our method along with other, complementary methods could strengthen inferences

based on routinely collected epidemiological data and open up new possibilities to make use of

even more data types, such as serological data, prevalence surveys, and pathogen genetic data

[35,37,38].

Conclusions

We demonstrated the limitations of routinely collected surveillance data for the inference of

individual-level transmission networks of *P. falciparum*. We identified a tendency to

overestimate local transmission using routinely collected surveillance data, especially when

outbreaks overlapped in space and time. Using both real data from Eswatini and simulated data,

we identified strong sensitivities of our inferences to the epidemiological setting, the choice of

data types included, and assumptions about the accuracy of travel-history data. Our results

indicated that using spatial and temporal data and believing travel histories yielded the most

plausible estimates of transmission when applied to the Eswatini surveillance data. However, our

simulation sweep demonstrated that the accuracy of our inferences strongly depended on the

accuracy of the travel-history data when the travel-history data were assumed to be accurate.

These sensitivities to the choice of data types and assumptions about the accuracy of travel-

history data could have important programmatic implications if outputs of transmission network

inferences are operationalized. Although this study was specific to *P. falciparum*, the results of

our analyses indicate that future studies inferring transmission networks of *P. falciparum*, or any

pathogen, should carefully consider the epidemiological setting and the choice of data types and

assumptions that inform the model and should validate them using simulated data.

Methods

Bayesian framework for estimating transmission linkages

Our goal was to obtain probabilistic estimates of a transmission network N that defines

transmission linkages among a set of known cases. The transmission network is defined as a

directed, acyclic graph comprised of a set of directed edges represented as $N = \{N_{i,j}\}$ for all i, j.

Each $N_{i,j}$ indicates that case i is hypothesized to contain parasites that are the most direct

observed ancestors of the parasites contained in case j. In addition, at least one edge denoted $N_{u,j}$

must exist in the network, indicating that the parasites contained in case j have no ancestors

among the parasites contained in any known local case and are instead contained in some

unknown case u from some source population s, such that it is denoted u_s. To illustrate this

terminology, an example transmission network is depicted in Fig 8.

Fig 8. Schematic of a hypothetical transmission network. A hypothetical transmission

network is presented along with the corresponding notation. In the schematic, white circles

denote unobserved cases, and black circle denote observed cases. Arrows represent transmission

between two cases.

To estimate N, we used spatial, temporal, and travel-history data about all cases, denoted

as $\vec{X}_s$, X_t, and X_h, respectively. We did so within a Bayesian statistical framework, meaning that

we sought to estimate the joint posterior probability density,

463
$$\Pr\left(N, \Theta \middle| X_t, \vec{X}_s, X_h\right) = \frac{\Pr\left(X_t, \vec{X}_s, X_h \middle| N, \Theta\right) \Pr\left(N, \Theta\right)}{\Pr\left(X_t, \vec{X}_s, X_h\right)}, \quad (1)$$

of the transmission network defined by N and the model parameters Θ conditional on the data $\vec{X}_s$,

X_t, and X_h. The first term in the numerator of eq. (1) is the likelihood of N and Θ conditional on

the data. The second term in the numerator is the prior probability of N and Θ. The term in the

denominator is the probability of the data, which is an intractable quantity to calculate directly

given that it would require evaluation of an extremely high-dimensional integral over N and Θ.

To address this, we used a Markov chain Monte Carlo algorithm to draw random samples of N

and Θ from the posterior distribution specified in eq. (1).

The most critical piece of our inference framework is the likelihood, which we define as a

function of each case j as

$$\mathcal{L}(N, \Theta | X_t, \vec{X}_s, X_h) = \prod_j \Pr(X_{t,j}, \vec{X}_{s,j}, X_{h,j} | N_{.,j}, \Theta). \quad (2)$$

Below, we define the probability of the data associated with each known case j as a function of different assumptions that are possible about how case j is connected to the rest of the transmission network.

Scenario 1: Local transmission between known cases i and j

When case i contains parasites that are immediate ancestors of the parasites contained in case j, we represent its contribution to the likelihood as

$$\Pr(X_{t,j}, \vec{X}_{s,j}, X_{h,j} | N_{i,j}, \Theta) = \Pr(X_{t,j} | N_{i,j}, \Theta) \Pr(\vec{X}_{s,j} | X_{t,j}, N_{i,j}, \Theta) \Pr(X_{h,j} | N_{i,j}, \Theta), \quad (3)$$

which is the product of the probabilities of the temporal, spatial, and travel-history data given the network and model parameters. This formulation assumes that those data are generated independently for each individual, with the exception of a dependence of the spatial data on the temporal data.

Probability of the temporal data. To characterize the time elapsed between two cases resulting from local transmission, we used a model of the generation and serial intervals for *P. falciparum* malaria by Huber *et al*. [25]. The generation interval represents the time between infection of a primary and secondary case, whereas the serial interval represents the time between detection of those cases. Because the timing of infection per se (i.e., an infectious mosquito inoculating a

susceptible human) is typically unknown, we focused on the serial interval as the most apropos

temporal quantity relating cases.

In deriving the probability of a given length of the serial interval, Huber *et al.* [25]

convolved a random variable representing variability in the generation interval (*GI*) with a

random variable representing variability in the time between infection with *P. falciparum* and

detection by surveillance – i.e., the infection to detection period (*IDP*). That framework yielded

504
$$\Pr\left(SI_{i,j} = -a + b + c\right) = \sum_{a}\sum_{b}\sum_{c} \Pr(IDP_i = a)\Pr\left(GI_{i,j} = b\right)\Pr\left(IDP_j = c\right), \quad (4)$$

as the probability of a serial interval of length $SI_{i,j}$. We allowed for different models of the serial

interval depending upon differences in the *GI* and *IDP* for different types of primary and

secondary cases. For instance, we assumed that symptomatic cases present in a clinic some

number of days after infection as informed by empirical data from Zanzibar [25]. For an

asymptomatic infection, we assumed that detection occurred through active surveillance at a

randomly drawn day among all days where its asexual parasitemia exceeds a detection threshold

[25]. The choice of *IDP* for both the primary and secondary case informs the probability of two

cases separated in time by $SI_{i,j} = X_{t,j} - X_{t,i}$ days.

Probability of the spatial data. Following Reiner *et al.* [6], we assumed that a simple two-

dimensional Wiener diffusion process determines the location of secondary cases relative to the

location of their associated primary case. It follows that, for a given diffusion coefficient *D* with

units $km^2 day^{-1}$ and generation interval $GI_{i,j}$, the two-dimensional location $\vec{X}_{s,j}$ of the secondary

case *j* is described by a bivariate normal distribution with probability density

$$f\left(\vec{X}_{s,j} \mid \vec{X}_{s,i}, D, GI_{i,j}, N_{i,j}, \Theta\right) = \frac{1}{2\pi\sigma^2\left(GI_{i,j}\right)} e^{-\frac{\|\vec{X}_{s,j}-\vec{X}_{s,i}\|}{2\sigma^2\left(GI_{i,j}\right)}}, \quad (5)$$

where $\sigma^2\left(GI_{i,j}\right) = DGI_{i,j}$. This formulation assumes that each spatial dimension is independent,

that the variance scales linearly with the generation interval, and that movement is isotropic

across a continuous landscape.

One complication to eq. (5) is that the generation interval $GI_{i,j}$ is unobserved and,

therefore, cannot take on a fixed value. Instead, we must use data about the serial interval $SI_{i,j}$ to

inform our generative model for $\vec{X}_{s,j}$. To do so, we take advantage of the property of normal

random variables that the sum of two or more random variables is itself a normal random

variable [39]. This property allows us to recast eq. (5) as a function of SI rather than GI by

computing the appropriate σ^2 as

$$\sigma^2(SI) = \int \sigma^2(GI) \Pr(GI|SI) \, dGI, \quad (6)$$

535 which is effectively a weighted sum of the spatial variances associated with a given GI

536 proportional to the probability of each GI conditional on the observed SI. This results in

$$f\left(\vec{X}_{s,j} \mid \vec{X}_{s,i}, D, SI_{i,j}, N_{i,j}, \Theta\right) = \frac{1}{2\pi\sigma^2\left(SI_{i,j}\right)} e^{-\frac{\|\vec{X}_{s,j}-\vec{X}_{s,i}\|}{2\sigma^2\left(SI_{i,j}\right)}}, \quad (7)$$

540 as the probability density of the spatial data that we assume.

541   In the event that case i has missing spatial data, we cannot compute the spatial likelihood

542   of eq. (7). To address this, we define a latent unobserved quantity $\tilde{X}_{s,i}$, which represents the

543   unknown location of case i. We then integrate over the uncertainty in $\tilde{X}_{s,i}$,

545
$$f\left(\vec{X}_{s,j}\middle|D, SI_{i,j}, N_{i,j}, \Theta\right) = \int f\left(\vec{X}_{s,j}\middle|\tilde{X}_{s,i}, D, N_{i,j}, \Theta\right) f\left(\tilde{X}_{s,i}\middle|\vec{X}_{s,j}, D\right) d\,\tilde{X}_{s,i}, \quad (8)$$

to compute the probability density of case j with known spatial location $\vec{X}_{s,j}$ arising from case i

with unknown spatial location $\tilde{X}_{s,i}$. Equation (8) is computed as the product of the probability

density of the location of a known case j conditional on an unknown location $\tilde{X}_{s,i}$ and the

probability density of spatial separation $\vec{X}_{s,j} - \tilde{X}_{s,i}$ conditional on the diffusion coefficient D for

all $\tilde{X}_{s,i}$. Because we assume that movement is isotropic, eq. (8) is a two-dimensional Gaussian

integral, simplifying to

554
$$f\left(\vec{X}_{s,j}\middle|D, SI_{i,j}, N_{i,j}, \Theta\right) = \frac{1}{4\pi\sigma^2(SI_{i,j})}. \quad (9)$$

In the event that case j has missing spatial data and case i has known spatial data, the latent

unobserved quantity becomes $\tilde{X}_{s,j}$. We then integrate over the uncertainty in $\tilde{X}_{s,j}$ and calculate

$f\left(\vec{X}_{s,j}\middle|D, SI_{i,j}, N_{i,j}, \Theta\right)$ using eq. (8-9).

Probability of the travel-history data. Although we assume in this scenario that a person's

infection was locally acquired, our model must still be capable of explaining the travel-history

data $X_{h,j}$. We define a probability τ_l that case j reported travel (i.e., $X_{h,j} = 1$) even though they

were not infected during that period of travel, such that

565
$$\Pr(X_{h,j}|N_{i,j}, \Theta) = \begin{cases} \tau_l, & X_{h,j} = 1 \\ 1 - \tau_l, & X_{h,j} = 0 \end{cases}. \quad (10)$$

567   In the event that case j has missing travel-history data, we cannot compute the travel-

568 history likelihood of eq. (10). To address this, we defined a latent unobserved quantity $\tilde{X}_{h,j}$,

569 which represents the unknown travel history of case j. We then sum across the uncertainty in

570 $\tilde{X}_{h,j}$,

572
$$\Pr(X_{h,j} = \text{NA}|N_{i,j}, \Theta) = \Pr(\tilde{X}_{h,j} = 1)\tau_l + (1 - \Pr(\tilde{X}_{h,j} = 1))(1 - \tau_l), \quad (11)$$

to compute the probability that case j was locally acquired given an unknown travel history. In

eq. (11), $\Pr(\tilde{X}_{h,j} = 1)$ was computed as the proportion of cases with a positive travel history

among all cases with known travel-history data.

Taken together with the probabilities of the temporal and spatial data described above,

the product of these three probabilities constitutes the entirety of the contribution of a case j

infected by a known local case i to the overall likelihood of N and Θ.

Scenario 2: Importation of local case j from source population s

In the event of $N_{u_s,j}$, we represent the contribution of such a case to the overall likelihood of N

and Θ as the product of the probabilities of its temporal, spatial, and travel-history data under

similar assumptions as in Scenario 1. The key difference in this scenario is that there is no

information about the unknown source case that gave rise to case *j*.

Probability of the temporal data. Because the person containing parasites that are the direct

ancestors of those in case *j* is unobserved and does not have an $X_{t,i}$, we are unable to compute the

probability of the temporal data as described in Scenario 1. It is important though to obtain a

probability comparable to that from Scenario 1 as a reference point for determining whether it is

more likely that a given case arose from some other known local case or from an unknown case

u_s from source population *s*. To do so, we consider the variable $\tilde{X}_{t,u_s}$, which is a latent variable

describing the timing of when u_s would have been detected, had it been detected.

Because u_s is not observed, we considered it to be asymptomatic and untreated. We then

calculated the probability of the timing of a known case *j* arising from an unknown case u_s as

$$\Pr(X_{t,j}|N_{u_s,j},\Theta) = \int \Pr(X_{t,j}|\tilde{X}_{t,u_s}, N_{u_s,j}, \Theta) \Pr(\mathrm{SI} = X_{t,j} - \tilde{X}_{t,u_s}) \, d\tilde{X}_{t,u_s}, \quad (12)$$

by integrating over uncertainty in $\tilde{X}_{t,u_s}$. We represented this as the product of the probability of

the timing of a known case *j* conditional on an unknown time of detection $\tilde{X}_{t,u_s}$ and the

probability of the serial interval $X_{t,j} - \tilde{X}_{t,u_s}$ for all $\tilde{X}_{t,u_s}$. In equation (12), we did not distinguish

between symptomatic and asymptomatic cases *j* because the calculation is identical; only the

serial interval distributions differ.

Probability of the spatial data. Without an $\tilde{X}_{t,u_s}$ for the unobserved case u_s, we lacked

information on the serial interval between it and case j. Consequently, we were unable to use the

probability from eq. (7) in that particular form. Instead, we computed the spatial variance as a

function of the diffusion coefficient alone, yielding

$$\sigma^2(D) = \int D \, GI \, \Pr(GI) \, dGI. \quad (13)$$

612   Equation (13) integrates across all possible generation intervals and simplifies to $D\mathbb{E}[GI]$, the

613   product of the diffusion coefficient and the expectation of the generation interval.

614   We applied this spatial variance to the unobserved latent variable $\tilde{X}_{s,u_s}$, which represents

615   the unknown location of the unobserved case u_s. We integrated over uncertainty in $\tilde{X}_{s,u_s}$ to

616   compute the probability density,

$$f(X_{s,j}|D, N_{u_s,j}, \Theta) = \int f(X_{s,j}|\tilde{X}_{s,u_s}, D, N_{u_s,j}, \Theta) f(\tilde{X}_{s,u_s}|X_{s,j}, D) d\tilde{X}_{s,u_s}, \quad (14)$$

620   of the location of a known case j arising from an unknown source case u_s with unknown location

621    $\tilde{X}_{s,u_s}$. This is represented as the product of the probability density of the location of a known case

622   j conditional on an unknown location $\tilde{X}_{s,u_s}$ and the probability density of spatial separation

623    $X_{s,j} - \tilde{X}_{s,u_s}$ conditional on the diffusion coefficient D for all $\tilde{X}_{s,u_s}$. As in eq. (9), we treated eq.

624   (14) as an evaluation of the Gaussian integral, evaluating to

$$f\left(X_{s,j}|D, N_{u_s,j}, \Theta\right) = \frac{1}{4\pi D \mathbb{E}[GI]}. \quad (15)$$

In eq. (15), D is the diffusion coefficient and $\mathbb{E}[GI]$ is the expectation of the generation interval.

Probability of the travel-history data. We considered the travel history $X_{h,j}$ to be a binary

variable with a value of 1 indicating reported travel to an area with known or assumed malaria

transmission within a timeframe consistent with the person having become infected there. After

defining the probability τ_s that $X_{h,j} = 1$ conditional on $N_{u_s,j}$, it follows that

$$\Pr\left(X_{h,j}|N_{u_s,j}, \Theta\right) = \begin{cases} \tau_s, & X_{h,j} = 1 \\ 1 - \tau_s, & X_{h,j} = 0 \end{cases}, \quad (16)$$

which constitutes the contribution of the travel history of such a case to the overall likelihood of

N and Θ. If the travel history of case j is unknown, an analogous calculation to eq. (11) is made

using τ_s.

Bayesian inference

Markov Chain Monte Carlo Algorithm. To avoid evaluating the high-dimensional integral

over N and Θ, we drew samples of N and Θ from their posterior distribution defined by eq. (1)

using a Metropolis-Hastings Markov chain Monte Carlo (MCMC) method [40,41]. To begin the

chain, N and Θ were initialized to $N^{(1)}$ and $\Theta^{(1)}$, and each subsequent step i in the chain was

646   denoted $N^{(i)}$ and $\Theta^{(i)}$. At each step, states N' and Θ' were proposed with $\Pr\left(\left(N^{(i)}, \Theta^{(i)}\right) \rightarrow\right.$

647    $\left.(N', \Theta')\right)$. Proposed states were accepted with probability

649
$$\alpha_{\text{update}} = \min\left[1, \frac{\pi(N', \Theta')\,\Pr\left((N', \Theta') \rightarrow \left(N^{(i)}, \Theta^{(i)}\right)\right)}{\pi(N^{(i)}, \Theta^{(i)})\,\Pr\left((N^{(i)}, \Theta^{(i)}) \rightarrow (N', \Theta')\right)}\right], \quad (17)$$

651   where $\pi(N, \Theta)$ is the product of the likelihood $\Pr\left(\vec{X}_s, X_t, X_h | N, \Theta\right)$ of N and Θ conditional on the

652   data and the assumed prior probability $\Pr(N, \Theta)$ of N and Θ. After a random draw R from a

653   uniform distribution, the chain was updated according to

655
$$N^{(i+1)}, \Theta^{(i+1)} = \begin{cases} N', \Theta', & R \leq \alpha \\ N^{(i)}, \Theta^{(i)}, & R > \alpha \end{cases}. \quad (18)$$

657   To reduce the probability of the chain becoming stuck at a local maximum, we employed

658   Metropolis-coupled Markov chain Monte Carlo (MC³) [42]. Implementing MC³ involved

659   running multiple chains in parallel, with $\pi_c(N, \Theta)$ in chain c raised to the power β_c according to

661
$$\beta_c = 1 + \lambda(c - 1), \quad (19)$$

where $\lambda > 0$ is a temperature increment parameter that governs the degree to which each chain is

"heated." As a result of setting $\beta_1 = 1$, $\pi_1(N, \Theta)$ is directly proportional to the joint posterior

distribution and is referred to as the master or "cold" chain. This algorithm effectively flattens

the likelihood in the heated chains by setting $\beta_c > 1$, allowing them to explore the parameter

space more freely and to encounter alternative high-density regions more readily than the cold

chain would alone. At a pre-defined frequency, two randomly selected chains i and j were

allowed to swap parameter sets according to a swap probability

$$\alpha_{\text{swap}} = \min\left[1, \frac{\pi\left(N^{(j)}, \Theta^{(j)}\right)^{\beta_i} \pi\left(N^{(i)}, \Theta^{(i)}\right)^{\beta_j}}{\pi(N^{(i)}, \Theta^{(i)})^{\beta_i} \pi(N^{(j)}, \Theta^{(j)})^{\beta_j}}\right], \quad (20)$$

where $\pi(N, \Theta)$ is the same as it was in eq. (17). A swap into the master chain only occurred if it

was from one of the two randomly selected chains and $R \leq \alpha_{\text{swap}}$. We recorded a total of 100

million samples from the posterior distribution, discarding the first 50 million samples as burn-in

and thinning the chain every 10,000 samples between each recorded sample.

Proposals. Proposals made by the MC^3 algorithm involved changes to the parameters (i.e., D, τ_s,

and τ_l) and changes to the transmission network topology. Each proposal occurred with a fixed

probability, where the sum of these proposal probabilities was equal to one.

Proposals to change parameters involved updating D, τ_s, or τ_l. To update the value of D, a

new value was drawn from a normal distribution with mean set to the current value of the

parameter and variance set to 2.5. Values of D proposed must be strictly nonnegative, so we

rejected any proposed D that was less than zero and assigned $\alpha_{\text{update}} = 0$. Similarly, new values

of τ_s and τ_l were chosen according to normal distributions with means set to their current

parameter value and variance set to 0.25. Because τ_s and τ_l are probabilities, we rejected any

proposed value that fell outside the range [0,1] and assigned $\alpha_{\text{update}} = 0$.

Changes proposed to the network topology involved the addition or removal of an

ancestor from a randomly selected node. We assigned a uniform probability of proposing case a

as an ancestor to a randomly selected case i, such that proposals to the network topology are

uninformed by spatial and temporal data. Furthermore, we defined the proposal probability of

removing case a as an ancestor to a randomly selected case i as

$$\Pr(\text{remove } a) = \bar{A}_i^{-1}, \quad (21)$$

where $\bar{A}_i$ represents the size of set A_i of all ancestors to case i. Proposed changes to the network

are then accepted according to eq. (17).

Prior assumptions. We placed strong priors on τ_s and τ_l, because we assumed that travel

histories were mostly, but not completely, accurate. We used a beta-distributed prior on τ_s, with

parameters $\alpha_{\tau_s} = 12$ and $\beta_{\tau_s} = 3$, which resulted in a mean of 0.8 and a variance of 0.01 for this

prior distribution. We also used a beta distributed prior on τ_l, with parameters $\alpha_{\tau_l} = 3$ and $\beta_{\tau_l} =$

12, which resulted in a mean of 0.2 and a variance of 0.01. We assumed a uniform prior on D

over the interval $[10^{-3}, \infty)$ and an even prior across all possible network configurations,

meaning that those prior probabilities canceled out in eqs. (17) and (20).

Assessing convergence. For D, τ_s, and τ_l, we assessed convergence using the Gelman-Rubin

statistic [43], with values below 1.1 indicating convergence. For the transmission network N, we

assessed convergence by calculating correlation coefficients of case-level probabilities across

five chains from independent realizations of the MC^3 algorithm, for a total of 10 pairwise

comparisons across the five chains. The two case-level probabilities that we considered were the

posterior probability that each case was infected by an unknown case u_s from a source population

and the posterior probability that each case j was infected by each other case i. Higher values of

these correlation coefficients provided stronger support for convergence.

Ethical Considerations

Ethical approval was obtained from the Eswatini Ministry of Health, the University of

California, San Francisco, and the University of Notre Dame (IRB 19-06-5408). All data were

analyzed anonymously.

Acknowledgments

The authors thank the National Malaria Elimination Programme as well as Nontokozo Mngadi

and Deepa Pindolia from the Clinton Health Access Initiative for their support in the collection

of the surveillance data used in this study. We also thank Brooke Whittemore for her support in

data management.

721 ## References

1. Wallinga J, Teunis P. Different epidemic curves for severe acute respiratory syndrome

reveal similar impacts of control measures. Am J Epidemiol. 2004;160: 509–516.

doi:10.1093/aje/kwh255

2. White LF, Archer B, Pagano M. Estimating the reproductive number in the presence of

spatial heterogeneity of transmission patterns. Int J Health Geogr. 2013;12: 35.

doi:10.1186/1476-072X-12-35

3.   Métras R, Baguelin M, Edmunds WJ, Thompson PN, Kemp A, Pfeiffer DU, et al.

Transmission Potential of Rift Valley Fever Virus over the Course of the 2010 Epidemic in

South Africa. Emerg Infect Dis. 2013;19: 916–924. doi:10.3201/eid1906.121641

4.   Backer JA, Wallinga J. Spatiotemporal Analysis of the 2014 Ebola Epidemic in West

Africa. Lessler J, editor. PLoS Comput Biol. 2016;12: e1005210.

doi:10.1371/journal.pcbi.1005210

5.   Salje H, Cummings DAT, Lessler J. Estimating infectious disease transmission distances

using the overall distribution of cases. Epidemics. 2016;17: 10–18.

doi:10.1016/j.epidem.2016.10.001

6.   Reiner RC, Le Menach A, Kunene S, Ntshalintshali N, Hsiang MS, Perkins TA, et al.

Mapping residual transmission for malaria elimination. eLife. 2015;4.

doi:10.7554/eLife.09520

7.   Lau MSY, Dalziel BD, Funk S, McClelland A, Tiffany A, Riley S, et al. Spatial and

temporal dynamics of superspreading events in the 2014-2015 West Africa Ebola epidemic.

Proc Natl Acad Sci USA. 2017;114: 2337–2342. doi:10.1073/pnas.1614595114

8.   Lau MSY, Gibson GJ, Adrakey H, McClelland A, Riley S, Zelner J, et al. A mechanistic

spatio-temporal framework for modelling individual-to-individual transmission—With an

application to the 2014-2015 West Africa Ebola outbreak. Ferrari M (Matt), editor. PLoS

Comput Biol. 2017;13: e1005798. doi:10.1371/journal.pcbi.1005798

9.   Jombart T, Cori A, Didelot X, Cauchemez S, Fraser C, Ferguson N. Bayesian

Reconstruction of Disease Outbreaks by Combining Epidemiologic and Genomic Data.

Tanaka MM, editor. PLoS Comput Biol. 2014;10: e1003457.

doi:10.1371/journal.pcbi.1003457

10. Didelot X, Gardy J, Colijn C. Bayesian Inference of Infectious Disease Transmission from

Whole-Genome Sequence Data. Molecular Biology and Evolution. 2014;31: 1869–1879.

doi:10.1093/molbev/msu121

11. Mollentze N, Nel LH, Townsend S, le Roux K, Hampson K, Haydon DT, et al. A Bayesian

approach for inferring the dynamics of partially observed endemic infectious diseases from

space-time-genetic data. Proc Biol Sci. 2014;281: 20133251. doi:10.1098/rspb.2013.3251

12. Ypma RJF, van Ballegooijen WM, Wallinga J. Relating Phylogenetic Trees to

Transmission Trees of Infectious Disease Outbreaks. Genetics. 2013;195: 1055–1062.

doi:10.1534/genetics.113.154856

13. Sturrock HJW, Hsiang MS, Cohen JM, Smith DL, Greenhouse B, Bousema T, et al.

Targeting Asymptomatic Malaria Infections: Active Surveillance in Control and

Elimination. PLoS Med. 2013;10: e1001467. doi:10.1371/journal.pmed.1001467

14. Bousema T, Griffin JT, Sauerwein RW, Smith DL, Churcher TS, Takken W, et al. Hitting

Hotspots: Spatial Targeting of Malaria for Control and Elimination. PLoS Med. 2012;9:

e1001165. doi:10.1371/journal.pmed.1001165

15. Bejon P, Williams TN, Nyundo C, Hay SI, Benz D, Gething PW, et al. A micro-

epidemiological analysis of febrile malaria in Coastal Kenya showing hotspots within

hotspots. eLife. 2014;3: e02130. doi:10.7554/eLife.02130

16.   Cohen JM, Moonen B, Snow RW, Smith DL. How absolute is zero? An evaluation of

historical and current definitions of malaria elimination. Malar J. 2010;9: 213.

doi:10.1186/1475-2875-9-213

17.   Cohen JM, Le Menach A, Pothin E, Eisele TP, Gething PW, Eckhoff PA, et al. Mapping

multiple components of malaria risk for improved targeting of elimination interventions.

Malar J. 2017;16: 459. doi:10.1186/s12936-017-2106-3

18.   Churcher TS, Cohen JM, Novotny J, Ntshalintshali N, Kunene S, Cauchemez S. Measuring

the path toward malaria elimination. Science. 2014;344: 1230–1232.

doi:10.1126/science.1251449

19.   Routledge I, Chevéz JER, Cucunubá ZM, Rodriguez MG, Guinovart C, Gustafson KB, et

al. Estimating spatiotemporally varying malaria reproduction numbers in a near elimination

setting. Nat Commun. 2018;9: 2476. doi:10.1038/s41467-018-04577-y

20.   Routledge I, Lai S, Battle KE, Ghani AC, Gomez-Rodriguez M, Gustafson KB, et al.

Tracking progress towards malaria elimination in China: Individual-level estimates of

transmission and its spatiotemporal variation using a diffusion network approach. Pascual

M, editor. PLoS Comput Biol. 2020;16: e1007707. doi:10.1371/journal.pcbi.1007707

21.   Marshall JM, Bennett A, Kiware SS, Sturrock HJW. The Hitchhiking Parasite: Why Human

Movement Matters to Malaria Transmission and What We Can Do About It. Trends in

Parasitology. 2016;32: 752–755. doi:10.1016/j.pt.2016.07.004

22.   Campbell F, Cori A, Ferguson N, Jombart T. Bayesian inference of transmission chains

using timing of symptoms, pathogen genomes and contact data. Pitzer VE, editor. PLoS

Comput Biol. 2019;15: e1006930. doi:10.1371/journal.pcbi.1006930

23.   Wood SN. Generalized additive models: an introduction with R. Boca Raton, FL: Chapman

& Hall/CRC; 2006.

24.   R Development Core Team. R: A Language and Environment for Statistical Computing.

Vienna, Austria: R Foundation for Statistical Computing; 2017.

25.   Huber JH, Johnston GL, Greenhouse B, Smith DL, Perkins TA. Quantitative, model-based

estimates of variability in the generation and serial intervals of Plasmodium falciparum

malaria. Malaria Journal. 2016;15. doi:10.1186/s12936-016-1537-6

26.   Cohen JM, Dlamini S, Novotny JM, Kandula D, Kunene S, Tatem AJ. Rapid case-based

mapping of seasonal malaria transmission risk for strategic elimination planning in

Swaziland. Malar J. 2013;12: 61. doi:10.1186/1475-2875-12-61

27.   Hsiang MS, Ntuku H, Roberts KW, Dufour M-SK, Whittemore B, Tambo M, et al.

Effectiveness of reactive focal mass drug administration and reactive focal vector control to

reduce malaria transmission in the low malaria-endemic setting of Namibia: a cluster-

randomised controlled, open-label, two-by-two factorial design trial. The Lancet. 2020;395:

1361–1373. doi:10.1016/S0140-6736(20)30470-0

28.   Sturrock HJ, Cohen JM, Keil P, Tatem AJ, Le Menach A, Ntshalintshali NE, et al. Fine-

scale malaria risk mapping from routine aggregated case data. Malar J. 2014;13: 421.

doi:10.1186/1475-2875-13-421

29.   Wesolowski A, Stresman G, Eagle N, Stevenson J, Owaga C, Marube E, et al. Quantifying

travel behavior for infectious disease research: a comparison of data from surveys and

mobile phones. Sci Rep. 2015;4: 5678. doi:10.1038/srep05678

30.   Marshall JM, Wu SL, Sanchez C. HM, Kiware SS, Ndhlovu M, Ouédraogo AL, et al.

Mathematical models of human mobility of relevance to malaria transmission in Africa. Sci

Rep. 2018;8: 7713. doi:10.1038/s41598-018-26023-1

31.   Blumberg S, Lloyd-Smith JO. Inference of R0 and Transmission Heterogeneity from the

Size Distribution of Stuttering Chains. Ferguson N, editor. PLoS Comput Biol. 2013;9:

e1002993. doi:10.1371/journal.pcbi.1002993

32.   Blumberg S, Lloyd-Smith JO. Comparing methods for estimating R0 from the size

distribution of subcritical transmission chains. Epidemics. 2013;5: 131–145.

doi:10.1016/j.epidem.2013.05.002

33.   Wesolowski A, Eagle N, Tatem AJ, Smith DL, Noor AM, Snow RW, et al. Quantifying the

Impact of Human Mobility on Malaria. Science. 2012;338: 267–270.

doi:10.1126/science.1223467

34.   Weiss DJ, Nelson A, Gibson HS, Temperley W, Peedell S, Lieber A, et al. A global map of

travel time to cities to assess inequalities in accessibility in 2015. Nature. 2018;553: 333–

336. doi:10.1038/nature25181

35.   Wesolowski A, Taylor AR, Chang H-H, Verity R, Tessema S, Bailey JA, et al. Mapping

malaria by combining parasite genomic and epidemiologic data. BMC Med. 2018;16: 190.

doi:10.1186/s12916-018-1181-9

36.   Guzzetta G, Marques-Toledo CA, Rosà R, Teixeira M, Merler S. Quantifying the spatial

837   spread of dengue in a non-endemic Brazilian metropolis via transmission chain

838   reconstruction. Nature Communications. 2018;9. doi:10.1038/s41467-018-05230-4

839   37.   Weiss DJ, Lucas TCD, Nguyen M, Nandi AK, Bisanzio D, Battle KE, et al. Mapping the

840   global prevalence, incidence, and mortality of Plasmodium falciparum, 2000–17: a spatial

841   and temporal modelling study. The Lancet. 2019;394: 322–331. doi:10.1016/S0140-

842   6736(19)31097-9

843   38.   Greenhouse B, Smith DL, Rodríguez-Barraquer I, Mueller I, Drakeley CJ. Taking Sharper

844   Pictures of Malaria with CAMERAs: Combined Antibodies to Measure Exposure Recency

845   Assays. The American Journal of Tropical Medicine and Hygiene. 2018;99: 1120–1127.

846   doi:10.4269/ajtmh.18-0303

847   39.   Lemons DS, Langevin P. An introduction to stochastic processes in physics: containing

848   "On the theory of Brownian motion" by Paul Langevin, translated by Anthony Gythiel.

849   Baltimore: Johns Hopkins University Press; 2002.

850   40.   Metropolis N, Rosenbluth AW, Rosenbluth MN, Teller AH, Teller E. Equation of State

851   Calculations by Fast Computing Machines. The Journal of Chemical Physics. 1953;21:

852   1087–1092. doi:10.1063/1.1699114

853   41.   Hastings WK. Monte Carlo Sampling Methods Using Markov Chains and Their

854   Applications. Biometrika. 1970;57: 97. doi:10.2307/2334940

42. Altekar G, Dwarkadas S, Huelsenbeck JP, Ronquist F. Parallel Metropolis coupled Markov

chain Monte Carlo for Bayesian phylogenetic inference. Bioinformatics. 2004;20: 407–415.

doi:10.1093/bioinformatics/btg427

43. Gelman A, Rubin DB. Inference from Iterative Simulation Using Multiple Sequences.

Statistical Science. 1992;7: 457–472. doi:10.1214/ss/1177011136

Supporting Information

S1 Fig. Likelihood profile of the diffusion coefficient conditioning on the true network. The

likelihood profile for two inference settings that incorporate spatial data are shown as a function

of the diffusion coefficient. The green line corresponds to the default inference setting, the purple

line corresponds to the setting in which spatial and temporal data are used and the travel history

is believed, and the blue line corresponds to the setting in which spatial and temporal data are

used and the travel history was ignored. Black bars denote the true value of the diffusion

coefficient, dark grey shapes denote the region with the maximum likelihood, and light grey

shapes denote the region within 10 log likelihood units of the maximum likelihood.

S2 Fig. Comparison of individual-level importation probabilities on Eswatini data under

default settings. Pairwise scatter plots of importation probabilities are shown for each pairing of

the 5 independent replicates to assess convergence. The correlation between importation

probabilities for each pair of replicates is reported in red.

S3 Fig. Comparison of transmission linkage probabilities on Eswatini data under default

settings. Pairwise scatter plots of the probability of each transmission linkage are shown for each

pairing of the 5 independent replicates to assess convergence. The correlation between

transmission linkage probabilities for each pair of replicates is reported in red.

S4 Fig. Comparison of individual-level R_c estimates on Eswatini data under default settings. Pairwise scatter plots of individual-level R_c are show for each pairing of the five independent replicates to assess convergence. The correlation between individual-level R_c values is reported in red.

S5 Fig. Comparison of the prior and posteriors of τ_s and τ_l from the Eswatini surveillance

data using spatial and temporal data and estimating the accuracy of the travel history. The

prior (gray shape), the analytical posterior distribution (black line), and the numerical posterior

distribution from MC3 (green histogram) are plotted for τ_s and τ_l.

S6 Fig. Comparison of the prior and posteriors of τ_s and τ_l from the Eswatini surveillance data using temporal data and estimating the accuracy of the travel history. The prior (gray shape), the analytical posterior distribution (black line), and the numerical posterior distribution from MC3 (pink histogram) are plotted for τ_s and τ_l.

S7 Fig. Comparison of the prior and posteriors of τ_s from simulated data using spatial and

temporal data and estimating the accuracy of the travel history. The prior (gray shape), the

analytical posterior distribution (black line), and the numerical posterior distribution from MC3

(green histogram) are plotted for τ_s.

S8 Fig. Univariate relationships between accuracy metrics and simulation parameters

using spatial and temporal data and estimating the accuracy of the travel history.

Scatterplots of the relationship between five accuracy metrics and the simulation parameters are

reported. Imported classification refers to the proportion of imported cases that are correct

classified as imported, local classification refers to the proportion of locally acquired cases that

are correctly classified as locally acquired, and case classification refers to the proportion of all

cases that are correctly classified as imported or locally acquired. Transmission linkage is the

proportion of locally acquired cases for which the true parent is correctly identified, and outbreak

is the proportion of locally acquired cases for which the inferred parent belongs to the same

outbreak.

S9 Fig. Univariate relationships between accuracy metrics and simulation parameters

using spatial and temporal data only. Scatterplots of the relationship between five accuracy

metrics and the simulation parameters are reported. Imported classification refers to the

proportion of imported cases that are correct classified as imported, local classification refers to

the proportion of locally acquired cases that are correctly classified as locally acquired, and case

classification refers to the proportion of all cases that are correctly classified as imported or

locally acquired. Transmission linkage is the proportion of locally acquired cases for which the

true parent is correctly identified, and outbreak is the proportion of locally acquired cases for

which the inferred parent belongs to the same outbreak.

S10 Fig. Univariate relationships between accuracy metrics and simulation parameters

using spatial and temporal data and believing the travel history. Scatterplots of the

relationship between five accuracy metrics and the simulation parameters are reported. Imported

classification refers to the proportion of imported cases that are correct classified as imported,

local classification refers to the proportion of locally acquired cases that are correctly classified

as locally acquired, and case classification refers to the proportion of all cases that are correctly

classified as imported or locally acquired. Transmission linkage is the proportion of locally

acquired cases for which the true parent is correctly identified, and outbreak is the proportion of

locally acquired cases for which the inferred parent belongs to the same outbreak.

S1 File. Full description of the analyses. This file contains a full description of the analyses run

on both the Eswatini data sets and the simulated data sets.

S2 File. Analytical solution for the posterior distribution of τ_s and τ_l. The analytical solutions

for τ_s and τ_l are derived using conjugate-prior relationships and compared to the empirical

posterior distributions obtained using MC^3.