Preliminary report on the development of a colorimetric loop-mediated isothermal amplification diagnostic assay for deer tick virus

Brandon G. Roy[1]* and Eric J. Ryndock[2]

[1]Biochemistry and Molecular Biology, Lebanon Valley College, 101 North College Ave, Annville, PA 17003.

[2]Biology Department, Millersville University, 40 Dilworth Road, Millersville, PA 17551.

*To whom correspondence should be addressed: bgr36@cornell.edu (Brandon G. Roy)

Abstract: Deer tick virus (DTV) is an emerging pathogen in North America. This virus can cause nervous system complications such as encephalitis in humans. Further, no data has been surmounted around long-term effects of infection from DTV patients across variable age groups. Diagnostic tools of DTV used by government laboratories are based on RT-PCR using patient serum or ticks. This paper explores the feasibility of a colorimetric loop-mediated isothermal amplification (LAMP) assay to create a point-of-care diagnostic methodology for use in field and in primary care. LAMP consists of six primers that bind to target DNA and amplifies variable length nucleotide strands that can be visualized through side reactions or via electrophoresis. First, a viable LAMP primer set, and a primer set that dimerizes and amplifies DNA regardless of compatibility were created in silico and validated in vitro. Then, a specific LAMP assay was developed. Our findings showed his method can be performed within 30 minutes and can measure with limits of detection comparable to PCR.

Introduction

Deer tick virus (DTV) has a single-stranded RNA genome and belongs to the genus *Flavivirus* in the family *Flaviviridae*. Other members of this virus genus and family are dengue virus, Zika virus and West Nile virus. The genome length of DTV is roughly 10,800 bases and it known to cause tick-bourne encephalitis (Artsob et al., 2001).

The increasing presence of DTV throughout North America is coupled with various tick species (CDC, 2019). DTV is also referred to as the second lineage Powassan virus (POW2), deviating from a native strain in Europe and Russia. POW1, or Powassan virus lineage I, has a variable genetic diversity along the theoretical divergence cutoff at 84% nucleotide identity to DTV, 94% amino acid identity, and is estimated to have diverged around 200 years ago (Pesko et al., 2010). This cutoff provided the difference between the two Powassan viruses on the cladogram of *Flaviviridae* (Telford III, 1997). Patient testing for DTV is usually performed in response to severe cases where symptoms arrive suddenly. Most people tested for DTV have symptoms of acute encephalitis or overactive inflammation of the brain, which leads to death in some cases without proper medical treatment (Solomon et al., 2018; Tavakoli et al., 2009). Remaining cases go unbeknownst because testing measures are not approached from typical or asymptomatic standpoints as other tick-borne infections. Current diagnostic testing is limited to the Center of Disease Control and its partnering laboratories using either reverse transcription-polymerase chain reaction (RT-PCR) or immunohistochemical techniques (CDC, 2019; Morozova et al., 2002). A sample of serum or cerebral spinal fluid is collected and sent to one of the offsite diagnostic centers, further complicating the testing process and taking up to a few days for results to become available.

A more time effective detection method than RT-PCR would be beneficial for point-of-care testing (POCT), as well as for field surveys of tick samples that are performed by state game commissions. Three different species of ticks, i.e., *Ixodes cookie, Ixodes marxi,* and *Ixodes scapularis*, have the potential to transmit DTV from their main vertebrate hosts, i.e., woodchucks, squirrels, and white-footed mice, to humans (Brackney et al., 2010). White-tailed deer is also a known host to *Ixodes scapularis.* DTV has no known pathology on these intermediate mammal hosts, but concern is comprehensible for humans (CDC, 2019). The main tick vector that raises human concern is *Ixodes scapularis.* This tick species has a parasitic relationship with white-tailed deer, *Odocoileus virginianus,* and other species of deer that humans encounter frequently in rural areas (Brackney et al., 2010). This host relationship is unique to DTV and unlike the previously mentioned POW1 strain. The prevalence of outdoor recreation and leisure activities give exposure to the tick vectors and potential pathogens to be transmitted.

Monitoring the presence of DTV provides the first steps to surveying disease spread and to developing mitigation efforts. A rapid DTV detection assay like colorimetric loop-mediated isothermal amplification (LAMP) would prove useful for field surveys and POCT. LAMP is an alternative nucleic acid amplification technique that has gained popularity in the past decade (Anupama et al., 2019; Chen et al., 2014; Lamb et al., 2018; Lau et al., 2015; Luo et al., 2011; Shin et al., 2018; Tomlinson, 2008). LAMP is a nucleotide amplification process that occurs through the binding of specially designed primers at approximately 60°C and will generate variable ladder structures that can be visualized in two manners. Colorimetric LAMP does not require expensive and sophisticated equipment or visualization techniques other than the naked eye. In addition, it is inexpensive and quick. For example, the Colorimetric Warmstart® MasterMix (New England Biolabs, Cat# M1800S) LAMP technique, utilized in this paper, is up to five times quicker than PCR and changes color for visual assessment of the results (Nagamine et al., 2002). The colorimetric test starts out as pink and turns a bright yellow color

if the result is positive. Color change is dependent on the emission of protons through nucleotide assimilation on target DNA (Mitra et al., 2019). The amplified DNA can then be confirmed through electrophoresis on an agarose gel followed by staining with ethidium bromide or Coomassie Blue® to display banding, validating the presence of DTV in the tick sample tested. In this paper we present our preliminary data on the development of a colorimetric LAMP detection of DTV and discuss perspectives of our research for POCT.

Figure 1. General schematic of LAMP proliferation into ladder structures. Each arrow indicates a new step of the process and arrows indicate polymerase writing genetic code. Three prime and five prime ends are indicated on each side of the strands of DNA as well as the LAMP products. Horizontally positioned arrows indicate polymerase activity and amplification. The initial double stranded DNA is representative of the target cDNA that was cloned into a plasmid. F1, F2, and F3 represent the forward primer sequences and are contained within the FIP, F3, and LF primers. F1c, F2c, and F3c represent coding sequences of target DNA. This same principle applies to the B1, B2, B3, B1c, B2c, B3c, BIP and LB sequences in a similar manner to create the end loop product of LAMP.

Materials and Methods

Nucleotide Sequence Analysis and Homology Analysis for DTV Strains

The greatest amount of genetic variance between strains of DTV were estimated through a maximum composite likelihood analysis method. The greatest amount of conservation was determined to be at the RNA-dependent RNA Polymerase (RdRp) coding site using the codon based Z-test. Two bioinformatics methods, i.e., the codon-based Z-test, and the UPGMA phylogenetic tree method, were used to analyze the RNA-dependent RNA polymerase (RdRp) coding region of the DTV genome of four strains and determine nucleotide sequence conservation. These tests were performed using the software suite MEGA7: Molecular Evolutionary Genetic Analysis version 7.0 (Kumar et al., 2015) using DTV accession numbers MG196295.1, EU338403.1, NC003687.1, and HM991145.1 retrieved from GenBank, an open-source genetic sequence database operated by the National Institute of Health (Soloman et al., 2018; Tavakoli et at., 2008, Unpublished Accession, 2010).

Codon Based Z-Test

Alignment tools were utilized to investigate the homology between the conserved sequences of DTV. Positive selection for the resistance to mutation was analyzed through the codon-based Z-test as defined by Equation 1. When analyzing the homology, the null hypothesis is regarded such that the sequences are not similar and only by rejecting the null hypothesis when $p < 0.05$, can we assume that the sequences are considered homologous.

$$Z = \frac{dN - dS}{\sqrt{(\mathrm{Var}(dS) - \mathrm{Var}(dN))}}$$

Equation 1. The Z-test for codon homology testing across multiple DTV genomes. Variables denoted in the equation are defined as the number of nonsynonymous substitutions per nonsynonymous site (dN), number of synonymous substitutions per synonymous site (dS), and the respective variances of these variables (Var(dN) and Var(dS)).

Primer Design

Primers for the LAMP assay were designed through the PrimerExplorerV5 LAMP Primer Designer using the following criteria: (1) genomic regions with a GC ratio of 50-60%, (2) no more than four successive G's or C's to prevent primers from binding too tightly to the target DNA strand, and (3) stability of F3 and B3 primer ends be lower than -4.00 to create the most stable 5' and 3' loop structures for LAMP proliferation (Wang et al., 2015). The region of the viral genome that was selected for primer design was NS5a that codes the non-structural polymerase.

Primer Constructs

Below are the two primer constructs evaluated in this paper.

Primer	Sequence (5' to 3')	Len.
F3	GTACCTCGACTTGCCAACC	19
B3	TGCGGGTAGGAAGTAGTGC	19
FIP	ATGTCCCGGAAAGGTTTGGACA-TACGACCACTCACCTCTGC	41
BIP	TGGGCATTCCTACACCCGCT-ACACTCGTCTTCCCGTGAA	39
LF	CTACCCAAACTGCTGCGTC	19
LB	AGTCACCCCAAGTGTCGGAC	20

Table 1. Primer candidate 1 sequences generated using the PrimerExplorerV5 software with respective loop primers (LF and LB).

Primer	Sequence (5' to 3')	Len.
F3	TAAGGATGTGGGCGAGTGG	19
B3	TGGAAGTAGTTGAGGAGCCA	20
FIP	GAACGCCCATCCTTCATCACGA-TCACAGCCTGGGAAGAAGT	41
BIP	ACGAGTGTCGCCAGGATGTG-CATACGCTTTGGACAGGCAG	40
LF	TCATGGAAATGGTGTGAGCAGAAGG	25
LB	ATGGAGCGTGCGGGAGACA	19

Table 2. Primer candidate 2.0 generated using the PrimerExplorerV5 software with respective loop primers (LF and LB).

Figure 2. General reaction scheme for a colorimetric LAMP assay. Reagents are listed below first reaction vessel on left and diverge based on using either 1 µL of sample or water control. The results from this procedure include the stagnant pink color for the negative control, and either a color change to yellow, indicating a positive result, or remain pink indicating a negative result for the experimental DNA.

LAMP primers were determined using the PrimerExplorerV5 software and the NC003687.1 accession (Notomi et al., 2000). The following primers have been designed using the PrimerExplorerV5 software and have been experimentally tested in the laboratory. Primer candidates 1 and 2 are displayed with annealing temperatures and dimerization rates in Tables 1 and 2, respectively. Four primers are designed within the first iteration of software for binding to a target sequence. Figure 1 depicts the amplification process that occurs to create the loop products generated by *Bst* polymerase. The inner primers denoted Forward Inner Primer (FIP) and Backward Inner Primer (BIP) are the central component of amplification and loop structure formation. Two additional primers F3 and B3, complementary to F3c and B3c respectively, can bind to the target DNA and displace the amplification product. This allows a repetitive, rapid amplification process. The final two primers, Loop Front (LF) and Loop Back (LB) are not pictured in Figure 1 but are able to further amplify and accelerate the loop creation process by providing templates. The reaction is still possible without LF and LB, but a greater limit of detection can be achieved using loop primers. The constructs for LF and LB were generated by inserting the initial primer file into the PrimerExplorerV5 software and following the same protocol.

Loop-Mediated Isothermal Amplification

For the purposes of this experimentation, the optimal temperature for primer candidate 1 was determined to be 59.3℃ and the optimal time was twenty minutes. The Colorimetric Warmstart® MasterMix contains nucleotides, *Bst* polymerase, 16.0 mM Magnesium Sulfate, and pH sensitive indicator phenol red. The general reaction scheme for colorimetric based LAMP assays is seen in Figure 2. After the addition of genetic material and molecular grade water, the reaction will proceed in either a hot water bath or a thermocycler held at constant temperature. Results are visualized by the naked eye to be pink (negative) or yellow (positive).

	Sample Vessel	Control
Warmstart® Colorimetric LAMP 2X Master Mix	12.5 µL	12.5 µL
LAMP Primer Mix 10X	2.5 µL	2.5 µL
Target Genetic Sample	1 µL	N/A
Molecular Grade H₂O	9 µL	10 µL

Table 3. Reagents added to reaction vessels to perform the colorimetric LAMP assay.

Transformation and Purification of Plasmid cDNA

A plasmid was created (GenScript) containing a 500 bp cDNA of DTV's RdRp, with flanking *EcoRI* and *XhoI* restriction enzyme sites. The plasmid was transformed into DH5α chemically competent *E. coli* for 2 minutes at 37°C and plated on nutrient agar plates containing ampicillin. DNA stocks were harvested from single clones by plasmid miniprep kits (Qiagen). The desired cDNA was digested from the vector by EcoRI and XhoI, run on an agarose gel, and purified from the gel with a gel extraction kit (Qiagen). The cDNA from this purification process was used for the experimentation as the DTV RdRp coding strand sample.

Discussion

From the analysis methods, we concluded that these sequences are statistically significant ($p < 0.05$) to move forward with the RdRp targeted primer design. A 2,000 base-pair sequence was chosen from the DTV NS5a coding region GenBank accession NC003687.1, which includes the coding region for the RdRp. The codon-based test resulted in rejecting the null hypothesis for comparing the four different accessions, with $p < 0.05$ for each entry. An evolutionary analysis showed the branching of four accessions from GenBank including one complete genome and three RdRp sequences and is visualized in Figure 3. The maximum divergence was 19.22% across all accessions, and primer development should represent the ideal areas of conservation. Alignment was performed against the full genome to match the RdRp sequences.

Primer candidate 1 displayed potential and consistency within LAMP assays. Dimerization events were unlikely to occur and provided consistent results. In Figure 4 typical results of this primer set are displayed. When LAMP is run with these primers, a colorimetric change is visualized, and the corresponding products are visualized following electrophoresis on an agarose gel. Controls of nonspecific DNA and water provide negative results as expected. These are desirable traits with LAMP primers to avoid false positives or false negatives. Additional primer sets were designed to confirm our design strategies. The optimal temperature at which this primer candidate function was 59.3ºC (Figure 4.D).

The second primer set displayed dimerization concerns after looking closely at the dimerization rate of the 5' end of the F3 primer. This observation was denoted after consistent false positives were obtained with primer candidate 2. The instability of these primers was directly correlated to the instability of the F3 primer being more positive than dG = -4.00 (Table 2). This observation can be regarded as a learning guide for future primer design experimentation and allowed for a consistency check with the first iteration generated.

Figure 3. Evolutionary Relationships of Powassan Virus Lineage II, DTV in the United States.[2,3] The evolutionary history was inferred using the UPGMA method [14]. The optimal tree with the sum of branch length = 0.19223306 is shown. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The evolutionary distances were computed using the Maximum Composite Likelihood method [15] and are in the units of the number of base substitutions per site. This analysis involved 4 nucleotide sequences accessed through GenBank. Codon positions included were 1st+2nd+3rd+Noncoding. All ambiguous positions were removed for each sequence pair (pairwise deletion option). There was a total of 10840 positions in the final dataset. Evolutionary analyses were conducted in MEGA X [16]

Table 4. Codon-based Test of Neutrality for analysis between sequences. The probability of rejecting the null hypothesis of strict-neutrality (dN = dS) (below diagonal) is shown. Values of P less than 0.05 are considered significant at the 5% level and are highlighted. The test statistic (dN - dS) is shown above the diagonal. dS and dN are the numbers of synonymous and nonsynonymous substitutions per site, respectively. The variance of the difference was computed using the analytical method. Analyses were conducted using the Nei-Gojobori method [17]. This analysis involved 4 nucleotide sequences. All ambiguous positions were removed for each sequence pair (pairwise deletion option). There was a total of 3556 positions in the final dataset. Evolutionary analyses were conducted in MEGA X [16]

	EU338403.1	MG196295.1	NC003687.1	HM991145.1
EU338403.1	---	-2.5830	-3.7536	4.4586
MG196295.1	0.0110	---	-15.7194	-3.2327
NC003687.1	0.0003	0.0000	---	-3.2986
HM991145.1	0.0000	0.0016	0.0013	---

Figure 4. (A) Colorimetric reaction vessels for Primer 1.0 with the following reaction vessel containments: 1 – Primer w/DNA, 2 – Primer w/DNA, 3 – Primer w/DNA, 4 – Primer w/o DNA, 5 – Primer w/o DNA, 6 – No Primer/DNA, 7 – No Primer/DNA. **(B) Ethidium bromide electrophoresis gel with samples of panel (A).** The containments of each lane correspond with the reaction vessels as indicated. **(C) Temperature gradient assay for LAMP primer set 1.0.** Each of the reaction vessels (left) were also run with negative controls, not pictured. Reaction vessel numbers correspond with the ethidium bromide gel pictured in box (D), with the exception of vessel (1) and (8). From vessel (1) to (8), the temperature gradient follows the order (in ºC): 70.8, 69.3, 67.8, 64.8, 59.3, 57.8, 56.3, and 54.8.

Future primer sets will be compared to these initial two primer candidates and put through a comparative analysis for accuracy and limit of detection. An additional primer candidate has already been synthesized and experimentation shows additional promise. The efforts of this research are to bring attention to this pathosystem and the lack of robust diagnostic testing for downstream health effects.

DTV is a recent pathogen of interest and needs additional detection methods to monitor the long-term effects and spread. In the creation of a LAMP detection system point-of-care and field diagnostic, additional tools can be enacted to understand the spread of this virus. Primer 1.0 was an excellent candidate for the detection of *in vitro* genetic material of the virus. Our second primer candidate was faulty but further improved primer design hypotheses and tactics. Further validation is currently being investigated for newer primer sets. Additional experimentation will be performed with RNA expression samples, tick samples spiked with DNA and RNA, and additional collaborations with state environmental agencies for tick testing. Through these efforts we hope to further promote mitigation tactics and reduce the number of infections in North America.

Acknowledgments

We would like to thank the Lebanon Valley College Biology Department for financially supporting this research, as well as the Beta Beta Beta Biological Honors Society National Research Grant for the 2019-2020 grant received for this project.

Conflict of Interests

The authors hereby state no conflict of interests.

References

[1] Anupama, K.P., Chakraborty, A., Karunasagar, Iddya, Karunasagar, Indrani, Maiti, B., 2019. Loop-mediated isothermal amplification assay as a point-of-care diagnostic tool for Vibrio parahaemolyticus: recent developments and improvements. Expert Review of Molecular Diagnostics 19, 229–239. https://doi.org/10.1080/14737159.2019.1571913

[2] Artsob, H., Karabatsos, N., Kuno, G., Tsuchiya, K.R., Chang, G.J., 2001. Genomic sequencing of deer tick virus and phylogeny of powassan-related viruses of North America. The American Journal of Tropical Medicine and Hygiene 65, 671–676. https://doi.org/10.4269/ajtmh.2001.65.671

[3] Brackney, D.E., Brown, I.K., Nofchissey, R.A., Fitzpatrick, K.A., Ebel, G.D., 2010. Homogeneity of Powassan virus populations in naturally infected Ixodes scapularis. Virology 402, 366–371. https://doi.org/10.1016/j.virol.2010.03.035

[4] CDC, 2019. Powassan Virus (National Center for Emerging and Zoonotic Infectious Diseases), Division of Vector-Borne Diseases. Center for Disease Control.

[5] Chen, Q., Yuan, L., Wan, J., Chen, Y., Du, C., 2014. Colorimetric detection of hepatitis E virus based on reverse transcription loop mediated isothermal amplification (RT-LAMP) assay. Journal of Virological Methods 197, 29–33. https://doi.org/10.1016/j.jviromet.2013.11.006

[6] Lamb, L.E., Bartolone, S.N., Tree, M.O., Conway, M.J., Rossignol, J., Smith, C.P., Chancellor, M.B., 2018. Rapid Detection of Zika Virus in Urine Samples and Infected Mosquitos by Reverse Transcription-Loop-Mediated Isothermal Amplification. Scientific Reports 8. https://doi.org/10.1038/s41598-018-22102-5

[7] Lau, Y.-L., Lai, M.-Y., Teoh, B.-T., Abd-Jamil, J., Johari, J., Sam, S.-S., Tan, K.-K., AbuBakar, S., 2015. Colorimetric Detection of Dengue by Single Tube Reverse-Transcription-Loop-Mediated Isothermal Amplification. PLOS ONE 10, e0138694. https://doi.org/10.1371/journal.pone.0138694

[8] Luo, L., Nie, K., Yang, M.-J., Wang, M., Li, J., Zhang, C., Liu, H.-T., Ma, X.-J., 2011. Visual Detection of High-Risk Human Papillomavirus Genotypes 16, 18, 45, 52, and 58 by Loop-Mediated Isothermal Amplification with Hydroxynaphthol Blue Dye. Journal of Clinical Microbiology 49, 3545–3550. https://doi.org/10.1128/JCM.00930-11

[9] Mitra, D., Kimov, I.K., Waldeisen, J.R., 2019. Colorimetric Detection of Nucleic Acid Amplification. 10,253,357B2.

[10] Morozova, O.V., Dobrotvorsky, A.K., Livanova, N.N., Tkachev, S.E., Bakhvalova, V.N., Beklemishev, A.B., Cabello, F.C., 2002. PCR Detection of Borrelia burgdorferi Sensu Lato, Tick-Borne Encephalitis Virus, and the Human Granulocytic Ehrlichiosis Agent in Ixodes persulcatus Ticks from Western Siberia, Russia. Journal of Clinical Microbiology 40, 3802–3804. https://doi.org/10.1128/JCM.40.10.3802-3804.2002

[11] Nagamine, K., Hase, T., Notomi, T., 2002. Accelerated reaction by loop-mediated isothermal amplification using loop primers. Molecular and Cellular Probes 16, 223–229. https://doi.org/10.1006/mcpr.2002.0415

[12] Pesko, K.N., Torres-Perez, F., Hjelle, B.L., Ebel, G.D., 2010. Molecular epidemiology of Powassan virus in North America. Journal of General Virology 91, 2698–2705. https://doi.org/10.1099/vir.0.024232-0

[13] Shin, D.-S., Heo, G.-I., Son, S.-H., Oh, C.-S., Lee, Y.-K., Cha, J.-S., 2018. Development of an Improved Loop-Mediated Isothermal Amplification Assay for On-Site Diagnosis of Fire Blight in Apple and Pear. The Plant Pathology Journal 34, 191–198. https://doi.org/10.5423/PPJ.FT.03.2018.0055

[14] Solomon, I.H., Spera, K.M., Ryan, S.L., Helgager, J., Andrici, J., Zaki, S.R., Vaitkevicius, H., Leon, K.E., Wilson, M.R., DeRisi, J.L., Koo, S., Smirnakis, S.M., De Girolami, U., 2018. Fatal Powassan Encephalitis (Deer Tick Virus, Lineage II) in a Patient With Fever and Orchitis Receiving Rituximab. JAMA Neurol 75, 746. https://doi.org/10.1001/jamaneurol.2018.0132

[15] Tavakoli, N.P., Wang, H., Dupuis, M., Hull, R., Ebel, G.D., Gilmore, E.J., Faust, P.L., 2009. Fatal Case of Deer Tick Virus Encephalitis. New England Journal of Medicine 360, 2099–2107. https://doi.org/10.1056/NEJMoa0806326

[16] Telford III, S., 1997. A New Tick-borne Encephalitis-like Virus Infecting New England Deer Ticks, Ixodes dammini. Emerg. Infect. Dis. 3, 165–170. https://doi.org/10.3201/eid0302.970209

[17] Tomlinson, J., 2008. Potential of LAMP for detection of plant pathogens. CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources 3. https://doi.org/10.1079/PAVSNNR20083066

[18] Wang, D.-G., Brewster, J., Paul, M., Tomasula, P., 2015. Two Methods for Increased Specificity and Sensitivity in Loop-Mediated Isothermal Amplification. Molecules 20, 6048–6059. https://doi.org/10.3390/molecules20046048