SARS-CoV-2 induces inflammasome-dependent pyroptosis and downmodulation of HLA-DR in human monocytes

André C. Ferreira[1,2,10#*,] Vinicius Cardoso Soares[1,9#], Isaclaudia G. de Azevedo-Quintanilha[1], Suelen da Silva Gomes Dias[1], Natalia Fintelman-Rodrigues[1,10], Carolina Q. Sacramento[1,10], Mayara Mattos[1,10], Caroline S. de Freitas[1,10], Jairo R. Temerozo[3,4], Lívia Teixeira[1], Eugenio Damaceno Hottz[1,5], Ester A Barreto[1], Camila R. R. Pão[1], Lohanna Palhinha[1], Milene Miranda[6], Dumith Chequer Bou-Habib[3,4], Fernando A. Bozza[7,8], Patrícia T. Bozza[1] , Thiago Moreno L. Souza[1,10*]

– Laboratório de Imunofarmacologia, Instituto Oswaldo Cruz (IOC), Fundação Oswaldo Cruz (Fiocruz), Rio de Janeiro, RJ, Brazil.

– Laboratório de Pesquisa Pré-clínica - Universidade Iguaçu, Nova Iguaçu, RJ, Brazil.

– Laboratório de Pesquisas sobre o Timo, IOC, Fiocruz, Rio de Janeiro, RJ, Brazil.

– National Institute for Science and Technology on Neuroimmunomodulation, Oswaldo Cruz Institute, Fiocruz, Rio de Janeiro, RJ, Brazil.

– Laboratório de Imunotrombose, Departamento de Bioquímica, Universidade Federal de Juiz de Fora, Juiz de Fora, MG, Brazil.

– Laboratório de Vírus Respiratório e do Sarampo, IOC, Fiocruz, Rio de Janeiro, RJ, Brazil.

– Instituto Nacional de Infectologia Evandro Chagas, Fiocruz, Rio de Janeiro, RJ, Brazil.

– Instituto D'or de Pesquisa e Ensino, Rio de Janeiro, RJ, Brazil.

- Program of Immunology and Inflammation, Federal University of Rio de Janeiro, UFRJ, Rio de Janeiro, RJ, Brazil.

- Centro De Desenvolvimento Tecnológico Em Saúde (CDTS), Fiocruz, Rio de Janeiro, Brasil.

# These authors contributed equally to this work

* corresponding author

Abstract

Infection by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has been associated with leukopenia and uncontrolled inflammatory response in critically ill patients. A better comprehension of SARS-CoV-2-induced monocyte death is essential for the identification of therapies capable to control the hyper-inflammation and reduce viral replication in patients with COVID-19. Here, we show that SARS-CoV-2 induces inflammasome activation and cell death by pyroptosis in human monocytes, experimentally infected and from patients under intensive care. Pyroptosis was dependent on caspase-1 engagement, prior to IL-1ß production and inflammatory cell death. Monocytes exposed to SARS-CoV-2 downregulate HLA-DR, suggesting a potential limitation to orchestrate the immune response. Our results originally describe mechanisms by which monocytes, a central cellular component recruited from peripheral blood to respiratory tract, succumb to control severe 2019 coronavirus disease (COVID-19).

Author summary

Since its emergence in China in late 2019, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has caused thousands of deaths worldwide. Currently, the number of individuals infected with SARS-CoV-2 and in need of antiviral, anti-inflammatory, anticoagulant and more invasive treatments has overwhelmed the health systems worldwide. In our study, we found that SARS-CoV-2 is capable of inducing inflammatory cell death in human monocytes, one of the main cell types responsible for anti-SARS-CoV-2 immune response. As a consequence of this intracellular inflammatory mechanism (inflammasome engagement), an exacerbated production of inflammatory mediators occurs. The infection also decreases the expression of HLA-DR in monocytes, a molecule related to the orchestration of the immune response in case of viral infections. We also demonstrated that the HIV-1 protease inhibitor, atazanavir (ATV), prevented the uncontrolled inflammatory response, cell death and reduction in HLA-DR expression in SARS-CoV-2-infected monocytes. Our study provides relevant information on the effects of SARS-CoV-2 infection on human monocytes, as well as on the effect of ATV in preventing these pathological effects on the host.

Introduction

Severe acute respiratory coronavirus 2 (SARS-CoV-2), the etiological agent of the 2019 coronavirus disease (COVID-19), emerged in China, causing a major public health burden in decades. Patients with COVID-19 may develop an asymptomatic or mild disease or be affected by the life threatening acute respiratory distress syndrome (ARDS), which is characterized by elevated serum levels of proinflammatory mediators, the cytokine storm (1–3). In the respiratory tract of patients with severe COVID-19, monocytes/macrophages may be the main source of uncontrolled levels of the pro-inflammatory mediators TNF-α and IL-6 (4). Plasmatic levels of IL-6 have been associated with mortality, intensive care admission and hospitalization, representing a poor prognostic factor for COVID-19 (5). The uncontrolled inflammation promoted by SARS-CoV-2 in severe COVID-19 is not acute, it negatively associates with viral loads in nasopharyngeal swabs and represents an important event from 7 to 10 days after onset of illness (6,7). The SARS-CoV-2-induced cytokine storm associates with intense cell death (8,9).

There are various mechanisms involved in cell death, which are differently engaged from development to responses to infection (10). For certain diseases, such as COVID-19, in which the immunopathogenesis mechanisms associate with poor clinical outcomes, controlling the way cells collapse to infection is vital for the host (10). For example, cell death from necrosis, which can occur from necroptosis to pyroptosis, tends to exacerbate inflammation due to the rupture of the cellular plasma membrane. Pyroptosis, in particular, begins with the activation of the inflammasome, an intracellular structure that involves several intracellular molecules such as caspase-1(10,11), leads to leakage of the cytoplasmic content, favors inflammatory infiltrate (11) and amplifies of the inflammatory response (12). These mechanisms of cell death contrasts with apoptosis, a more controlled process that maintain the host's homeostasis.

In severe COVID-19, the cytokine storm associates with high levels of tissue insult, judged by increased levels lactate dehydrogenase (LDH) and D-dimer in the plasma (6–8). Moreover, high LDH levels and leukopenia in severe COVID-19 points out that white cells loses the integrity of plasma membrane (6–8). Among these cells,

monocytes should orchestrate the equilibrium between innate and adaptive immune responses, which may be presumably affected during cytokine storm. Indeed, severe immune dysregulation with low antigen presenting capacity is evidenced by reduced expression of HLA-DR in monocytes during COVID-19, which is strongly associated with severe respiratory failure (4).

The leukopenia of patients with severe COVID-19 seem to precede the cytokine storm (13–18). Moreover, in other virus-induced cytokine storm episodes in the respiratory tract, such as induced by influenza A virus, monocytes and macrophages are severely affected (13). Thus, we hypothesized that the monocyte cell death induced by SARS-COV-2 exacerbates the production of inflammatory cytokines, as well as impairs the immune balance in the hosts. In fact, we found that SARS-CoV-2 triggers the activation of the inflammasome, leading to pyroptosis in human monocytes, by experimental or natural infection. Pyroptosis was dependent on caspase-1 engagement, prior to IL-1ß production and dysregulation of cytokine release. Monocytes that survive after SARS-CoV-2 challenge downregulate HLA-DR, being less likely to properly orchestrate the immune response. Finally, we show that the reproposed antiviral drug atazanavir (ATV) could block this deleterious loop in favor of the host.

Material and Methods

Reagents

Atazanavir (ATV) and Ribavirin were received as donations from Instituto de Tecnologia de Fármacos (Farmanguinhos, Fiocruz). The antiviral Lopinavir/ritonavir (4:1 proportion) was pruchased from AbbVie (Ludwingshafen, Germany). ELISA assays were purchased from R&D Bioscience. Lipopolysacchadides - LPS, adenosine triphosphate (ATP), the specific inhibitor of caspase-1 (AC-YVAD-CMK), pan-caspase inhibitor (ZVAD-FMK), RIPK1 (Necrostatin-1 – Nec-1) and IL-1 receptor (IL-1RA) were all purchased from Sigma-Aldrich (St. Louis, MO, USA). All small molecule inhibitors were dissolved in 100 % dimethylsulfoxide (DMSO) and subsequently diluted at least 10^4-fold in culture or reaction medium before each assay. The final DMSO concentrations showed no cytotoxicity. The materials for cell culture were purchased from Thermo Scientific Life Sciences (Grand Island, NY), unless otherwise mentioned.

Cells and Virus

African green monkey kidney (Vero, subtype E6) cells were cultured in DMEM
high glucose supplemented with 10 % fetal bovine serum (FBS; HyClone, Logan, Utah)
and 100 U/mL penicillin, and 100 μg/mL streptomycin (P/S). Vero cells were incubated
at 37°C in 5 % CO_2 atmosphere.

Human primary monocytes were obtained through plastic adherence of peripheral
blood mononuclear cells (PBMCs), which were obtained from buffy coat preparations of
healthy donors by density gradient centrifugation (Ficoll-Paque, GE Healthcare). In brief,
PBMCs (2.0 x 10^6 cells) were plated onto 48-well plates (NalgeNunc) in RPMI-1640
without serum for 2 to 4 h; then, non-adherent cells were removed by washing and the
remaining monocytes were maintained in DMEM with 5% human serum (HS; Millipore)
and penicillin/streptomycin.

SARS-CoV-2 was isolated and expanded on Vero E6 cells from a nasopharyngeal
swab of a confirmed case from Rio de Janeiro, Brazil. Experiments were performed after
one passage in cell culture, when Vero E6 cells with DMEM plus 2% FBS in 150 cm^2
flasks were incubated at 37 °C in 5 % CO_2 atmosphere. Cytopathic effect was observed
daily and peaked 4 to 5 days after infection. All procedures related to virus culture were
handled at biosafety level 3 (BSL3) multiuser facility, according to WHO guidelines.
Virus titers were determined as the tissue culture infectious dose at 50% (TCID50/mL).
Virus stocks were kept in -80 °C ultralow freezers. The virus strain was sequenced to
confirm the virus identity and its complete genome is publicly deposited (GenBank
accession no. MT710714).

Yield-reduction assay

Human primary monocytes were infected with multiplicity of infection (MOI) of
0.01 at density of 2-8 x 10^5 cells/well in 48-well culture plates, depending on the total cell
number from each donor. After 1 h at 37 °C, cells were washed, and various
concentrations of compounds were added in DMEM with 2% FBS. After 48 h, the

supernatants were harvested and virus replication was quantified by real time RT- PCR

and infectious titers by TCID50/mL.

Virus titration

Monolayers of Vero cells (2×10^4 cell/well) in 96-well culture plates were infected

with log- based dilutions of the supernatants containing SARS-CoV-2 for 1 h at 37°C.

The cells were washed and fresh medium with 2% FBS was added. After 3 to 5 days, the

cytopathic effects were scored in at least 10 replicates per dilution by independent readers,

who were blind with respect to source of the supernatant. Reed and Muench scoring

method was employed to determine TCID50/mL.

Flow cytometer analysis

For flow cytometry analysis, monocytes were diluted in labelling buffer (10^6

cells/mL). Then, 100 µL of cell samples were marked with 5 µL of AnnexinV and PI for

15 minutes for cell death evaluation. Around 10,000 gated events were acquired using

FACSCalibur and the analysis was performed using the CellQuest software. Monocytes

were gated through cell size (foward light scatter, FSC) and granularity (side light scatter,

SSC) analysis.

Human monocytes were stained for caspase-1 activity with FAM-YVAD-FMK

(fluorescent-labeled inhibitor of caspase-1 [FLICA] and FAM-FLICA Caspase-3/7

activity or HLA-DR APC.H7 or IgG APC.H7. Caspase-1 and caspase-3/7 activity was

determined via flow cytometry (FACSCalibur) by detecting FLICA fluorescence and

expression of HLA-DR as mean fluorescence intensity (MFI) value for each sample.

Acquisition of data was set to count a total of 10,000 events, and the FLOWJO software

package was used to analyze the data.

Microscopic analysis

Human primary monocytes were plated on glass coverslips at density of $2\text{-}8 \times 10^5$

cells/well in 48-well plates. Infection was performed for 2 h at 37 °C and then fresh

medium with 2% FBS was added. After 24 h, the cells were washed with Binding buffer

and stained with PI (0.5 µg/mL) for 5 minutes. Next, the cells were fixed with 3.7%

formaldehyde for 30 minutes at room temperature. The nuclei were stained with DAPI

(1μg/mL) for 5 min and the coverslips were mounted using an antifade mounting medium

(VECTASHIELD). Fluorescence was analyzed by fluorescence microscopy with an x100

objective lens (Olympus, Tokyo, Japan).

Measurements Inflammatory Mediators and cell death marker

The levels of IL-1ß, TNF-α, IL-6, IL-8 and LDH were quantified in the culture

supernatants from infected and uninfected monocytes using ELISA kits, according to the

manufacturer's intructions (R&D System).

Extracellular lactate dehydrogenase (LDH) was quantified using Doles® kit

according to manufacturer's instructions. In brief, cell culture supernatants were

centrifuged at 5,000 rpm for 1 minute, to remove cellular debris, and then 25 μL were

placed into 96-well plates and incubated with 5 μL of ferric alum and 100 μL of LDH

substrate for 3 minutes at 37 °C. Nicotinamide adenine dinucleotide (NAD, oxidized

form) was added followed by the addition of a stabilizing solution. After 10 minutes, the

reaction was read in a spectrophotometer at 492 nm.

Western blot assay

Cellular extracts of $1x10^6$ cells were homogenized in the RIPA lysis buffer in the

presence of proteinase inhibitor cocktail (Roche), and the protein levels were measured

by BCA protein assay kit. A total of 20 μg of protein was loaded onto a 10% sodium

dodecyl sulfate polyacrylamide gel (SDS-PAGE) for separation by electrophoresis and

the protein bands were then transferred to a polyvinylidene difluoride membranes

(ImmobilonP-SQ, Millipore). The membranes were blocked with 5 % albumin diluted in

Tris-buffered saline containing 0.05 % of Tween 20 for 2 hours at room temperature and

incubated with the specific primary antibodies (Cell Signaling Technology), to detect pro-

caspase-1 and cleaved-caspase-1, after overnight incubation at 4 °C. After washing,

membranes were incubated with secondary antibodies (IRDye® 800CW Goat-anti-

Mouse and IRDye® 680LT Goat anti-Rabbit IgG Antibody, LI-COR, Lincoln) for 30

minutes at RT. The protein bands were visualized by digital fluorescence

(Odyssey® CLx Imaging System), and protein density was analyzed by the ImageJ

software. All the data were normalized by β-actin expression quantification.

Human subjects.

We prospectively enrolled severe COVID19 RT-PCR-confirmed cases, as well as

SARS-CoV-2-negative healthy controls. Blood samples were obtained from 12 patients

with severe COVID-19 within 72 hours from intensive care unit (ICU) admission in two

reference centers (Instituto Estadual do Cérebro Paulo Niemeyer and Hospital Copa Star,

Rio de Janeiro, Brazil). Severe COVID-19 was defined as critically ill patients presenting

viral pneumonia on computed tomography scan and in mechanical ventilation. All

patients had SARSCoV-2 confirmed diagnostic through RT-PCR of nasal swab or

tracheal aspirates. Peripheral vein blood was also collected from 8 SARS-CoV-2-negative

healthy control participants as tested by RT-PCR on the day of blood sampling. The

characteristics of severe (n = 12), and control (n = 8) participants are presented in Table

1. Severe COVID-19 patients usually present older age and higher prevalence of

comorbidities as obesity, cardiovascular diseases and diabetes as in previously reported

patient cohorts. In the present study, the SARS-CoV-2-negative control group was

designed to include subjects of older age and chronic non-communicable diseases, so it

is matched with critically ill COVID-19 patients (Table 1). Patients with acute respiratory

distress syndrome (ARDS) were managed with neuromuscular blockade and a protective

ventilation strategy that included low tidal volume (6 mL/kg of predicted body weight)

and limited driving pressure (less than 16 cmH2O) as well as optimal PEEP calculated

based on the best lung compliance and PaO_2/FiO_2 ratio. In those with severe ARDS and

PaO_2/FiO_2 ratio below 150 despite optimal ventilatory settings, prone position was

initiated. Our management protocol included antithrombotic prophylaxis with enoxaparin

40 to 60 mg per day. Patients did not receive routine steroids, antivirals or other anti-

inflammatory or anti-platelet drugs. The SARS-CoV-2- negative control participants

were not under anti-inflammatory or anti-platelet drugs for at least two weeks. All clinical

information were prospectively collected using a standardized form - ISARIC/WHO

Clinical Characterization Protocol for Severe Emerging Infections (CCPBR).

Ethics statement.

Experimental procedures involving human cells from healthy donors were

performed with samples obtained after written informed consent and were approved by

the Institutional Review Board (IRB) of the Oswaldo Cruz Foundation/Fiocruz (Rio de

Janeiro, RJ, Brazil) under the number 397-07. The National Review Board approved the

study protocol (CONEP 30650420.4.1001.0008), and informed consent was obtained

from all participants or patients' representatives.

Statistical analysis

The assays were performed in blinded way. They were performed by one

professional, codified and read by another fellow. All experiments were carried out at

least three independent times, including a minimum of two technical replicates in each

assay. The equations to fit the best curve were generated based on R^2 values ≥ 0.9.

Student's T-test was used to access statistically significant P values <0.05.

Results

SARS-CoV-2 promotes pyroptosis in human monocytes

To characterize the mechanism by which SARS-CoV-2 triggers monocyte death,

human primary monocytes were infected with SARS-CoV-2 or treated with LPS+ATP.

Next, cell death was analyzed by quantifying the LDH activity in the culture supernatant,

and by AnnexinV/propidium iodide (PI) cell labeling through for flow cytometry and

fluorescence microscopy. As shown in Figure 1A, SARS-CoV-2 increased LDH levels

similarly to the positive control, LPS+ATP (Figure 1A). Flow cytometry and fluorescence

microscopy images demonstrated higher percentages of PI$^+$ cells in the infected cultures,

as well as a significant increase in mean fluorescence intensity (MFI). Both, LDH leak

and PI labeling characterize cell membrane disruption in human monocytes after SARS-

CoV-2 infection (Figure 1B-E). These data suggest that SARS-CoV-2 infection can

induce cell death characterized by loss of plasma membrane integrity suggestive of

pyroptosis cell death.

SARS-CoV-2 induces pyroptosis in monocytes through inflammasome activation

To gain insights on the mechanisms of monocyte cell death in SARS-CoV-2

infection, we accessed caspase-1 activation - a key events that require inflammasome's

proteolytic activity. Cells stimulated with LPS+ATP were used as a positive control of

inflammasome activation. We observed that SARS-CoV-2 induced pro-caspase-1

cleavage, similarly to positive control (Figure 2A and 2B). To confirm these results, cells

were labeled with FAM-YVAD-FLICA (as an indicative of caspase-1 activation) and

analyzed by flow cytometry. We found that SARS-CoV-2 infection induced the
activation of caspase-1 in monocytes in the same magnitude to the positive control group
(Figure 2C and 2D). Moreover, SARS-CoV-2-induced caspase-1 activation was a
specific event, since we did not observe activation of the apoptotic caspases-3 and -7 in
infected monocytes (Figure 2E and 2F). As caspase-1 activity is critical for the production
of IL-1β (19,20), we measured the levels of this cytokine in our experiments. Our data
show that SARS-CoV-2 infection was able to increase IL-1β production, this
phenomenon being prevented with pretreatment with AC-YVAD-CMK, and not altered
with a necroptosis inhibitor. (Figure 2G). Importantly, inhibition of IL-1R engagement
reduced SARS-CoV-2-mediated caspase-1 activation and LDH release (Figure S1),
suggesting that inflammasome-dependent IL-1β secretion amplify caspase-1 activation
and pyroptosis in SARS-CoV-2 infection.

Inflammasome activation amplify pro-inflammatory cytokines secretion in infected
monocytes

To evaluate the effect of the activation of inflammasome and pyroptosis on the
immune response, we quantified the production of key cytokines in the amplification of
the inflammatory process observed clinically in a patient during the course of the
infection with IL-6 and TNF-α. Remarkably, caspase-1 specific inhibition, but not pan-
caspase or RIPK1 inhibitor, also led to lower levels of the pro-inflammatory cytokines
IL-6 and TNF-α, highlighting the participation of the caspase-1-IL-1β axis in
inflammatory amplification during SARS-CoV-2 infection. (Figure 3A and 3B). To
confirm the effects in SARS-CoV-2 infection in the immune response, whether the
observed effect is specifically related to inflammasome activation, we evaluated the
production of IL-8, in which induction is independent of the inflammasomes pathway.
SARS-CoV-2 infection induced a higher IL-8 production, which was not prevented by
caspase-1 inhibition (Figure 3C). In addition, inhibition of IL-1R engagement reduced
SARS-CoV-2-mediated was able to significantly decrease the production of IL-1 β, IL-
6, TNF-α and did not alter the production of IL-8 (Figure S2).

Infection with SARS-CoV-2 decreases the expression of HLA class II in human monocytes independently of the activation of inflammasomes

To investigate the impact of inflammasome activation and cell death through pyroptosis on the orchestration of anti-SARS-CoV-2 immune response, cells were then marked with HLA-DR and analyzed by flow cytometry. We found that SARS-CoV-2 infection induced a decrease in HLA-DR expression in monocytes (Figure 4A and 4B). However, the inhibition of caspase-1 (Figure 4A and 4B) and IL-1R engagement (Figure S3) did not prevent the SARS-CoV-2-mediated decreasing of HLA expression. These results suggest that even in disassociation with the engagement of SARS-CoV-2-induced inflammasome activation, monocytes exposed to the new CoV will present limited ability to orchestrate the immune response.

Atazanavir prevented inflammasome-dependent pyroptosis

Given our findings that pyroptosis and decreased HLA-DR expression are triggered during SARS-CoV-2 replication, we tested whether atazanavir (ATV), an HIV protease inhibitor recently shown to possess antiviral activity against the new CoV in monocytes (17), could prevent these events. Flow cytometry analysis of SARS-CoV-2-infected monocytes treated with ATV demonstrated a significant reduction in caspase-1 activity, observed by the decreased FAM-YVAD-FLICA labeling (Figure 5A and 5B). Other orally available repurposed anti-SARS-CoV-2 drugs, such as lopinavir (LPV) and ribavirin (RBV), did not affect SARS-CoV-2-induced pyroptosis (Figure 5C). Moreover, treatments with ATV did not alter the activity of caspase-1, -3 and -7 in monocytes exposed to LPS and ATP, used as a positive control of pyroptosis (Figure S4), indicating a specific antiviral activity of this drug. Consistently, treatment with ATV also reduced the levels of IL-1β, IL-6 and TNF-α in SARS-CoV-2-infected monocytes, when compared to the untreated cells (Figure 5D-F). ATV did not interfere with the production of IL-8, which is independent of inflammasome engagement (Figure 5G).

We then confirmed whether ATV can prevent SARS-CoV-2-induced pyroptosis. Lower levels of LDH were detected in the supernatants of SARS-CoV-2-infected monocytes treated with ATV, when compared to infected and untreated cells (Figure 5H). The treatment with ATV was also able to prevent the decrease of HLA-DR expression in infected monocytes (Figure 5I and J). LPV and RBV did not alter the HLA-DR expression

in monocytes infected with SARS-Cov-2 (Figure S5). Since ATV inhibits the early proteolytic processing of viral antigens, an early event during SARS-CoV-2 replication, this drug represented the most upstream process pharmacologically inhibited in this investigation to prevent pyroptosis and HLA-DR knockdown.

Inflammasome activation and pyroptosis are risk factors in critically ill COVID-19 patients

To clinically validate our findings, we evaluated if monocytes obtained from critically ill patients with COVID-19 would also display signals of inflammasome engagement and pyroptosis. We observed that monocytes from COVID-19 patients had increased caspase-1 activation (Figure 6A and 6B) and significantly higher PI+ staining, when compared to monocytes from healthy donors (HD) (Figure 6C and 6D). Corroborating with our *in vitro* data, we also detected higher levels of IL-1β in the plasma of critically ill patients (Figure 6E). Therefore, the *in vitro* results from the previous sections stand on the shoulders of the clinical relevance of monocytes in patients with severe COVID-19.

SARS-CoV-2 infection induces inflammasome-dependent pyroptosis and downmodulation of HLA-DR monocytes, which can be prevented by atazanavir

A representative scheme to summarize the effect of SARS-CoV-2 infection upon in the increase of TNF-α and IL-6 levels, as part of an immunodysregulation promoted by pyroptosis. SARS-CoV-2-induced pyropotosis engages inflammasomes, to activate caspase-1 and IL-1ß. Release of IL-1ß amplify the activation of inflammasomes during the SARS-CoV-2 infection. In parallel, SARS-CoV-2 also induces the HLA-DR downregulation. ATV prevented SARS-CoV-2-induced deleterious effects on monocyte biology.

Discussion

COVID-19 has caused in less than 8 months over 800,000 deaths worldwide (21) and represent the major public health crisis of the beginning of 21st century, leading to an unpredictable impact in global economics (22,23). SARS-CoV-2 infection triggers an uncontrolled inflammatory response and marked leukopenia with consequent lung/respiratory dysfunction, which are the characteristics of the most severe

manifestations of the COVID-19 (24,25). Similarly, to other respiratory viruses (26-29),

SARS-CoV-2 induces a cytokine storm, characterized by an uncontrolled inflammatory

response mediated by monocytes/macrophages, when they should orchestrate the

antiviral immune response (30). In this work, we demonstrate that SARS-CoV-2 infection

triggers inflammasome activation, increases IL-1ß secretion by monocytes, resulting in

pyroptotic cell death, which could be a key event for SARS-CoV-2 pathogenesis in

critically ill patients (31). This deleterious immune dysregulation loop triggered by

SARS-CoV-2 may be impaired by ATV.

Our results demonstrate that SARS-CoV-2 leads to an intense cell death in human

monocytes, observed by the increase in LDH release in infected cultures, as well as by

the higher number of PI$^+$ cells when compared to uninfected controls. Even though the

pretreatment of infected monocytes with the pan-caspase inhibitor ZVAD-FMK

prevented IL-1β secretion, it was not able to prevent the release of LDH. Indeed, others

have demonstrated that the treatment with ZVAD-FMK induces necroptosis in diverse

cell types (32-34). Notably, as pyroptosis is an inflammatory and programmed cell death

mediated by inflammasome and inflammatory caspase-1 activation (10,11), we

investigated the engagement and activation of these structures in human monocytes

infected with SARS-CoV-2. Activation of caspase-1 and increased production of IL-1ß

were observed in infected monocytes. This is correlated to the activation of

inflammasomes, because the pretreatment of infected monocytes with YVAD prevented

caspase-1 activation. These *in vitro* results are in accordance with the literature and with

our finding described here that indicate the formation of inflammasomes in patients with

severe COVID-19 (35). We also showed that the release of IL-1ß could be promoted by

the activation of inflammasomes during the SARS-CoV-2 infection, because blockage of

IL-1ß receptors reduced caspase-1 activation and cell death. These results corroborate

with studies showing that the increase in IL-1ß production is associated with severe

COVID-19 (36-38). Under our experimental conditions, the increase in IL-1ß precedes

the unbalanced IL-6 release. These information should not be neglected when considering

biopharmaceuticals to tackle cytokine storm in severe COVID-19.

Monocytes that survived from SARS-CoV-2 infection displayed decreased HLA-

DR expression, which has also been described in monocytes isolated from COVID-19

patients (39,40). Our results suggest that this modulation of HLA expression is not

directly associated with the activation of inflammasomes, and may be involved by other
inflammatory pathways, since we and others have observed increased levels of IL-6 by
myeloid cells impairing HLA-DR membrane expression (41).

To stablish the clinical relevance of SARS-CoV-2-induced pyroptosis in
monocytes (41,41), we analyzed peripheral monocytes isolated from patients with severe
COVID-19. We found that the cells from the patients displayed higher caspase-1
activation, when compared with monocytes isolated from HD. Recent clinical data reveal
high levels of LDH and consistent leukopenia in critically ill COVID-19 patients
(6,7,35,42-44). Our data also demonstrate intense monocyte death in COVID-19 patients,
as detected through flow cytometry analyzes. Altogether, these data suggest that the
severity of COVID-19 may be associated with inflammasome activation in monocytes
that results in large amounts of IL-1ß and generates an excessive inflammatory response,
further characterized by high levels of IL-6 and TNF-α. Consistently, treatment with IL-
1RA has been associated with clinical and inflammatory improvements in critically ill
COVID-19 patients (45). These results are in line with clinical case reports that
demonstrate that monocytes/macrophages are key cells in the deleterious pro-
inflammatory events that characterize the most serious cases of COVID-19 (46-49).

To the best of our knowledge, effective therapies for COVID-19 should ideally
combine antiviral/anti-inflammatory drugs to reduce viral load and to mitigate the
cytokine storm. In the present study, we showed that ATV, an antiretroviral approved in
2003 for HIV-1 treatment and previously described by us as having anti-SARS-CoV-2
effects in different cell types (including monocytes)(50), to block new CoV-induced
pyroptosis and HLA-DR knockdown in monocytes. In addition, ATV inhibited the
release of LDH and the production of IL-1ß, f IL-6 and TNF-α from SARS-CoV-2-
infected monocytes, which are key players in the cytokine storm associated with COVID-
19 (51,52).

In this work, we originally describe that infection with SARS-CoV-2 can induce
pyroptotic cell death by inflammasome activation, which may be related to the intense
leukopenia and exacerbated inflammation seen in severe cases of the COVID-19. Since
there is no specific therapy for this disease, our results point out that ATV has a promising
therapeutic potential against SARS-CoV-2-induced cell death.

Acknowledgments and funding

We thank the Hemotherapy Service from Hospital Clementino Fraga Filho (Federal University of Rio de Janeiro, Brazil) for providing buffy-coats. This work was supported by grants from Fundação de Amparo a Pesquisa do Estado do Rio de Janeiro (FAPERJ), Inova Fiocruz, Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) and Mercosur Fund for Structural Convergence (FOCEM, Mercosur, grant number 03/11) granted for Thiago Moreno L. Souza, Patrícia T. Bozza, Fernando A. Bozza and Dumith Chequer Bou-Habib. Funding was also provided by CNPq, CAPES and FAPERJ through the National Institutes of Science and Technology Program (INCT) to Carlos Morel (INCT-IDPN) and to Wilson Savino (INCT-NIM). Thanks are due to Oswaldo Cruz Foundation/Fiocruz under the auspicious of Inova program. The funding sponsors had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, and in the decision to publish the results.

References

1. Conti P, Ronconi G, Caraffa A, Gallenga CE, Ross R, Frydas I, et al. Induction of pro-inflammatory cytokines (IL-1 and IL-6) and lung inflammation by COVID-19: anti-inflammatory strategies. Journal of biological regulators and homeostatic agents. 2020.

2. Long B, Brady WJ, Koyfman A, Gottlieb M. Cardiovascular complications in COVID-19. American Journal of Emergency Medicine. 2020.

3. Li J, Fan J-G. Characteristics and Mechanism of Liver Injury in 2019 Coronavirus Disease. J Clin Transl Hepatol. 2020.

4. Giamarellos-Bourboulis EJ, Netea MG, Rovina N, Akinosoglou K, Antoniadou A, Antonakos N, et al. Cell Host Microbe. 2020.

5. Du RH, Liang LR, Yang CQ, Wang W, Cao TZ, Li M, et al. Predictors of mortality for patients with COVID-19 pneumonia caused by SARSCoV- 2: A prospective cohort study. Eur Respir J. 2020.

457   6.   Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors
458   for mortality of adult inpatients with COVID-19 in Wuhan, China: a
459   retrospective cohort study. Lancet. 2020.

460   7.   Wölfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, Müller MA, et
461   al. Virological assessment of hospitalized patients with COVID-2019. Nature.
462   2020.

463   8.   Wang J, Jiang M, Chen X, Montaner LJ. Cytokine storm and leukocyte
464   changes in mild versus severe SARS-CoV-2 infection: Review of 3939
465   COVID-19 patients in China and emerging pathogenesis and therapy
466   concepts. Journal of Leukocyte Biology. 2020.

467   9.   Chen Z, John Wherry E. T cell responses in patients with COVID-19. Nature
468   Reviews Immunology. 2020.

469   10.   Danthi P. Viruses and the Diversity of Cell Death. Annual Review of
470   Virology. 2016.

471   11.   Jorgensen I, Miao EA. Pyroptotic cell death defends against intracellular
472   pathogens. Immunological Reviews. 2015.

473   12.   Man SM, Karki R, Kanneganti TD. Molecular mechanisms and functions of
474   pyroptosis, inflammatory caspases and inflammasomes in infectious diseases.
475   Immunological Reviews. 2017.

476   13.   Xu Z, Shi L, Wang Y, Zhang J, Huang L, Zhang C, et al. Pathological findings
477   of COVID-19 associated with acute respiratory distress syndrome. Lancet
478   Respir Med. 2020.

479   14.   Han Y, Zhang H, Mu S, Wei W, Jin C, Xue Y, et al. Lactate dehydrogenase,
480   a Risk Factor of Severe COVID-19 Patients. medRxiv. 2020.

481   15.   Terpos E, Ntanasis-Stathopoulos I, Elalamy I, Kastritis E, Sergentanis TN,
482   Politou M, et al. Hematological findings and complications of COVID-19.
483   American Journal of Hematology. 2020.

484   16.   Li YX, Wu W, Yang T, Zhou W, Fu YM, Feng QM, et al. [Characteristics of
485   peripheral blood leukocyte differential counts in patients with COVID-19].
486   Zhonghua nei ke za zhi. 2020.

487   17.   Ding X, Yu Y, Lu B, Huo J, Chen M, Kang Y, et al. Dynamic profile and
488   clinical implications of hematological parameters in hospitalized patients with
489   coronavirus disease 2019. Clin Chem Lab Med. 2020.

18. Singh A, Sood N, Narang V, Goyal A. Morphology of COVID-19-affected
cells in peripheral blood film. BMJ Case Reports. 2020.

19. Zalinger ZB, Elliott R, Weiss SR. Role of the inflammasome-related cytokines
Il-1 and Il-18 during infection with murine coronavirus. J Neurovirol. 2017.

20. Franchi L, Eigenbrod T, Muñoz-Planillo R, Nuñez G. The inflammasome: A
caspase-1-activation platform that regulates immune responses and disease
pathogenesis. Nature Immunology. 2009.

21. OPAS. COVID-19 (doença causada pelo novo coronavírus). Folha Inf. 2020.

22. Dong E, Du H, Gardner L. An interactive web-based dashboard to track
COVID-19 in real time. The Lancet Infectious Diseases. 2020.

23. Wissel BD, Van Camp PJ, Kouril M, Weis C, Glauser TA, White PS, et al. An
Interactive Online Dashboard for Tracking COVID-19 in U.S. Counties,
Cities, and States in Real Time. J Am Med Inform Assoc. 2020.

24. Lucas C, Wong P, Klein J, Castro TBR, Silva J, Sundaram M, et al.
Longitudinal analyses reveal immunological misfiring in severe COVID-19.
Nature. 2020.

25. Tay MZ, Poh CM, Rénia L, MacAry PA, Ng LFP. The trinity of COVID-19:
immunity, inflammation and intervention. Nature Reviews Immunology.
2020.

26. Teijaro JR. The role of cytokine responses during influenza virus pathogenesis
and potential therapeutic options. Curr Top Microbiol Immunol. 2014.

27. Szretter KJ, Gangappa S, Lu X, Smith C, Shieh W-J, Zaki SR, et al. Role of
Host Cytokine Responses in the Pathogenesis of Avian H5N1 Influenza
Viruses in Mice. J Virol. 2007.

28. Peschke T, Bender A, Nain M, Gemsa D. Role of Macrophage Cytokines in
Influenza A Virus lnfections. Immunobiology. 1993.

29. Woo PCY, Tung ETK, Chan KH, Lau CCY, Lau SKP, Yuen KY. Cytokine
profiles induced by the novel swine-origin influenza A/H1N1 virus:
Implications for treatment strategies. J Infect Dis. 2010.

30. Jamilloux Y, Henry T, Belot A, Viel S, Fauter M, El Jammal T, et al. Should
we stimulate or suppress immune responses in COVID-19? Cytokine and anti-
cytokine interventions. Autoimmunity Reviews. 2020.

31. Zheng Z, Peng F, Xu B, Zhao J, Liu H, Peng J, et al. Risk factors of critical & mortal COVID-19 cases: A systematic literature review and meta-analysis. Journal of Infection. 2020.

32. Siegmund D, Kums J, Ehrenschwender M, Wajant H. Activation of TNFR2 sensitizes macrophages for TNFR1-mediated necroptosis. Cell Death Dis. 2016.

33. Kim SJ, Li J. Caspase blockade induces RIP3-mediated programmed necrosis in Toll-like receptor-activated microglia. Cell Death Dis. 2013.

34. Wu YT, Tan HL, Huang Q, Sun XJ, Zhu X, Shen HM. ZVAD-induced necroptosis in L929 cells depends on autocrine production of TNFα mediated by the PKC-MAPKs-AP-1 pathway. Cell Death Differ. 2011.

35. Rodrigues S, de Sá KS, Ishimoto AY, Becerra A, Oliveira S, Almeida L, et al. Inflammasome activation in COVID-19 patients 1 2 Tamara. medRxiv. 2020;

36. Abbasifard M, Khorramdelazad H. The bio-mission of interleukin-6 in the pathogenesis of COVID-19: A brief look at potential therapeutic tactics. Life Sciences. 2020.

37. Yao Z, Zheng Z, Wu K, Zheng J. Immune environment modulation in pneumonia patients caused by coronavirus: SARS-CoV, MERS-CoV and SARS-CoV-2. Aging (Albany NY). 2020.

38. Zhang Y, Li J, Zhan Y, Wu L, Yu X, Zhang W, et al. Analysis of serum cytokines in patients with severe acute respiratory syndrome. Infect Immun. 2004.

39. Ong EZ, Chan YFZ, Leong WY, Lee NMY, Kalimuddin S, Haja Mohideen SM, et al. A Dynamic Immune Response Shapes COVID-19 Progression. Cell Host Microbe. 2020.

40. Laing AG, Lorenc A, del Molino del Barrio I, Das A, Fish M, Monin L, et al. A dynamic COVID-19 immune signature includes associations with poor prognosis. Nat Med [Internet]. 2020; Available from: https://doi.org/10.1038/s41591-020-1038-6.

41. Ohno Y, Kitamura H, Takahashi N, Ohtake J, Kaneumi S, Sumida K, et al. IL-6 down-regulates HLA class II expression and IL-12 production of human dendritic cells to impair activation of antigen-specific CD4+ T cells. Cancer Immunol Immunother. 2016.

42. Yap JKY, Moriyama M, Iwasaki A. Inflammasomes and Pyroptosis as
Therapeutic Targets for COVID-19. J Immunol. 2020.

43. Yuan J, Zou R, Zeng L, Kou S, Lan J, Li X, et al. The correlation between
viral clearance and biochemical outcomes of 94 COVID-19 infected
discharged patients. Inflamm Res. 2020.

44. Yan L, Zhang HT, Goncalves J, Xiao Y, Wang M, Guo Y, et al. An
interpretable mortality prediction model for COVID-19 patients. Nat Mach
Intell. 2020.

45. Cauchois R, Koubi M, Delarbre D, Manet C, Carvelli J, Blasco VB, et al. Early
IL-1 receptor blockade in severe inflammatory respiratory failure
complicating COVID-19. Proc Natl Acad Sci. 2020.

46. Qin C, Zhou L, Hu Z, Zhang S, Yang S, Tao Y, et al. Dysregulation of immune
response in patients with COVID-19 in Wuhan, China. Clin Infect Dis. 2020.

47. McGonagle D, Sharif K, O'Regan A, Bridgewood C. The Role of Cytokines
including Interleukin-6 in COVID-19 induced Pneumonia and Macrophage
Activation Syndrome-Like Disease. Autoimmunity Reviews. 2020.

48. Merad M, Martin JC. Pathological inflammation in patients with COVID-19:
a key role for monocytes and macrophages. Nature Reviews Immunology.
2020.

49. Park MD. Macrophages: a Trojan horse in COVID-19? Nat Rev Immunol.
2020.

50. Fintelman-Rodrigues N, Sacramento CQ, Lima CR, Silva FS da, Ferreira A,
Mattos M, et al. Atazanavir inhibits SARS-CoV-2 replication and pro-
inflammatory cytokine production. bioRxiv. 2020.

51. Hantoushzadeh S, Norooznezhad AH. Possible Cause of Inflammatory Storm
and Septic Shock in Patients Diagnosed with (COVID-19). Arch Med Res.
2020.

52. Hantoushzadeh S, Norooznezhad AH. Inappropriate Antibiotic Consumption
as a Possible Cause of Inflammatory Storm and Septic Shock in Patients
Diagnosed with Coronavirus Disease 2019 (COVID-19). Arch Med Res.
2020.

586
587
588
589

 FIGURE LEGENDS

Figure 1. SARS-Cov-2 induces monocyte cell death through pyroptosis. Human

monocytes were infected with SARS-CoV-2 (MOI 0.1) for 24 h. As a positive control,

monocytes were stimulated with LPS (500ng/mL) for 23 h and after this time, incubated

with ATP (2 mM) for 1 h. **(A)** Cell viability was assessed through the measurement of

LDH release in the supernatant of monocytes. **(B, C and D)** Monocytes` monolayers were

stained with propidium iodide (PI) plus Annexin V and pyroptotic cell death were

determine by flow cytometer analysis and **(E)** labeled with DAPI to visualize the nuclei

fluorescence microscopy. Images and Graph data are representative of six independent

experiments. Data are presented as the mean ± SEM [*] $P < 0.05$ *versus* control group

(MOCK).

Figure 2. SARS-CoV-2 induce inflammasome activation in human monocytes.

Human monocytes were pretreated with caspase-1 inhibitor (AC-YVAD-CMK – 1 µM),

pan-caspase inhibitor (ZVAD-FMK - 10 µM) or necroptosis inhibitor (Nec-1 - 25 µM)

for 1 h and infected with SARS-Cov-2 (MOI 0.1) for 24 h. Monocytes were stimulated

with LPS (500ng/mL) for 23 h and after this time were stimulated with ATP (2 mM) for

1 h as a positive control group for the formation of inflammasomes and pyroptosis

induction. **(A, B)** Monocytes` monolayers were used for determination of the levels of

pro-caspase-1 and cleaved caspase-1 by western blot analysis. **(C - F)** Monocytes were

stained with FAM-YVAD-FLICA and FAM-FLICA to determine the caspase-1 and

caspase-3/7 activity, respectively, by flow cytometry. **(G)** Cell culture supernatants were

collected for the measurement of the levels of IL-1β. Western blot, histogram and graph

data are representative of six independent experiments. Data are presented as the mean ±

SEM [#] $P < 0.05$ *versus* infected and untreated group.

Figure 3. Inflammasome activation amplify pro-inflammatory cytokines secretion

in infected monocytes. Human monocytes were pretreated with caspase-1 inhibitor (AC-

YVAD-CMK – 1 µM), pan-caspase inhibitor (ZVAD-FMK - 10 µM), necroptosis

inhibitor (Nec-1 - 25 µM) or with inhibitor of IL-1ß receptor (IL-1RA – 1 µM).

Monocytes were stimulated with LPS (500ng/mL) for 23 h and after this time were

stimulated with ATP (2 mM) for 1 h as a positive control group for the formation of

inflammasomes and pyroptosis induction. Cell culture supernatants were collected for the

measurement of the levels of **(A)** IL-6, **(B)** TNF-α and **(C)** IL-8. Graphs data are

representative of six independent experiments. Data are presented as the mean ± SEM [#]

P < 0.05 *versus* infected and untreated group.

Figure 4. Infection with SARS-CoV-2 decreases the expression of HLA in human

monocytes independently of the activation of inflammasomes. Human monocytes

were pretreated with caspase-1 inhibitor (AC-YVAD-CMK – 1 µM) for 1 h and infected

with SARS-Cov-2 for 24 h. **(A, B)** Monocytes were stained with HLA-DR APC.H7 or

Isotype control APC.H7 to determine the HLA-DR expression by flow cytometry. Mean

fluorescence intensity (MFI) value for each sample was represented in graphics.

Histogram and graphs datas are representative of six independent experiments. Data are

presented as the mean ± SEM [*] *P* < 0.05 *versus* uninfected group (MOCK).

Figure 5. Atazanavir prevented inflammasome-dependent pyroptosis. Human

monocytes were infected with SARS-Cov-2 and treated with atazanavir – ATV (10 µM),

ribavirina (10 µM) or Lopinavir (10 µM) for 24 h. **(A, B and C)** Monocytes were stained

with FAM-YVAD-FLICA or FAM-FLICA and HLA-DR APC.H7 or IgG APC.H7 to

determine the caspase-1 and caspase-3/7 activity and HLA-DR expression, respectively

and analyzed by flow cytometry. **(D – G)** Culture supernatants were collected and the

levels of IL-1β, IL-6, TNF-α and IL-8 were determined by ELISA. **(H)** Assessment of

cell viability through the measurement of LDH release in the supernatant of monocytes.

(I and J) Monocytes were stained with HLA-DR APC.H7 to determine the HLA-DR

expression by flow cytometry. Histogram and graphs datas are representative of six

independent experiments. Data are presented as the mean ± SEM [*] *P* < 0.05 *versus* control

group (MOCK); [#] *P* < 0.05 *versus* only infected group.

Figure 6. Inflammasome activation and pyroptosis are risk factors in critically ill

COVID-19 patients. Monocytes isolated from blood samples of critically ill patients

with COVID-19 and healthy donors. **(A and B)** monocytes were stained with FAM-

YVAD-FLICA or **(C and D)** propidium iodide (PI) and analyzed by flow cytometry. **(E)**

plasma was separated and the levels of IL-1β were determined by ELISA. Western blot,

histogram and graphs datas are representative of 9 critically ill patients and 8 healthy

donors. Data are presented as the mean ± SEM [*] *P* < 0.05 *versus* healthy donors (HD).

Figure 7. Representative scheme of the SARS-CoV-2 infection in activation of
inflammasome and pyroptosis in monocyte with downregulation of HLA-DR expression
and effects of ATV treatment in this phenomenon.

Table 1: Characteristics of COVID-19 patients and control subjects.

Characteristics[1]	Control (N=8)	Covid-19 (N=9)
Age, years	52.6 (47 – 60)	61 (33 – 79)
Sex, male	8 (100 %)	6 (66.7 %)
Respiratory support		
Oxygen supplementation	0 (0 %)	3 (33.3 %)
Mechanical ventilation	0 (0 %)	6 (66.7 %)
SAPS 3		59.25 (31 – 75)
PaO2/FiO2 ratio	-	210.75 (87 – 509.5)
Vasopressor	-	3 (33.3 %)
Time from symptom onset to blood sample, days	-	10 (6 – 18)
Status on Jun 30th		
Dead	-	5 (55.55 %)
Discharged	-	1 (11.1 %)
Hospitalized	-	3 (33.3 %)
Comorbidities		
Obesity	1 (11.1 %)	1 (11.1 %)
Hypertension	3 (33.3%)	3 (33.3 %)
Diabetes	0 (0%)	0 (0 %)
Cancer	0 (0%)	0 (0 %)
Chronic heart disease[2]	0 (0%)	0 (0 %)
Presenting symptoms		
Cough	0 (0 %)	3 (33,3 %)
Fever	0 (0 %)	3 (33,3 %)
Dyspnea	0 (0 %)	4 (44,4 %)
Headache	0 (0 %)	0 (0 %)
Anosmia	0 (0 %)	0 (0 %)
Laboratory findings on admission		
Lymphocyte count, cells/mm^3	-	1,355.8 (552 –2,564)
Platelet count, x 1000/mm^3	-	169.6 (92 – 278)
Leukocytes	-	8,036 (8.03-18,670)
C Reactive Protein, mg/L	-	14.78 (0.1 – 30.8)

FIGURE 1

FIGURE 2

771   **FIGURE 3**

FIGURE 4

FIGURE 5

FIGURE 6

FIGURE 7

    FIGURE S1

S1 Fig. SARS-CoV-2 induce inflammasome activation and pyroptosis in human
monocytes by IL-1β amplification. Human monocytes were pretreated with IL-1β
receptor (IL-1RA – 1 μM) for 1 h and infected with SARS-Cov-2 (MOI 0.1) for 24 h.
Cell viability was assessed through the measurement of LDH release in the supernatant
and monocytes were stained with FAM-YVAD-FLICA to determine the caspase-1 and
caspase-3/7 activity, respectively, by flow cytometry. Cell culture supernatants also were
collected for the measurement of the levels of IL-1β. Data are presented as the mean ±
SEM [#] $P < 0.05$ *versus* infected and untreated group; [*] $P < 0.05$ *versus* control group
(MOCK). Graphs are representative of four independent experiments.

 FIGURE S2

S2 Fig. Infection with SARS-CoV-2 decreases the expression of HLA in human

monocytes independently of the IL-1ß activation. Human monocytes were pretreated

with inhibitor of IL-1ß receptor (IL-1RA – 1 µM) for 1 h and infected with SARS-Cov-

2 for 24 h. Monocytes were stained with HLA-DR APC.H7 or IgG APC.H7 to determine

the HLA-DR expression by flow cytometry. Mean fluorescence intensity (MFI) value for

each sample was represented in graphics. Graphs datas are representative of four

independent experiments. Data are presented as the mean ± SEM [#] $P < 0.05$ *versus*

uninfected group (MOCK).

 FIGURE S3

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057 **S3 Fig. Evaluation of anti-inflammatory activity of ATV, LPV and ribavirin in LPS-induced monocytes.** Human monocytes were stimulated

1058 with LPS (500ng/mL) for 23 h and after this time were stimulated with ATP (2 mM) for 1 h as a positive control group for the formation of

1059 inflammasomes and pyroptosis induction. Simultaneously, cells were treated with atazanavir – ATV (10 µM), LPV (10 µM) or ribavirin (10 µM)

1060 for 24 h. Monocytes were stained with FAM-YVAD-FLICA (blue) or FAM-FLICA (green) to determine the caspase-1 and caspase-3/7 activity

1061 and HLA-DR expression, respectively, and analyzed by flow cytometry. Histograms are representative of six independent experiments.

1062
1063
1064
1065
1066 **FIGURE S4**

1096 **S4 Fig. Effects of LPV and ribavirin in preventing the SARS-CoV-2-related reduced**

1097 **expression of HLA-DR in monocytes.** Human monocytes were infected with SARS-

1098 Cov-2 and treated with LPV (10 µM) or ribavirina (10 µM) for 24 h. Monocytes were

1099 stained with HLA-DR APC.H7 or IgG APC.H7 to determine the HLA-DR expression by

1100 flow cytometry. Histogram data are representative of six independent experiments.

1101
1102
1103
1104
1105