County-level exposures to greenness and associations with COVID-19 incidence and mortality in the United States

Jochem O Klompmaker [a], Jaime E Hart [a, b], Isabel Holland [b], M Benjamin Sabath [c], Xiao Wu [c], Francine Laden [a, b, d], Francesca Dominici [c], Peter James [a, e]

[a] *Department of Environmental Health, Harvard T. H. Chan School of Public Health, 655 Huntington Avenue, Boston, Massachusetts 02115*
[b] *Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital, 181 Longwood Avenue, Boston, Massachusetts 02115*
[c] *Department of Biostatistics, Harvard T.H. Chan School of Public Health, 677 Huntington Avenue, Boston, Massachusetts 02115*
[d] *Department of Epidemiology, Harvard T. H. Chan School of Public Health, 677 Huntington Avenue, Boston, Massachusetts 02115*
[e] *Department of Population Medicine, Harvard Medical School and Harvard Pilgrim Health Care Institute, 401 Park Drive, Boston, Massachusetts 02215*

Corresponding author:
Jochem O Klompmaker, PhD
Postdoctoral fellow
Department of Environmental Health
Harvard T. H. Chan School of Public Health

Landmark Center, 401 Park Drive, Boston MA 02215
jklompmaker@hsph.harvard.edu

RUNNING TITLE: Greenness and COVID-19 incidence and mortality

KEY WORDS: green space, Coronavirus, epidemiology, environmental health

Declaration of interest
The authors declare they have no actual or potential competing financial interests.

Acknowledgement
This study was funded by R01HL150119 and R01ES028033.

Abstract

Background: COVID-19 is an infectious disease that has killed more than 175,000 people in the US. During a time of social distancing measures and increasing social isolation, green spaces may be a crucial factor to maintain a physically and socially active lifestyle while not increasing risk of infection.

Objectives: We evaluated whether greenness is related to COVID-19 incidence and mortality in the United States.

Methods: We downloaded data on COVID-19 cases and deaths for each US county up through June 7, 2020, from Johns Hopkins University, Center for Systems Science and Engineering Coronavirus Resource Center. We used April-May 2020 Normalized Difference Vegetation Index (NDVI) data, to represent the greenness exposure during the initial COVID-19 outbreak in the US. We fitted negative binomial mixed models to evaluate associations of NDVI with COVID-19 incidence and mortality, adjusting for potential confounders such as county-level demographics, epidemic stage, and other environmental factors. We evaluated whether the associations were modified by population density, proportion of Black residents, median home value, and issuance of stay-at-home order.

Results: An increase of 0.1 in NDVI was associated with a 6% (95% Confidence Interval: 3%, 10%) decrease in COVID-19 incidence rate after adjustment for potential confounders. Associations with COVID-19 incidence were stronger in counties with high population density and high median home values, and in counties with stay-at-home orders. Greenness was not associated with COVID-19 mortality in all counties; however, it was protective in counties with high percentages of Black residents, high median home value, and higher population density.

Discussion: Exposures to NDVI had beneficial impacts on county-level incidence of COVID-19 in the US and may have reduced county-level COVID-19 mortality rates, especially in densely populated counties.

Introduction

The global spread of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2), the
virus responsible for COVID-19, has caused a worldwide public health emergency. (Sohrabi et
al. 2020; WHO 2020a) The outbreak was declared a pandemic by the World Health Organization
(WHO) on March 11, 2020, (WHO 2020b) and as of August 25, 2020, 23.5 million cases of
COVID-19 had been documented worldwide, and more than 800,000 deaths had been recorded.
(Johns Hopkins Coronavirus Resource Center 2020) To date, there are few effective therapies
and no vaccine, therefore public health measures at the population level (e.g. social distancing
measures, stay-at-home orders, public education initiatives (Prem et al. 2020; Tammes 2020))
have been the primary approach for reducing transmission.

The coronavirus pandemic presents an unprecedented situation for the globe; however, this is not
the first time the world has confronted a large-scale infectious disease threat. Historical
approaches to combat infectious disease outbreaks provide crucial lessons that we can still apply
today. One of those approaches is the use of urban parks as a resilience measure. Frederick Law
Olmsted, who designed New York's Central Park, Boston's Emerald Necklace, and many other
major urban parks, championed the concept of "parks as lungs" and the espoused "two great
natural agents of disinfection: sunshine, and fall foliage".(Beveridge and Hoffman 1997)

The spread of infectious diseases, like COVID-19, is dependent on the duration of
infectiousness, transmissibility, and the contact rate.(Heederik et al. 2020; Delamater et al. 2019)
These three factors generally summarize the basic reproduction number (R0). R0 is affected by
numerous biological, socio-behavioral and environmental factors that influence pathogen
transmission.(Delamater et al. 2019) Green spaces may influence the contact rate and in turn the
reproduction number, as they provide a setting to obtain much needed physical activity and a
place for social interactions while remaining the recommended safe distance (six feet). Because
these activities take place outdoors, wind dilutes the amount of virus in the air substantially
(Qian et al. 2020), which greatly decreases transmission risk. In addition, theory (Ulrich 1984;
Kaplan & Kaplan 1989) and empirical evidence (Banay et al. 2019; Bezold et al. 2018) suggests
that living near green spaces allows us to restore our attention and decrease stress, leading to
lower incidence of depression, anxiety, and other negative psychological factors. During a time
of social distancing measures and increasing social isolation, urban green spaces may be a
crucial factor to maintain a physically and socially active lifestyle while not increasing risk of
infection.

To quantify whether greenness is related to COVID-19 incidence and mortality, we compiled
county level data on both greenness and COVID-19 outcomes. Furthermore, based on evidence
that there are large disparities in incidence and mortality rates, we evaluated whether the
relationship between greenness and incidence/mortality differed according to county-level
population density, percentage of black residents, median home value and issuance of stay-at-
home orders.

Methods

Data used in this study are publicly available and links to each of the data sources can be found
in Table S1.

COVID-19 incidence and mortality data

The Johns Hopkins University Center for Systems Science and Engineering Coronavirus Resource Center provides daily updates about COVID-19 death counts and cases for each country.(Dong et al. 2020) For the US, county level data is provided by the US Centers for Disease Control and Prevention (CDC) and State governments. The number of COVID-19 cases is the sum of the number of deaths and active cases. As of April 14, 2020, CDC case counts and death counts included both confirmed and probable cases and deaths in accordance with CDC guidelines.(CDC 2020)

We downloaded data on the cumulative number of COVID-19 cases and deaths for each county through June 7, 2020. County-level COVID-19 mortality/incidence rates were defined as the ratio of COVID-19 deaths/cases to county level population size.(Wu et al. 2020)

Greenness exposure

For each county, the Normalized Difference Vegetation Index (NDVI) was estimated using satellite imagery. The NDVI is calculated as the ratio between the red and near infrared values, and ranges from -1 to 1.(NASA 2020) Values close to 1 correspond to areas with complete coverage by live green vegetation, values close to zero correspond to areas without much live vegetation (e.g. rocks, sand) and negative values correspond to water. We used Landsat 8 (Collection 1 Tier 1 Operational Land Imager DN values, representing scaled, calibrated at sensor radiance (USGS 2020)) images for the entire US from April 1, 2020 up to May 31, 2020, to represent the exposure during the initial COVID-19 outbreak in the US. Landsat 8 images are generated every 16 days at 30m resolution. Using Google Earth Engine (https://earthengine.google.com/), cloud-free Landsat composites were created for the US. We calculated the spatially weighted mean April-May NDVI for each county in the US, after setting negative NDVI values to zero. County shapefiles were based on the US Census Bureau Tiger dataset of 2018.(US Census Bureau 2020)

Potential confounders

To adjust for potential confounding bias, we obtained data on several variables that might be linked to green space and COVID-19 incidence and mortality. We collected eleven county level Census variables from the 2000 Census (https://www.census.gov) and the 2010 5-year American Community Surveys (https://www.census.gov/programs-surveys/acs/): proportion of residents older than 65, proportion of residents aged 15-44, proportion of residents aged 45-64, proportion of Hispanic residents, proportion of Black residents, median household income, median home value, proportion of residents in poverty, proportion of residents with a high school diploma, population density, and proportion of residents that own their house. From the Behavioral Risk Factor Surveillance System (BRFSS) (https://www.countyhealthranki ngs.org/) we obtained the proportion of individuals that were obese and the proportion of current smokers in 2011, the most recent year available.

We used days since first COVID-19 case reported in a county as a proxy for stage of the COVID-19 outbreak. Further, we linked days since issuance of stay-at-home order (state-level), days since closure of non-essential businesses (state-level), and days since nursing home visitor ban (state-level) from the COVID-19 US State Policy Database (Raifman et al. 2020) to our data. Since the availability of adequate hospital resources might influence COVID-19 outcomes,

185 we collected county-level information on the number of hospital beds available in 2019 from the
186 Homeland Infrastructure Foundation-Level Data (HIFLD). In addition, we used state level
187 information on number of COVID-19 tests performed up to June 7, 2020 from the COVID
188 tracking project (https://covidtracking.com/).
189
Based on previous studies implicating relations between exposure to particulate matter less than
2.5 microns ($PM_{2.5}$), temperature and/or relative humidity and COVID-19 incidence and
mortality (Wu et al. 2020; Raines et al. 2020), we also adjusted our analyses for these factors.
Temperature and relative humidity data were available from Gridmet, and we created long-term
(2000-2016) summer (June-August) and winter (December-February) averages for each
county.(Abatzoglou 2013) $PM_{2.5}$ concentration estimates for 2000-2016 were derived from an
established exposure prediction model.(Van Donkelaar et al. 2019)
197
Statistical methods
We used negative binomial mixed models to evaluate associations of NDVI with COVID-19
incidence and mortality. We report mortality rate ratios (MRR) and incidence rate ratios (IRR),
i.e., exponentiated effect estimates from the negative binomial mixed model, and 95% CI per 0.1
unit NDVI increase. To evaluate effects of potential confounders, we specified a series of models
with increasing covariate adjustment. In model 1 we only included a population size offset and a
random intercept by state. In model 2 we additionally adjusted for degree of urbanization. In
model 3 we added all county-level SES covariates and BRFSS covariates. In model 4 we added
date since first COVID-19 case reported, date since issuance of stay-at-home order for each state,
number of hospital beds per unit population. In model 5 we additionally included temperature,
relative humidity and $PM_{2.5}$ (main model for COVID-19 mortality). The number of tests per unit
population was added to model 6 (main model for COVID-19 incidence). We used a general
additive mixed model with penalized cubic regression splines (with 2 degrees of freedom as the
upper limit) to evaluate whether the association of NDVI with COVID-19 mortality and
incidence was linear in the full cohort, in rural counties, and in urban counties. We carried out all
analyses in R statistical software and performed model fitting using the lme4 package or the
gamm4 package (for spline analyses).(Package "lme4"; Package "gamm4")
215
We evaluated whether associations of NDVI with COVID-19 deaths and cases were modified by
population density, proportion of black residents, median home value, and issuance of stay-at-
home order by adding an interaction term to the model. Significance of interaction terms were
tested by Chi-square tests between the models with and without the interaction terms. We
hypothesized that associations of NDVI with COVID-19 incidence and mortality were stronger
in densely populated counties, in counties with issuance of stay-at-home orders, in counties with
higher proportions of black residents, and in counties with lower median home values.
223
We conducted several sensitivity analyses to assess the robustness of the associations. We
excluded 27 counties comprising the New York metropolitan area (n = 3,062), as this area
experienced the most severe COVID-19 outbreak. We also conducted analysis excluding
counties with 10 or fewer confirmed COVID-19 cases. We additionally added days since closure
of non-essential businesses and days since nursing home visitor ban to our models. In addition,
we used county averages of NDVI with negative values excluded (instead of set to zero).
230

Results
Our study cohort consisted of 3,089 counties of which 2,297 counties reported more than 10 cases. The highest COVID-19 death rates were in New York, Illinois, Michigan, Florida, Louisiana, and California (Figure 1). COVID-19 incidence rates were more equally spread over the US. NDVI values were high along the West coast and in the South. The median COVID-19 death rate per 100,000 individuals was 2.8 and the median COVID-19 incidence rate per 100,000 individuals was 163.5 (Table 2). NDVI was moderately positively correlated with % Black, % smoke, and $PM_{2.5}$, and weakly negatively correlated with median household income (Figure S1).

In main models (model 5 for mortality and model 6 for incidence) we found an IRR of 0.94 (95% CI: 0.90, 0.97) and a MRR of 0.99 (95% CI: 0.94, 1.05) per 0.1 unit increase in NDVI. There was little impact of population density on estimates; however, effects were attenuated in models that included county-level SES, BMI and smoking. The additional inclusion of epidemic stage, timing of stay-at-home-orders, hospital beds per capita, long-term exposures to $PM_{2.5}$, weather and COVID test rate did not confound the association. Estimated IRR and MRR for all covariates included in the fully adjusted models can be found in Table S2. The overall exposure-response curve for COVID-19 incidence showed some small evidence of deviations from linearity, with a potential threshold effect around an NDVI of 0.5 (Figure 2). For urban counties, the curve for COVID-19 incidence was inverse and linear, while for rural counties increasing NDVI appeared beneficial at the lower end of the distribution only. Similar patterns, although slightly more pronounced, were observed for COVID-19 mortality.

The associations of NDVI with COVID-19 incidence and mortality were stronger in densely populated counties and counties with higher median home values (Figure 3). For COVID-19 incidence, but not for mortality, we found stronger associations for counties with issuance of stay-at-home orders. NDVI was also inversely associated with COVID-19 mortality in counties with the highest proportion of Black residents, while in the lowest quintiles of proportion of Black residents the association of NDVI with COVID-19 mortality was positive.

In sensitivity analyses, associations were robust to additional adjustment for days since closures of non-essential businesses or days since a nursing home visitor ban, exclusions of the NYC metro area, restriction to counties with at least 10 cases, or alternative procedure for calculating NDVI. (Figure S3).

Discussion
We observed that greenness in April-May of 2020 was inversely associated with COVID-19 incidence, especially in urban counties. An increase of 0.1 in NDVI was associated with a 6% decrease in COVID-19 incidence rate after adjustment for potential confounders at the county level. Associations with COVID-19 incidence were stronger in counties with high median home values, in more densely populated counties, and in counties with stay-at-home orders. NDVI was not associated with COVID-19 mortality in all counties; however, NDVI was protective in counties with high percentages of Black residents, high median home value, and higher population density.

Since the COVID-19 outbreak, social distancing policies and guidelines have led to more time spent at home and therefore people may be more dependent on their immediate surroundings.

Because gyms were closed in large parts of the US during this time period, people may have relied on parks to be physically active. Parks also provide places for social gatherings outdoors. Being outside might substantially reduce the chance of SARS-CoV-2 transmission. According to a study performed among 7,324 identified cases in China, only a single small outdoor outbreak was identified.(Qian et al. 2020)

We found stronger associations between NDVI and COVID-19 incidence. This seems plausible as neighborhood green space might affect contact rates and therefore COVID-19 incidence, while COVID-19 mortality also depends on available treatments and on host susceptibility, such as age and presence of chronic diseases. Several reviews showed inverse associations of greenness with a variety of diseases.(Fong et al. 2018; James et al.) A couple of studies also reported inverse associations with cardiovascular and respiratory disease mortality, even after adjustment for air pollution.(Crouse et al. 2017; Vienneau et al. 2017) This suggests that increased amounts of greenness could influence host susceptibility. For COVID-19 incidence, associations were stronger (and linear) in urban versus rural counties. This is in line with the literature on the health effects of green spaces, which suggest benefits of green space are stronger in urban areas.(Fong et al. 2018)

For COVID-19 incidence and mortality, we found stronger associations in densely populated counties and counties with high median home values. Median home value is likely related to health insurance and the ability to work from home, which affects COVID-19 incidence and mortality. The positive associations of NDVI with COVID-19 mortality in the lowest population density quintile could be because an increase in greenness in these areas is related to limited access to health care. Associations of NDVI with COVID-19 incidence were modified by state-level issuance of stay-at-home orders. Individuals living in states with stay-at-home orders might spend more time at home and are thus more dependent on their immediate surroundings, like greenness. Individuals living in states without stay-at-home orders might not practice social distancing and may differ in COVID-19 health risk perceptions. However, differences in associations could also be due to differences in epidemic stage (number of COVID-19 cases) in counties with and without stay-at-home orders.

Associations of NDVI with COVID-19 mortality, but not COVID-19 incidence, were modified by percentage Black, where NDVI was harmful in the counties with the lowest proportion of Black residents, but protective in all other quintiles. This finding may be related to higher observed rates of mortality among Black individuals, or may reflect the moderate correlation between percentage of Black residents and population density.

This study has several strengths. We used NDVI for April-May 2020, largely overlapping with the beginning of the COVID-19 outbreak in the US, allowing us to assess the impact of temporally relevant exposures on incidence and mortality. Associations of NDVI with COVID-19 incidence remained in analyses stratified by urban-rural status or population density, indicating that our associations are not a result of differences in urban-rural COVID-19 incidence or testing rates. We adjusted for several potentially important confounders, such as proportion Black residents, population density, and days since first COVID-19 case. We note that NDVI was moderately positively (Spearman rho > 0.40) correlated with % less than high school education, % Black residents, % current smokers, and $PM_{2.5}$, while these variables were all

323 positively associated with COVID-19 incidence and mortality. Further, sensitivity analyses
324 showed that associations were robust to exclusion of counties with 10 or fewer COVID-19 cases,
325 excluding all counties comprising the New York metropolitan area and additional adjustment for
326 physical distance closures.
327
We acknowledge that this study has several limitations. This is an ecological study with
aggregated data on county level. Ecological designs should not be used to make inferences about
individual risks even though they are valid for hypothesis-generating purposes. Although we
adjusted for several important confounders, such as days since first COVID-19 case reported and
days since stay-at-home order, it is possible that there is residual confounding by these factors.
Days since stay-at-home order is based on the start date of the issuance of the order. However, in
several states the stay-at-home order was ended/relaxed in (the end of) April or May (earlier than
June 7). Further, there are other state-level physical distance closures (e.g. day cares, K-12
schools, gyms) that we did not take into account. As additional adjustment for days since non-
essential business closure and days since nursing home visitor ban did not affect our
associations, we do not think that adjustments for additional closures would greatly impact our
findings. We also note that physical distance closures and face coverings requirements could
differ between counties within a state. We used a county-level vegetation index as a proxy for
green space access, which does not distinguish whether vegetation represents urban parks,
forests, agricultural land, or overgrown vacant lots. Further detail on park amenities, vegetation
species or typology, and park usage during the COVID-19 pandemic were unavailable at the
time of data collection, but would add to future analyses. Another major limitation is the
underreporting of COVID-19 cases and deaths. Widespread testing has been limited in most
areas of the US and differences in testing availability might differ between counties and could
have changed over time due to additional resources and increased recognition of the disease.
348
Despite these limitations, our findings suggest that exposures to NDVI had beneficial impacts on
county-level incidence of COVID-19 in the US and may have reduced county-level COVID-19
mortality rates, especially in areas of higher population density. Although casual relationships
cannot be drawn from ecological studies, our findings imply that keeping parks open,
maintaining funding for parks in light of coming surges of COVID-19 and future pandemics may
have important public health benefits.

References

Abatzoglou JT. 2013. Development of gridded surface meteorological data for ecological
applications and modelling. Int J Climatol 33:121–131; doi:10.1002/joc.3413.

Banay RF, James P, Hart JE, Kubzansky LD, Spiegelman D, Okereke OI, et al. 2019. Greenness
and Depression Incidence among Older Women. ehp.niehs.nih.gov 127;
doi:10.1289/EHP1229.

Beveridge C, Hoffman C. 1997. The Papers of Frederick Law Olmsted: Writings on Public Parks,
Parkways, and Park Systems. Supplementary Series. Vol. 1.

Bezold CP, Banay RF, Coull BA, Hart JE, James P, Kubzansky LD, et al. 2018. The Association
Between Natural Environments and Depressive Symptoms in Adolescents Living in the
United States. J Adolesc Heal 62:488–495; doi:10.1016/j.jadohealth.2017.10.008.

CDC (Centers for Disease Control and Prevention). Coronavirus Disease 2019 (COVID-19) | 2020
Interim Case Definition, Approved April 5, 2020. Available:
https://wwwn.cdc.gov/nndss/conditions/coronavirus-disease-2019-covid-19/case-
definition/2020/ [accessed 16 June 2020].

Crouse DL, Pinault L, Balram A, Hystad P, Peters PA, Chen H, et al. 2017. Urban greenness and
mortality in Canada's largest cities: a national cohort study. Lancet Planet Heal 1:e289–
e297; doi:10.1016/S2542-5196(17)30118-3.

Delamater PL, Street EJ, Leslie TF, Yang YT, Jacobsen KH. 2019. Complexity of the basic
reproduction number (R0). Emerg Infect Dis 25:1–4; doi:10.3201/eid2501.171901.

Dong E, Du H, Gardner L. 2020. An interactive web-based dashboard to track COVID-19 in real
time. thelancet.com; doi:10.1016/S1473-3099(20)30120-1.

Fong KC, Hart JE, James P. 2018. A Review of Epidemiologic Studies on Greenness and Health:
Updated Literature Through 2017. Curr Environ Heal reports 5:77–87; doi:10.1007/s40572-
018-0179-y.

Heederik DJJ, Smit LAM, Vermeulen RCH. 2020. Go slow to go fast: A plea for sustained
scientific rigor in air pollution research during the COVID-19 pandemic. Eur Respir J
2001361; doi:10.1183/13993003.01361-2020.

James P, Banay RF, Hart JE, Laden F. A Review of the Health Benefits of Greenness. Springer;
doi:10.1007/s40471-015-0043-7.

Johns Hopkins Coronavirus Resource Center. 2020. COVID-19 United States Cases by County.
Available: https://coronavirus.jhu.edu/us-map [accessed 17 July 2020].

Kaplan R & Kaplan S. 1989. The Experience of Nature: A Psychological Perspective - Rachel
Kaplan, Stephen Kaplan - Google Books. Available:
https://books.google.com/books?hl=en&lr=&id=7l80AAAAIAAJ&oi=fnd&pg=PR7&dq=The+
Experience+of+Nature:+A+psychological+perspective.+&ots=TpQ-
NDt34l&sig=4s4Rdvae2yJ5xlXfpe1nRscE_Wk#v=onepage&q=The Experience of Nature%3A
A psychological perspective.&f=false [accessed 17 July 2020].

NASA. 2020. Measuring Vegetation (NDVI & EVI). Available:
https://earthobservatory.nasa.gov/features/MeasuringVegetation/measuring_vegetation_
2.php [accessed 16 June 2020].

Package "gamm4". 2020. https://cran.r-project.org/web/packages/gamm4/gamm4.pdf

Package "lme4." 2020. https://cran.r-project.org/web/packages/lme4/lme4.pdf

Prem K, Liu Y, Russell TW, Kucharski AJ, Eggo RM, Davies N, et al. 2020. The effect of control

strategies to reduce social mixing on outcomes of the COVID-19 epidemic in Wuhan,
China: a modelling study. Lancet Public Heal 5:e261–e270; doi:10.1016/S2468-
2667(20)30073-6.

Qian H, Miao T, Liu L, Zheng X, Luo D, Li Y. 2020. Indoor transmission of SARS-Cov-2.
medrxiv.org; doi:10.1101/2020.04.04.20053058.

Raifman J, Nocka K, Jones D, Bor J, Lipson S, Jay J. 2020. COVID-19 US state policy database.

Raines KS, Doniach S, Bhanot G. 2020. The transmission of SARS-CoV-2 is likely comodulated by
temperature and by relative humidity.; doi:10.1101/2020.05.23.20111278.

Sohrabi C, Alsafi Z, O'Neill N, Khan M, Kerwan A, Al-Jabir A, et al. 2020. World Health
Organization declares global emergency: A review of the 2019 novel coronavirus (COVID-
19). Int J Surg 76:71–76; doi:10.1016/j.ijsu.2020.02.034.

Tammes P. 2020. Social distancing, population density, and spread of COVID-19 in England: a
longitudinal study. BJGP open; doi:10.3399/bjgpopen20X101116.

US census bureau. 2020. Geography Program. Available: https://www.census.gov/programs-
surveys/geography.html [accessed 16 June 2020].

USGS (United States Geological Survey). 2020. Landsat 8 Data Users Handbook. Available:
https://www.usgs.gov/media/files/landsat-8-data-users-handbook [accessed 16 June
2020].

Ulrich RS. 1984. View through a window may influence recovery from surgery. Science (80-)
224:420–421; doi:10.1126/science.6143402.

Van Donkelaar A, Martin R V., Li C, Burnett RT. 2019. Regional Estimates of Chemical
Composition of Fine Particulate Matter Using a Combined Geoscience-Statistical Method
with Information from Satellites, Models, and Monitors. Environ Sci Technol 53:2595–
2611; doi:10.1021/acs.est.8b06392.

Vienneau D, Hoogh K de, Faeh D, Kaufmann M, Wunderli JM, Röösli M. et al. 2017. More than
clean air and tranquillity: residential green is independently associated with decreasing
mortality. Elsevier.

WHO 2020a. Coronavirus Disease (COVID-19) Situation Reports. Available:
https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports/
[accessed 20 July 2020].

WHO 2020b. WHO Director-General's opening remarks at the media briefing on COVID-19 - 11
March 2020. Available: https://www.who.int/dg/speeches/detail/who-director-general-s-
opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020 [accessed 17 July
2020].

Wu X, Nethery RC, Sabath B, Braun D, Dominici F, James C. 2020. Exposure to air pollution and
COVID-19 mortality in the United States: A nationwide cross-sectional study.

457 **Table 1. Descriptive statistics of the full cohort (n = 3,089 U.S. counties) [a].**
458

Variable	Median (IQR)
COVID-19 death rate (per 100,000)	2.8 (13.4)
COVID-19 incidence rate (per 100,000)	163.5 (330.2)
NDVI (April-May 2020)	0.44 (0.27)
Population density (person/sq. mi.)	60.6 (208.7)
% in poverty	9.2 (6.1)
% owner occupied housing	76.7 (9.3)
% less than high school education	19.1 (13.2)
% Black	1.4 (7.8)
% Hispanic	3.1 (5.8)
% 65 years of age	15.6 (5.0)
% 45-64 years of age	26.5 (2.9)
% 15-44 years of age	37.9 (5.4)
Median home value ($1,000)	110.3 (67.7)
Median household income ($1,000)	47.7 (15.2)
% Obese	33.1 (7.2)
% Smoke	17.0 (4.8)
Days since stay-at-home order	68 (74)
Days since non-essential businesses closure	69 (74)
Days since nursing homes visitor ban	62 (83)
Days since first case	74 (13)
Rate of hospital beds (per 100,000)	50 (173)
Rate of tests (per 100,000)	5670.7 (2394.6)
Average summer temperature (K)	303.3 (5.0)
Average winter temperature (K)	280.2 (10.5)
Average summer relative humidity (%)	91.3 (6.7)
Average winter relative humidity (%)	88 (5.6)
$PM_{2.5}$ ($\mu g/m^3$)	8.8 (4.1)
Urban counties [NCHS classification ≤4 (n)]	1149
Counties with issuance of stay-at-home order (n)	2196
Counties with 10< cases (n)	2297

459
460
461 *[a] The number of COVID-19 cases and deaths are based on data from March 22, 2020 through*
462 *June 7, 2020.*
463
464
465
466

Figure 1. Maps of the US that show (A) the county-level number of COVID-19 cases per 1 million population in the United States up to and including June 7, 2020, (B) the county-level number of COVID-19 deaths per 1 million population in the United States up to and including June 7, 2020, and (C) county-level average NDVI (April-May 2020).

Figure 2. Exposure-response curves of the association of NDVI with COVID-19 incidence and COVID-19 mortality in the full cohort, in urban counties and in rural counties.

Figure 3. Associations of NDVI with COVID-19 incidence and COVID-19 mortality by strata. Main = main model, quintile 1 = lowest quintile, quintile 5 = highest quintile. Home value = median home value, Pop. density = population density, no stay-at-home = counties with no issuance of stay-at-home order, stay-at-home = counties with issuance of stay-at-home order.