

Covid-19 Social Distancing Interventions by State Mandate and their Correlation to Mortality in the United States

Short Title: Covid-19 Social Mandate Correlation to Mortality in the USA

Authors:

Sean McCafferty, MD, Arizona Eye Consultants, 6422 E. Speedway Blvd. Tucson, AZ 85710 (520) 327-3487 Corresponding author SJMccafferty66@hotmail.com

Sean Ashley, M.S. Intuor Technologies 6422 E. Speedway Blvd. Tucson, AZ 85710 (520) 325-9400 Sashley@email.arizona.edu

Précis:

The mean-effects of Covid-19 mandated social interventions were found to have no affect the maximum slope of the daily mortality nor the overall mortality but did correlate to the population density.

Abstract:

Background: Evaluate the correlation between U.S. state mandated social interventions and Covid-19 mortality using a retrospective analysis of Institute for Health Metrics and Evaluation (IHME) data.

Methods: Twenty-seven (27) states in the United States were selected on June 17, 2020 from IHME data which had clearly defined and dated establishment of statewide mandates for social distancing measures to include: School closures, Prohibition on mass gatherings, business closures, stay at home orders, severe travel restrictions, and closure of non-essential businesses. The state Covid-19 mortality prevalence was defined as total normalized deaths to the peak daily mortality rate. The state mortality prevalence was correlated to the total number of mandates-days from their date of establishment to the peak daily mortality date. The slope of the maximum daily mortality rate was also correlated to mandate-days.

Results: The standardized mortality per state to the initial peak mortality rate did not demonstrate a discernable correlation to the total mandate days ($R^2 = 0.000006$, $p = 0.995$). The standardized peak mortality rate per state suggested a slight correlation to the total mandate days ($R^2 = 0.053$, $p = 0.246$), but was not statistically significant. There was a significant correlation between standardized mortality and state population density ($R^2 = 0.524$, $p = 0.00002$).

Conclusions: The analysis appears to suggest no mandate effective reduction in Covid-19 mortality nor a reduction in Covid-19 mortality rate to its defined initial peak when interpreting the mean-effect of the mandates as present in the data. A strong correlation to population density suggests human interaction frequency does affect the total mortality and maximum mortality rate.

Trial Registration: Public data is not patient specific and made available for public download on IHME Websites which can be used, shared, modified or built upon by non-commercial users via the Creative Commons Attribution-NonCommercial 4.0 International License. <https://creativecommons.org/licenses/by-nc/4.0/>

Key words: Covid-19, social distancing, mandates, mortality, mortality rate, United States

Background:

Socially distancing policy has been theorized to effectively reduce the rate of transmission of contagious diseases.^{1,2,3} Mandating social distancing state-wide within the United States has been instituted to varying degrees during the Covid-19 pandemic. Reducing the maximum contagion transmission by mandating decreased social interactions has been theorized to reduce mortality by allowing for social/medical mobilization and proper resource allocation.^{3,4,5} Verification of this theoretical model has not been demonstrated, except analyses demonstrating marginal social distancing effectiveness in delaying the peak infection.⁶ The extensive global incidence of the Covid-19 pandemic and unique real-time record keeping presents an opportunity to evaluate the social distancing theory over a large population with considerably different social distancing measures instituted.

Methods:

The study was conducted using the Institute for Health Metrics and Evaluation (IHME) openly published data on Covid-19 infections by individual states in the United States, to include daily infections/deaths as well as onset and discontinuation dates of state mandated social interaction interventions. As this was the primary source of information used for predictive modeling and setting public policy, it was chosen for its accuracy and regular updates.⁷

The methods utilized are an extension of the recently submitted manuscript evaluating European countries in the same manner. The methods are discussed extensively in the available preprint manuscript: [“Covid-19 Social Distancing Interventions by State Mandate and Their Correlation to Mortality”](#).⁸

The purpose was to quantify the decrease in the infection/mortality or decrease in the maximum infection/mortality rate as a result of defined state mandated social distancing measures. Mortality was chosen to define endpoint peaks and rates of change over registered infections within a state. Covid-19 registered infections are beset by inaccuracies due to: Inaccurate testing, testing frequency, asymptomatic patients, test availability, and regional variations in testing criteria. Individual states routinely record the cause and time of death. The analyses account for the state’s stage in the epidemiological infectious cycle and to extend those techniques to the states in the USA while additionally accounting for population density.

IHME data accessed on June 17, 2020 at 1900 EST was used to select all U.S. states with more than a maximum mortality rate of 10 Covid-19 deaths per day. Each of the selected states independently instituted preventive social interaction law on a given date to include one of the six (6) possible mandates.

Considered forms of social mandates:

1. Public School Closures
2. Social Gathering Restrictions
3. Stay-at-Home Orders
4. Business Closures
5. Non-Essential Business Closures
6. Severe Travel Restrictions

States which did not maintain their state-wide mandated social distancing through the end of the examination period (initial peak mortality rate) were excluded. Twenty-Seven (27) States were included as listed in figure 1. All states mandated social distancing universally across their respective territory of the 6 variations listed above on a specific date provided by the IHME data set accessed on June 17, 2020 and maintained them through their initial peak infection rate.

Figure 1 captures population density, social mandate form and duration, and Covid-19 mortality and maximum mortality rate data on the 27 states of the USA included in the study. Covid-19 data was obtained from IHME.⁷ State median age data is obtained from StatsAmerica’s USA States in Profile website for median age in 2019.⁹ State population density was obtained from the United States Census Bureau 2010 summary.¹⁰

State	Date of Peak Death Rate	Mortality at Peak (deaths / million)	Peak Mortality Rate (deaths/million/day)	School Closure		Gathering Restrictions		Stay-at-Home Orders		Business Closures		Non-Essential Business Closures		Travel Restrictions		Total Mandate Days (days)	Covid-19 Data Sources	Average Population Density (persons / sq. mi.)	Median Age (years)
				Date Start	Mandate Days Before Peak	Date Start	Mandate Days Before Peak	Date Start	Mandate Days Before Peak	Date Start	Mandate Days Before Peak	Date Start	Mandate Days Before Peak	Date Start	Mandate Days Before Peak				
Arizona	14-May	83.1	2.34	19-Mar	59	30-Mar	45	30-Mar	45	30-Mar	45	30-Mar	45	N/A	0	230	IHME	64.08	38.2
California	26-Apr	43.5	2.84	19-Mar	38	11-Mar	46	19-Mar	38	19-Mar	38	19-Mar	38	N/A	0	198	IHME	253.64	37
Colorado	23-Apr	98.1	4.74	23-Mar	31	19-Mar	35	26-Mar	28	17-Mar	37	26-Mar	28	N/A	0	159	IHME	55.56	37.1
Connecticut	22-Apr	413	28.1	17-Mar	36	12-Mar	41	N/A	0	16-Mar	37	23-Mar	30	N/A	0	144	IHME	736.27	41.1
Florida	18-Apr	32.9	2.16	17-Mar	32	3-Apr	15	3-Apr	15	17-Mar	32	N/A	0	N/A	0	94	IHME	399.96	42.5
Georgia	18-Apr	61.8	3.45	18-Mar	31	24-Mar	25	6-Apr	15	24-Mar	25	N/A	0	N/A	0	96	IHME	184.61	37.1
Illinois	11-May	279.5	8.29	17-Mar	55	13-Mar	59	21-Mar	51	16-Mar	56	21-Mar	51	N/A	0	272	IHME	228.24	38.6
Indiana	22-Apr	102.1	3.15	19-Mar	34	12-Mar	41	25-Mar	28	16-Mar	37	24-Mar	29	N/A	0	169	IHME	187.91	37.9
Iowa	11-May	116.6	4.46	4-Apr	37	17-Mar	55	N/A	0	17-Mar	55	N/A	0	N/A	0	202	IHME	56.48	38.5
Louisiana	12-Apr	191.7	13.6	16-Mar	27	13-Mar	30	23-Mar	20	17-Mar	26	22-Mar	21	N/A	0	124	IHME	107.60	37.5
Maryland	23-Apr	133.8	9.26	16-Mar	38	16-Mar	38	30-Mar	24	16-Mar	38	23-Mar	31	N/A	0	169	IHME	622.80	39.1
Massachusetts	26-Apr	410	25.1	17-Mar	40	13-Mar	44	N/A	0	17-Mar	40	24-Mar	33	N/A	0	157	IHME	883.65	39.6
Michigan	15-Apr	195.9	14.4	16-Mar	30	13-Mar	33	24-Mar	22	16-Mar	30	23-Mar	23	N/A	0	138	IHME	176.64	39.9
Minnesota	11-May	103.7	4.05	18-Mar	54	28-Mar	44	28-Mar	44	17-Mar	55	N/A	0	N/A	0	197	IHME	70.83	38.3
Mississippi	18-May	178.3	5.25	19-Mar	60	24-Mar	55	27-Apr	21	24-Mar	55	27-Apr	21	N/A	0	212	IHME	63.43	38
Missouri	24-Apr	43.6	2.45	23-Mar	32	23-Mar	32	6-Apr	18	23-Mar	32	N/A	0	N/A	0	114	IHME	89.28	38.9
New Jersey	18-Apr	445.7	32.2	18-Mar	31	16-Mar	33	23-Mar	28	16-Mar	33	21-Mar	28	N/A	0	153	IHME	1207.77	40.1
New York	11-Apr	568.1	50.8	18-Mar	24	12-Mar	30	22-Mar	20	16-Mar	26	22-Mar	20	N/A	0	120	IHME	412.80	39.2
North Carolina	11-May	54.5	1.63	14-Mar	58	14-Mar	58	3-Apr	38	17-Mar	55	3-Apr	38	N/A	0	247	IHME	215.72	39.1
Ohio	11-May	117.4	3.76	16-Mar	56	12-Mar	60	23-Mar	49	15-Mar	57	30-Mar	42	N/A	0	264	IHME	286.07	39.5
Pennsylvania	30-Apr	191	9.92	17-Mar	44	1-Apr	29	4-Apr	26	18-Mar	43	23-Mar	38	N/A	0	180	IHME	286.12	40.8
Rhode Island	18-May	402.9	13.4	16-Mar	63	17-Mar	62	7-Apr	41	17-Mar	62	N/A	0	N/A	0	228	IHME	1024.72	40.1
South Carolina	1-May	46.9	1.92	16-Mar	46	18-Mar	44	7-Apr	24	18-Mar	44	N/A	0	N/A	0	158	IHME	171.28	39.9
Texas	8-May	34.8	1.05	19-Mar	50	21-Mar	48	9-Apr	29	21-Mar	48	N/A	0	N/A	0	175	IHME	111.00	35
Virginia	8-May	91.4	3.31	16-Mar	53	15-Mar	54	30-Mar	39	17-Mar	52	24-Mar	45	N/A	0	243	IHME	216.14	38.6
Washington	8-Apr	59	2.63	13-Mar	26	11-Mar	28	23-Mar	16	16-Mar	23	25-Mar	14	N/A	0	107	IHME	114.59	37.8
Wisconsin	11-Apr	22.8	1.86	18-Mar	24	17-Mar	25	25-Mar	17	17-Mar	25	25-Mar	17	N/A	0	108	IHME	107.51	39.8

Figure 1: Table of population density, median age, social mandate, and Covid-19 mortality data on 27 states of the USA used in this analysis. Fields with “N/A” indicate the state did not implement that particular form of social mandate.

The defined endpoints for the analysis included the date of the initial peak mortality in deaths per day for each state. A state’s peak mortality was defined as the highest recorded daily deaths over a seven-day moving-average which was followed by a seven-or-more day decline in mortality with no other discernable peaks (using the same criteria) at the time of accessing the data. The maximum daily mortality rate was used as an easily defined universal milestone in any infectious disease progression to examine the total viral mortality up to the maximum mortality rate.

Additionally, the maximum slope of the Covid-19 mortality was determined by evaluating the maximum of the derivative of the mortality curve. Specifically, this slope was defined as the total recorded mortality five days after the peak-mortality-rate minus the total recorded mortality five days before the peak-mortality-rate divided by the 10-day interval. Both the mortality-at-peak and peak-mortality-rate were normalized by dividing by the population of the selected state.

Equation 1: Formula to calculate the estimate for the peak-mortality-rate.

$$\text{Peak Mortality Rate Estimate} = \frac{(\text{Mortality 5 Days After Peak}) - (\text{Mortality 5 Days Before Peak})}{10 \text{ days}}$$

All data was analyzed with Matlab and Microsoft Excel.¹¹ General linear mixed effects (GLME) was used to examine combined effects of the multiple variables. Specifically, Microsoft Excel was used for the bivariate linear regressions and ANOVA summaries for both mortality-at-peak and peak-mortality-rate while Matlab was used to generate the 95% confidence interval bounds on those linear regressions as well as the separate multivariate GLME analyses.

The clinical study was conducted within the ethical principles contained in Declaration of Helsinki, Code of Federal Regulations (CRF), Obligations of Clinical Investigators (21 CFR 812). All data was public and anonymous so no IRB was needed.

Similar to the previous study on European countries, the study conducted two high-level bivariate analyses on this data: one for each output of interest – peak-mortality-rate (PMR) and mortality-at-peak (MAP) – against the *total mandate days*.⁸ The *total mandate days* is defined the same way as it was in the previous study which is the summation from each form of social-mandate of the total number of days prior to the initial peak that form of social-mandate was implemented.⁸

Equation 2: Formula on how Total Mandate Days are calculated from each form of social-mandate and the number of days prior to the initial peak the mandates were implemented for.

$$\begin{aligned} \text{Total Mandate Days} = & \text{Days}_{\text{PriorToPeak}}^{\text{SchoolClosure}} + \text{Days}_{\text{PriorToPeak}}^{\text{Gatherings}} + \text{Days}_{\text{PriorToPeak}}^{\text{StayAtHome}} \\ & + \text{Days}_{\text{PriorToPeak}}^{\text{Business}} + \text{Days}_{\text{PriorToPeak}}^{\text{NonEssentBus}} + \text{Days}_{\text{PriorToPeak}}^{\text{TravelRestrict}} \end{aligned}$$

After the bivariate analyses against total mandate days, we then conducted bivariate analyses with the same response variables against population density instead. After the bivariate analyses, two multivariate analyses were conducted. These multivariate analyses used the same response variables as in the previous bivariate studies.

Equation 3: GLME model for peak-mortality-rate studied in the multivariate analysis.

$\text{Peak Mortality Rate} = \mu_{PMR}$	Overall mean peak mortality rate
$+ \alpha_{\text{StayAtHome}} \cdot \text{Days}_{\text{PriorToPeak}}^{\text{StayAtHome}}$	Fixed-effect from stay-at-home orders
$+ \alpha_{\text{SchoolClosure}} \cdot \text{Days}_{\text{PriorToPeak}}^{\text{SchoolClosure}}$	Fixed-effect from school closure
$+ \alpha_{\text{Gatherings}} \cdot \text{Days}_{\text{PriorToPeak}}^{\text{Gatherings}}$	Fixed-effect from social gathering restrictions
$+ \alpha_{\text{Business}} \cdot \text{Days}_{\text{PriorToPeak}}^{\text{Business}}$	Fixed-effect from general business closure
$+ \alpha_{\text{NonEssentBus}} \cdot \text{Days}_{\text{PriorToPeak}}^{\text{NonEssentBus}}$	Fixed-effect from non-essential business closure
$+ \alpha_{\text{TravelRestrict}} \cdot \text{Days}_{\text{PriorToPeak}}^{\text{TravelRestrict}}$	Fixed-effect from travel restrictions
$+ \alpha_{\text{GeneVar}} \cdot \text{PopDens}$	Fixed-effect from population density
$+ \alpha_{\text{MedAge}} \cdot \text{MedianAge}$	Fixed-effect from median age
$+ \epsilon_{PMR}$	Random Gaussian error

Equation 4: GLME model for mortality-at-peak studied in the multivariate analysis.

$Mortality\ at\ Peak = \mu_{MAP}$	Overall mean mortality at peak
$+ \beta_{StayAtHome} \cdot Days_{PriorToPeak}^{StayAtHome}$	Fixed-effect from stay-at-home orders
$+ \beta_{SchoolClosure} \cdot Days_{PriorToPeak}^{SchoolClosure}$	Fixed-effect from school closure
$+ \beta_{Gatherings} \cdot Days_{PriorToPeak}^{Gatherings}$	Fixed-effect from social gathering restrictions
$+ \beta_{Business} \cdot Days_{PriorToPeak}^{Business}$	Fixed-effect from general business closure
$+ \beta_{NonEssentBus} \cdot Days_{PriorToPeak}^{NonEssentBus}$	Fixed-effect from non-essential business closure
$+ \beta_{TravelRestrict} \cdot Days_{PriorToPeak}^{TravelRestrict}$	Fixed-effect from travel restrictions
$+ \beta_{GeneVar} \cdot PopDens$	Fixed-effect from population density
$+ \beta_{MedAge} \cdot MedianAge$	Fixed-effect from median age
$+ \varepsilon_{MAP}$	Random Gaussian error

These models were then regressed onto the data from **Figure 1**. Note that where certain social distancing mandates were not implemented, the “Days Prior to Peak” values were coded as zero in modeling to avoid singular data matrices during the regression analysis. The travel restrictions mandate was removed from the analysis since no states implemented strict travel restrictions in the time of this study according to **Figure 1**.


Results:

Results from Bivariate Analysis, MAP vs Total Mandate Days:

Twenty-Seven states were selected June 17, 2020 to examine their collective correlation between standardized mortality and total mandate-days of state directed social distancing directives. All states maintained social distancing directives through the study endpoint.⁷ States were found to have a significant diversity in total mandated intervention over time (mean = 171 total-mandate-days, std-dev = 50 total-mandate-days). The total population studied was 292 million. All states had statistically similar age distributions (mean = 38.6 years-old, std-dev = 1.5 years-old). The population density varied significantly (mean = 304 Pop./square mile, std-dev = 306 Pop./square mile)

Results for the bivariate analysis of mortality-at-peak against total mandate days is captured in **Figure 2**. The correlation of the standardized mortality-at-peak with total-mandate-days of social-distancing mandates prior to the peak was found to be statistically insignificant ($R^2 = 2E-06$ with $p\text{-val} = 0.9946$).


Figure 2: Standardized Covid-19 mortality on the day of peak-mortality-rate correlated to days under state-mandated social distancing directives prior to the peak.



Results from Bivariate Analysis, PMR vs Total Mandate Days:

Results for the bivariate analysis of peak-mortality-rate against total mandate days is captured in **Figure 3**. The correlation of the standardized peak-mortality-rate with total-mandate-days of social-distancing mandates prior to the peak was found to be statistically insignificant ($R^2 = 0.0534$ with $p\text{-val} = 0.2463$).


Figure 3:-Standardized Covid-19 peak-mortality-rate correlated to days under state-mandated social distancing directives prior to the peak.



Results from Bivariate Analysis, MAP vs Population Density:

Results for the bivariate analysis of mortality-at-peak against average population density is captured in **Figure** . The correlation of the mortality-at-peak with average population density was found to be statistically significant ($R^2 = 0.5238$ with $p\text{-val} = 1.99E-05$).


Figure 4: Standardized Covid-19 mortality on the day of peak-mortality-rate correlated to state average.



Results from Bivariate Analysis, PMR vs Population Density:

Results for the bivariate analysis of peak-mortality-rate against average population density is captured in **Figure 4**. The correlation of the standardized peak-mortality-rate with total-mandate-days of social-distancing mandates prior to the peak was found to be statistically significant ($R^2 = 0.3814$ with $p\text{-val} = 0.0006$).

Figure 4: Standardized Covid-19 peak-mortality-rate correlated to state average population density.



Results from Multivariate Analysis for MAP:

Results for the multivariate analysis of mortality-at-peak on the date of initial peak-mortality-rate are captured in **Figure 5**. In this study, only one modeled effect was found to be statistically significant at the 5% significance level; this was a state's average population density (p -val = 0.0004).

Figure 5: Matlab console output from GLME regression observing mortality-at-peak as the response variable using the mathematical model from Equation 4.

Generalized linear mixed-effects model fit by ML

Model information:

Number of observations	27
Fixed effects coefficients	8
Random effects coefficients	0
Covariance parameters	1
Distribution	Gaussian
Link	Identity
FitMethod	Laplace

Formula:

Linear Mixed Formula with 7 predictors.

Model fit statistics:

AIC	BIC	LogLikelihood	Deviance
341.88	353.55	-161.94	323.88

Fixed effects coefficients (95% CIs):

Name	Estimate	SE	tStat	DF	pValue	Lower	Upper
'(Intercept)'	-171.45	689.16	-0.24879	19	0.80619	-1613.9	1271
'Days_SchoolClosure'	-1.7793	4.4186	-0.40269	19	0.69167	-11.028	7.469
'Days_Gatherings'	4.4052	4.4709	0.98532	19	0.33685	-4.9524	13.763
'Days_StayAtHome'	-0.56839	1.8435	-0.30833	19	0.76119	-4.4268	3.29
'Days_BusinessClosure'	-2.7449	5.6195	-0.48847	19	0.63081	-14.507	9.0167
'Days_NonEssentBus'	0.72053	1.3313	0.54123	19	0.59464	-2.0659	3.5069
'AvgPopDens'	0.32657	0.075464	4.3274	19	0.00036298	0.16862	0.48452
'MedianAge'	6.1787	17.807	0.34698	19	0.73242	-31.092	43.45

Random effects covariance parameters:

Group: Error

Name	Estimate
'sqrt(Dispersion)'	97.407

Results from Multivariate Analysis for PMR:

Results for the multivariate analysis of peak-mortality-rate are captured in **Figure 6**. In this study, only one modeled effect was found to be statistically significant at the 5% significance level; this was a state's average population density (p-val = 0.0025).

Figure 6: Matlab console output from GLME regression observing standardized peak-mortality-rate as the response variable using the mathematical model from **Equation 3**.

```

Generalized linear mixed-effects model fit by ML

Model information:
  Number of observations      27
  Fixed effects coefficients    8
  Random effects coefficients  0
  Covariance parameters       1
  Distribution                 Gaussian
  Link                         Identity
  FitMethod                    Laplace

Formula:
  Linear Mixed Formula with 7 predictors.

Model fit statistics:
  AIC      BIC      LogLikelihood      Deviance
  207.15   218.82   -94.577                189.15

Fixed effects coefficients (95% CIs):
  Name                Estimate      SE        tStat      DF      pValue      Lower      Upper
  '(Intercept)'      -5.2635     56.854    -0.092579  19      0.92721    -124.26    113.73
  'Days_SchoolClosure' -0.23278    0.36453   -0.63858   19      0.53072    -0.99576   0.53019
  'Days_Gatherings'   0.29588    0.36884    0.80219   19      0.43236    -0.47611   1.0679
  'Days_StayAtHome'  -0.028495   0.15208   -0.18736   19      0.85336    -0.34681   0.28982
  'Days_BusinessClosure' -0.35576    0.4636    -0.7674    19      0.45228    -1.3261    0.61456
  'Days_NonEssentBus' 0.067505    0.10983    0.61464    19      0.54608    -0.16237   0.29738
  'AvgPopDens'        0.021662    0.0062257 3.4795     19      0.0025095  0.0086316  0.034693
  'MedianAge'         0.49308     1.4691     0.33564    19      0.74082    -2.5817    3.5679

Random effects covariance parameters:
Group: Error
  Name                Estimate
  'sqrt(Dispersion)'  8.0359
  
```

Discussion:

The analysis appears to show no statistically significant reduction in neither the slope of the Covid-19 mortality rate ($p\text{-val} = 0.995$) nor in Covid-19 mortality ($p\text{-val} = 0.246$) to its defined initial infection peak. There was a significant correlation to both the overall mortality and the maximum mortality rate to state population density. The correlation to population density suggests that the proximity and frequency of social interactions directly affect the mortality rate and overall mortality. However, the evaluation of state mandates suggests that they were unable to reduce the frequency of social interactions to be effective. The results corroborate the findings of the previous study conducted using the same methodology with the IHME data for 12 European countries.⁸

One of the advantages in this analytical approach is in the design of the social-mandate input variables to be more robust to possible biases caused by different states being in different epidemiologic infectious stages of Covid-19. It is worth noting that quite a few of the states involved in this study have since experience a second wave of infections not included in this analysis. This may be a successive superposition of infectious cycles within the state in sequestered populations or an effect of behavioral changes within the population. As of the drafting of the manuscript on July 19, 2020, Six States; Arizona, California, Florida, Mississippi, South Carolina, Texas have had secondary peaks of maximum infection rates which eclipsed the primary peak used in this analysis. Only their initial peaks were the subject of this study and they were not revised to the new peak nor excluded. Mandates and social behaviors changed substantially during the interim between the first maximum and could confound the analysis. Furthermore, the strong correlation to population density demonstrating a likely effect of personal interaction frequency which assists in validating the methodology. It is also worth noting that start dates of social-mandate state implementation does not necessarily imply a certain degree of compliance of the population with enacted mandates.

Conclusions:

How early before the initial peak an investigated form of social-mandate was implemented by a USA state was found to be statistically insignificant in its effects on peak-mortality-rate and mortality-at-peak in both the bivariate studies and the multivariate studies. (The lowest associated $p\text{-value}$ with social-mandate terms across all studies above was $p\text{-val} > 0.2463$ in the bivariate study of studying standardized Covid-19 peak-mortality-rate against total-mandate-days.) However, the average population density was found to be a statistically significant factor for both peak-mortality-rate and mortality-at-peak of a state in both the bivariate and multivariate studies. (The largest associated $p\text{-value}$ with average population density across all studies above was $p\text{-val} = 0.0025$ in the multivariate study with peak-mortality-rate.)

What we could conclude from this study is that the timing of social-mandates alone is not enough to explain the variability in Covid-19 peak-mortality-rate and mortality-at-peak numbers between states. However, a state's average population density is a significant factor and should be accounted for in one way or another in current and future investigations to explain the variability in Covid-19 peak-mortality-rate and mortality-at-peak numbers between states.

Factors which influence the effectiveness of social distancing interventions include the reproduction number (R_0) of the virus, the mortality rate, and the mandate effectiveness of the isolation.^{12,13} A reduction of the slope of the maximum infection rate without a change in mortality is consistent with

theory of viral infectious epidemiology.¹⁴ Reductions in total mortality due to reduced healthcare system or advancements in medical care did not materialize in this analysis. However, the time from the probable first infection to the peak of the infection rate was likely only around 3 months. It is possible that if the viral mortality estimated at 0.6% to 1.2% or the viral reproduction R_0 , estimated between 2.8 and 3.3, were significantly higher may have produced a correlation.^{11,12} Mandated interventions which were significantly more isolating may have produced a correlation to the mortality.

List of Abbreviations:

- FDA
 - Food and Drug Administration
- CFR
 - Code of Federal Regulations
- IHME
 - Institute of Health Metrics and Evaluation
- GLME
 - General Linear Mixed-Effects
- PMR
 - Peak-mortality-rate due to Covid-19, standardized by state total population, as calculated from **Equation 1** using data from IHME
- MAP
 - Mortality-at-peak due to Covid-19, standardized by state total population as reported in referenced IHME data

Declarations:

Ethics approval and Consent to participate:

- This clinical study was conducted in accordance with the ethical principles contained within Declaration of Helsinki, Protection of Human Volunteers (21 CFR 50), Institutional Review Boards (21 CFR 56), and Obligations of Clinical Investigators (21 CFR 812). No IRB was required with the use of depersonalized public data.

Consent for publication:

- Not Applicable. Depersonalized data made available for download on IHME Websites can be used, shared, modified or built upon by non-commercial users via the Creative Commons Attribution-NonCommercial 4.0 International License. <https://creativecommons.org/licenses/by-nc/4.0/>

Availability of data and materials:

- Depersonalized data made available for download on IHME Websites can be used, shared, modified or built upon by non-commercial users via the Creative Commons Attribution-NonCommercial 4.0 International License. <https://creativecommons.org/licenses/by-nc/4.0/>

Funding:

- This study was not funded.

Competing interests:

- None.

Authors' Contributions:

- Sean McCafferty – Design, drafting, analysis, and interpretation
- Sean Ashley – Drafting, analysis, and interpretation

Acknowledgements:

- None.

References

- 1 Reluga T. Game theory of social distancing in response to an epidemic. *PLoS Comput Biol.* 2010;6(5):e1000793. doi:10.1371/journal.pcbi.1000793
- 2 Caley P, Philp D, McCracken K. Quantifying social distancing arising from pandemic influenza. *Journal of The Royal Society Interface* Volume 5, Issue 23 <https://doi.org/10.1098/rsif.2007.1197>
- 3 Kelso, J.K., Milne, G.J. & Kelly, H. Simulation suggests that rapid activation of social distancing can arrest epidemic development due to a novel strain of influenza. *BMC Public Health* 2009;9:117 <https://doi.org/10.1186/1471-2458-9-117>
- 4 Kissler S, Tedijanto C, Lipsitch M, Grad Y. Social distancing strategies for curbing the COVID-19 epidemic. doi: <https://doi.org/10.1101/2020.03.22.20041079>
- 5 Rashid H, Ridda I, King C, Begun M, Tekin H, Wood J, Booy R. Evidence compendium and advice on social distancing and other related measures for response to an influenza pandemic. *Paediatric Respiratory Reviews* 2015;16:1526-42 <https://doi.org/10.1016/j.prrv.2014.01.003>.
- 6 Marsland R, Mehta P. Data-driven modeling reveals a universal dynamic underlying the COVID-19 pandemic under social distancing. arXiv:2004.10666
- 7 Institute for Health Metrics and Evaluation (IHME) <http://www.healthdata.org/covid> June 17, 2020, July 19, 2020
- 8 McCafferty S, Ashley S. Covid-19 Social Distancing Interventions by State Mandate and Their Correlation to Mortality. *Research Square* (pre-print), June 1, 2020. [Online]. <https://www.researchsquare.com/article/rs-30828/v1>.
- 9 StatsAmerica, "USA States in Profile: Median Age in 2019," Indiana Business Research Center, 9 July 2020. [Online]. http://www.statsamerica.org/sip/rank_list.aspx?rank_label=pop46&ct=S09. [Accessed 9 July 2020].
- 10 United States Census Bureau, "United States Summary: 2010, Population and Housing Unit Counts, 2010 Census of Population and Housing," pp. V-2, 1& 41 (Tables 1 & 18), 2012. <https://www.census.gov/data/tables/2010/dec/density-data-text.html>

- 11 MATLAB. (2020). *Version 9.6.0.1072779 (R2019a)*. Natick, Massachusetts: The MathWorks Inc.
- 12 Natick, Massachusetts, 2020.11 Viceconte G, Petrosillo N. Covid-19 R0: Magic Number or Conundrum? *Infect Dis Rep.* 2020;12:8616
- 13 Russel T, Hallewell J, Jarvis C, et al. Estimating the Infection and Case Fatality Ratio (Covid-19) using Age Adjusted Data from the Outbreak on the Diamond Princess Cruise ship, February 2020. *Euro Surveill.* 2020;25:pii=2000256
- 14 Daley G, Gani J. *Epidemic Modeling: An Introduction*. NY: 2005: Cambridge University Press

Figures Legends:

Figure 1: Table of population density, median age, social mandate, and Covid-19 mortality data on 27 states of the USA used in this analysis. Fields with “N/A” indicate the state did not implement that particular form of social mandate.

Figure 2: Standardized Covid-19 mortality on the day of peak-mortality-rate correlated to days under state-mandated social distancing directives prior to the peak.

Figure 3: Standardized Covid-19 peak-mortality-rate correlated to days under state-mandated social distancing directives prior to the peak.

Figure 4: Standardized Covid-19 mortality on the day of peak-mortality-rate correlated to state average population density.

Figure 5: Standardized Covid-19 peak-mortality-rate correlated to state average population density.

Figure 6: Matlab console output from GLME regression observing mortality-at-peak as the response variable using the mathematical model from **Equation 4**.

Figure 7: Matlab console output from GLME regression observing standardized peak-mortality-rate as the response variable using the mathematical model from **Equation 3**.

Equation 1: Formula on how to calculate the estimate for the peak-mortality-rate.

Equation 2: Formula on how Total Mandate Days are calculated from each form of social-mandate and the number of days prior to the initial peak the mandates were implemented for.

Equation 3: GLME model for peak-mortality-rate studied in the multivariate analysis.

Equation 4: GLME model for mortality-at-peak studied in the multivariate analysis.