

Short communication

**A Comparison of Covid-19 Patient Characteristics Before versus After Partial Lockdown
in Vietnam**

Hoang-Long Vo^{1,2,*}, Hao Anh Si Nguyen¹, Thao Phuong Thi Nguyen², Khanh Ngoc Nguyen¹, Kim-Duy Vu¹, Vu Dam Trong Anh¹, Tham Thi Nguyen¹, Men Thi Hoang^{3,4}, Trang Huyen Thi Nguyen^{3,4,*}, Giang Thu Vu⁵, Huong Thi Le¹

¹Institute for Preventive Medicine and Public Health, Hanoi Medical University, Hanoi 100000, Vietnam

²Institute of Health Economics and Technology, Hanoi 100000, Vietnam

³Institute for Global Health Innovations, Duy Tan University, Da Nang 550000, Vietnam

⁴Faculty of Medicine, Duy Tan University, Da Nang 550000, Vietnam

⁵Center of Excellence in Evidence-based Medicine, Nguyen Tat Thanh University, Ho Chi Minh City 700000, Vietnam

*** Correspondence:**

Hoang-Long Vo, M.D., vohoanglonghmu@gmail.com or drvohoanglonghmu@daihocyhanoi.edu.vn

Trang Huyen Nguyen, M.Sc., trangbn.18@gmail.com or nguyenthuyentrang46@duytan.edu.vn

Word count: 2702

Running Title: Characteristics Before versus After Partial Lockdown

What is already known on this topic

- Most of Covid-19 confirmed patients in Vietnam were acquired overseas.
- In Vietnam, Covid-19 epidemic had the low estimated reproduction ratio (R_0) and SARS-CoV-2 spread was on the downward trend before July 25.

What this study adds

- Covid-19 confirmed patients were older in the time after lockdown end compared to in the period before lockdown end by a median of 5 years.
- All discharged Covid-19 patients and no Covid-19 death were in the phase before lockdown, while post lockdown period had still remained significant patients being under treatment, especially reported first fatalities.
- Continuous volume of people returning Vietnam from abroad through two stages during the Covid-19 epidemic.
- Our analysis might be considered as a prompt reference source for further in-depth surveys to understand the adaptive models of Covid-19 patients among different provinces.

0 **Abstract**

1 **Background.** Given the Covid-19' sudden rapid spread in the community since 25 July 2020, an
2 updated analysis of Covid-19 cases was conducted to examine the differences of characteristics
3 of Covid-19 patients before versus after partial lockdown end in Vietnam.

4 **Methods.** Data of 569 Covid-19 patients confirmed SARS-CoV-2 infection from 23 January to
5 31 July was collected from available official databases. We divided Covid-19 situation timeline
6 into two main periods, before lockdown end (23 January - 22 April) and after lockdown end (23
7 April - 31 July).

8 **Results.** We found significant variations in the distribution of Covid-19 patients among different
9 provinces between two periods. Covid-19 confirmed patients were older in the time after
10 lockdown end compared to in the period before lockdown end by a median of 5 years. All
11 discharged Covid-19 patients and no Covid-19 death were in the phase before lockdown, while
12 post lockdown period had still remained significant patients being under treatment, especially
13 reported first fatalities. The number of Covid-19 patients who returned from other countries
14 (excluding China) slightly increased through two stages ($p > 0.05$), partially showing that the
15 continuous volume of people returning Vietnam from abroad during the Covid-19 epidemic.

16 **Conclusions.** Our analysis indicated demographic and epidemiological disparity of Covid-19
17 patients before versus after loosening the national partial lockdown in Vietnam. It is important to
18 suggest that, proactive efforts in Covid-19 control after partial lockdown end will be effective
19 when the measures to closely control and monitor repatriation and immigration via the borders of
20 Vietnam are strictly enforced.

21 **Keywords:** SARS-CoV-2; Covid-19; patient characteristics; Comparison; Vietnam

22 **Introduction**

23 Since the beginning of the coronavirus disease 2019 (Covid-19) crisis, the lockdown has been
24 considered a key policy response in many countries to prevent the further spread of the severe
25 acute respiratory syndrome coronavirus 2 (SARS-CoV-2) which causes the disease. The
26 implementation of the lockdown order, however, has varied in countries and territories around
27 the world, thanks to the unpredictable situation of the pandemic [1]. Vietnam is one of the few
28 countries achieving effective control of Covid-19 outbreak after the implementation of lockdown
29 strategy [2]. Under Directive No. 16/CT-TTg issued in March 31, 2020, a “stay-at-home” order
30 was imposed nationwide to curb the spread of the contagion, limit the number of infections and
31 casualties, and alleviate the pressure on the health service providers. The effectiveness of such
32 outbreak control was evidenced by the fact that there were only 59 new infected cases confirmed
33 during the first 14-day nationwide social distancing (from April 1 to 14, 2020), equaled to 40%
34 of the two weeks prior to the national lockdown [3]. Of these, there were 30 confirmed cases in
35 isolation zones and 29 confirmed cases in the community [3]. Within 99 days since April 15,
36 Vietnam had confirmed no community transmission despite extensive testing. While in many
37 regions of the world strict Covid-19 lockdown orders had still been in effect, the Government of
38 Vietnam relaxed social distancing rules for almost all provinces and cities on 23 April [4]. The
39 early success in Vietnam’ epidemic control has mainly attributed to the national emergency
40 response across the whole sociopolitical system, in particular, 22-day nationwide lockdown.

41 However, the strike of no Covid-19 community transmission has been broken on July 25 when a
42 new case were reported in Da Nang City [5]. This disruption made the Covid-19 pandemic in
43 Vietnam more unforeseeable than ever [6], [7], [8]. Immediately, contact tracing efforts as well
44 as epidemiologic investigations have been strengthened under the strong guidance of the
45 Vietnamese Prime Minister, yet the source of the infection is not known in several new cases [8],

46 [9]. With the estimation of the reproduction ratio (R_0) for 257 first Covid-19 patients in Vietnam
47 (between 23/1/2020 and 10/4/2020), a Huy G. Nguyen' early analysis revealed the downward
48 trend of the epidemic in this country [10]. However, the potential SARS-CoV-2 transmission in
49 the community is not reflective of current reality in each stage under Government' strict
50 restrictions. An initial understanding of the characteristics of Covid-19 patients detected before
51 and after lockdown end in Vietnam is needed to suggest prompt and efficient actions in the
52 worst-case scenarios for in Vietnam. Given the Covid-19' rapid spread, an updated analysis of
53 cases was conducted to examine the differences of characteristics of Covid-19 patients before
54 versus after lockdown end in Vietnam.

55 **Methods**

56 In this paper, we collected data of Covid-19 patients who were confirmed SARS-CoV-2
57 infection from 23 January to 31 July, 2020. We divided Covid-19 situation timeline from 23
58 January to 31 July 2020 into two main periods, before lockdown end (23 January – 22 April,
59 2020) and after lockdown end (23 April – 31 July, 2020). We selected cut-off point of Covid-19
60 situation timeline in Vietnam on 23 April, 2020 (23 April, 2020 is the day that the Vietnam
61 Government decided to loosen the national lockdown [4]).

62 **Data sources**

63 Analysis data in this paper was extracted from two sources, official database of the Ministry of
64 Health of Vietnam (MOH) (<https://ncov.moh.gov.vn/>) and the website of Our World in Data
65 (<https://ourworldindata.org/coronavirus#coronavirus-country-profiles>) [11]. A total of 569
66 Covid-19 confirmed patients were consecutively detected from 23 January to 31 July, 2020
67 (Figure 1).

68 **Case definition for Covid-19**

69 *Covid-19 positive case* and *Covid-19 suspected case* were defined according to the Decision No.
70 322 / QD-BYT of February 6, 2020 of the Minister of Health for the period before lockdown end
71 [12] and according to the Decision 1344 / QD-BYT 2020 Guidance on diagnosis and treatment
72 of COVID-19 [13] for the period after lockdown. The Vietnam' tests were reported by the WHO
73 and MOH approved laboratories [12], [13].

74 *Covid-19 recovered case* before and after lockdown end was defined that, a patient who
75 recovered from Covid-19 and discharged from the health facility met the following criteria: 1) no
76 fever for at least three days; 2) a good general health condition; (3) two samples taken at least 1
77 day apart having negative test results for SARS-CoV-2 [12], [13].

78 *Covid-19 death case* is a patient with confirmed Covid-19 infection whose death resulted from
79 clinically compatible illness, unless there is an other certain cause of mortality that cannot be
80 related to Covid-19 disease (e.g., trauma). No period of full recovery was recorded between the
81 illness and death.

82 **Variables**

83 The variables in this analysis were selected on the basis of the availability of above official
84 databases. The variables related to the Covid-19 patient characteristics were divided into
85 subgroups as following: mean age, age group (0-10 years/11–20 years/21–30 years/31–40
86 years/41–50 years/51–60 years/60 years and over), gender (male/female), nationality
87 (Vietnam/others), the history of returning from China (yes/no), the history of returning from
88 other countries (excluding China) (yes/no), and treatment status (under treatment/ discharged/
89 died).

90 **Data analysis**

91 Vietnam' Covid-19 situation timeline was considered as a variable with two main values: (code
92 1) before lockdown end and (code 2) after lockdown end. We used both descriptive and
93 analytical method. While age data was expressed as mean \pm standard deviation (SD) with
94 interquartile ranges (IQR), categorical variables were presented as frequency with percentage
95 (%). We estimated the overall proportion of Covid-19 confirmed cases and its percentage before
96 and after lockdown end according to selected demographic and epidemiological characteristics.
97 Chi-square tests and Mann–Whitney U tests were utilized to compare patient' characteristics
98 between two periods. All statistical analyses in this paper were conducted by Stata® 15
99 (StataCorp LLC, USA). The level of statistical significance was set at 0.05.

100 **Results**

101 Among the 569 patients from 23 January to 31 July 2020, there were 268 patients by 22 April
102 and 301 patients between 23 April and 31 July (Figure 1). Figures 2 illustrated the significant
103 variation in the distribution of Covid-19 patient among different provinces between two periods.
104 The demographic and epidemiological characteristics of the patients are shown in Table 1.
105 Patients had a median age of 38.56 (SD 16.61) years, and the patients in the time after lockdown
106 end were older than those in the period before lockdown end by a median of 5 years. Overall,
107 those aged 21–30 years are recorded with the highest number of Covid-19 confirmed cases in
108 both periods (35.82% (n = 96) vs. 27.91% (n = 84)). Almost 90% (n = 241) of patients aged 59
109 years or younger before lockdown, while this figure after lockdown end was nearly 85% (n =
110 254). There was significant difference in the patient gender between two periods, 45.52% male
111 before lockdown and 61.79% male after lockdown.
112 Of 10 Covid-19 patients returned from China, 9 patients (3.36%) were in the first period and 1
113 (0.33%) was in the second period, while no difference between two periods was observed

114 according to the number of patients returned from other countries. Besides, the Covid-19 patient'
115 nationality between two periods was also significantly varied (Table 1).

116 Compared to the phase before lockdown with 268 discharged Covid-19 patients (100%) and no
117 Covid-19 death, post lockdown period had only 34.88% patients cured and discharged from the
118 facilities with 5 fatalities (1.66%) and 191 those being under treatment (63.46%) (Table 1).

Table 1. Characteristics of the Covid-19 patients detected before versus after lockdown end, 23 January to 31 July 2020 (N = 569)

Characteristics	All patients	Before lockdowns end	After lockdowns end	p
	N = 569 192 (Jan 23–Jul 31)	N = 268 91 days (Jan 23–Apr 22)	N = 301 101 days (Apr 23–Jul 31)	
Gender				<0,001*
Male	308 (54.13)	122 (45.52)	186 (61.79)	
Female	261 (45.87)	146 (54.48)	115 (38.21)	
Age, mean±SD (IQR)	38.56±16.61 (0 – 90)	35.89±15.93 (0 – 88)	40.93±16.86 (0 – 90)	<0,001**
Age group				<0,001*
0–10 years	13 (2.28)	6 (2.24)	7 (2.33)	
11–20 years	44 (7.73)	32 (11.94)	12 (3.99)	
21–30 years	180 (31.63)	96 (35.82)	84 (27.91)	
31–40 years	101 (17.75)	36 (13.43)	65 (21.59)	
41–50 years	86 (15.11)	44 (16.42)	42 (13.95)	
51–60 years	71 (12.48)	27 (10.07)	44 (14.62)	
60 years and over	74 (13.01)	27 (10.07)	47 (15.61)	
Nationality				<0,001*
Vietnamese	498 (87.52)	220 (82.09)	278 (92.36)	
Others	71 (12.48)	48 (17.91)	23 (7.64)	
Had returned from China				0.006*
Yes	10 (1.76)	9 (3.36)	1 (0.33)	
No	559 (98.24)	259 (96.64)	300 (99.67)	
Had returned from other countries (excluding China)				0.844
Yes	333 (58.52)	158 (58.96)	175 (58.14)	
No	236 (41.48)	110 (41.04)	126 (41.86)	
Treatment status				<0,001*
Under treatment	191 (33.57)	0 (0.00)	191 (63.46)	
Discharged	373 (65.55)	268 (100.00)	105 (34.88)	
Died	5 (0.88)	0 (0.00)	5 (1.66)	

* Chi-square test, ** Mann–Whitney U test




Figure 1. The incidence of confirmed Covid-19 cases and its cumulative numbers, 23 January to 31 July 2020

Discussion

At the time of writing this article, the first 569 consecutive confirmed cases of Covid-19 through 31 July 2020 are utilized to compare according to two stages, before and after lockdown end in Vietnam. Among the 569 patients, the a median age was 38.56 years, which was significantly lower compared to recent reports in other countries such as 62.2 years in 393 patients in New York City, U.S [14] and 50.1 years in 198 patients in Shanghai, China [15]. Age structure of each country may be the best explanation for the difference in the confirmed case number through two phases. The age distribution of cases was revealed with strong and important role on Covid-19 case fatality [16]. In Vietnam, as compared to before lockdown end, the stage after lockdown had a higher proportion of elder patients. The Covid-19 patient distribution in general is quite similar amongst age groups and those aged 21–30 years are recorded with the highest number of Covid-19 confirmed cases in both periods, implying that SARS-CoV-2 infection risk considerations are not focused solely on any age group. Those in later stage are also patients having pre-existing comorbidities, contributing to the partial explanation of 5 first fatality cases before 31 July. In this analysis, it was observed that significant difference in the gender was shown between two stages (45.52% male patients before lockdown end vs 61.79% male patients after lockdown end).

Despite the more and more increasing global Covid-19 patient number with the overload of the health system, Vietnam has still been responding well to the Covid-19 pandemic with the mobilization of the entire political system. Significant difference was observed in the nationality of Covid-19 patients between two phases. During the initial phase of the study timeline, 17.91% patients (n=48) were residents of other countries, which decreased to 7.64% patients (n=23) during phase after the lockdown end. This point is mainly because, in the time after lockdown

end, most of them are Vietnamese citizens flown home from abroad [17]. In our analysis, the Vietnamese cases increase in the community after lockdown may be associated with the reopening of domestic tourism in the “new normal” condition after the first Covid-19 wave and the illegal border entry [4], [18], [19]. Current situation requires that, Vietnam’ proactive efforts in Covid-19 control will be effective when volume, speed, and reach of travel in tourism are well managed and the measures to closely control and monitor repatriation and immigration via its borders are prioritized. Despite having a long borderline with China, Vietnam only recorded 9 confirmed patients returned from China at the time before lockdown end and only a case after lockdown who had illegally immigrated to Vietnam. On the other hand, we found, the number of Covid-19 patients who returned from other countries (excluding China) slightly increased through two stages, however, this difference was not statistically significant. This result partially shows that the continuous volume of people returning Vietnam from abroad during the Covid-19 epidemic. Such the detection of the Covid-19 cases from overseas is the effort of not only the entire political system in immigration control and management of people returning from abroad in general and but also the Vietnam Border Defense Force in particular. The more clear proof is that, over 16,000 people illegally entering Vietnam have been prevented by the Vietnam Border Defence Force since the beginning of 2020 and, particularly, over 2,400 people have been arrested in July once trespassing on the trail [19].

Our study showed that the difference in the treatment status of Covid-19 patients between before and after lockdown end. Vietnam’ early success in the Covid-19 patient treatment was proved by the absolute figure of 268 discharged Covid-19 patients (100%) before lockdown end. Although the number of the recovered patients, overall, continued to slightly increase from 23 April to 25 July after lockdown end, yet the number of new cases in this phase gradually decreased within

three months before Covid-19 outbreak point documented in the community in Da Nang on 25 July. Importantly, all new Covid-19 instances from 23 April to 25 July after lockdown were Vietnamese people who returned from abroad and this risk group was then required to isolate for 14 days [20]. When Vietnam was retaining 99 days without any cases in the community, various countries across the world have been faced a rapid growth in the number of new and fatal cases of Covid-19 [21]. Since 25 July after lockdown, almost patients being treatment in Vietnam have been confirmed SARS-CoV-2 infection, which known as the second wave of Covid-19. In such emerging Covid-19 situation, the community transmission cases reported to be linked to the hospitals in Da Nang where the elder patients with chronic severe diseases such as cancer, chronic kidney disease and dialysis, heart failure, diabetes, or hypertension were being treated [22]. Therefore, compared to the phase before lockdown end (all discharged Covid-19 patients (100%) and no death), the post-lockdown time recorded the worse situation with mostly patients being treated ($n = 191$, 63.46%) and first 5 fatalities (1.66%). It should be noted that five fatal cases were identified at the moment of Vietnam' second Covid-19 outbreak after the national lockdown implementation for three months. After synthesizing current analysis data, two main reasons associated with Vietnamese Covid-19 deaths were suggested by the Vietnamese experts. Firstly, the new coronavirus strain detected in second wave of Covid-19 infections there had a faster speed of infection compared to the earlier strain. However, many uncertainties have remained concerning the virus-host interaction and the evolution of the epidemic, and its harmfulness compared to the strain previously existed in Vietnam has been not yet known [22], [23]. Secondly, the presence of multiple coexisting severe illness was reported in Covid-19 fatalities [24].

The present results should be viewed with the limitations. First, because the paper' secondary analysis data was collected from two publicly available datasets, the information of the exposure history, clinical symptoms or signs, radiologic assessments, laboratory testing as well as the other risks could be unknown. Second, the challenges in national large-scale coronavirus testing resulted in the underestimated true number of Covid-19 infections. Finally, due the ongoing Covid-19 pandemic with the more complicated situation in Vietnam, the current estimates are temporary at the time of this writing.

Conclusions

Our short communication illustrated demographic and epidemiological disparity of Covid-19 patients before versus after loosening the national lockdown in Vietnam. The Covid-19 patient distribution in general is quite similar amongst age groups and these patients were largely recorded in the 21–30-year-old group in both periods. Most of the Covid-19 patients are residents of Vietnam and Vietnamese patient number has still increased rapidly during phase after the lockdown end. Importantly after the lockdown period, Vietnam' proactive efforts in Covid-19 control after lockdown end will be effective when the measures to closely control and monitor repatriation and immigration via its borders are strictly enforced.

Author Contributions

Conceptualization: H-LV; Data analysis: H-LV, HNSA and DKV; Methodology: H-LV and TPTN; Supervision: MTH, THN, GTV and HLT; Writing – original draft: H-LV, TPTN, KNN, VATD and TTN; Writing – review and editing: H-LV, MTH, THN, GTV and HLT.

Funding support

This research is supported by Vingroup Innovation Foundation (VINIF) in project code VINIF.2020.COVID-19.DA03.

Acknowledgments

This research is supported by Vingroup Innovation Foundation (VINIF) in project code VINIF.2020.COVID-19.DA03.

Conflicts of Interest

The authors declare that this study received funding from Vingroup Innovation Foundation (VINIF). The funder was not involved in the study design, collection, analysis, interpretation of data, the writing of this article or the decision to submit it for publication.

References

1. *National responses to the COVID-19 pandemic*. 2020 [cited 2020 31/07]; Available from: https://en.wikipedia.org/wiki/National_responses_to_the_COVID-19_pandemic.
2. Todd Pollack, et al. *Emerging COVID-19 success story: Vietnam's commitment to containment*. 2020 [cited 2020 27/07]; This article is one of a series focused on identifying and understanding Exemplars in the response to the Coronavirus pandemic. It is hosted by the Exemplars in Global Health (EGH) platform]. Available from: <https://ourworldindata.org/covid-exemplar-vietnam>.
3. Duc Tuan. *Prime Minister agreed to continue implementing Directive 16 at least until 22/4 in some localities*. 2020 [cited 2020 01/08]; Available from: <http://baochinhphu.vn/Tin-noi-bat/Thu-tuong-nhat-tri-tiep-tuc-thuc-hien-Chi-thi-16-it-nhat-den-224-tai-mot-so-dia-phuong/393140.vgp>.
4. Khanh Vu, Phuong Nguyen, and James Pearson. *After aggressive mass testing, Vietnam says it contains coronavirus outbreak*. 2020 [cited 2020 28/07]; Available from: <https://www.reuters.com/article/us-health-coronavirus-vietnam-fight-insi/after-aggressive-mass-testing-vietnam-says-it-contains-coronavirus-outbreak-idUSKBN22B34H>.
5. *Vietnam, Which Avoided Coronavirus for 99 Days, Records First COVID-19 Death*. [cited 2020 01/08]; Available from: <https://time.com/5874146/vietnam-first-coronavirus-death/>.
6. Ministry of Health of Vietnam. *Prime Minister: Calculate closely before deciding on social distance*. 2020 [cited 2020 02/08]; Available from: https://moh.gov.vn/tin-noi-bat/-/asset_publisher/3Yst7YhbKA5j/content/thu-tuong-tinh-toan-chat-che-truoc-khi-quyet-inh-gian-cach-xa-hoi?inheritRedirect=false&redirect=https%3A%2F%2Fmoh.gov.vn%3A443%2Ftin-noi-bat%3Fp_p_id%3D101_INSTANCE_3Yst7YhbKA5j%26p_p_lifecycle%3D0%26p_p_state%3Dnormal%26p_p_mode%3Dview%26p_p_col_id%3Drow-1-column-1%26p_p_col_count%3D1.
7. Ministry of Health of Vietnam. *Tightening the entire system, putting forces in Da Nang*. 2020 [cited 2020 02/08]; Available from: https://moh.gov.vn/tin-noi-bat/-/asset_publisher/3Yst7YhbKA5j/content/siet-chat-toan-bo-he-thong-don-luc-dap-dich-o-a-nang?inheritRedirect=false&redirect=https%3A%2F%2Fmoh.gov.vn%3A443%2Ftin-noi-bat%3Fp_p_id%3D101_INSTANCE_3Yst7YhbKA5j%26p_p_lifecycle%3D0%26p_p_state%3Dnormal%26p_p_mode%3Dview%26p_p_col_id%3Drow-1-column-1%26p_p_col_count%3D1.
8. Ministry of Health of Vietnam. *The Prime Minister instructed: Speed up the tracking of the source of COVID-19 infection*. 2020 [cited 2020 02/08]; Available from: https://moh.gov.vn/hoat-dong-cua-lanh-dao-bo/-/asset_publisher/TW6LTp1ZtwaN/content/thu-tuong-chi-ao-tang-toc-truy-vet-nguon-lay-covid-19?inheritRedirect=false&redirect=https%3A%2F%2Fmoh.gov.vn%3A443%2Fhoat-dong-cua-lanh-dao-bo%3Fp_p_id%3D101_INSTANCE_TW6LTp1ZtwaN%26p_p_lifecycle%3D0%26p_p

- [state%3Dnormal%26p_p_mode%3Dview%26p_p_col_id%3Drow-0-column-2%26p_p_col_count%3D2.](#)
9. L.Anh and T.Duong. *COVID-19 Prevention of community infection: Many measures are applied for the first time*. 2020 [cited 2020 02/08]; Available from: <https://tuoitre.vn/phong-chong-covid-19-lay-nhiem-cong-dong-nhieu-bien-phap-lan-dau-ap-dung-20200727000320695.htm>.
 10. Nguyen, H.G. and T.V. Nguyen, *An epidemiologic profile of COVID-19 patients in Vietnam*. medRxiv, 2020.
 11. Roser, M., et al., *Coronavirus pandemic (COVID-19)*. Our World in Data, 2020.
 12. Ministry of Health of Vietnam, *Decision 322 / QD-BYT guides the diagnosis and treatment of acute respiratory infections*. 2020.
 13. Ministry of Health of Vietnam, *Decision 1344 / QD-BYT 2020 Guidance on diagnosis and treatment of COVID-19*. 2020.
 14. Parag Goyal, e.a., *Clinical Characteristics of Covid-19 in New York City*. NEJM Group Public Health Emergency Collection, 2020.
 15. Cao, M., et al., *Clinical features of patients infected with the 2019 novel coronavirus (COVID-19) in Shanghai, China*. MedRxiv, 2020.
 16. Sudharsanan, N., et al., *The Contribution of the Age Distribution of Cases to COVID-19 Case Fatality Across Countries: A 9-Country Demographic Study*. Annals of Internal Medicine, 2020.
 17. Quang Minh. *Nearly 7,000 Vietnamese return home due to coronavirus fear in foreign countries*. 2020 [cited 2020 31/07]; Available from: <http://news.chinhphu.vn/Home/Nearly-7000-Vietnamese-return-home-due-to-coronavirus-fear-in-foreign-countries/20203/39268.vgp>.
 18. Minh Chien. *Chinese people illegally enter Vietnam to travel, find work and go to Cambodia to gamble*. 2020 [cited 2020 29/07]; Available from: <https://nld.com.vn/thoi-su/nguoi-trung-quoc-nhap-canh-trai-phep-vao-viet-nam-de-du-lich-tim-viec-va-sang-campuchia-danh-bai-20200803185032684.htm>.
 19. Chi Tue and Tien Thang. *Preventing 16,000 people from illegally entering Vietnam*. 2020 [cited 2020 31/07]; Available from: <https://tuoitre.vn/ngan-chan-16-000-nguoi-nhap-canh-trai-phep-vao-viet-nam-20200731183650665.htm>.
 20. Ministry of Health of Vietnam, *Decision 322/QD-BYT 2020; Guidelines for Diagnosis and Treatment of acute respiratory infections caused by a new Corona Virus (2019-nCoV)*. 2020.
 21. World Health Organization. *WHO Coronavirus Disease (COVID-19) Dashboard*. 2020 [cited 2020 30/07]; Available from: <https://covid19.who.int/>.
 22. Vo Thu. *The third COVID-19 patient in Vietnam died*. 2020 [cited 2020 31/07]; Available from: <https://ncov.moh.gov.vn/web/guest/-/benh-nhan-covid-19-thu-3-o-viet-nam-tu-vong>.
 23. Phuong Nguyen and Khanh Vu. *'Act now and act fast': Vietnam says every city at risk of coronavirus*. 2020 [cited 2020 29/07]; Available from: <https://www.reuters.com/article/us-health-coronavirus-vietnam/vietnam-says-every-city-province-now-at-risk-of-virus-infection-idUSKCN24U0C7>.
 24. Duong Hai. *Prof. Dr. Nguyen Van Kinh: "The new virus strain causing COVID-19 spreads rapidly, but its virulence remains unchanged"*. 2020 [cited 2020 31/07];

Available from: <https://ncov.moh.gov.vn/web/guest/-/gs-ts-nguyen-van-kinh-chung-virus-moi-gay-covid-19-lay-lan-nhanh-nhung-oc-luc-khong-oi->.