Full Title: 12 Novel Clonal Groups of *Leptospira* Infecting Humans in Multiple

Contrasting Epidemiological Contexts in Sri Lanka

Jayasundara JMDD[1,2], Senavirathna RMISK[1,3], Warnasekara YPJN[1], Gamage CD[4],

Siribaddana S[5], Kularatne SAM[6], Matthias M[7], Mariet JF[8], Picardeau M[8], Agampodi

SB[1,7], Vinetz JM[7]

[1] Leptospirosis Research Laboratory, Department of Community Medicine, Faculty of

Medicine and Allied Sciences, Rajarata University of Sri Lanka

[2] Department of Microbiology, Faculty of Medicine and Allied Sciences, Rajarata

University of Sri Lanka

[3] Department of Biochemistry, Faculty of Medicine and Allied Sciences, Rajarata

University of Sri Lanka

[4] Department of Microbiology, Faculty of Medicine, University of Peradeniya, Sri Lanka

[5] Department of Medicine, Faculty of Medicine and Allied Sciences, Rajarata University of

Sri Lanka

[6] Department of Medicine, Faculty of Medicine, University of Peradeniya, Sri Lanka

[7] Yale University School of Medicine, New Haven, Connecticut, USA

[8] Institut Pasteur, Biology of Spirochetes unit, Paris, France

Abstract

Leptospirosis is a ubiquitous disease and a major clinical challenge owing to the multitude

of clinical presentations and manifestations that are possibly attributable to the diversity of

Leptospira, the understanding of which is key to study the epidemiology of this emerging

global disease threat. Sri Lanka is a hotspot for leptospirosis with high levels of endemic

disease as well as annual epidemics. We carried out a prospective study of *Leptospira*

diversity in Sri Lanka, covering the full range of climatic zones, geography, and clinical

severity. Samples were collected for leptospiral culture from 1192 patients from 15 of 25

districts in Sri Lanka over two and half years period. Twenty-five isolates belonging to four

pathogenic *Leptospira* species were identified: *L. interrogans*, *L. borgpetersenii*, *L. weilii*,

and *L. kirschneri*. At least six serogroups were identified among the isolates: Autumnalis

(6), Pyrogenes (4), Icterohaemorrhagiae (2), Celledoni (1), Grippotyphosa (2) and Bataviae

(1). Seven isolates did not agglutinate using available antisera panels, suggesting new

serogroups. Isolates were sequenced by Illumina. These data add 25 new core genome

sequence types and were clustered in 15 clonal groups, including 12 new clonal groups. *L.*

borgpetersenii was found only in the dry zone and *L. weilii* only in the wet zone. Acute

kidney injury and cardiovascular involvement were seen only with *L. interrogans*

infections. Thrombocytopenia and liver impairment were seen in both *L. interrogans* and *L.*

borgpetersenii infections. The inadequate sensitivity of culture isolation to identify

infecting *Leptospira* species underscores the need for culture-independent typing methods

for *Leptospira*.

Author Summary

There is a huge diversity in pathogenic *Leptospira* species worldwide, and our knowledge

of the currently circulating species is deficient owing to limited isolation and identification

of *Leptospira* species from endemic countries. This prospective study reveals the wide

pathogen diversity that causes human leptospirosis in Sri Lanka, representing four species,

more than eight serogroups, and fifteen clonal groups. Further, the different geographic and

climatic zone distributions and clinical manifestations observed underscores the need for

prospective studies to expand the molecular epidemiological approaches to combat

leptospirosis

Introduction

Introduction

Leptospirosis is caused by a group of pathogenic *Leptospira* species of the phylum

Spirochetes and is considered one of the commonest zoonotic diseases worldwide [1][2].

Leptospira spp. have the ability to colonize proximal convoluted tubules of kidney tissue of

various mammals (including rodents), birds and marsupials, and the hosts excrete the

bacteria to the environment via urine [3][4]. Humans are incidental hosts who acquire the

disease by direct contact with urine or tissues of reservoir animals or, more frequently by

indirect contact with contaminated water sources [5][6]. The incidence of leptospirosis is

estimated to be 1.03 million annually worldwide, with 58,900 deaths [1]. The majority of

tropical countries in Oceania, Southeast Asia, the Caribbean region, central and eastern

sub-Saharan Africa, and South Asia are estimated to have substantial morbidity and

mortality that is attributable to leptospirosis.[7]

Understanding the diversity of infecting *Leptospira* has been a major global focus,

especially in recent years. The phenomenal changes in *Leptospira* classification backed by

next-generation sequencing methods and whole-genome sequencing have led to the

identification of 43 new *Leptospira* species during the period 2018 to 2020 [8–11]. , the

more robust classification of *Leptospira* strains beyond the species level using core-genome

multi-locus sequence typing (cgMLST) [9] and single nucleotide polymorphism typing

methods has rapidly expanded our knowledge of the molecular epidemiology of

Leptospira. However, the goal of reducing the global burden of this deadly disease will

require an enhanced understanding of pathogen types and applications and linkage to

disease distribution, transmission, clinical presentations, and outcomes.

The global leptospirosis disease burden study [1] has highlighted Sri Lanka as a

hyperendemic country with an estimated morbidity of 300.6 and mortality of 17.98 per

100,000 population. The disease incidence tends to be higher during the rainy seasons, i.e.,

the southwest and northeast monsoons. Cases, however, are not confined exclusively to the

wet zone and are reported in the dry zone as well, where the majority of residents are

engaged in farming activities. Outbreaks have also occurred in the dry zone following

extreme weather events like flooding [12].

As in many other endemic countries, understanding *Leptospira* diversity in Sri Lanka is

limited because of a lack of knowledge of the circulating pathogenic species and serovars.

Studies that utilized culture-based isolation of *Leptospira* species were carried out in Sri

Lanka during the period from 1950 to 1970 in the wet zone only. Several pathogenic strains

of the species *L. interrogans* [13][14][15], *L. borgpetersenii* [16], *L. kirschneri* [17], and *L.*

santarosai were detected during that time [18][19][20]. Since the 1970s, no culture-based

isolation studies were reported until 2018, when two human isolates belonging to *L.*

interrogans were recovered from the wet zone [21]. Despite the availability of next-

generation sequencing methods for many years, whole-genome sequencing data for Sri

Lankan isolates were not available until recently [22].

A systematic review published in 2016 revealed the large diversity of *Leptospira* strains in

Sri Lanka based on historical data [20]. Being an island with a high leptospirosis disease

burden makes Sri Lanka an ideal location to study pathogen diversity linked with

epidemiological and clinical patterns of the disease. Low-passage isolates from human

sources with high-resolution genetic typing in a place with high pathogen diversity would

enhance our global knowledge of leptospirosis. This study was designed to provide a

comprehensive understanding of the circulating pathogenic *Leptospira* species and types

responsible for human leptospirosis in Sri Lanka, covering different clinical presentations

and geographical locations as well as an epidemic and endemic disease over 3 years.

Methods

The present study was embedded in a larger clinical-epidemiological study on leptospirosis,

in Sri Lanka and the study protocol was published elsewhere [23]. Specific details related

to *Leptospira* diversity and methods, in brief, are given here.

Study setting

This study was carried out at several locations in Sri Lanka that differed in mean

temperature, rainfall, elevation, ecology, human activities, and leptospirosis endemicity.

The main data collection sites were the Teaching Hospital Anuradhapura (THA) and

Teaching Hospital Peradeniya (THP). THA is in the dry zone located at a low elevation

with low humidity, high temperature, large rice paddy fields, water reservoir–based

irrigation systems, and low endemicity for the disease. THP is in the wet zone located at

high elevation, low temperature, rainfall-based farming activities, and high endemicity.

Samples for *Leptospira* diversity assessment were collected through two approaches. First,

as a part of the main study described in Agampodi et al. 2019, prospective data and sample

collection was done in THA and THP. Also, during an outbreak of leptospirosis in 2017,

we set up the same procedure at Base Hospital Awissawella and Provincial General

Hospital Rathnapura. These two wet-zone areas have high endemicity, representing low

and intermediate elevations. As a part of the service component of this study, we offered

diagnostic services to all requesting physicians and also collected additional culture

samples. This resulted in sample collection from District General Hospital Kegalle, Base

Hospital Karawanella, Sri Jayawardanapura General Hospital, and General Hospital

Polonnaruwa (GHP)—again representing different [locations OR geographical locations].

These study sites (Fig 1) represent seven districts belonging to four provinces of the

country, and the patients who visited these hospitals came from all nine provinces.

Fig 1. Locations of the seven hospitals involved in the study

Study samples

Culture collection was done from three types of patients. Acute undifferentiated febrile

(temperature >38°C) patients who presented to adult wards (age >13 years) of THA, THP,

Provincial General Hospital Rathnapura and Base Hospital Awissawella (both outpatient

department and hospitalized patients) were included as possible cases of leptospirosis.

Probable cases of clinical leptospirosis were included from GHP. Culture samples from

Provincial General Hospital Rathnapura, Sri Jayawardanapura General Hospital and Base

Hospital Karawanella were included only if they came from clinically confirmed cases of

leptospirosis. Physician-diagnosed probable or definite acute bacterial meningitis or lower

respiratory tract infections (e.g., consolidated lobar pneumonia), traumatic or post-operative

fever per physician discretion, fever owing to nosocomial infections, and any patient with

confirmed diagnosis as a cause for the fever were excluded.

Sample collection and isolation of Leptospira

Blood (7 ml) was collected from all eligible patients. Bedside inoculation of 2 and 4 drops

(100–400 µl) was done into two tubes with 9 ml Ellinghausen-McCullough-Johnson-Harris

(EMJH) semisolid medium with added antibiotics (5-fluorouracil and neomycin). These

cultures were kept at room temperature (usually 28–32°C) until transfer to the

Leptospirosis Research Laboratory of the Faculty of Medicine and Allied Sciences,

Rajarata University, Sri Lanka, and then incubated at 30°C until the cultures become

positive or for 6 months.

In brief, EMJH semisolid media were prepared by adding 2.3 g of EMJH base (Difco), 1.5

g bacteriological agar, and 100 mg sodium pyruvate into 785 ml distilled water and

adjusting the pH to 7.4. The media were autoclaved, and once cooled to ~50°C, 100 ml

Leptospira enrichment media and 100 ml fetal bovine serum was added. To suppress the

growth of possible contaminating bacteria, 5-fluorouracil (100 µg/ml, final) and neomycin

(25 µg/ml) were added. Each inoculated medium was inspected under a dark-field

microscope to check for the presence of motile spirochetes; this was done initially after 3

weeks and then every month. However, samples were inspected before 3 weeks if

quantitative PCR of the corresponding whole-blood sample indicated a positive reaction.

When positive growth was detected, subcultures were made into liquid and semisolid

media, and an aliquot was fixed with 5% dimethyl sulfoxide and stored at –80°C. Isolates

were subcultured in liquid media 2 weekly and on semisolid media 3 monthly. Certain

isolates required weekly subculture into liquid media to maintain viability. None of the

isolates became contaminated during the subculture process, although two positive original

clinical samples were contaminated with bacilli. For those two samples, subcultures were

made into liquid media and subsequently filtered through a 0.2 µm pore-size microfilter to

overcome the problem of contamination.

16S rRNA gene-based species identification

Species-level identification for all isolates was first done using nested PCR targeting the

16S rRNA gene [24][25][26]. Amplification was carried out in 50 µl containing 25 µl

HotStarTaq® master mix, 0.5 µl each primer (1 µM), 10 µl template DNA, and 15 µl

molecular-grade water. The thermal cycler conditions were done as reported

previously[26]. The 16S rRNA region obtained with the amplified product was sequenced

using an ABI 3500 genetic analyzer with an 8-capillary system (Applied Biosystems®).

The target gene segment was 292 bp. All amplified 16 rRNA sequences were submitted to

the NCBI database under Bio Project number PRJNA528695. The 16S rRNA sequences of

the 25 *Leptospira* isolates were aligned using Muscle Mega 7 software [27]. A

phylogenetic tree was constructed with Mega 7 using maximum likelihood analysis with

the General Time Reversible model. Validation of the statistical confidence of each node

was based on 1000 bootstrap replicates.

Next-generation sequencing, cgMLST and phylogenetic tree

DNA was extracted from culture using the PureLink® Genomic DNA Mini kit (Invitrogen,

Dublin, Ireland) and Wizard® Genomic DNA Purification Kit (Promega, Southampton,

UK) according to manufacturer instructions. NGS was performed using Nextera XT DNA

Library Preparation kit and the NextSeq 500 sequencing systems (Illumina, San Diego, CA,

USA) at the Mutualized Platform for Microbiology (P2M) at Institut Pasteur. The data were

analyzed using the CLC Genomics Workbench 9 software (Qiagen, Hilden, Germany) (S2

Table). cgMLST typing was performed for strain taxonomy using a scheme based on 545

highly conserved genes with BIGSdb (http://bigsdb.pasteur.fr/leptospira), and a

phylogenetic tree was generated using cgMLST with Interactive Tree of Life v3, and

GrapeTree [28] was used to visualize the core genomic relationships among the isolates and

the previously reported Sri Lankan isolates [9][29][30].

Serotyping of new isolates

Serotyping of newly isolated *Leptospira* strains was done at the Pasteur Institute, France,

using a standard battery of rabbit antisera raised against 24 reference serovars representing

the main serogroups [31][32].

 Ethics statement

 Written informed consent was obtained from all patients before sample collection. This

 study is approved by the Ethics Review Committee of the Faculty of Medicine and Allied

 Sciences, Rajarata University of Sri Lanka. Protocol No. 2015/EC/98

 Results

 From June 2016 through January 2019, we acquired blood cultures from 1192 patients.

 Patients were from 14 districts of Sri Lanka, representing all 9 provinces (Fig 2).

 Fig 2. Distribution of probable exposure sites/residence of the patients recruited for

 the study.

 The majority of patients were male (n = 985, 82.6%). The mean age of the sample was 43.4

 years (SD 14.7). Most of the cultures (96%) were obtained from hospital inpatients, and the

 remaining 4% (48 cultures) were from outpatients. Of the 1192 cultures, 80 (6.7%) were

 received from hospitals where only typical clinical cases of leptospirosis were sampled

 (from Sri Jayewardenepura General Hospital, Base Hospital Karawanella and Provincial

 General Hospital Rathnapura). Another 107 (9.0%) were from GHP, where patients

 represented only probable cases of leptospirosis. The remaining patients (n = 1047) were

 from THA, THP, Base Hospital Awissawella and Provincial General Hospital Rathnapura,

 representing patients who presented with acute undifferentiated fever.

 Of the 1192 patients, 25 isolates had been identified by January 2019. Among the acute

 undifferentiated febrile patients, 1.5% (16/1047) had culture-positive leptospirosis; among

 the probable and clinically confirmed cases of leptospirosis, culture positivity was 4.7%

 (5/107) and 5.0% (4/80), respectively. The incubation period required to detect positive

growth (assessed with dark-field microscopy) varied between 1 and 17 weeks. For each

patient, both 2 drops and 4 drops of blood-inoculated media gave positive results. Fig 3

shows the range of incubation period required for the 25 positive *Leptospira* cultures. The

median duration was 15 weeks for the first 9 isolates and was 6 weeks for the remaining 16

isolates.

Fig 3. Distribution of incubation period for culture isolation among 25 Leptospira

cultures

Leptospira was isolated from one female and 24 male patients who presented with fever.

Only three patients were from outpatient departments, and the rest were inpatients. Of the

three outpatients, two were later admitted to a hospital owing to increased disease severity.

Species identification using 16S rRNA led to inconclusive results for certain isolates owing

to inadequate length of the PCR product of the target DNA. However, all nucleotides

obtained from 16S rRNA sequencing were deposited in GenBank (see Table 2 for

accession numbers). cgMLST analysis revealed that the 25 isolates represented four

species: *L. interrogans* (15 isolates, 60%), *L. borgpetersenii* (7 isolates, 28%), *L. weilii* (2

isolates, 8%), and *L. kirschneri* (1 isolate, 4%) (Fig 4).

Fig 4 Phylogenetic tree showing the distribution of species and serogroups of the

isolates from the present study along with the previously reported Sri Lankan isolates

and other species.

Interpretable data for serogroup assay was available for only 19 isolates. Seven samples

resulted   in   no   agglutination,   probably   owing   to   new,   previously   unreported

serogroups/serovars. The assay revealed that the 19 seropositive isolates represented at

least six serogroups, namely Autumnalis, Pyrogenes, Icterohaemorrhagiae, Grippotyphosa,

Celledoni, and Bataviae (Table 2).

Table 2. Putative species, serogroups, and cgMLST data for the *Leptospira* isolates

Strain ID	Species	Serogroup	GenBank accession number of the 16S rrs amplicon	cgMLST	Cluster40
FMAS_KW1	*L. interrogans*	Pyrogenes	MN250143	784	10
FMAS_KW2	*L. interrogans*	Autumnalis	MN250144	631	291
FMAS_AW1	*L. interrogans*	Autumnalis	MN250145	555	74
FMAS_RT1	*L. weilii*	No agglutination	MN250146	556	262
FMAS_AW2	*L. interrogans*	Autumnalis	MN250147	567	269
FMAS_AW3	*L. interrogans*	Pyrogenes	MN250148	557	9
FMAS_RT2	*L. interrogans*	Autumnalis	MN250149	569	271
FMAS_PD1	*L. interrogans*	Pyrogenes	MN250150	558	263
FMAS_PD2	*L. weilii*	Celledoni	MN250151	559	264
FMAS_KG1	*L. interrogans*	Bataviae	MN250152	560	265
FMAS_KG2	*L. interrogans*	Pyrogenes	MN250153	561	263
FMAS_AP1	*L. interrogans*	Autumnalis	MN250154	562	266
FMAS_AP2	*L. borgpetersenii*	No agglutination	MN250155	563	267
FMAS_AP3	*L. borgpetersenii*	No agglutination	MN250156	575	267
FMAS_AP4	*L. borgpetersenii*	No agglutination	MN250157	564	267
FMAS_AP5	*L. interrogans*	Pyrogenes	MN250158	565	10
FMAS_AP6	*L. interrogans*	Pyrogenes	MN250159	785	321

FMAS_AP7	*L. interrogans*	Autumnalis	MN250160	786	266
FMAS_PN1	*L. borgpetersenii*	No agglutination	MN250161	787	267
FMAS_PN2	*L. interrogans*	Icterohaemorrhagiae	MN250162	788	322
FMAS_PN3	*L. interrogans*	Autumnalis	MN250163	789	266
FMAS_PN4	*L. borgpetersenii*	No agglutination	MN250164	790	267
FMAS_AP8	*L. borgpetersenii*	No agglutination	MN250165	791	267
FMAS_AP9	*L. borgpetersenii*	No agglutination	MN250166	792	267
FMAS_PN5	*L. kirschneri*	Grippotyphosa	MN250167	793	323

The isolates were classified using genome-based taxonomy based on clusters created at the

40-mismatch level. The predominant clonal group (CG) was CG267 (7 *L. borgpetersenii*

isolates among the 25 total isolates). This was followed by CG266 (3/25), CG10 (2/25),

and CG263 (2/25). The two *L. weilii* isolates clustered in [different clonal groups (CG262,

CG264). GrapeTree analysis revealed that the seven *L. borgpetersenii* isolates formed a

distinct cluster separated from previously reported isolates of the same species. However,

L. interrogans isolates had a different lineage. The distribution of serogroups closely

matched that of the genetic clusters (Fig 5).

Fig 5. Genome GrapeTree showing the core-genome relationship among the 25 new

and 9 previously isolated *Leptospira* strains from Sri Lanka

Culture-positive leptospirosis patients were distributed widely in the study areas (Fig 6). *L.*

borgpetersenii was exclusively isolated from patients in areas of the dry zone at low

elevation, with hot and dry conditions. The single isolate of *L. kirschneri* was from the

same setting. In contrast, *L. weilii* was isolated from patients in the wet zone, whereas *L.*

interrogans was isolated from patients in all geographical areas (Fig 6). The serogroup

distribution also revealed a specific pattern for all nonagglutination isolates, which mainly

were from the dry zone at low elevation. The most frequent serogroup Autumnalis, was

observed in all geographical settings. All *L. borgpetersenii* isolates were in CG267, the

majority of which failed to agglutinate with rabbit sera (some slightly agglutinated with

serovars Grippotyphosa and Louisiana).

Fig 6. Geographic distribution of *Leptospira* species, serogroups and clonal groups

Clinical profile of culture-positive patients

Supplementary Table 1 presents complete demographic and clinical profiles for culture-

positive patients. All culture-positive patients were later contacted to collect additional

data. Each patient's clinical records were also retrieved from the corresponding hospital.

No fatalities were reported for patients with culture-positive leptospirosis. Among

hospitalized patients, the median duration of hospital stay was 4 days (interquartile range,

3–5 days), and the longest stay was 12 days followed by 9 days (both patients had acute

kidney injury and required hemodialysis). Another two had renal involvement with

elevated serum creatinine but not acute kidney failure. Twelve patients had elevated

serum glutamic-oxaloacetic transaminase and serum glutamic-pyruvic transaminase, and

three other patients had elevated serum bilirubin. Thrombocytopenia was common in those

12 patients, with a platelet count <100,000 per microliter. Two patients who underwent

hemodialysis also had cardiac involvement with hypotension. Each patient who had a

severe complication was infected with *L. interrogans*. Infection with *L. interrogans* or *L.*

borgpetersenii was associated with thrombocytopenia and liver involvement.

Other diagnostic tests for culture-positive patients

For the 25 *Leptospira*-positive patients, whole-blood and serum samples were acquired for

20 and 15 patients in the hospitals. Sera included 3 paired and 12 single samples, and only

three patient samples yielded positive results for the microscopic agglutination test. PCR

for whole-blood samples revealed 14 positives, 3 probable (only 1 of 3 replicates was

positive), and 3 negatives.

Discussion

This is the first study to report information on the diversity of pathogenic *Leptospira*

species in Sri Lanka and probably one of the few prospective studies on *Leptospira* disease

diversity in literature representing all geographical regions of an entire country. Also, we

report here the first isolation of *L. weilii* in Sri Lanka, the existence of which was suggested

based on molecular studies of clinical samples [33]. This study also provides the first

evidence of serogroups Celledoni and Bataviae circulating in Sri Lanka. Moreover, except

for *L. santarosai*, our study identified all pathogenic *Leptospira* species that were reported

to have existed in Sri Lanka during the 1960s and 70s (as reviewed by Naotunna et al. in

2015), confirming the breadth of *Leptospira* diversity in Sri Lanka [20]. Recently reports

indicated that *L. interrogans*, *L. borgpetersenii* and *L. kirschneri* are also the most common

circulating species in other tropical regions of the world [34][35][36][37][38][39].

However, the geographical distribution of certain *Leptospira* species is limited, such as *L.*

santarosai, which has been mainly reported in South America [40][41][42][43], and *L.*

weilii in Asia [39][44][45][46]. Only limited reports have described the existence of *L.*

weilii outside Asia, where cattle are the dominant reservoir host [46][47][48][49].

We identified 15 distinct clonal groups of *Leptospira*, underscoring the diversity of

pathogenic *Leptospira* circulating in Sri Lanka. The distribution of certain clonal groups

revealed a geographical demarcation between the dry and wet zones. The presence of

diverse clonal groups and the observed demarcation of the limited geographical distribution

of certain species suggests that different environmental drivers of leptospirosis operate in

distinct ways in these climatic zones. The predominance of *L. borgpetersenii* and *L.*

kirschneri over *L. interrogans* has been reported in both humans and cattle in the African

continent and nearby islands such as Mayotte, with possible cattle-to-human transmission

[50][51][52]. Moreover, certain rat species also excrete serovars of *L. borgpetersenii*

[53][54]. Notably, areas where *L. borgpetersenii* was found in our study, cattle and buffalo,

are commonly used in paddy farming activities.

In our present study, *L. interrogans* was the predominant species identified in patients

residing in the wet zone. The emergence of a single dominant clone that caused an outbreak

of leptospirosis following a flood was reported in Thailand] [36]. In contrast with that

study, we observed diverse clonal groups of *L. interrogans* as the cause of human

leptospirosis during floods, probably owing to disease transmission from several different

reservoir hosts.

The diversity of circulating serogroups in Sri Lanka that has been known since the 1960s

has been preserved, and 19 isolates belong to 8 different serogroups, and the non-

agglutinated cultures might reflect newly emerged serovars. Serological and molecular

assays done on the veterinary field has already identified the role of rodents, cattle and dogs

as reservoir hosts in Sri Lanka[55][56][57]. However, the observed diversity in serogroups

offers evidence for a wider range of reservoir hosts even though Sri Lanka is a small island.

Although culture isolation is required for in-depth molecular epidemiological studies, our

present study highlights the constraints faced for culture isolation of *Leptospira*, for which

both a high level of skill and procedural optimization are required. *Leptospira* spp. are

fastidious organisms, and their growth requirements differ from those of many other

bacterial genera. *Leptospira* tend to have a relatively long incubation period, as the lag

period during *in vitro* culture range from days to several weeks [58]. Other culture-isolation

studies have reported an incubation period of ~3 weeks, but with a wide range of duration

[50][59][60]. We attribute the relatively lengthy incubation period of 15 weeks required for

the first nine cultures in our study mainly to the lack of skills and optimization required for

culture procedures during the first phase of the study. Isolation is essential for genomic and

vaccine studies on *Leptospira*, and for that purpose a specific skill set is mandatory.

Although culture isolation has a 100% positive predictive value for diagnosing

leptospirosis, its sensitivity has been consistently <10% in most studies [36][61][62].

Similarly, our study yielded low sensitivity, which can be attributed to a few possible

causes: Because the study population consisted of patients with acute febrile illness, their

fever may have been caused by another unrelated illness; Use of antibiotics before culture;

also, performing blood cultures during the late phase of illness and infection with fastidious

Leptospira spp. may account for the low sensitivity during culture isolation.

This large collection of pathogenic *Leptospira* isolates from clinical samples will be a great

addition to the global knowledge base for leptospirosis. Whole-genome sequencing and

genomic analysis of this set of isolates will reveal the pathogenic diversity and evolution of

Leptospira species, in comparison to archived *Leptospira* isolated from Sri Lanka more

than 50 years ago. The first three isolates from this study are already published [REFs?],

and the isolate FMAS_AW1 is now listed as the reference genome for *L. interrogans* in the

NCBI genome database (https://www.ncbi.nlm.nih.gov/genome/?term =

Leptospira+interrogans).

Whole genome sequencing and comparative genomic analysis of this collection will

facilitate ongoing studies on identifying the putative virulent genes, pathogenic

mechanisms with specific host adaptations, horizontal gene transfer mechanisms, and

microbial resistance as shown in studies on the diversity and epidemiology of other

microorganisms [63][64][65][66][67][68].

Acknowledgement

This work was supported by The National Institute of Allergy and Infectious Diseases of

the National Institutes of Health, Award Number U19AI115658. The content is solely the

responsibility of the authors and does not necessarily represent the official views of the

National Institutes of Health. The funding body had no role in the design of the study or the

collection, analysis, or interpretation of data or in writing the manuscript and publication.

We would like to thank the co-applicants of the main study, Prof.Sisira Siribaddana and

Prof.S.A.M Kularatne for their support for the clinical study. We would also like to thank

Ms. Thilakanjali Gamage, Mr. K.M.R. Premathilaka, Mr. S.K. Senevirathna, and Mr.

Milinda Perera for technical assistance, Mr. Shalka Srimantha and Ms. Chamila Kappagoda

for culture maintenance and laboratory support, and Dr. Muditha Abeykoon, Dr. Chamida

Wickramasinghe, and Dr. Shanika Gamage for additional culture collections. We also

thank all the physicians and healthcare staff in the various participating hospitals, the team

of core facility P2M (Institut Pasteur, Mutualized Platform for Microbiology) for genomic

sequencing and members of the National Reference Center for Leptospirosis (Institut

Pasteur) for technical assistance with the cultures of *Leptospira*.

References

1.   Costa F, Hagan JE, Calcagno J, Kane M, Torgerson P, Martinez-Silveira MS, et al.

Global Morbidity and Mortality of Leptospirosis: A Systematic Review. PLoS Negl

Trop Dis. 2015;9: e0003898. doi:10.1371/journal.pntd.0003898

2.   Pappas G, Papadimitriou P, Siozopoulou V, Christou L, Akritidis N. The

globalization of leptospirosis: worldwide incidence trends. International Journal of

Infectious Diseases. 2008. pp. 351–357. doi:10.1016/j.ijid.2007.09.011

3.   Levett PN. Leptospira and Leptospirosis. Systematics of Leptospiraceae. Adler B,

editor. Curr Top Microbiol Inmunol. 2015;387: 11–20. doi:10.1007/978-3-662-

45059-8

4.   Human leptospirosis: guidance for diagnosis, surveillance and control. Rev Inst Med

Trop Sao Paulo. 2003;45: 292–292. doi:10.1590/s0036-46652003000500015

5.   Levett PN. Systematics of leptospiraceae. Curr Top Microbiol Immunol. 2015;387:

11–20. doi:10.1007/978-3-662-45059-8_2

6.   Haake DA, Levett PN. Leptospirosis in humans. Curr Top Microbiol Immunol.

2015;387: 65–97. doi:10.1007/978-3-662-45059-8_5

7.   Torgerson PR, Hagan JE, Costa F, Calcagno J, Kane M, Martinez-Silveira MS, et al.

Global Burden of Leptospirosis: Estimated in Terms of Disability Adjusted Life

Years. PLoS Negl Trop Dis. 2015. doi:10.1371/journal.pntd.0004122

8.   Thibeaux R, Iraola G, Ferrés I, Bierque E, Girault D, Soupé-Gilbert ME, et al.

Deciphering the unexplored Leptospira diversity from soils uncovers genomic

evolution to virulence. Microb genomics. 2018;4. doi:10.1099/mgen.0.000144

9.   Guglielmini J, Bourhy P, Schiettekatte O, Zinini F, Brisse S, Picardeau M. Genus-

wide Leptospira core genome multilocus sequence typing for strain taxonomy and

global surveillance. PLoS Negl Trop Dis. 2019;13.

doi:10.1371/journal.pntd.0007374

10.   Vincent AT, Schiettekatte O, Goarant C, Neela VK, Bernet E, Thibeaux R, et al.

Revisiting the taxonomy and evolution of pathogenicity of the genus Leptospira

through the prism of genomics. Martins EAL, editor. PLoS Negl Trop Dis. 2019;13:

e0007270. doi:10.1371/journal.pntd.0007270

11.   Casanovas-Massana A, Hamond C, Santos LA, de Oliveira D, Hacker KP,

Balassiano I, et al. Leptospira yasudae sp. Nov. and Leptospira stimsonii sp. nov.,

two new species of the pathogenic group isolated from environmental sources. Int J

Syst Evol Microbiol. 2020. doi:10.1099/ijsem.0.003480

12.   Bandaranayaka AK, Vinetz JM, Perera M, Dahanayaka NJ, Weerawansa P,

Priyankara S, et al. Regional Differences of Leptospirosis in Sri Lanka: Observations

from a Flood-Associated Outbreak in 2011. PLoS Negl Trop Dis. 2014;8: e2626.

doi:10.1371/journal.pntd.0002626

13.   Nityananda K. Isolation of Leptospira in Ceylon. Ceylon Med J. 1962;7: 95–6.

Available: https://www.cabdirect.org/cabdirect/abstract/19632705150

14.   Nityananda K SC. A New Leptospiral Serotype in the Icterohaemorrhagiae

Serogroup from Ceylon. Ceylon J Med Sci. 1972;21: 9–13. Available:

http://archive.cmb.ac.lk:8080/research/bitstream/70130/3637/1/A_

New_Leptospiral_Serotype.PDF

15.   Nityananda K, Sulzer CR. A new serotype of Leptospira belonging to the autumnalis

serogroup. J Trop Med Hyg. 1971;74: 184–6. Available:

http://www.ncbi.nlm.nih.gov/pubmed/5570126

16.   Nityananda K, Sulzer CR. A new leptospiral serotype in the Javanica serogroup from

Ceylon. Trop Geogr Med. 1969;21: 207–209. Available:

http://archive.cmb.ac.lk:8080/research/handle/70130/3637

17.   Kokovin IL, Chernukha IG. [Serological classification of leptospirae of the

grippotyphosa serogroup--new serological type ratnapura]. Zh Mikrobiol Epidemiol

Immunobiol. 1970;47: 102–5. Available:

http://www.ncbi.nlm.nih.gov/pubmed/4251597

18.   Nityananda K, Harvey T. Leptospirosis_in_Ceylon-Epidemiological and Laboratory

Studies. Ceylon J Med Sci. 1971;20: 5–14.

19.   Kmety E, Dikken H. Classification of the species Leptospira interrogans and history

of its serovars. University Press Groningen; 1993. p. 104. Available:

https://books.google.lk/books/about/Classification_of_the_species_Leptospira.html?

id=EUBWAAAAYAAJ&redir_esc=y

20.   Naotunna C, Agampodi SB, Agampodi TC. Etiological agents causing leptospirosis

in Sri Lanka: A review. Asian Pacific Journal of Tropical Medicine. 2016.

doi:10.1016/j.apjtm.2016.03.009

21.   Nisansala GGT, Muthusinghe D, Gunasekara TDCP, Weerasekera MM, Fernando

SSN, Ranasinghe KNP, et al. Isolation and characterization of Leptospira interrogans

from two patients with leptospirosis in Western Province, Sri Lanka. J Med

Microbiol. 2018 [cited 30 Aug 2018]. doi:10.1099/jmm.0.000800

22.   Seneviratha I, Jayasundara D, Lefler JP, Chaiboonm KL, Warnasekara J, Agampodi

S, et al. Complete Genome Sequence of Leptospira interrogans Strains

FMAS_KW1, FMAS_KW2 and FMAS_AW1 Isolated from Leptospirosis Patients

from Karawanalla and Awissawella, Sri Lanka. J Genomics. 2020;8: 49–52.

474   doi:10.7150/jgen.43953

23.   Agampodi S, Warnasekara J, Jayasundara D, Senawirathna I, Gamage C, Kularatne

S, et al. Study protocol: Characterising the clinical, epidemiological and aetiological

aspects of leptospirosis in Sri Lanka: A hospital based clinico-epidemiological study.

BMJ Open. 2019;9: 1–8. doi:10.1136/bmjopen-2018-027850

24.   Gonçalves AT, Paiva C, Melo-Mota F, Vieira ML, Carreira T, Nunes MS, et al. First

isolation of human Leptospira strains, Azores, Portugal. Int J Infect Dis. 2010;14:

e148–e153. doi:10.1016/j.ijid.2009.12.004

25.   Merien F, Amouriaux P, Perolat P, Baranton G, Girons IS. Polymerase chain

reaction for detection of Leptospira spp. in clinical samples. J Clin Microbiol.

1992;30: 2219–2224. doi:10.1128/jcm.30.9.2219-2224.1992

26.   Esteves LM, Bulhões SM, Branco CC, Carreira T, Vieira ML, Gomes-Solecki M, et

al. Diagnosis of Human Leptospirosis in a Clinical Setting: Real-Time PCR High

Resolution Melting Analysis for Detection of Leptospira at the Onset of Disease. Sci

Rep. 2018;8. doi:10.1038/s41598-018-27555-2

27.   Kumar S, Stecher G, Tamura K. MEGA7: Molecular Evolutionary Genetics

Analysis Version 7.0 for Bigger Datasets. Mol Biol Evol. 2016;33: 1870–1874.

doi:10.1093/molbev/msw054

28.   Zhou Z, Alikhan NF, Sergeant MJ, Luhmann N, Vaz C, Francisco AP, et al.

Grapetree: Visualization of core genomic relationships among 100,000 bacterial

pathogens. Genome Res. 2018;28: 1395–1404. doi:10.1101/gr.232397.117

29.   Jolley KA, Cj Maiden M. BIGSdb: Scalable analysis of bacterial genome variation at

the population level. 2010. doi:10.1186/1471-2105-11-595

30.   Letunic I, Bork P. Interactive tree of life (iTOL) v3: an online tool for the display

and annotation of phylogenetic and other trees. Nucleic Acids Res. 2016;44.

doi:10.1093/nar/gkw290

31.   Kmety E, Dikken H. Classification of the species Leptospira interrogans and history

of its serovars. 1993. p. 104.

32.   Bourhy P, Collet L, Clément S, Huerre M, Ave P, Giry C, et al. Isolation and

characterization of new Leptospira genotypes from patients in Mayotte (Indian

Ocean). PLoS Negl Trop Dis. 2010;4. doi:10.1371/journal.pntd.0000724

33.   Agampodi SB, Matthias MA, Moreno AC, Vinetz JM. Utility of quantitative

polymerase chain reaction in leptospirosis diagnosis: Association of level of

leptospiremia and clinical manifestations in Sri Lanka. Clin Infect Dis. 2012;54:

1249–1255. doi:10.1093/cid/cis035

34.   Nalam K, Ahmed A, Devi SM, Francalacci P, Baig M. Genetic Affinities within a

Large Global Collection of Pathogenic Leptospira: Implications for Strain

Identification and Molecular Epidemiology. PLoS One. 2010;5: 12637.

doi:10.1371/journal.pone.0012637

35.   Bourhy P, Collet L, Lernout T, Zinini F, Hartskeerl RA, Van Der Linden H, et al.

Human Leptospira isolates circulating in Mayotte (Indian Ocean) have unique

serological and molecular features. J Clin Microbiol. 2012;50: 307–311.

doi:10.1128/JCM.05931-11

36.   Thaipadungpanit J, Wuthiekanun V, Chierakul W, Smythe LD, Petkanchanapong W,

Limpaiboon R, et al. A Dominant Clone of Leptospira interrogans Associated with

an Outbreak of Human Leptospirosis in Thailand. [cited 3 Mar 2019].

doi:10.1371/journal.pntd.0000056

37.   Zarantonelli L, Suanes A, Meny P, Buroni F, Nieves C, Salaberry X, et al. Isolation

of pathogenic Leptospira strains from naturally infected cattle in Uruguay reveals

high serovar diversity, and uncovers a relevant risk for human leptospirosis. PLoS

Negl Trop Dis. 2018;12. doi:10.1371/journal.pntd.0006694

38.   Wuthiekanun V, Sirisukkarn N, Daengsupa P, Sakaraserane P, Sangkakam A,

Chierakul W, et al. Clinical diagnosis and geographic distribution of leptospirosis,

Thailand. Emerg Infect Dis. 2007;13: 124–126. doi:10.3201/eid1301.060718

39.   Kurilung A, Chanchaithong P, Lugsomya K, Niyomtham W, Wuthiekanun V,

Prapasarakul N. Molecular detection and isolation of pathogenic Leptospira from

asymptomatic humans, domestic animals and water sources in Nan province, a rural

area of Thailand. Res Vet Sci. 2017;115: 146–154. doi:10.1016/j.rvsc.2017.03.017

40.   Storck CH, Postic D, Lamaury I, Perez JM. Changes in epidemiology of

leptospirosis in 2003-2004, a two El Niño Southern Oscillation period, Guadeloupe

archipelago, French West Indies. Epidemiol Infect. 2008;136: 1407–1415.

doi:10.1017/S0950268807000052

41.   Peláez Sanchez RG, Lopez JÁ, Pereira MM, Naranjo MA, Agudelo-F Lórez P.

Genetic diversity of leptospira in northwestern Colombia: First report of leptospira

santarosai as a recognised leptospirosis agent. Mem Inst Oswaldo Cruz. 2016;111:

737–744. doi:10.1590/0074-02760160245

42.   Moreno LZ, Miraglia F, Marvulo MFV, Silva JCR, Paula CD, Costa BLP, et al.

Characterization of Leptospira santarosai serogroup Grippotyphosa serovar Bananal

isolated from Capybara (Hydrochaeris hydrochaeris) in Brazil. J Wildl Dis. 2016;52:

688–693. doi:10.7589/2015-09-245

43.   Valverde M de los A, Goris MGA, González V, Anchia ME, Díaz P, Ahmed A, et al.

New serovars of Leptospira isolated from patients in Costa Rica: Implications for

public health. J Med Microbiol. 2013;62: 1263–1271. doi:10.1099/jmm.0.058545-0

44.   Xu Y, Zheng H, Zhang Y, Wang Y, Zhang J, Li Z, et al. Genomic analysis of a new

serovar of Leptospira weilii serogroup Manhao. Front Microbiol. 2017;8.

doi:10.3389/fmicb.2017.00149

45. Haake DA, Dundoo M, Cader R, Kubak BM, Hartskeerl RA, Sejvar JJ, et al.

Leptospirosis, Water Sports, and Chemoprophylaxis. Clin Infect Dis. 2002;34: e40–

e43. doi:10.1086/339942

46. Kurilung A, Chanchaithong P, Lugsomya K, Niyomtham W, Wuthiekanun V,

Prapasarakul N. Molecular detection and isolation of pathogenic Leptospira from

asymptomatic humans, domestic animals and water sources in Nan province, a rural

area of Thailand. Res Vet Sci. 2017;115: 146–154.

doi:10.1016/J.RVSC.2017.03.017

47. Roberts MW, Smythe L, Dohnt M, Symonds M, Slack A. Serologic-based

investigation of leptospirosis in a population of free-ranging eastern grey kangaroos

(Macropus giganteus) indicating the presence of Leptospira weilii serovar Topaz. J

Wildl Dis. 2010;46: 564–569. doi:10.7589/0090-3558-46.2.564

48. Slack AT, Symonds ML, Dohnt MF, Corney BG, Smythe LD. Epidemiology of

Leptospira weilii serovar Topaz infections in Australia. Commun Dis Intell.

2007;31: 216–222.

49. Corney BG, Slack AT, Symonds ML, Dohnt MF, McClintock CS, McGowan MR, et

al. Leptospira weilii serovar Topaz, a new member of the Tarassovi serogroup

isolated from a bovine source in Queensland, Australia. Int J Syst Evol Microbiol.

2008;58: 2249–2252. doi:10.1099/ijs.0.64884-0

50. Bourhy P, Collet L, Clément S, Huerre M, Ave P, Giry C, et al. Isolation and

characterization of new Leptospira genotypes from patients in Mayotte (Indian

Ocean). PLoS Negl Trop Dis. 2010;4. doi:10.1371/journal.pntd.0000724

51. Feresu SB. DNA relatedness of Leptospira strains isolated from beef cattle in

Zimbabwe. Int J Syst Bacteriol. 1999;49: 1111–1117. doi:10.1099/00207713-49-3-

1111

52.   Mgode GF, Machang'u RS, Goris MG, Engelbert M, Sondij S, Hartskeerl RA. New

Leptospira serovar Sokoine of serogroup Icterohaemorrhagiae from cattle in

Tanzania. Int J Syst Evol Microbiol. 2006;56: 593–597. doi:10.1099/ijs.0.63278-0

53.   Benacer D, Mohd Zain SN, Sim SZ, Mohd Khalid MKN, Galloway RL, Souris M, et

al. Determination of Leptospira borgpetersenii serovar Javanica and Leptospira

interrogans serovar Bataviae as the persistent Leptospira serovars circulating in the

urban rat populations in Peninsular Malaysia. Parasites and Vectors. 2016;9: no

pagination. doi:10.1186/s13071-016-1400-1

54.   Eslabão MR, Kremer FS, Ramos RTJ, Da Silva AL da C, Azevedo VA de C, Pinto L

da S, et al. Genome of leptospira borgpetersenii strain 4E, a highly virulent isolate

obtained from mus musculus in southern Brazil. Mem Inst Oswaldo Cruz. 2018;113:

137–141. doi:10.1590/0074-02760170111

55.   Nityananda K, Harvey T. Leptospirosis in Ceylon epidemiological and laboratory

studies. Ceylon J Med Sci. 1971;20: 5–14. Available:

https://www.researchgate.net/publication/235020299_Leptospirosis_in_Ceylon_-

_Epidemiological_and_Laboratory_Investigation

56.   Gamage CD, Koizumi N, Muto M, Nwafor-Okoli C, Kurukurusuriya S, Rajapakse

JRPV, et al. Prevalence and Carrier Status of Leptospirosis in Smallholder Dairy

Cattle and Peridomestic Rodents in Kandy, Sri Lanka. Vector-Borne Zoonotic Dis.

2011;11: 1041–1047. doi:10.1089/vbz.2010.0153

57.   Gamage CD, Koizumi N, Perera AKC, Muto M, Nwafor-Okoli C, Ranasinghe S, et

al. Carrier status of leptospirosis among cattle in Sri Lanka: A zoonotic threat to

public health. Transbound Emerg Dis. 2014;61: 91–96. doi:10.1111/tbed.12014

58.   Cameron CE. Leptospiral structure, physiology, and metabolism. Curr Top

599  Microbiol Immunol. 2015;387: 21–41. doi:10.1007/978-3-662-45059-8_3

600 59. Wuthiekanun V, Chierakul W, Limmathurotsakul D, Smythe LD, Symonds ML,

601  Dohnt MF, et al. Optimization of Culture of Leptospira from Humans with

602  Leptospirosis. J Clin Microbiol. 2007;45: 1363–1365. doi:10.1128/JCM.02430-06

603 60. Girault D, Soupé-Gilbert ME, Geroult S, Colot J, Goarant C. Isolation of Leptospira

604  from blood culture bottles. Diagn Microbiol Infect Dis. 2017;88: 17–19.

605  doi:10.1016/j.diagmicrobio.2017.01.014

606 61. Meny P, Menéndez C, Quintero J, Hernández E, Ríos C, Balassiano IT, et al.

607  Characterization of Leptospira isolates from humans and the environment in

608  Uruguay. Rev Inst Med Trop Sao Paulo. 2017;59. doi:10.1590/S1678-

609  9946201759079

610 62. Matthias MA, Ricaldi JN, Cespedes M, Diaz MM, Galloway RL, Saito M, et al.

611  Human leptospirosis caused by a new, antigenically unique Leptospira associated

612  with a Rattus species reservoir in the Peruvian Amazon. PLoS Negl Trop Dis.

613  2008;2. doi:10.1371/journal.pntd.0000213

614 63. Jorge S, Kremer FS, De Oliveira NR, Navarro GDOSV, Guimarães AM, Sanchez

615  CD, et al. Whole-genome sequencing of leptospira interrogans from southern Brazil:

616  Genetic features of a highly virulent strain. Mem Inst Oswaldo Cruz. 2018;113: 80–

617  86. doi:10.1590/0074-02760170130

618 64. Beriwal S, Padhiyar N, Bhatt D, Pandit PD, Ansari A, Lata KS, et al. LeptoDB: an

619  integrated database of genomics and proteomics resource of Leptospira. Database.

620  2018;2018. doi:10.1093/database/bay057

621 65. Llanes A, Restrepo CM, Rajeev S. Whole genome sequencing allows better

622  understanding of the evolutionary history of leptospira interrogans serovar hardjo.

623  PLoS One. 2016;11. doi:10.1371/journal.pone.0159387

624   66.   Xu Y, Zhu Y, Wang Y, Chang Y-F, Zhang Y, Jiang X, et al. Whole genome

625   sequencing revealed host adaptation-focused genomic plasticity of pathogenic

626   Leptospira. Sci Rep. 2016;6: 20020. Available: https://doi.org/10.1038/srep20020

627   67.   Kurilung A, Keeratipusana C, Suriyaphol P, Hampson DJ, Prapasarakul N. Genomic

628   analysis of Leptospira interrogans serovar Paidjan and Dadas isolates from carrier

629   dogs and comparative genomic analysis to detect genes under positive selection.

630   BMC Genomics. 2019;20: 168. doi:10.1186/s12864-019-5562-z

631   68.   Llanes A, Restrepo CM, Riesgo-Ferreiro P, Rajeev S. Genomic Variability among

632   Field Isolates and Laboratory-Adapted Strains of Leptospira borgpetersenii Serovar

633   Hardjo. Int J Microbiol. 2018;2018. doi:10.1155/2018/2137036

636   **Fig 1. Locations of seven hospitals used for the study**

Fig 2. Distribution of probable exposure sites/ residence of the patients recruited for

the study.

Fig 3. Distribution of incubation period for culture isolation among 25 *Leptospira*

cultures

Fig 4. Phylogenetic Tree showing the species and serogroup distribution of the isolates from the present study with previously reported Sri Lankan isolates and other species.

Fig 5. Genome GrapeTree showing the core-genome relationship among 25 new and

nine previously isolated Leptospira strains from Sri Lanka

655 **Fig 6. Geographical distribution of *Leptospira* species, serogroups and clonal groups**