Bronchoscopy on Intubated COVID-19 Patients is Associated with Low Infectious Risk to Operators at a High-Volume Center Using an Aerosol-minimizing Protocol

Catherine A. Gao,*[1] Joseph I. Bailey,[1] James M. Walter,[1] John M. Coleman,[1] Elizabeth S. Malsin,[1] A. Christine Argento,[1] NU COVID Investigators,[2] Richard G. Wunderink,[1] Sean B. Smith[1]

* Corresponding author: catherine.gao@northwestern.edu
[1] Division of Pulmonary and Critical Care, Department of Medicine, Northwestern University Feinberg School of Medicine, Chicago IL 60611
[2] Multiple departments, Northwestern University Feinberg School of Medicine, Chicago IL 60611

ABSTRACT

Background: The coronavirus disease 2019 (COVID-19) pandemic raised concern for exposure to healthcare providers through aerosol generating procedures, such as bronchoalveolar lavage (BAL). Current society guidelines recommended limiting use of BAL to reduce operators' risk for infection, yet data on the infection rate for providers after BAL is sparse. Since March 2020, our institution used a modified protocol to perform over 450 BALs on intubated COVID-19 patients. We therefore sought to describe the subsequent infectious risks to providers associated with BAL.

Methods: Fifty-two pulmonary and critical care providers (faculty and fellows) at our tertiary-care, urban medical center were surveyed. Survey participants were asked to provide the number of BALs on COVID-19 patients they performed, the number of weeks they cared for intensive care unit (ICU) patients with COVID-19, and the results of any SARS-CoV-2 testing that they received. Participants were asked to assess the difficulty of BAL on intubated COVID-19 patients as compared to routine ICU BAL using a numeric perceived difficulty score ranging from 1 (easier) to 10 (harder).

Results: We received forty-seven responses from fifty-two surveyed (90% response rate), with 2 declining to participate. Many respondents (19/45, 42%) spent >5 weeks on an ICU service with COVID-19

patients. The number of BALs performed by providers ranged from 0 to >60. Sixteen of the 35 providers (46%) who performed BALs underwent at least one nasopharyngeal (NP) swab to test for SARS-CoV-2, but none were positive. Twenty-seven of the 35 providers (77%) who performed BALs underwent SARS-CoV-2 serology testing, and only one (3.7%) was positive. Respondents indicated occasionally not being able to follow aerosol-minimizing steps but overall felt BALs in COVID-19 patients was only slightly more difficult than routine bronchoscopy.

Discussion: At a high-volume center having performed >450 BALs on intubated COVID-19 patients with aerosol-limiting precautions, our survey of bronchoscopists found no positive NP SARS-CoV-2 tests and only one positive antibody test result. While the optimal role for COVID-19 BAL remains to be determined, these data suggest that BAL can be safely performed in intubated COVID-19 patients if experienced providers take precautions to limit aerosol generation and wear personal protective equipment.

INTRODUCTION

The coronavirus disease 2019 (COVID-19) pandemic has caused tremendous amounts of morbidity and mortality. Bacterial co-infections in viral pneumonia have been well described, especially in the setting of influenza pandemics, and have been associated with significant mortality (Martin-Loeches et al., 2017). Preliminary studies from COVID-19 have also shown found bacterial co-infection, with especially high rates in those with fatal outcomes (Zhou et al., 2020), suggesting that identification of bacterial co-infection may affect COVID-19 outcomes (Cox et al., 2020). Preliminary data on a subset of our population found high coinfection rates (Grant et al., 2020). Given the long average duration of ventilation in COVID-19 patients (Madhok & Mihm, 2020), they are also at high risk for secondary infection with nosocomial bacterial organisms (Koenig & Truwit, 2006).

Therefore, an accurate diagnosis of bacterial superinfection of ventilated COVID-19 patients is important for management, especially to guide appropriate antibiotic stewardship (Pickens & Wunderink, 2019; Wunderink et al., 2020). Bronchoalveolar lavage (BAL) with quantitative cultures can be a helpful diagnostic test in the management of severe pneumonia (Torres et al., 2017). We have a longstanding practice of examining lower respiratory tract samples with rapid diagnostic platforms to inform the diagnosis and treatment of severe pneumonia, as well as research the alveolar microenvironment in severe pneumonia (Reyfman et al., 2019; Walter et al., 2018).

The standard in our intensive care unit (ICU) had been nonbronchoscopic BAL performed by respiratory therapists, with occasional bronchoscopic BAL for more challenging cases. Since the primary method of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) transmission is aerosolization of respiratory droplets (Shereen et al. 2020), nonbronchoscopic BAL was felt to be contraindicated. Endotracheal aspirates have been shown to be inferior to BAL (Fagon et al., 2000; Fujitani et al., 2009) When done correctly for diagnostic purposes, this maneuver also involves a long break in the closed ventilator circuit (Ranzani et al., 2019), and thus is not preferred. Bronchoscopic BAL was therefore felt to be the safest procedure for determining the presence and etiology of bacterial infections in intubated COVID-19 patients.

There is concern that bronchoscopy exposes healthcare workers by generating aerosols (Tran et al., 2012). Society and panel guidelines discourage routine use of bronchoscopy in patients with COVID-19, though this is only based on expert opinion (Wahidi et al., n.d., 2020). A paucity of data informs the infectious risk to providers performing BAL in COVID-19 patients. In a single-center report of 101 COVID-19 BALs, one of two bronchoscopists became sick with COVID-19 and had to be replaced by a third, who remained uninfected for the duration of the study (Torrego et al., 2020). Some studies where COVID-19 BAL was performed mentioned low provider infection rates, though details are not provided (Kon et al., 2020).

We therefore designed a safety-modified protocol for BALs in patients intubated with respiratory failure from known or suspected COVID-19. To-date in our tertiary-care, urban academic medical ICU, we have performed over 450 bronchoscopies on such patients. Given our high volume of BALs, we designed the following study to assess the incidence of SARS-CoV-2 infection and seropositivity among bronchoscopists.

METHODS

We surveyed all clinical faculty and fellows in the Pulmonary and Critical Care Division at Northwestern Memorial Hospital (NMH) in Chicago, IL. (Survey available as Supplementary 1.) Survey participants were asked to provide the number of COVID-19 BALs that they personally performed or directly participated in, the number of weeks they cared for ICU patients with COVID-19, and the results of any SARS-CoV-2 testing that they had received. Participants were asked to assess the difficulty of BALs on intubated COVID-19 patients compared to routine ICU BALs using a numeric perceived difficulty score ranging from 1 (easier) to 10 (harder). They were also queried regarding COVID-19 exposures outside of work. No identifiers were collected, and all respondents were offered the choice to decline to participate. This study was reviewed and deemed exempt by our institution's Institutional Review Board.

SARS-CoV-2 testing

Nasopharyngeal (NP) testing for SARS-CoV-2 by polymerase chain reaction (PCR) (Becton and Dickinson, Becton, Dickinson and Company, Franklin Lakes, NJ; Cepheid, Sunnyvale, CA; and an

institutionally-developed platform) was offered to providers *ad hoc* at the discretion of our hospital's

infection control team, either in response to providers' symptoms, after a known exposure without PPE, or

routine pre-procedure screening for the provider's own medical care. Serology (Abbott Architect SARS-

CoV IgG, Abbott Park, IL) testing was offered to all medical staff by NMH starting in June 2020.

COVID-19 bronchoscopy protocol

Bronchoscopies were performed in the ICU on intubated patients with respiratory failure from

either known or suspected COVID-19. The decision to perform bronchoscopy was at the discretion of the

ICU team, in some cases to establish a diagnosis or in others to exclude bacterial co-

infection. Bronchoscopies were performed by a pulmonary ICU attending, interventional pulmonology

(IP) attending, and/or Pulmonary/IP fellows. Nurses and respiratory therapists were not in the rooms

during the actual bronchoscopy. PPE included an N95 mask, eye protection, gloves, gown, and hair

protection. Bronchoscopy was performed with a disposable Ambu® aScope™ (Ambu Inc., Columbia,

MD) bronchoscope. Patients were sedated per ICU protocols or at the proceduralist's discretion.

Administration of cisatracurium (0.1 mg/kg) in order to minimize coughing during bronchoscopy was

recommended. The endotracheal tube was clamped while the ventilatory circuit was manipulated to

accommodate scope placement, and the inspiratory limb of the ventilator was transiently disconnected

during the brief break in circuit.

Statistical Analyses

Survey results were exported from Qualtrics (Provo, UT) and compiled in Microsoft Excel

(Version 15.39 for Mac, Microsoft, Redmond, WA). Graphs were generated in GraphPad Prism 8

(version 8.4.3, GraphPad Software LLC), Adobe Illustrator (Version CC for Mac, Adobe Inc, San Jose,

CA), and Python using Seaborn v0.10.1. Not all continuous data were normally distributed, and so

median values with interquartile ranges (IQR) were reported. Non-parametric analyses included Wilcoxon

rank sum for categorical and Spearman's rank correlation for continuous variables. Kruskal-Wallis rank

testing was used to compare values across multiple categories.

RESULTS

We surveyed 52 clinical Pulmonary and Critical Care faculty and fellows. Forty-five respondents (90% response rate) agreed to participate: 18 fellows and 27 attendings (Table 1, Figure 1). The majority (35/45, 78%) had performed at least one bronchoscopy. The most common range of bronchoscopies performed by a respondent was 10-30 bronchoscopies, which 14/45 (31%) respondents selected. Two respondents reported performing bronchoscopy without full PPE under emergent situations, and 12 reported performing bronchoscopies without being able to follow the full protocol.

Many respondents (42%) spent >5 weeks on an ICU service with COVID-19 patients. The number of weeks on ICU service correlated to bronchoscopy volume (Spearman r=0.66, p<0.05). The median perceived difficulty score was 6 (IQR 6-7), with fellows finding the procedure more difficult than attendings (p<0.05). The perceived difficulty score did not correlate with either bronchoscopy volume (p=0.5) or time spent on ICU service with COVID-19 patients (p=0.18). Comments included that respondents felt these bronchoscopies were quite safe given numerous precautions instituted and adequate availability of PPE.

Sixteen of the 35 providers (46%) who performed bronchoscopies underwent at least one nasopharyngeal swab for SARS-CoV-2. No respondents reported a positive test. Five respondents indicated having a respiratory illness that prompted their testing. Eight respondents indicated their SARS-CoV-2 testing was for screening purposes only.

Twenty-seven of the 35 providers (77%) who performed bronchoscopies underwent SARS-CoV-2 serology testing. One individual (3.7%) had two sets of serology testing, with one result being positive and another result being negative. This individual spent >5 weeks on an ICU service with COVID-19 patients, performed an estimated 10-30 bronchoscopies, but did not have any febrile respiratory symptoms.

DISCUSSION

Bronchoalveolar lavage plays an important role in the evaluation of acute respiratory failure in the ICU. Current society guidelines recommend a limited role for bronchoscopy in the care of COVID-19 patients to minimize the potential infectious risk to beside providers of aerosol-generating procedures

(Wahidi et al., 2020). These guidelines are based on expert opinion and appropriately emphasize provider safety, but data regarding the infectious risk of BAL to providers have been lacking. Our survey is small and limited to a single center, but our data suggest that the risk of transmitting COVID-19 to providers performing BAL may be low. No provider developed COVID-19, and only one had positive serology.

This survey was not a formal study of our specific bronchoscopy protocol. However, we hypothesized that careful technique during bronchoscopy can limit infectious risk. Principles of our protocol included: 1) limiting the number of providers in the room, 2) minimizing aerosol generation by with clamping the endotracheal tube and disconnecting the inspiratory limb of the ventilator during manipulations, 3) minimizing cough by neuromuscular blockade or heavy sedation and instillation of lidocaine into the tracheobronchial tree, and 4) use of a disposable bronchoscope. We also had a small number of highly skilled providers (our IP team) performing a high number of the BALs, ensuring adherence to protocol. Fortunately, most providers felt that the bronchoscopies following the protocol were only slightly more difficult than routine ICU bronchoscopy. We suspect that other centers may have similar protocols, and a more scientific exploration of techniques would be needed to comment on protocol efficacy.

Our group has supported adhering to evidence-based critical care during the COVID-19 pandemic rather than making practice changes solely in response to uncertainties associated with the COVID pandemic (Singer et al., 2020). Medical provider safety is of paramount importance and precluded relying on our standard of care nonbronchoscopic BAL performed by respiratory therapists. However, our providers already utilize BAL during their routine critical care practice, and were likely to be able to safely continue that practice while using appropriate PPE and minimizing aerosolization. Our results demonstrate that assumption was accurate.

Our study has several limitations. First, this is a retrospective survey of providers and their recollection of practices. We did not directly pull exact bronchoscopy numbers and test results from the electronic medical records so as to respect our colleagues' privacy, and instead looked for voluntary responses. However, collateral data suggests overestimation of bronchoscopy numbers is very unlikely. Adherence to the modified bronchoscopy protocol was not monitored. We were not able to capture a 100% response rate for the survey but obtained results higher than most physician survey response rates

(Kellerman & Herold, 2001). While serology testing was offered to all providers per hospital protocol, only a subset of providers actually got tested. NP testing was performed *ad hoc* based upon symptoms or exposure. Since neither form of testing was mandated across the survey cohort, both may in fact under-represent the true infectious risk. It is also possible that asymptomatic infections were missed if providers did not seek testing in a short time period after performing said procedures.

In summary, this is the first granular report of infectious risks from BAL from a large-volume center that routinely incorporates bronchoscopy as part of critical care for patients with COVID-19 respiratory failure. While additional research is needed to inform the optimal use of invasive sampling of the lower respiratory tract to improve outcomes for patients with COVID-19, our data suggest that BAL can be routinely incorporated into the ICU care of these patients with minimal infectious risk to providers.

Table 1: Survey Results

	Total	Attendings	Fellows
Total Surveyed Providers	52	31	21
Responded to survey	47		
Agreed to complete survey	45	27	18
Weeks of COVID-19 ICU Service			
0	9	7	2
1	2	2	0
2	3	1	2
3	4	4	0
4	8	6	2
>5	19	7	12
No answer			
Number of Bronchoscopies			
0	10	8	2
1 - 10	9	6	3
10 - 30	15	9	6
30 - 60	5	0	5
>60	6	4	2
No answer			
Perceived Difficulty Scores (median, IQR)	6 (6-7)	6 (6-7)	6.5 (5.25-7)

Figure 1: Survey Response Rates and COVID-19 Test Results

Figure 2: Weeks of COVID-19 ICU Service and Bronchoscopy Volume

Funding:

Dr. Wunderink is supported by a NIH grant U19AI135964 and a GlaxoSmithKline Distinguished Scholar in Respiratory Health grant from the CHEST Foundation.

Conflicts of Interest: The authors report none.

Author contributions:

CAG and SBS had full access to all of the data and took responsibility for the integrity of the data and accuracy of data analysis.

Concept and design: CAG, JIB, JMC, AAC, ESM, RGW, SBS

Survey: CAG, JIB, RGW, ACA

Compiling data: CAG, JIB

Statistical analysis: CAG, SBS

Drafting of manuscript: CAG, JIB

Revision of manuscript: CAG, JIB, JMW, JMC, RGW, SBS

Supervision: RGW, ACA, SBS

Acknowledgements: We thank the many individuals who took care of the COVID-19 patients; this list of physicians and APPs is recognized by their inclusion in the NU COVID Investigators. We thank all the nurses, respiratory therapists, social workers, physical therapists, and providers who cared for these patients.

Works Cited

Cox, M. J., Loman, N., Bogaert, D., & O'Grady, J. (2020). Co-infections: potentially lethal and unexplored
in COVID-19. *The Lancet. Microbe*, *1*(1), e11.

Fagon, J. Y., Chastre, J., Wolff, M., Gervais, C., Parer-Aubas, S., Stéphan, F., Similowski, T., Mercat, A.,
Diehl, J. L., Sollet, J. P., & Tenaillon, A. (2000). Invasive and noninvasive strategies for
management of suspected ventilator-associated pneumonia. A randomized trial. *Annals of
Internal Medicine*, *132*(8). https://doi.org/10.7326/0003-4819-132-8-200004180-00004

Fujitani, S., Cohen-Melamed, M. H., Tuttle, R. P., Delgado, E., Taira, Y., & Darby, J. M. (2009).
Comparison of semi-quantitative endotracheal aspirates to quantitative non-bronchoscopic
bronchoalveolar lavage in diagnosing ventilator-associated pneumonia. *Respiratory Care*, *54*(11).
https://pubmed.ncbi.nlm.nih.gov/19863828/

Grant, R. A., Morales-Nebreda, L., Markov, N. S., Swaminathan, S., Guzman, E. R., Abbott, D. A.,
Donnelly, H. K., Donayre, A., Goldberg, I. A., Klug, Z. M., Borkowski, N., Lu, Z., Kihshen, H.,
Politanska, Y., Sichizya, L., Kang, M., Shilatifard, A., Qi, C., Christine Argento, A., … for The NU
SCRIPT Study Investigators. (2020). *Alveolitis in severe SARS-CoV-2 pneumonia is driven by
self-sustaining circuits between infected alveolar macrophages and T cells* (p.
2020.08.05.238188). https://doi.org/10.1101/2020.08.05.238188

Kellerman, S. E., & Herold, J. (2001). Physician response to surveys. A review of the literature. *American
Journal of Preventive Medicine*, 20(1), 61–67.

Koenig, S. M., & Truwit, J. D. (2006). Ventilator-associated pneumonia: diagnosis, treatment, and
prevention. *Clinical Microbiology Reviews*, *19*(4), 637–657.

Kon, Z. N., Smith, D. E., Chang, S. H., Goldenberg, R. M., Angel, L. F., Carillo, J. A., Geraci, T. C.,
Cerfolio, R. J., Montgomery, R. A., Moazami, N., & Galloway, A. C. (2020). Extracorporeal
Membrane Oxygenation Support in Severe COVID-19. *The Annals of Thoracic Surgery*.
https://doi.org/10.1016/j.athoracsur.2020.07.002

Madhok, J., & Mihm, F. G. (2020). Rethinking sedation during prolonged mechanical ventilation for
COVID-19 respiratory failure. *Anesthesia and Analgesia*.
https://doi.org/10.1213/ANE.0000000000004960

Martin-Loeches, I., Schultz M, J., Vincent, J. L., Alvarez-Lerma, F., Bos, L. D., Solé-Violán, J., Torres, A., & Rodriguez, A. (2017). Increased incidence of co-infection in critically ill patients with influenza. *Intensive Care Medicine*, *43*(1). https://doi.org/10.1007/s00134-016-4578-y

Pickens, C. I., & Wunderink, R. G. (2019). Principles and Practice of Antibiotic Stewardship in the ICU. In *Chest* (Vol. 156, Issue 1, pp. 163–171). https://doi.org/10.1016/j.chest.2019.01.013

Ranzani, O. T., Senussi, T., Idone, F., Ceccato, A., Bassi, G. L., Ferrer, M., & Torres, A. (2019). Invasive and non-invasive diagnostic approaches for microbiological diagnosis of hospital-acquired pneumonia. *Critical Care*, 23(1), 1–11.

Reyfman, P. A., Walter, J. M., Joshi, N., Anekalla, K. R., McQuattie-Pimentel, A. C., Chiu, S., Fernandez, R., Akbarpour, M., Chen, C. I., Ren, Z., Verma, R., Abdala-Valencia, H., Nam, K., Chi, M., Han, S., Gonzalez-Gonzalez, F. J., Soberanes, S., Watanabe, S., Williams, K. J. N., … Misharin, A. V. (2019). Single-Cell Transcriptomic Analysis of Human Lung Provides Insights into the Pathobiology of Pulmonary Fibrosis. *American Journal of Respiratory and Critical Care Medicine*, *199*(12). https://doi.org/10.1164/rccm.201712-2410OC

Singer, B. D., Jain, M., Budinger, G. R. S., & Wunderink, R. G. (2020). A Call for Rational Intensive Care in the Era of COVID-19. *American Journal of Respiratory Cell and Molecular Biology*, *63*(1). https://doi.org/10.1165/rcmb.2020-0151LE

Torrego, A., Pajares, V., Fernández-Arias, C., Vera, P., & Mancebo, J. (2020). Bronchoscopy in Patients with COVID-19 with Invasive Mechanical Ventilation: A Single-Center Experience. *American Journal of Respiratory and Critical Care Medicine*, *202*(2), 284–287.

Torres, A., Niederman, M. S., Chastre, J., Ewig, S., Fernandez-Vandellos, P., Hanberger, H., Kollef, M., Li Bassi, G., Luna, C. M., Martin-Loeches, I., Paiva, J. A., Read, R. C., Rigau, D., Timsit, J. F., Welte, T., & Wunderink, R. (2017). International ERS/ESICM/ESCMID/ALAT guidelines for the management of hospital-acquired pneumonia and ventilator-associated pneumonia: Guidelines for the management of hospital-acquired pneumonia (HAP)/ventilator-associated pneumonia (VAP) of the European Respiratory Society (ERS), European Society of Intensive Care Medicine (ESICM), European Society of Clinical Microbiology and Infectious Diseases (ESCMID) and Asociación Latinoamericana del Tórax (ALAT). *The European Respiratory Journal: Official*

Journal of the European Society for Clinical Respiratory Physiology, *50*(3). https://doi.org/10.1183/13993003.00582-2017

Tran, K., Cimon, K., Severn, M., Pessoa-Silva, C. L., & Conly, J. (2012). Aerosol Generating Procedures and Risk of Transmission of Acute Respiratory Infections to Healthcare Workers: A Systematic Review. *PloS One*, *7*(4). https://doi.org/10.1371/journal.pone.0035797

Wahidi, M. M., Lamb, C., Murgu, S., Musani, A., Shojaee, S., Sachdeva, A., Maldonado, F., Mahmood, K., Kinsey, M., Sethi, S., Mahajan, A., Majid, A., Keyes, C., Alraiyes, A. H., Sung, A., Hsia, D., & Eapen, G. (2020). American Association for Bronchology and Interventional Pulmonology (AABIP) Statement on the Use of Bronchoscopy and Respiratory Specimen Collection in Patients with Suspected or Confirmed COVID-19 Infection. *Journal of Bronchology & Interventional Pulmonology*. https://doi.org/10.1097/LBR.0000000000000681

Wahidi, M. M., Shojaee, S., Lamb, C. R., Ost, D., Maldonado, F., Eapen, G., Caroff, D. A., Stevens, M. P., Ouellette, D. R., Lilly, C., Gardner, D. D., Glisinski, K., Pennington, K., & Alalawi, R. (n.d.). The Use of Bronchoscopy During the Coronavirus Disease 2019 Pandemic: CHEST/AABIP Guideline and Expert Panel Report. *Chest*. https://doi.org/10.1016/j.chest.2020.04.036

Walter, J. M., Helmin, K. A., Abdala-Valencia, H., Wunderink, R. G., & Singer, B. D. (2018). Multidimensional assessment of alveolar T cells in critically ill patients. *JCI Insight*, *3*(17). https://doi.org/10.1172/jci.insight.123287

Wunderink, R. G., Srinivasan, A., Barie, P. S., Chastre, J., Cs, D. C., Douglas, I. S., Ecklund, M., Evans, S. E., Evans, S. R., Gerlach, A. T., Hicks, L. A., Howell, M., Hutchinson, M. L., Hyzy, R. C., Kane-Gill, S. L., Lease, E. D., Metersky, M. L., Munro, N., Niederman, M. S., … Weiss, C. H. (2020). Antibiotic Stewardship in the Intensive Care Unit. An Official American Thoracic Society Workshop Report in Collaboration with the AACN, CHEST, CDC, and SCCM. *Annals of the American Thoracic Society*, *17*(5). https://doi.org/10.1513/AnnalsATS.202003-188ST

Zhou, F., Yu, T., Du, R., Fan, G., Liu, Y., Liu, Z., Xiang, J., Wang, Y., Song, B., Gu, X., Guan, L., Wei, Y., Li, H., Wu, X., Xu, J., Tu, S., Zhang, Y., Chen, H., & Cao, B. (2020). Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*, *395*(10229). https://doi.org/10.1016/S0140-6736(20)30566-3

Supplementary

NMH COVID-19 ICU Bronchoscopy protocol:

Bronchoscopy in the COVID ICU will be performed for diagnostic and therapeutic purposes, including, but not limited to:
- Diagnostic evaluation of newly intubated patient, including COVID rule-out testing
- Evaluation of possible VAP
- Airway clearance

Bronchoscopy can be performed at the discretion of the ICU attending, but generally policy will be that off-hour bronchoscopy performed by a fellow without an attending should be limited to emergent situations such as mucous plugging. The following protocol will be used for bronchoscopy and will be performed by 1-2 member(s) of the IP service or the COVID ICU attending. The COVID ICU fellow will participate if willing and available (per fellow's preference).

Given the longer circuit break with NBBAL, these will not be performed on COVID-19 patients.

The nurse will help with printing order labels, pre-procedural sedation including administration of neuromuscular blockade but then will leave the room.
RT may help with gathering equipment but does not need to be present in the room.

Role 1: primary bronchoscopist (fellow, attending, IP attending) – this person is responsible for assessing the clinical situation, consenting the family, ensuring all necessary equipment are ready, ordering lab tests, ensuring adequate sedation, performing the bronchoscopy, cleaning the equipment afterwards, and ensuring samples delivered to the lab.

Role 2: secondary bronchoscopist (supervising IP attending or COVID attending or fellow) – this person will assist the primary bronchoscopist, silence ventilator alarms, assist in circuit manipulation, instill saline for lavage, withdraw BAL, and connect lukens trap.

Order checklist:
- Adequate sedation
- Cisatricurium 0.2mg/kg (Pharmacist may often have, otherwise call 9th floor pharmacy)
- Labwork
  - BAL cell count and differential
  - BAL amylase
  - BAL respiratory culture (normal, +/- fungal, AFB per provider suspicion)
  - Lower Respiratory Tract Panel (BioFire Pneumonia Panel)
  - SARS-CoV-2 test (even if already know status, to monitor for clearance/reinfection)
  - Cytology per provider suspicion
  - Galactomannan, PJP DFA, etc – per provider suspicion
  - Extra patient label for research specimen

PPE checklist:
- N95 mask (ensure adequate fit) + covering surgical mask
- Gown, gloves (discard after procedure)
- Faceshield or goggles (wipe down after procedure)

Equipment checklist:
- Ambu scope (large/orange if concern for mucus plugging, but check ETT size; regular/green otherwise)
- Ambu tower (ensure adequately charged)
- Drape or chuck to lay down supplies on
- Scope adaptor for ETT

- Clamp for ETT
- Scope lubricant
- Extra suction tubing, if far from the bed
- 4x 30cc syringes (slip tip preferred – if luer lock, ensure slip tip adaptors available; often in the scope bag)
- Normal saline (500cc bottles)
- Lukens trap
- Orange specimen cup + labels
- Research Eppendorf tube + patient sticker
- Extra specimen bags outside room for double-bagging specimens

Procedure steps:
1. Outside the room, Primary Bronchoscopist times out with nursing and ensures consent has been signed, orders placed and labels printed. Sedation adjusted to goal RASS-4 with cisatraciurium 0.2mg/kg administered at provider's discretion. FiO2 increased to 100%.
2. As able, Primary Bronchoscopist preps the bronch equipment outside the room
   a. Drawing up 30cc normal saline x4
   b. Connecting AmbuScope to Monitor
   c. Lubricating AmbuScope
   d. Pre-loading AmbuScope onto scope adaptor
   e. Applying labels to specimen cups and research tube
3. Primary Bronchoscopist and Secondary Bronchoscopist enter room; Nursing and RT available but are outside the room.
   a. Primary Bronchoscopist responsible for narrating steps out loud, stands at the side of the patient; Secondary Bronchoscopist stands next to first, closer to the ventilator
   b. Primary Bronchoscopist ensures equipment set up, suction connected and functioning, sedation is adequate and vitals are stable to tolerate the procedure
   c. Primary Bronchoscopist clamps ETT
   d. Secondary Bronchoscopist disconnects inspiratory limb from the vent distal to the filter
   e. Primary Bronchoscopist places the adaptor (pre-loaded with the scope) onto the ETT
   f. Secondary Bronchoscopist reconnects the inspiratory limb
   g. Primary Bronchoscopist unclamps the ETT
   h. Primary Bronchoscopist performs inspection, toileting secretions as needed, wedges into target lobe
   i. Secondary Bronchoscopist instills saline in 30cc aliquots, 120cc recommended, draws back and discards first 5 cc, draws back more sample if able, then connects the lukens trap (goal >40cc return)
   j. Primary Bronchoscopist suctions sample into lukens trap
   k. Secondary Bronchoscopist disconnect lukens trap, hooks back up to wall suction
   l. Primary Bronchoscopist cleans up any remaining secretions, pulls scope back to edge of adaptor, then clamps ETT
   m. Secondary Bronchoscopist disconnects inspiratory limb from the vent distal to the filter
   n. Primary Bronchoscopist removes adaptor and scope in one motion
   o. Secondary Bronchoscopist reconnects the inspiratory limb
   p. Primary Bronchoscopist unclamps the ETT
   q. Sample placed into orange specimen cup from lukens trap (have had several break), with 10-15cc placed in Eppendorf tube for research team
      i. Samples bagged first in room, then placed in another bag held by someone outside the room
      ii. Orange specimen cup + labels delivered to 7th floor lab (can tube if double-bagged)
      iii. Research specimen - The MICU research team can be contacted by phone at 62752 or by pager at 59285. If they are not readily available to pick up the specimen, it can be left in the specimen fridge in the dirty utility room in the 9th floor MICU.

r. Disposable equipment placed in red biohazard bag for disposal
s. Monitor and pole wiped down before leaving room, returned to RT room
t. Primary Bronchoscopist returns FiO2 to pre-procedure level (assuming tolerated) and ensures hemodynamics acceptable
u. Both proceduralists doff PPE and wash hands; surgical mask over N95 should be discarded but N95 can be reused; goggles/faceshields wiped down and reused
v. Primary Bronchoscopist documents procedure note

Special Circumstances:
- Prone positioning: continue with standard procedure
- iNO: continue with standard procedure
- Brushings: sometimes requested by research team, who will provide brushes, brush cutter (wire cutter can also be found in bronch suite), research tube/medium for brush to be cut into

Please do not hesitate to contact the Interventional Pulmonary team with questions.

Protocol updated 8-29-2020 by Catherine Gao, from version 4-5-20 from COVID-ICU-Guidelines.

Supplementary: Survey:

Title of Research Study: Provider safety during COVID-19 bronchoscopies

IRB Study Number: STU00213164, version 7-24-20

Principal Investigator: Sean Smith

Supported By: This research is supported by Divison of Pulmonary and Critical Care.

Key Information about this research study: The following is a short summary of this study to help you

decide whether to be a part of this study. The purpose of this study is to evaluate the safety of

bronchoscopists in COVID-19 patients. You will be asked to complete a survey on the number of

bronchoscopies you have performed on COVID patients, the number of COVID tests you have received,

and the results of your COVID testing. We expect that you will be in this research study for the duration

of this survey.

The primary risk of participation is potentially being identified based on how many bronchoscopies you

have performed and the small cohort of participants. The main benefit of participation is publicizing the

risks/safety of bronchoscopy on COVID patients. Why am I being asked to take part in this research

study?

We are asking you to take part in this research study because you have potentially bronched a COVID-19

patient.

How many people will be in this study? We expect about 40 people here will be in this research study.

What should I know about a research study?

·   Whether or not you take part is up to you.

·   You can choose not to take part.

·   You can agree to take part and later change your mind.

·   Your decision will not be held against you.

If you say that "Yes, you want to be in this research," here is what you will do:You will be asked to

complete a survey on the number of bronchoscopies you have performed on COVID patients, the number

of COVID tests you have received, and the results of your COVID testing. Will being in this study help me in any way?

We cannot promise any benefits to you from your taking part in this research. However, possible benefits include publicizing the risks/safety of bronchoscopy on COVID patients

Is there any way being in this study could be bad for me?

The primary risk of participation is potentially being identified based on how many bronchoscopies you have performed and the small cohort of participants.

What happens if I do not want to be in this research or if I say "Yes", but I change my mind later?

Participation in research is voluntary. You can decide to participate or not to participate.

You can decide not to participate in this research or you can start and then decide to leave the research at any time and it will not be held against you. To do so, simply exit the survey. Any data already collected will not be saved.

What happens to the information collected for the research?

Efforts will be made to limit the use and disclosure data gathered to people who have a need to review this information. We cannot promise complete secrecy. Organizations that may inspect and copy your information include the IRB and other representatives of this institution.

This survey is being hosted by Qualtrics and involves a secure connection. Upon receiving results of your survey, any possible identifiers will be deleted. You will be identified only by a unique subject number. All information will be kept on a password protected file only accessible by the research team. The results of the research study may be published, but your name will not be used.

Data Sharing

De-identified data from this study may be shared with the research community at large to advance science and health. We will remove or code any personal information that could identify you before files are shared with other researchers to ensure that, by current scientific standards and known methods, no one will be able to identify you from the information we share. Despite these measures, we cannot guarantee anonymity of your personal data.

What else do I need to know?

If you agree to take part in this research study, we will provide you with the survey questions.

Who can I talk to?

If you have questions, concerns, or complaints talk to the Principal Investigator Sean Smith, sean@northwestern.edu, or Catherine Gao, catherine.gao@northwestern.edu

This research has been reviewed and approved by an Institutional Review Board ("IRB"). You may talk to them at (312) 503-9338 or irb@northwestern.edu if:

·   Your questions, concerns, or complaints are not being answered by the research team.

·   You cannot reach the research team.

·   You want to talk to someone besides the research team.

·   You have questions about your rights as a research participant.

·   You want to get information or provide input about this research.

Do you agree to participate?

    I agree or I decline

What is your role?

    Attending or Fellow

How many weeks did you spend on the COVID ICU services?

    0, 1, 2, 3, 4, >5, other:

How many bronchoscopy procedures did you perform on COVID positive patients? Please provide the exact number if you have this available, or select an estimate from the ranges offered:

    0, 1-10, 10-30, 30-60, >60, other:

Were you ever tested for COVID via nasopharyngeal swab?

Was this for symptoms or for screening or in response to symptoms?

If yes, were any of these tests positive?

Were you ever tested for COVID-19 by antibody serology testing?

If yes, did you have detectable COVID antibodies?

Did you have any febrile respiratory illness during the COVID-19 pandemic (starting Jan 1 2020 to now)?

Yes/No

Did you ever perform a bronchoscopy without access to the recommended COVID-19 personal protective equipment – N95, eye protection, gown, and gloves?

If yes, estimate how many procedures were performed without recommended PPE:

Did you ever perform a bronchoscopy on a COVID positive patient without following the droplet minimization steps of clamping the endotracheal tube and disconnecting the ventilator? Yes/No

If yes, estimate how many procedures were performed without following droplet mitigation procedures.

Did you have any other high risk COVID exposures outside of work?

Compared to a typical, non-COVID bronchoscopy in intubated patients, how much more difficult were the COVID bronchoscopies (eg, donning/doffing PPE, clamping ETT, disconnecting inspiratory limb of the ventilator)?

(1 – much easier, 5 – same, 10 – much more difficult; sliding scale)

Other questions or comments: (free text)

NU COVID Investigators

Name	ORCID	Email	Affiliation	
A. Christine Argento	0000-0002-3807-7690	cargento@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Ajay A. Wagh	0000-0003-3402-0272	awagh@medicine.bsd.uchicago.edu	UChicago Division of Pulmonary and Critical Care Medicine	
Alexandra C. McQuattie-Pimentel	0000-0003-4028-9905	amhanrahan@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Alexis Rose Wolfe	0000-0001-9321-892X	alexis.wolfe@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Ankit Bharat	0000-0002-1248-0457	ankit.bharat@nm.org	Division of Thoracic Surgery	Division of Pulmonary and Critical Care Medicine
Anne R. Levenson	0000-0002-7587-1885	anne.levenson@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Anthony M. Joudi	0000-0003-1232-3256	anthony.joudi@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Arjun Sinha	0000-0001-9794-8209	arjun.sinha@northwestern.edu	Division of Cardiology	
Ashley N. Budd	0000-0002-2879-9294	ashley.budd@nm.org	Department of Anesthesiology	
Benjamin D. Singer	0000-0001-5775-8427	benjamin-singer@northwestern.edu	Division of Pulmonary and Critical Care Medicine	Department of Biochemistry and Molecular Genetics
Betty Tran	0000-0001-6794-7088	betty.tran@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Catherine A. Gao	0000-0001-5576-3943	catherine.gao@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Chiagozie O. Pickens	0000-0002-3400-9228	Chiagozie-Ononye@northwestern.edu	Division of Pulmonary and	

			Critical Care Medicine	
Chitaru Kurihara	0000-0003-3536-4675	Chitaru.kurihara@northwestern.edu	Division of Thoracic Surgery	
Christopher J. Soriano	0000-0003-3510-8866	christopher.soriano2@nm.org	Division of Organ Transplantation	
Clara J Schroedl	0000-0003-3280-4997	c-schroedl@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Daniel Meza	0000-0001-8154-1295	daniel.meza@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
David Alexander Kidd	0000-0003-0411-9511	david.kidd@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
David W. Kamp	0000-0001-5796-1206	d-kamp@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Elizabeth S. Malsin	0000-0002-3294-6752	elizabeth.malsin@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Emily M. Leibenguth	0000-0001-7809-7947	emily.leibenguth@nm.org	Division of Pulmonary and Critical Care Medicine	
Eric P. Cantey	0000-0002-7065-7169	Eric.cantey@northwestern.edu	Division of cardiology	
Gabrielle Y. Liu	0000-0003-2049-423	gabrielle.liu@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Giang T. Quach	0000-0002-4558-3348	giang.quach@northwestern.edu	Division of Trauma and Surgical Critical care	
GR Scott Budinger	0000-0002-3114-5208	s-buding@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Jacqueline M. Kruser	0000-0003-3258-1869	jacqueline.kruser@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
James M. Walter	0000-0001-7428-3101	james.walter@northwestern.edu	Division of Pulmonary and Critical Care Medicine	

Jane E. Dematte	0000-0003-4907-9525	j-dematte@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Jason A. Bonomo	0000-0002-6214-2587	jason.bonomo@northwestern.edu	Division of Cardiology	
John M. Coleman	0000-0003-1246-3814	jcolema2@nm.org	Division of Pulmonary and Critical Care Medicine	
Joseph Isaac Bailey	0000-0002-4225-3108	joseph.bailey1@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Joseph S. Deters	0000-0002-3522-3824	joseph.deters@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Joseph Sun			Department of Anesthesiology	
Justin A. Fiala	0000-0001-5196-7076	justin.fiala@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Kaitlyn Vitale		kaitlyn.vitale@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Kara M. Joseph		kjoseph9@bwh.harvard.edu	Department of Anesthesiology	
Katharine Secunda	0000-0003-0310-5783	k-secunda@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Khalilah L. Gates	0000-0002-5965-8136	k-gates@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Kristy Todd	0000-0001-8210-8522	kristy.todd2@nm.org	Division of Pulmonary and Critical Care Medicine	
Lindsey D. Gradone	0000-0001-8145-2997	lindsey.gradone@nm.org	Division of Pulmonary and Critical Care Medicine	
Lindsey N. Textor	0000-0001-7699-9429	lindsey.textor@nm.org	Division of Pulmonary and Critical Care Medicine	

Lisa F. Wolfe	0000-0001-9044-4147	lwolfe@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Luisa Morales-Nebreda	0000-0002-7126-0518	luisamorales@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Madeline L Rosenbaum	0000-0001-9866-9850	madeline.rosenbaum@nm.org	Division of Pulmonary and Critical Care Medicine	
Manu Jain	0000-0003-1534-5629	m-jain@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Marc A. Sala	0000-0002-2900-0595	marc.sala@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Mark Saine	0000-0002-3516-4254	mark.saine@nm.org	Division of Thoracic Surgery	
Marysa V. Leya	0000-0002-6583-5389	Marysa-Leya@northwestern.edu	Division of Cardiology	
Michael J. Alexander	0000-0001-6271-3426	michael.alexander@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Michael J. Cuttica	0000-0002-0084-8760	m-cuttica@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Michelle Hinsch Prickett	0000-0002-1119-0079	micprick@nm.org	Division of Pulmonary and Critical Care Medicine	
Natalie Jensema		natalie.jensema@nm.org		
Paul A. Reyfman	0000-0002-6435-6001	paul.reyfman@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Peter H. S. Sporn	0000-0002-1006-9437	p-sporn@northwestern.edu	Division of Pulmonary and Critical Care Medicine	Medical Service, Jesse Brown Veterans Affairs Medical Center
Rachel B. Kadar	0000-0002-2583-3637	rachel.kadar@northwestern.edu	Section of Critical Care Medicine,	

			Department of Anesthesiology	
Rachel M. Kaplan	0000-0003-0038-9231	rachel-kaplan1@northwestern.edu	Division of Cardiology	
Rade Tomic	0000-0003-2394-1287	rade.tomic@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Radhika Patel	0000-0002-1255-5818	radhika.patel@nm.org	Division of Pulmonary and Critical Care Medicine	
Rafael Garza-Castillon	0000-0003-4970-6584	rgarza@nm.org	Division of Thoracic Surgery	
Ravi Kalhan	0000-0003-2443-0876	r-kalhan@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Richard G. Wunderink	0000-0002-8527-4195	r-wunderink@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Romy Lawrence		romy.lawrence@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Ruben J. Mylvaganam	0000-0002-7203-7609	ruben.mylvaganam@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Samuel S. Kim	0000-0001-8889-478X	samuel.kim2@nm.org	Division of Thoracic Surgery	
Sanket Thakkar	0000-0002-0026-6533	sthakkar@nm.org	Division of Thoracic Surgery	
Sean B. Smith	0000-0001-8796-3966	sean@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
SeungHye Han	0000-0001-5625-6337	shan@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Sharon R. Rosenberg	0000-0002-0260-2545	sharon.rosenberg@nm.org	Division of Pulmonary and Critical Care Medicine	
Susan R. Russell	0000-0003-3006-1018	s-russell4@northwestern.edu	Division of Pulmonary and Critical Care Medicine	

Sydney M. Hyder	0000-0002-6395-9062	sydney.hyder@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Taylor A. Poor	0000-0002-2191-5037	taylor-poor@northwestern.edu	Division of Pulmonary and Critical Care Medicine	
Theresa A. Lombardo	0000-0002-7468-7541	theresa.lombardo@nm.org	Division of Pulmonary and Critical Care Medicine	