The impact of high frequency rapid viral antigen screening on COVID-19 spread and

outcomes: a validation and modeling study

Authors: Beatrice Nash[1,2*], Anthony Badea[1,3*], Ankita Reddy[1,4*], Miguel Bosch[1,5], Nol Salcedo[1],

Adam R. Gomez[1], Alice Versiani[6], Gislaine Celestino Dutra Silva[6], Thayza Maria Izabel Lopes

dos Santos[6], Bruno H. G. A. Milhim[6], Marilia M Moraes[6], Guilherme Rodrigues Fernandes

Campos[6], Flávia Quieroz[6], Andreia Francesli Negri Reis[6], Mauricio L. Nogueira[6], Elena N.

Naumova[7], Irene Bosch[1,8], Bobby Brooke Herrera[1,9†]

*These authors contributed equally to this work.

Affiliations:

[1]E25Bio, Inc., Cambridge, MA, USA

[2]Department of Computer Science, Harvard University School of Engineering and Applied

Sciences, Cambridge, MA, USA

[3]Department of Physics, Harvard University, Cambridge, MA, USA

[4]Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

[5]InfoGeosciences LLC, Houston, TX, USA

[6]Faculdade de Medicina de São José do Rio Preto (FAMERP), São José do Rio Preto, Brazil

[7]Division of the Nutrition Epidemiology and Data Science, Friedman School of Nutrition Science

and Policy, Tufts University, Boston, MA, USA

[8]Department of Medicine, Mount Sinai School of Medicine, New York, NY, USA

[9]Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of Public Health,

Boston, MA, USA

†Corresponding Author. BBH, email: bbherrera@e25bio.com

Abstract

High frequency screening of populations has been proposed as a strategy in facilitating

control of the COVID-19 pandemic. Here we use computational modeling, coupled with clinical

data from a rapid antigen test, to predict the impact of frequent rapid testing on COVID-19 spread

and outcomes. Using patient nasopharyngeal or nasal swab specimens, we demonstrate that the

sensitivity/specificity of two rapid antigen tests compared to quantitative real-time polymerase

chain reaction (qRT-PCR) are 80.0%/91.1% and 84.7%/85.7%, respectively; moreover, sensitivity

correlates directly with viral load. Based on COVID-19 data from three regions in the United States

and São José do Rio Preto, Brazil, we show that high frequency, strategic population-wide rapid

testing, even at varied accuracy levels, diminishes COVID-19 infections, hospitalizations, and

deaths at a fraction of the cost of nucleic acid detection via qRT-PCR. We propose large-scale

antigen-based surveillance as a viable strategy to control SARS-CoV-2 spread and to enable

societal re-opening.

INTRODUCTION

The COVID-19 pandemic has taken an unprecedented toll on lives, wellbeing, healthcare systems, and global economies. As of 25 September 2020, there have been more than 32.1 million confirmed cases globally with more than 980,000 confirmed deaths (*1*). However, these numbers and the current mapping of disease spread present an incomplete picture of the outbreak largely due to the lack of adequate testing, particularly as undetected infected cases are the main source of disease spread (*2–7*). It is estimated that the reported detection rate of actual COVID-19 cases is only 1-2% (*5*). As of September 2020, the United States and Brazil remain the top two countries with the highest number of COVID-19 cases and deaths worldwide. As countries begin to re-open their economies, a method for accessible and frequent surveillance of COVID-19, with the necessary rapid quarantine measures, is crucial to prevent the multiple resurgences of the disease.

The current standard of care rightfully places a strong focus on the diagnostic limit of detection, yet frequently at the expense of both cost and turnaround time. This situation has contributed to limited population testing largely due to a dearth of diagnostic resources. Quantitative real-time polymerase chain reaction (qRT-PCR) is the gold-standard method for clinical diagnosis, with high sensitivity and specificity, but these tests are accompanied by the need for trained personnel, expensive reagents and instrumentation, and a significant amount of time to execute. Facilities offering qRT-PCR sometimes require a week or longer to complete and return the results to the patient. During this waiting period the undiagnosed individual may spread the infection and/or receive delayed medical treatment. Moreover, due to the cost and relative inaccessibility of qRT-PCR in both resource-limited and abundant settings, large-scale screening using qRT-PCR at frequent intervals remains impractical to identify infected but asymptomatic or mildly symptomatic infections. Numerous studies have reported asymptomatic COVID-19 cases

as well as a variation in viral load within and between individuals at different time points,

suggesting the need for more frequent testing for informative surveillance.

Technologies alternate to qRT-PCR, such as rapid viral antigen detection, clustered

regularly interspaced short palindromic repeats (CRISPR), and loop-mediated isothermal

amplification (LAMP) of SARS-CoV-2 provide potential large-scale screening applications, yet

their implementation is stymied by requirements for qRT-PCR-like accuracy before they can reach

the market (*8*). In countries such as India, where the qRT-PCR resources would not be sufficient

to cover monitoring of the population, the use of rapid antigen tests is well underway(*9, 10*). In

early May 2020, the United States Food and Drug Administration (FDA) authorized the first

antigen test for the laboratory detection of COVID-19, citing a need for testing beyond molecular

and serological methods. Antigen testing detects the viral proteins rather than nucleic acids or

human antibodies, allowing for detection of an active infection with relative ease of sample

collection and assay. These rapid assays – like other commercially-available rapid antigen tests -

can be mass-produced at low prices and be administered by the average person without a laboratory

or instrumentation. These tests also take as little as 15 minutes to determine the result, enabling

real-time surveillance and/or diagnosis. Although antigen tests usually perform with high

specificities (true negative rate), their sensitivity (true positive rate) is often lower when compared

to molecular assays. While qRT-PCR can reach a limit of detection as low as 10^2 genome copies

per mL, rapid antigen testing detects viral protein that is assumed to correlate with approximately

10^5 genome copies per mL (*11*).

We hypothesize that frequent antigen-based rapid testing even with lower sensitivities

compared to qRT-PCR - along with appropriate quarantine measures - can be more effective at

decreasing COVID-19 spread than less frequent molecular testing of symptomatic individuals.

Keeping in mind the realities of daily testing in resource-limited regions, we also hypothesize that

testing frequency can be adjusted according to the prevalence of the disease; that is, an uptick in

reported cases should be accompanied by more frequent testing. During the viral incubation period,

high infectivity correlates with a high viral load that can be detected by either qRT-PCR or rapid

antigen testing (*12–16*). Rapid tests thus optimize diagnosis for the most infectious individuals.

Studies also point to the relatively small window of time during an individual's incubation period

in which the qRT-PCR assay is more sensitive than rapid tests (*12*).

In this study we report the clinical validation of two direct antigen rapid tests for detection

of SARS-CoV-2 spike glycoprotein (S) or nucleocapsid protein (N) using retrospectively collected

nasopharyngeal or nasal swab specimens. Using the clinical performance data, we develop a

modeling system to evaluate the impact of frequent rapid testing on COVID-19 spread and

outcomes using a variation of a SIR model, which has been previously used to model COVID-19

transmission (*17–23*). We build on this model to incorporate quarantine states and testing protocols

to examine the effects of different testing regimes. This model distinguishes between undetected

and detected infections and separates severe cases, specifically, those requiring hospitalization,

from those less so, which is important for disease response systems such as intensive care unit

triaging. We simulate COVID-19 spread with rapid testing and model disease outcomes in three

regions in the United States and São José do Rio Preto, Brazil - the site of the clinical validation

study - using publicly available data. To date, COVID-19 modeling describes the course of disease

spread in response to social distancing and quarantine measures, and a previous simulation study

has shown that frequent testing with accuracies less than qRT-PCR, coupled with quarantine

process and social distancing, are predicted to significantly decrease infections (*12, 17, 23–27*).

This is the first modeling system using publicly-available data to simulate how potential public

health strategies based on testing performance, frequency, and geography impact the course of

COVID-19 spread and outcomes. Our findings suggest that a rapid test, even with sensitivities

lower than molecular tests, when strategically administered 2-3 times per week, will reduce

COVID-19 spread, hospitalizations, and deaths at a fraction of the cost of nucleic acid testing via

qRT-PCR. Modern surveillance systems should be well equipped with rapid testing tools to ensure

that disease tracking and control protocols are effective and well-tailored to national, regional, and

community needs.

RESULTS

Accuracy of Direct Antigen Rapid Tests Correlate with Viral Load Levels

Rapid antigen tests have recently been considered a viable source for first-line screening,

although concerns about the accuracy of these tests persist. We clinically validated two different

direct antigen rapid tests for the detection of either N or S from SARS-CoV-2 in retrospectively

collected nasal or nasopharyngeal swab specimens. Of the total number of nasal swab specimens

evaluated by qRT-PCR for amplification of SARS-CoV-2 N, S, and ORF1ab genes, 100 tested

positive and 90 tested negative (Table 1, Table S1). The overall sensitivity and specificity of the

rapid antigen test for detection of SARS-CoV-2 N, evaluated across the nasal swab specimens,

was 80.0% and 91.1%, respectively. Of the total number of nasopharyngeal swab specimens

evaluated by qRT-PCR for amplification of SARS-CoV-2 N, RNA-dependent RNA polymerase

(RdRp), and envelope (E) genes, 72 tested positive and 49 tested negative (Table 2, Table S1). The

overall sensitivity and specificity of the rapid antigen test for detection of SARS-CoV-2 S,

evaluated across the nasopharyngeal swab specimens was 84.7% and 85.7%, respectively.

Altogether, our data demonstrate that the sensitivity of the rapid antigen tests are positively

correlated to the viral load level (Table S1).

The Ct value indirectly quantifies the viral RNA copy number related to the viral load of

the sample for the specific assay (*28, 29*). Ct values represent the number of qRT-PCR cycles at

which generated fluorescence crosses a threshold during the linear amplification phase; Ct values

are therefore inversely related to the viral load. Notably, the sensitivity of both rapid antigen tests

increases as Ct value decreases (Table S1). The sensitivity of the rapid antigen test for detection

of SARS-CoV-2 N increased from 80.0% at Ct values <40 to 95.8% at Ct values <20. Similarly,

the sensitivity of the rapid antigen test for detection of SARS-CoV-2 S increased from 84.7% at

Ct values <35 to 100.0% at Ct values <15. Taken together, the clinical data shows that the rapid

antigen test performs with increasing accuracy for individuals with a higher viral load, and

potentially the most infectious (*13–16*).

An Enhanced Epidemiological SIDHRE-Q Model

We propose an enhanced epidemiological modeling system, *SIDHRE-Q,* a variant of the

classical SIR model in order to expand our clinical validation study and to understand the effects

of using frequent rapid tests such as the rapid antigen test on COVID-19 outbreak dynamics. The

changes we make to the basic model to encompass the unique characteristics of the COVID-19

pandemic are similar to those presented by Giordano et al. *(16)* (Figure 1, Figure S1). The

differential equations governing the evolution of the *SIDHRE-Q* model and descriptions of the

parameter values are provided in the material and methods section (Equation 2, Table 3).

An individual that begins in **S** may either transition to a Quarantine Uninfected (**Q-U**) state

via a false positive result or to an Infected Undetected (**I**) state via interaction with an infected

individual. Should an individual in **S** move into **Q-U**, they are quarantined for 14 days before

returning to **S**, a time period chosen based on current knowledge of the infectious period of the

disease. One could also conceive of an effective strategy in which individuals exit quarantine after

producing a certain number of negative rapid tests in the days following their initial positive result

or confirm their negative result using qRT-PCR.

Given that those diagnosed are predominantly quarantined, individuals in **I** interact more

with the **S** population than do those in Infected Detected (**D**). Therefore, the infectious rate for **I** is

assumed to be significantly larger than for **D**. Furthermore, a region's ability to control an outbreak

is directly related to how quickly and effectively people in **I** test into **D**, reducing their

infectiousness through quarantine. This study, in particular, highlights the critical role frequency

of testing, along with strict quarantine, has in mitigating the spread of the disease and provides

specific testing strategies based on rapid tests we predict to be highly effective.

In this model, we assume that individuals receive a positive diagnosis before developing

severe symptoms and that those with symptoms severe enough to be potentially fatal will go to the

hospital. If an individual develops symptoms, we assume they are tested daily until receiving a

positive result; hence, before severe symptoms develop, they will be diagnosed with high

probability. Those who do not develop symptoms are tested according to the frequency of tests

administered to the general population. Therefore, there is no modeled connection between **I** and

H or between **I** and **E**. Removing these assumptions would have negligible impact on the results

as these flows are very small.

Should an individual test positive and transition to **D**, they may either develop serious

symptoms requiring care or recover. Those who develop serious symptoms and transition to state

H will then transition to either **R** or **E**. The recovered population is inevitably tested, as infected

individuals may recover without being detected. Therefore, the Quarantined Recovered (**Q-R**)

state is introduced with the same connections to **R** as the connections between **S** and **Q-U**. Though

the reinfection rate of SARS-CoV-2 has been a point of recent debate, it is assumed that the number

of re-infected individuals is small (*30–34*). Therefore, individuals cannot transition from **R** to **S**,

hence the separately categorized quarantined populations.

We considered several variations and extensions of the *SIDHRE-Q* model. In simulations,

we tested additional states, such as those in the *SIDARTHE* model, which include distinctions

between symptomatic and asymptomatic cases for both detected and undetected populations (*17*).

Incorporating information about the correlations between viral load and infectivity and sensitivity

were also considered. Altogether, our modeling system has been well tuned to predict the impact

of high frequency rapid testing on COVID-19 spread and outcomes.

Frequent Rapid Testing with Actionable Quarantining Dramatically Reduces Disease

Spread

In order to demonstrate how strategies could affect the disease spread in different

geographies and demographics, we used surveillance data obtained from regions of varying

characteristics: the state of Massachusetts (MA), New York City (NYC), Los Angeles (LA), and

São José do Rio Preto (SJRP), Brazil, the site of the rapid antigen test clinical validation study.

These regions are also selected in our study due to the readily available surveillance data provided

by the local governments. We fit the model to the data from each region starting 1 April 2020. At

this time point the disease reportedly is most advanced in NYC and least advanced in SJRP, Brazil

with estimated cumulative infection rates of 7.11% and 0.12%, respectively.

After calibrating the *SIDHRE-Q* model, the disease spread is observed with varying

validated rapid antigen test performances and frequencies (Figure 2A). Sensitivity (the ratio of true

positives to the total number of positives) and specificity (the ratio of true negatives to the total

number of negatives) compared to gold-standard qRT-PCR were used as measures of test

accuracy.

The rapid test frequency is varied while maintaining an accuracy of 80% sensitivity and

90% specificity, comparable to our clinical data collected in SJRP, Brazil. These testing scenarios

are then compared to symptomatic testing, in which individuals receive a rapid test only when

presenting symptoms, via either a rapid test or qRT-PCR. Since the primary testing regiment

deployed in MA, LA, NYC and SJRP, Brazil is qRT-PCR-based and focused on symptomatic

individuals, the symptomatic testing protocol via qRT-PCR is directly estimated from the data to

be the rate v (Table 3).

The difference between the qRT-PCR and rapid test simulations (red and orange lines,

respectively) is therefore only sensitivity of testing (Figure 2A). We assumed that test outcome

probability is a function only of whether an individual is infected and independent of other factors;

one can consider this a lower bound on effectiveness of a strategy, as sensitivity and infectivity

are often positively correlated with antigen testing.

To better understand the effect of rapid testing frequency and performance on healthcare

capacity and mortality rates, we simulate the testing strategy with 30%-90% sensitivity each with

80% or 90% specificity against the symptomatic testing strategy (Figure S2).

As per our hypothesis, frequency and symptom-based testing dramatically reduced

infections, simultaneous hospitalizations, and total deaths when compared to the purely symptom-

based testing regiments, and infections, hospitalization, and death were reduced as frequency

increased. Although testing every day was clearly most effective, even testing every fourteen days

with an imperfect test gave an improvement over symptomatic testing with qRT-PCR. While the

strategy works best when implemented at the very beginning of an outbreak, as demonstrated by

the results in SJRP, Brazil, it also works to curb an outbreak that is already large, as demonstrated

by the results in NYC. The difference between frequencies is more noticeable when the testing

strategy is applied to the outbreak in NYC, leading us to hypothesize that smaller outbreaks require

a lower testing frequency than larger ones; note the difference between the dependence on

frequency to curb a small initial outbreak in SJRP, Brazil versus a large one in NYC (Figure 2B).

For test performance of 80% sensitivity and 90% specificity, the percent of the population

that has been infected in total from the beginning of the outbreak to mid-July drops from 18%

(MA), 11% (LA), 26% (NYC), and 11% (SJRP, Brazil) to 3%, 2%, 12%, and 0.26%, respectively,

using a weekly rapid testing and quarantine strategy (with regards to predictions of overall

infection rates, other studies based on seroprevalence and epidemiological predictions have

reached similar conclusions (*35*, *36*)). If testing is increased to once every three days, these

numbers drop further to 1.6% (MA), 1.4% (LA), 9.4% (NYC), and 0.19% (SJRP, Brazil) (Table

S2).

To further examine the relationship between frequency and sensitivity, we modeled the

maximum number of individuals in a given state over the 105-day time period for four geographic

regions (Figure 2B, Figure S3). In all four geographic regions, as frequency of testing increases,

the total infections, maximum simultaneous hospitalizations, and total deaths converge to small

percentages regardless of the sensitivity at high frequencies. It is clear that the difference in

frequency required to achieve the same result using tests of differing sensitivities is very small.

For example, we predict that for the outbreak in LA, a testing strategy started on 1 April of every

10 days using a test of sensitivity 90% would have resulted in 2.5% of the population having been

infected, while using a test of sensitivity 30% would require a strategy of every 5 days to achieve

the same number. Thus, we conclude that frequency is more important than sensitivity in curbing

the spread, and a large range of sensitivities prove effective when testing sufficiently often (Figure

S3). How frequently, exactly, depends on the specific outbreak and what stage it is in, which leads

us to the location-based deployment strategy discussed in a later section. Frequency of testing can

be significantly reduced to effectively contain the disease once the initial outbreak has been

controlled; it is clear that this takes only a matter of weeks (Figure 2A).

On the other hand, according to the specificity of the rapid test and the quarantine duration,

larger testing frequency result in a larger percent of the population quarantined (Figure 2A).

Assuming a 90% rapid test specificity and 14-day quarantine duration, for the 1-, 3- and 7-day

frequencies almost 60%, 38% and 20% of the population, respectively, would be quarantined. This

figure may be reduced with additional rules for exiting quarantine early, such as after

complementary testing. An example of such a strategy is that individuals who test positive are

required to either quarantine for two weeks or produce two consecutive negative rapid tests in the

two days following their positive result. Assuming 80% sensitivity and 90% specificity, those

individuals will reenter the public while still infected with probability 0.04. If uninfected, that

individual will exit quarantine after two days with probability 0.81. However, a compromise

between the reduction of infections and the proportion of the population in quarantine would be

part of the planning for the appropriate testing protocol in each community or region.

Additionally, while high frequency may be necessary to contain a large outbreak initially,

relatively infrequent testing, such as every one or two weeks, is sufficient to keep controlled

outbreaks small, while reducing the number of quarantined individuals to less than 10% of the

population using a two-week mandatory quarantine.

A County-Based Testing Strategy Offers a Cost-effective Approach to Large-scale COVID-

19 Surveillance

To examine the effects of resource-strategic testing schemes, we modeled the COVID-19

prevalence by varying testing frequency across counties of California. For this analysis, only

California was analyzed because of the accessibility of the county level data. In this scheme, the

percent of active infected detected individuals in a county determines the frequency of testing. We

define thresholds for the number of active detected infections that, when hit, initiate testing

protocols of different frequencies depending on the threshold hit. We first tested evenly spaced

thresholds for the number of detected active infections up to 1% of the population, but later adopted

thresholds that were determined according to Equation 1. In Equation 1, D = population of state **D**

at the time of testing. T = number of active infections which, if reached, initiates everyday testing.

The days between tests are rounded to the closest integer value.

$$\text{Days between tests} = \max\left(1, 2\log_2(T/D) + 1\right)$$

(1)

The days between tests are chosen such that the detected active infections should remain

near to or below T. If the initial detected active infections are greater than T, then the testing

frequency of 1 will cause infections to rapidly drop. Both the threshold at which everyday testing

begins and the coefficient of log_2T/D can be modified to produce a strategy that is more or less

frequent in testing or resource effective; a range of days between tests from 14 days to 1 day are

used (Figure 3). A scan over different choices of T is shown in Figures S5 and S6; the threshold

we choose in Figure 3 is 0.05% because it is successful in curbing the outbreak within the time

period we consider. While these choices work for the epidemic in California at the point we start

our simulations, 10 April, they do not necessarily reflect the most resource effective choices

everywhere. Our analysis could be redone to select the best strategy in other states or in the country

as a whole.

Using a rapid test with a sensitivity of 80% and specificity of 90%, the county-based testing

with threshold 0.05% reduces the active infections from 0.94% to 0.0005%, while the uniform

strategy with tests administered every 7 days results in double the number of active infections

(Figure 3). As the threshold is reduced, the total cost increases while the cumulative infections,

maximum percentage hospitalized, and cumulative deaths all decrease (Figure S4).

Strategy B in Figure 3 consists of qRT-PCR testing uniformly applied to the highlighted

population with a frequency of once weekly. The average cost per person per day is just under $15.

Despite this frequency and the accuracy of qRT-PCR, the strategy does not succeed in curbing the

spread as fast as strategy A, which uses a testing sensitivity and specificity of 80% and 90%,

respectively, and testing frequency that vary between counties depending on the proportion of their

population that is currently infected. The total cost for strategy A is estimated at a fraction of the

other at $1.53 per person per day.

DISCUSSION

In this study we examine the potential effects of a novel testing strategy to limit the spread

of SARS-CoV-2 utilizing rapid antigen test screening approaches. Our clinical data and *SIDHRE-*

Q modeling system demonstrate that 1) frequent rapid testing even at a range of accuracies is

effective at reducing COVID-19 spread, 2) rapid antigen tests are a viable source for this strategy

and diagnose the most infectious individuals, and 3) strategic geographic-based testing can

optimize disease control with the amount of available resources. The information from a diagnostic

test itself is of tremendous value, as it can prompt the necessary quarantine measures to prevent

spread, guide proper care and triage, and provide crucial disease-tracking information. Diagnostic

testing in the United States and abroad, however, has been a significant public health hurdle. The

public has witnessed and experienced symptomatic individuals being denied testing due to

shortages, and few testing structures for asymptomatic or mildly symptomatic individuals – a

significant source of disease spread. Though several factors contributed to the stymied early

response measures, such as lockdown and quarantine protocols and adherence, severe testing

bottlenecks were a significant culprit (*37–39*). Early control measures have been shown to decrease

lives lost by several orders of magnitude (*40*). These challenges, though exacerbated during the

early months of the pandemic, remain at the forefront of the public health crises.

Diagnosis of SARS-CoV-2 infection by qRT-PCR is the current standard of care, yet

remains expensive and requires a laboratory and experienced personnel for sample preparations

and experimentation. Significantly, the turnaround time for results can be up to 10 days (*41*). On

an individual scale, this leaves the public in limbo, preventing people from either leaving

quarantine if they are negative, or delaying critical care and infecting others if they are positive.

On a societal level, this current testing scheme yields incomplete surveillance data on which

response efforts such as societal reopening and hospital management depend. Though qRT-PCR

is considered the gold-standard diagnostic method because of its high sensitivity and specificity,

the logistical hurdles render it unrealistic for large-scale screening.

As qRT-PCR remains impractical for this strategy, and rapid tests are facing regulatory

challenges because they do not perform with qRT-PCR-like accuracy, rapid test screening is either

nonexistent in several countries or symptom-based. Even under best-case assumptions, findings

have shown that symptom and risk-based screening strategies miss more than half of the infected

individuals (*42*). Some have argued that the need for widespread testing is overstated due to the

variability in test sensitivity and specificity (*43*). Here, we present alternative large-scale

diagnostic tools to qRT-PCR, and show that test performance, though valuable, is secondary to

widespread test frequency, which is enabled by accessibility and turnaround time. Furthermore,

test affordability is essential for the successful implementation in communities most affected by

infection and will to speed up the safe opening and functioning of the viral sectors of the economy.

Giordano et al. has modeled the evolution of SARS-CoV-2 spread, introducing a diagnosed

state to elucidate the importance of population-wide testing (*17*). Mina et al. has examined how

various test sensitivities and frequencies affect the reproductive number (*12*). We build upon these

findings to show how in affected United States and Brazil regions, population-wide frequent and

rapid testing schemes, with sensitivities ranging from 30%-90%, can be more effective in curbing

the pandemic than a PCR-based scheme. Integrating real-world surveillance and clinical data into

our modeling system has allowed us to incorporate regional differences - such as variances in

healthcare access, state health policy and adherence, state GDP, and environmental factors - under

the same model. Significantly, our findings hold true across Massachusetts, New York City, Los

Angeles, and São José do Rio Preto, Brazil. We also present the economic considerations of these

testing regimes, showing that widespread rapid testing is more cost efficient than less frequent

qRT-PCR testing. In line with these economic considerations, our model demonstrates the

effectiveness of a geographic-based frequent testing regime, in which high disease prevalence

areas receive more frequent testing than low disease prevalence areas.

Since COVID-19 is known to affect certain demographics differently, modeling would

benefit from incorporating demographic information correlated with disease progression and

spread to define sub-models and sets of parameters accordingly. Age, pre-existing conditions, job

types, and density of population are examples of possible categories, each of which influence the

risk of contracting and/or dying from COVID-19. Further studies would benefit from incorporating

these ideas to better understand the effectiveness of rapid testing on identifying potential super

spreading events. Future public health prevention programs should use the proposed modeling

system to develop and test scenarios for precision testing and prevention.

Our findings also point to low-cost tools for implementation of this testing strategy, such

as a rapid antigen-based test for the detection of SARS-CoV-2 proteins. We show that the rapid

antigen tests perform with a range of accuracies under which disease spread can be dramatically

mitigated under our model. Notably, the sensitivity is correlated to the individual's viral load,

effectively diagnosing those who are potentially the most infectious with the highest accuracy. Our

findings are significant because rapid antigen tests are cheaper than qRT-PCR, can be mass

produced to millions per day, present results within 15 minutes, and can be administered by a

nonexpert without a lab or special equipment.

There are several policy implications for these findings. First, our model supports that

systems of high frequency rapid testing should be implemented as a first-line screening method.

This can be first enabled by a more holistic regulatory evaluation of rapid diagnostics, such that

policy emphasizes accessibility and turnaround time even under a range of accuracies. One can

imagine a less accurate, though rapid method of first-line screening in schools, public

transportation, and airports, or even at home, and a qRT-PCR-based method for second-line

screening (testing those who present severe symptoms or have been in contact with infected

individuals, testing in a clinical setting, etc). Second, our cost analysis and rapid antigen test data

present a viable and potentially more cost-effective method for screening. Third, our county-based

testing scheme presents a possible method for wide-scale screening while optimizing resources.

Future studies should investigate how this selective testing strategy can be applied to different

location scales to further inform health policy. Moreover, though our models analyze regions in

the United States and Brazil, similar testing strategies can be considered globally in both resource

limited and abundant settings due to the higher accessibility of rapid tests compared to qRT-PCR.

We emphasize that integral to the effectiveness of diagnostic schemes is 1) the proper

adherence to quarantine measures and 2) the combined use of a variety of diagnostic methods

including nucleic acid, antigen, and antibody tests. According to these models, rapid antigen tests

are an ideal tool for first-line screening. Clinical molecular tests such as qRT-PCR are vital to the

diagnostic landscape, particularly to re-test suspected cases that were negative on the rapid test.

Because rapid tests present a higher rate of false negatives, methods such as qRT-PCR remain

integral to second-line screening. Antibody tests provide important information for immunity and

vaccination purposes as well as epidemiological surveillance. This model also assumes that

individuals will quarantine themselves before being tested and for 14 days following a positive

diagnostic result.

Our simulations combined with real-world data demonstrate a robust modeling system and

elucidates the significance of this novel testing strategy. However, there are important limitations

to be considered. Differences in disease reporting between the geographical regions and the

incomplete nature of COVID-19 surveillance data, often due to the lack of testing, are not

considered in the model. It is imperative that the testing results, hospitalization and death statistics,

and changes in protocol are reported in real-time to scientists and policy makers so that models

can be accurately tuned as the pandemic develops. The model also does not take into account

infrastructural limitations such as hospital capacity. Though the rapid antigen test offers several

advantages such as affordability, fast turnaround time, and ease of mass production, we are also

assuming that there are systems in place to implement frequent and safe low-cost screening across

different communities and settings.

Our model underscores the need for a point-of-care or at-home test for frequent screening,

particularly as lockdown restrictions ease. Regulatory agencies such as the FDA could work

towards regulating rapid tests to alternative standards other than comparison to high sensitivity

molecular diagnostics, as our model shows that frequency and scale of testing may overcome lower

sensitivities. Rather, we could refocus policy to implement first-line screening that optimizes

accuracy with efficiency and equitability.

MATERIAL AND METHODS

Development of Direct Antigen Rapid Tests for the Detection of SARS-CoV-2

We developed a direct antigen rapid test for the detection of the nucleocapsid protein or

spike glycoprotein from SARS-CoV-2 in nasal or nasopharyngeal swab specimens as previously

described (*44*). Briefly, the rapid antigen tests are immunochromatographic format with a visual

readout using anti-N or anti-S mouse monoclonal antibodies (E25Bio, Inc., Cambridge, MA, USA)

that are either coupled to 40 nm gold nanoparticles (Abcam, Cambridge, UK) or adsorbed to

nitrocellulose membranes (Sartorius, Goettingen, Germany). Each rapid antigen test has a control

area adjacent to the paper absorbent pad; the control is an anti-mouse Fc domain antibody (Leinco

Technologies, Fenton, MO, USA) that will capture any of the antibody-conjugated gold

nanoparticles to generate a control visual signal. A visual signal at the test area reflects SARS-

CoV-2 N or S that is "sandwiched" between an anti-N or anti-S antibody adsorbed to the

nitrocellulose membrane and a second anti-N or anti-S antibody covalently coupled to visible gold

nanoparticles.

Validation of Direct Antigen Rapid Test for the Detection of SARS-CoV-2

In a retrospective study of nasal swab specimens form human patients, we compared the

accuracy of the rapid antigen test for detection of SARS-CoV-2 N to the viral loads of individuals.

Nasal swab specimens (n=190) were tested following approved human subjects use protocols. The

nasal swab specimens were banked frozen from suspected patients submitted to PATH for routine

COVID diagnosis. Prior to using the rapid test, the nasal swab specimens were validated by qRT-

PCR using the FDA EUA ThermoFisher/AppliedBiosystems TaqPATH COVID-19 Combo Kit

(ThermoFisher, Waltham, MA USA). The primary study under which the samples and data were

collected received ethical clearance from the PATH Research Ethics Committee, protocol number

00004244. The nasal swab specimens were de-identified, containing no demographic data, prior

to analysis.

Additionally, in a retrospective study of nasopharyngeal swab specimens from human

patients, we compared the accuracy of the rapid antigen test to the viral load of individuals.

Nasopharyngeal swab specimens (n = 121) were tested in Brazil following approved human

subjects use protocols. The age of study participants ranged from 1 to 95 years with an overall

median of 37 years (interquartile range, 27–51 years), and 62% were female. The demographic

summary of the patients are included in Table S3. The nasopharyngeal swab specimens were

banked refrigerated or frozen samples from suspected patients submitted to the lab for routine

COVID diagnosis. Prior to using the rapid test, the nasopharyngeal swab samples were validated

by qRT-PCR using GeneFinder[TM] COVID-19 Plus Real*Amp* Kit (OSANGHealtcare, Anyang-si,

Gyeonggi-do, Republic of Korea I). The primary study under which the samples and data were

collected received ethical clearance from the Faculdade de Medicina de São José do Rio Preto

(FAMERP), protocol number 31588920.0.0000.5415. All excess samples and corresponding data

were banked and de-identified prior to the analyses.

A nasopharyngeal swab specimen (1 mL) was concentrated using a Vivaspin 500

centrifugal concentrator (Sartorius, Goettingen, Germany) at 12,000 x g for 10 minutes. The

concentrated nasopharyngeal swab specimen retentate was transferred to a collection tube and the

rapid antigen test was inserted into the tube with the retentate and allowed to react for 15 minutes.

After processing of the rapid antigen test, the visual positive or negative signal was documented.

Data for Modeling

As of August 2020, the United States and Brazil have the highest number of confirmed

COVID-19 cases and deaths worldwide, with both countries reporting their first case on 26

February 202) (*1*). Although several affected US regions could have been modeled, we look at

data from Massachusetts, New York, and Los Angeles: these regions each contained "hotspots",

or areas of surging COVID-19 cases, at different points in time during the pandemic and have

publicly available government-provided surveillance data. Our model is fit using data over 105

days beginning on April 1 for Figure 2 and 105 days beginning on April 10 for Figure 3 (see

"Modeling Parameters" in Methods). In order to understand the various testing proposals on a

global scale, we performed our clinical study in and expanded the modeling study to Brazil. The

specific data we use to fit our model are cumulative confirmed cases, total deaths, and number of

daily hospitalizations due to COVID-19. This surveillance data was retrieved from government-

provided online databases (*45–51*).

Modeling Parameters

Equation 2 below provides the exact differential equations governing the model.

$$
\begin{aligned}
d\mathbf{S} &= &-\mathbf{S}\left(\alpha\mathbf{I} + \eta\mathbf{D} + \gamma\right) && +\psi\mathbf{Q_U} \\
d\mathbf{I} &= &-\mathbf{I}\left(\varepsilon + \lambda + \nu\right) && +\mathbf{S}\left(\alpha\mathbf{I} + \eta\mathbf{D}\right) \\
d\mathbf{D} &= &-\mathbf{D}\left(\frac{\mathbf{D}+\mathbf{I}}{\mathbf{D}}\mu + \rho\right) && +\mathbf{I}(\nu + \varepsilon) \\
d\mathbf{H} &= &-\mathbf{H}\left(\sigma + \tau\right) && +\mu\left(\mathbf{D} + \mathbf{I}\right) \\
d\mathbf{E} &= & && +\tau\mathbf{H} \\
d\mathbf{R} &= &-\gamma\mathbf{R} && +\rho\mathbf{D} + \lambda\mathbf{I} + \sigma\mathbf{H} + \psi\mathbf{Q_R} \\
d\mathbf{Q_U} &= &-\psi\mathbf{Q_U} && +\gamma\mathbf{S} \\
d\mathbf{Q_R} &= &-\psi\mathbf{Q_R} && +\gamma\mathbf{R}
\end{aligned}
$$

(2)

In order to determine the values of the parameters defining the flows between states, we use a least

squares regression performed at seven day intervals in the datasets to which we fit. This allows

the model to take into account the time dependent nature of the parameters, which rely on factors

such as social distancing regulations and changes in testing capacity. We also fit window sizes

between 1 and 21 days and find that while the fit degrades with larger window size, the overall

shape of the fits do not change. We choose seven days assuming policy changes take a week to

become effective and that reasonable parameters can be expected to change within this time period.

Also, the seven day window size accounts for the fact that often data is not reported as diligently

over the weekend. Time series of the values of the parameters for the geographic locations

discussed in this paper are included in Figure S5.

Given the restrictions on data available for the populations of various states, varying all of

the parameters results in an over parameterized system. Therefore, a subset of the model

parameters are fit while the others are either extracted from other sources; see Table 3.

The fitting procedure minimizes the sum of the squared residuals of the total cases, current

daily hospitalizations, cumulative deaths, and percentage of total infected individuals currently

hospitalized. The first three are present in the data sets while the latter is derived from the estimates

of the ratio between infected undetected to infected detected individuals from the CDC Laboratory

Seroprevalence Survey Data (*52*). While this ratio changes over time, the percentage of infected

individuals developing severe symptoms should remain roughly constant throughout the course of

the epidemic in the different locations studied.

We consider the data sets for outbreaks in MA, NYC, LA, and SJRP, Brazil (*45–50*). While

each location has testing and fatality information dating back to January, hospitalization data was

not included until late March (for NYC and SJRP) and April (for MA and LA). Hence we begin

our fitting procedure and testing strategy on 1 April for each of the data sets; by this point, the

outbreak is advanced in NYC, substantial in MA, non-negligible, but far from its peak, in LA, and

in early stages in SJRP, Brazil. Starting simulations at various stages of the outbreak allows one

to see the difference in results between when a testing strategy is administered.

In order to determine the effectiveness of the county-based strategy when applied to the

state of California, we also fit all of the counties in California with a population greater than 1.5%

of that of the entire state and with greater than zero deaths. The results do not depend on these

selections, but instead suggest a practical criteria to administer limited resources. The fitting is

done starting 10 April for these counties, as at this point the outbreak is sufficiently well-

documented in each to successfully model. For the county-level data we compute a seven day

running average of each of the data sets to which we then fit in order to smooth out fluctuations in

the data, likely due to reporting, which are more significant here than in the other data sets

considered, as the county populations are smaller and hence discrepancies impact the smoothness

of the data more. The fits for each of the counties can be found in Figure S6.

As one can see from Figure 1, these data sets are particularly not smooth, which indicates

inefficiencies in reporting. Additionally, it is difficult to gauge their consistency within the dates

provided or to compare between locations, as reporting mechanisms changed over time within the

same locations. Despite this lack of consistency, our model and fitting mechanism was successful

in reproducing the progress of the outbreak in each data set studied.

The authors confirm that the data supporting the findings of this study are available within the

article and/or its supplementary materials; any other data will be made available upon request. Our

code can be found on github: https://github.com/badeaa3/COVID19_Rapid_Testing. The code is

written using python with the packages scipy, numpy, lmfit, matplotlib and plotly (53–57).

H2: Supplementary Materials

Results

Table S1. Data summary of direct antigen rapid test (DART) for detection of SARS-CoV-2 nucleocapsid protein and DART for detection of SARS-CoV-2 spike glycoprotein performance in comparison to qRT-PCR results.

Table S2. Summary of results of COVID-19 outcomes in 3 US Regions and Brazil as a result of Frequent Rapid Testing Protocol using SIDHRE-Q Model. Total Infected, Maximum Hospitalized, and Total Deaths are shown for Massachusetts, Los Angeles, New York, and São Jose do Rio Preto, Brazil under a qRT-PCR protocol (symptomatic testing) and a Rapid Testing protocol (once every three days with test performance of 80% sensitivity and 90% specificity).

Table S3. Demographic and clinical summary of patients evaluated by the SARS-CoV-2 Direct Antigen Rapid Test (DART). N response, N or mean of positive, and % or standard deviation for each group is presented. All samples (n=131) collected and tested in São José do Rio Preto, Brazil.

Figure S1. Graphical scheme displaying the relationships between the stages of quarantine and infection in *SIDHRE-Q* model: Q-U, quarantine uninfected; **S**, susceptible (uninfected); **I**, infected undetected (pre-testing and infected); **D**, infected detected (infection diagnosis through testing); **H**, hospitalized (infected with life threatening symptom progression); **R**, recovered

(healed); **E**, extinct (dead); and **Q-R**, quarantine recovered (healed but in quarantine by false

positive testing).

Figure S2. COVID-19 Outcomes as a result of Frequent Rapid Testing Protocol with

variable test performances using _SIDHRE-Q_ Model. The Cumulative Detected Infected,

Hospitalized, Deceased, Active Infections, Recovered, and Quarantined are modeled over 105

days (top to bottom) using reported data from 4 global regions: Massachusetts, Los Angeles,

New York City, and São José do Rio Preto in Brazil (left to right). The COVID-19 population

spread and outcomes are modeled under a Rapid Testing Protocol with variable testing

frequencies ranging from 1-21 days between tests, and variable test performances: 90%

specificity with 90% sensitivity (A), 70% sensitivity (B), 50% sensitivity (C), and 30%

sensitivity (D); and 80% specificity with 90% sensitivity (E), 70% sensitivity (F), 50%

sensitivity (G), and 30% sensitivity (H). This protocol is compared to a symptom-based Rapid

Testing protocol and a symptom-based qRT-PCR protocol.

Figure S3. Effect of Rapid Testing Protocol under variable testing sensitivities and

increasing frequency under the _SIDHRE-Q_ Model. The Cumulative Infections, Maximum

Simultaneously Hospitalized, and Deceased populations are modeled for Massachusetts, Los

Angeles, New York City, and São José do Rio Preto in Brazil. The effect of increasing frequency

of testing is modeled for various testing sensitivities (30%-90%) with an 80% specificity.

Figure S4. Effect of County Based Rapid Testing strategy on COVID-19 outcomes in

California. This protocol varies testing frequency in accordance to the number of recorded

609 cases; the threshold for number of active infections which, if reached, signals to commence

610 everyday testing, the highest frequency considered. A Rapid Test with a 80% sensitivity and

611 90% sensitivity is used in this deployment strategy. Shown is the total cost per person per day

612 versus the cumulative infections, maximum simultaneously hospitalized, and cumulative deaths

613 with varied thresholds for all of CA is shown. The County Based Rapid Testing strategy is

614 compared to uniform testing, which distributes the same number of total tests used in the county

615 strategy, albeit evenly across each county. The effects of uniform testing are modeled for both a

616 Rapid Testing protocol and a qRT-PCR protocol.

618 **Figure S5**. **Time series of the four fitted parameters α, ν, μ, and τ (left to right) for MA, LA,**

619 **NYC, and SJRP (top to bottom).** See Table 2 in the Methods section for an explanation of the

620 parameters. The values are extracted every seven days from data provided by the respective

621 regions. The parameters vary significantly over time and location. Flat points occur during the

622 seven day windows where the parameters are held constant. The fitting procedure is also outlined

623 in the Methods section.

625 **Figure S6. Time series of the three fitted pieces of data Cumulative Cases, Daily**

626 **Hospitalized, and Cumulative Deaths (left to right) for each county receiving testing in CA;**

627 Ventura (2A), Stanislaus (2B), Santa Clara (2C), San Joaquin (2D), San Francisco (2E), San

628 Diego (2F), San Bernardino (2G), Sacramento (2H), Orange (2I), Los Angeles (2J), Kern (2K),

629 Fresno (2L), Alameda (2M). The counties included satisfy two requirements: population greater

630 than 1.5% of the total CA population and nonzero total number of deaths at each point in time.

631 The fitting procedure is outlined in the Methods section.

References and Notes

633 1. Coronavirus Disease (COVID-19) Situation Reports, (available at

634   https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports).

635 2. T. F. Menkir, T. Chin, J. A. Hay, E. Surface, P. Martinez de Salazar, C. Buckee, M. J. Mina, K. Khan, A.

636   Watts, M. Lipsitch, R. Niehus, Estimating the number of undetected COVID-19 cases exported internationally

637   from all of China. *medRxiv* (2020), doi:10.1101/2020.03.23.20038331.

638 3. B. Ivorra, M. R. Ferrández, M. Vela-Pérez, A. M. Ramos, Mathematical modeling of the spread of the

639   coronavirus disease 2019 (COVID-19) taking into account the undetected infections. The case of China.

640   *Commun Nonlinear Sci Numer Simul*, 105303 (2020).

641 4. M. Salathé, C. L. Althaus, R. Neher, S. Stringhini, E. Hodcroft, J. Fellay, M. Zwahlen, G. Senti, M. Battegay,

642   A. Wilder-Smith, I. Eckerle, M. Egger, N. Low, COVID-19 epidemic in Switzerland: on the importance of

643   testing, contact tracing and isolation. *Swiss Med Wkly*. **150**, w20225 (2020).

644 5. H. Lau, T. Khosrawipour, P. Kocbach, H. Ichii, J. Bania, V. Khosrawipour, Evaluating the massive

645   underreporting and undertesting of COVID-19 cases in multiple global epicenters. *Pulmonology* (2020),

646   doi:10.1016/j.pulmoe.2020.05.015.

647 6. J. D. Silverman, N. Hupert, A. D. Washburne, Using influenza surveillance networks to estimate state-specific

648   prevalence of SARS-CoV-2 in the United States. *Science Translational Medicine* (2020),

649   doi:10.1126/scitranslmed.abc1126.

650 7. D. Böhning, I. Rocchetti, A. Maruotti, H. Holling, Estimating the undetected infections in the Covid-19

651   outbreak by harnessing capture-recapture methods. *Int. J. Infect. Dis.* **97**, 197–201 (2020).

652 8. Y. H. Baek, J. Um, K. J. C. Antigua, J.-H. Park, Y. Kim, S. Oh, Y. Kim, W.-S. Choi, S. G. Kim, J. H. Jeong, B.

653   S. Chin, H. D. G. Nicolas, J.-Y. Ahn, K. S. Shin, Y. K. Choi, J.-S. Park, M.-S. Song, Development of a reverse

654   transcription-loop-mediated isothermal amplification as a rapid early-detection method for novel SARS-CoV-2.

655   *Emerg Microbes Infect*. **9**, 998–1007 (2020).

9.   L. Leo, Mylab gets commercial approval from ICMR for Covid-19 antigen rapid testing kit. *Livemint* (2020),

(available at https://www.livemint.com/news/india/mylab-gets-commercial-approval-from-icmr-for-covid-19-

antigen-rapid-testing-kit-11595434040321.html).

10.   S. Dey, Coronavirus testing: Rapid antigen tests now make up nearly half of daily checks | India News - Times

of India. *The Times of India*, (available at https://timesofindia.indiatimes.com/india/rapid-antigen-tests-now-

make-up-nearly-half-of-daily-checks/articleshow/77340459.cms).

11.   C. B. F. Vogels, A. F. Brito, A. L. Wyllie, J. R. Fauver, I. M. Ott, C. C. Kalinich, M. E. Petrone, A. Casanovas-

Massana, M. Catherine Muenker, A. J. Moore, J. Klein, P. Lu, A. Lu-Culligan, X. Jiang, D. J. Kim, E. Kudo, T.

Mao, M. Moriyama, J. E. Oh, A. Park, J. Silva, E. Song, T. Takahashi, M. Taura, M. Tokuyama, A.

Venkataraman, O.-E. Weizman, P. Wong, Y. Yang, N. R. Cheemarla, E. B. White, S. Lapidus, R. Earnest, B.

Geng, P. Vijayakumar, C. Odio, J. Fournier, S. Bermejo, S. Farhadian, C. S. Dela Cruz, A. Iwasaki, A. I. Ko,

M. L. Landry, E. F. Foxman, N. D. Grubaugh, Analytical sensitivity and efficiency comparisons of SARS-

CoV-2 RT-qPCR primer-probe sets. *Nat Microbiol* (2020), doi:10.1038/s41564-020-0761-6.

12.   D. B. Larremore, B. Wilder, E. Lester, S. Shehata, J. M. Burke, J. A. Hay, M. Tambe, M. J. Mina, R. Parker,

Test sensitivity is secondary to frequency and turnaround time for COVID-19 surveillance. *medRxiv* (2020),

doi:10.1101/2020.06.22.20136309.

13.   Z. Shen, F. Ning, W. Zhou, X. He, C. Lin, D. P. Chin, Z. Zhu, A. Schuchat, Superspreading SARS events,

Beijing, 2003. *Emerging Infect. Dis.* **10**, 256–260 (2004).

14.   J. S. M. Peiris, C. M. Chu, V. C. C. Cheng, K. S. Chan, I. F. N. Hung, L. L. M. Poon, K. I. Law, B. S. F. Tang,

T. Y. W. Hon, C. S. Chan, K. H. Chan, J. S. C. Ng, B. J. Zheng, W. L. Ng, R. W. M. Lai, Y. Guan, K. Y. Yuen,

HKU/UCH SARS Study Group, Clinical progression and viral load in a community outbreak of coronavirus-

associated SARS pneumonia: a prospective study. *Lancet*. **361**, 1767–1772 (2003).

15.   X. He, E. H. Y. Lau, P. Wu, X. Deng, J. Wang, X. Hao, Y. C. Lau, J. Y. Wong, Y. Guan, X. Tan, X. Mo, Y.

Chen, B. Liao, W. Chen, F. Hu, Q. Zhang, M. Zhong, Y. Wu, L. Zhao, F. Zhang, B. J. Cowling, F. Li, G. M.

Leung, Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat. Med.* **26**, 672–675 (2020).

16. J. Bullard, K. Dust, D. Funk, J. E. Strong, D. Alexander, L. Garnett, C. Boodman, A. Bello, A. Hedley, Z.

Schiffman, K. Doan, N. Bastien, Y. Li, P. G. Van Caeseele, G. Poliquin, Predicting infectious SARS-CoV-2

from diagnostic samples. *Clin. Infect. Dis.* (2020), doi:10.1093/cid/ciaa638.

17. G. Giordano, F. Blanchini, R. Bruno, P. Colaneri, A. Di Filippo, A. Di Matteo, M. Colaneri, Modelling the

COVID-19 epidemic and implementation of population-wide interventions in Italy. *Nat. Med.* **26**, 855–860

(2020).

18. S. Choi, M. Ki, Estimating the reproductive number and the outbreak size of COVID-19 in Korea. *Epidemiol*

Health. **42**, e2020011 (2020).

19. Y. Y. Wei, Z. Z. Lu, Z. C. Du, Z. J. Zhang, Y. Zhao, S. P. Shen, B. Wang, Y. T. Hao, F. Chen, [Fitting and

forecasting the trend of COVID-19 by SEIR(+CAQ) dynamic model]. *Zhonghua Liu Xing Bing Xue Za Zhi*. **41**,

470–475 (2020).

20. Z. Yang, Z. Zeng, K. Wang, S.-S. Wong, W. Liang, M. Zanin, P. Liu, X. Cao, Z. Gao, Z. Mai, J. Liang, X. Liu,

S. Li, Y. Li, F. Ye, W. Guan, Y. Yang, F. Li, S. Luo, Y. Xie, B. Liu, Z. Wang, S. Zhang, Y. Wang, N. Zhong, J.

He, Modified SEIR and AI prediction of the epidemics trend of COVID-19 in China under public health

interventions. *J Thorac Dis*. **12**, 165–174 (2020).

21. S. Cao, P. Feng, P. Shi, [Study on the epidemic development of COVID-19 in Hubei province by a modified

SEIR model]. *Zhejiang Da Xue Xue Bao Yi Xue Ban*. **49**, 178–184 (2020).

22. R. Huang, M. Liu, Y. Ding, Spatial-temporal distribution of COVID-19 in China and its prediction: A data-

driven modeling analysis. *J Infect Dev Ctries*. **14**, 246–253 (2020).

23. A. Godio, F. Pace, A. Vergnano, SEIR Modeling of the Italian Epidemic of SARS-CoV-2 Using Computational

Swarm Intelligence. *Int J Environ Res Public Health*. **17** (2020), doi:10.3390/ijerph17103535.

24. M. Gatto, E. Bertuzzo, L. Mari, S. Miccoli, L. Carraro, R. Casagrandi, A. Rinaldo, Spread and dynamics of the

COVID-19 epidemic in Italy: Effects of emergency containment measures. *Proc. Natl. Acad. Sci. U.S.A.* **117**,

10484–10491 (2020).

25. C. Hou, J. Chen, Y. Zhou, L. Hua, J. Yuan, S. He, Y. Guo, S. Zhang, Q. Jia, C. Zhao, J. Zhang, G. Xu, E. Jia,

The effectiveness of quarantine of Wuhan city against the Corona Virus Disease 2019 (COVID-19): A well-

mixed SEIR model analysis. *J. Med. Virol.* **92**, 841–848 (2020).

26. T. Zhou, Q. Liu, Z. Yang, J. Liao, K. Yang, W. Bai, X. Lu, W. Zhang, Preliminary prediction of the basic

reproduction number of the Wuhan novel coronavirus 2019-nCoV. *J Evid Based Med*. **13**, 3–7 (2020).

27. C. Reno, J. Lenzi, A. Navarra, E. Barelli, D. Gori, A. Lanza, R. Valentini, B. Tang, M. P. Fantini, Forecasting

COVID-19-Associated Hospitalizations under Different Levels of Social Distancing in Lombardy and Emilia-

Romagna, Northern Italy: Results from an Extended SEIR Compartmental Model. *J Clin Med*. **9** (2020),

doi:10.3390/jcm9051492.

28. F. Yu, L. Yan, N. Wang, S. Yang, L. Wang, Y. Tang, G. Gao, S. Wang, C. Ma, R. Xie, F. Wang, C. Tan, L.

Zhu, Y. Guo, F. Zhang, Quantitative Detection and Viral Load Analysis of SARS-CoV-2 in Infected Patients.

Clin. Infect. Dis. **71**, 793–798 (2020).

29. S. N. Rao, D. Manissero, V. R. Steele, J. Pareja, A Narrative Systematic Review of the Clinical Utility of Cycle

Threshold Values in the Context of COVID-19. *Infect Dis Ther*, 1–14 (2020).

30. J. Alizargar, Risk of reactivation or reinfection of novel coronavirus (COVID-19). *J. Formos. Med. Assoc.* **119**,

1123 (2020).

31. D. Batisse, N. Benech, E. Botelho-Nevers, K. Bouiller, R. Collarino, A. Conrad, L. Gallay, F. Goehringer, M.

Gousseff, D. C. Joseph, A. Lemaignen, F.-X. Lescure, B. Levy, M. Mahevas, P. Penot, B. Pozzetto, D. Salmon,

D. Slama, N. Vignier, B. Wyplosz, Clinical recurrences of COVID-19 symptoms after recovery: viral relapse,

reinfection or inflammatory rebound? *J. Infect.* (2020), doi:10.1016/j.jinf.2020.06.073.

32. W. Deng, L. Bao, J. Liu, C. Xiao, J. Liu, J. Xue, Q. Lv, F. Qi, H. Gao, P. Yu, Y. Xu, Y. Qu, F. Li, Z. Xiang, H.

Yu, S. Gong, M. Liu, G. Wang, S. Wang, Z. Song, Y. Liu, W. Zhao, Y. Han, L. Zhao, X. Liu, Q. Wei, C. Qin,

Primary exposure to SARS-CoV-2 protects against reinfection in rhesus macaques. *Science* (2020),

doi:10.1126/science.abc5343.

33. M. Ota, Will we see protection or reinfection in COVID-19? *Nat. Rev. Immunol.* **20**, 351 (2020).

34. A. Victor Okhuese, Estimation of the Probability of Reinfection With COVID-19 by the Susceptible-Exposed-

Infectious-Removed-Undetectable-Susceptible Model. *JMIR Public Health Surveill.* **6**, e19097 (2020).

35. Y. Gu, COVID-19 Projections Using Machine Learning. *COVID-19 Projections Using Machine Learning*,

(available at https://covid19-projections.com/).

36. D. Stadlbauer, J. Tan, K. Jiang, M. Hernandez, S. Fabre, F. Amanat, C. Teo, G. A. Arunkumar, M. McMahon,

J. Jhang, M. Nowak, V. Simon, E. Sordillo, H. van Bakel, F. Krammer, *medRxiv*, in press,

doi:10.1101/2020.06.28.20142190.

37. A. Goodnough, M. D. Shear, The U.S.'s Slow Start to Coronavirus Testing: A Timeline. *The New York Times*

(2020), (available at https://www.nytimes.com/2020/03/28/us/coronavirus-testing-timeline.html).

38. M. D. Shear, A. Goodnough, S. Kaplan, S. Fink, K. Thomas, N. Weiland, The Lost Month: How a Failure to

Test Blinded the U.S. to Covid-19. *The New York Times* (2020), (available at

https://www.nytimes.com/2020/03/28/us/testing-coronavirus-pandemic.html).

39. S. Kaplan, K. Thomas, Despite Promises, Testing Delays Leave Americans 'Flying Blind.' *The New York*

Times (2020), (available at https://www.nytimes.com/2020/04/06/health/coronavirus-testing-us.html).

40. W. M. de Souza, L. F. Buss, D. da S. Candido, J.-P. Carrera, S. Li, A. E. Zarebski, R. H. M. Pereira, C. A.

Prete, A. A. de Souza-Santos, K. V. Parag, M. C. T. D. Belotti, M. F. Vincenti-Gonzalez, J. Messina, F. C. da

Silva Sales, P. dos S. Andrade, V. H. Nascimento, F. Ghilardi, L. Abade, B. Gutierrez, M. U. G. Kraemer, C. K.

V. Braga, R. S. Aguiar, N. Alexander, P. Mayaud, O. J. Brady, I. Marcilio, N. Gouveia, G. Li, A. Tami, S. B.

de Oliveira, V. B. G. Porto, F. Ganem, W. A. F. de Almeida, F. F. S. T. Fantinato, E. M. Macário, W. K. de

Oliveira, M. L. Nogueira, O. G. Pybus, C.-H. Wu, J. Croda, E. C. Sabino, N. R. Faria, Epidemiological and

clinical characteristics of the COVID-19 epidemic in Brazil. *Nature Human Behaviour*. **4**, 856–865 (2020).

41. S. Mervosh, M. Fernandez, 'It's Like Having No Testing': Coronavirus Test Results Are Still Delayed. *The*

New York Times (2020), (available at https://www.nytimes.com/2020/08/04/us/virus-testing-delays.html).

42. K. Gostic, A. C. Gomez, R. O. Mummah, A. J. Kucharski, J. O. Lloyd-Smith, Estimated effectiveness of

symptom and risk screening to prevent the spread of COVID-19. *eLife*. **9**, e55570 (2020).

43. T. Zitek, The Appropriate Use of Testing for COVID-19. *West J Emerg Med*. **21**, 470–472 (2020).

44. I. Bosch, H. de Puig, M. Hiley, M. Carré-Camps, F. Perdomo-Celis, C. F. Narváez, D. M. Salgado, D.

Senthoor, M. O'Grady, E. Phillips, A. Durbin, D. Fandos, H. Miyazaki, C.-W. Yen, M. Gélvez-Ramírez, R. V.

Warke, L. S. Ribeiro, M. M. Teixeira, R. P. Almeida, J. E. Muñóz-Medina, J. E. Ludert, M. L. Nogueira, T. E.

Colombo, A. C. B. Terzian, P. T. Bozza, A. S. Calheiros, Y. R. Vieira, G. Barbosa-Lima, A. Vizzoni, J.

Cerbino-Neto, F. A. Bozza, T. M. L. Souza, M. R. O. Trugilho, A. M. B. de Filippis, P. C. de Sequeira, E. T. A.

Marques, T. Magalhaes, F. J. Díaz, B. N. Restrepo, K. Marín, S. Mattar, D. Olson, E. J. Asturias, M. Lucera, M.

Singla, G. R. Medigeshi, N. de Bosch, J. Tam, J. Gómez-Márquez, C. Clavet, L. Villar, K. Hamad-Schifferli, L.

Gehrke, Rapid antigen tests for dengue virus serotypes and Zika virus in patient serum. *Sci Transl Med*. **9**

(2017), doi:10.1126/scitranslmed.aan1589.

45. Massachusetts Department of Public Health, COVID-19 Response Reporting. *Mass.gov*, (available at

https://www.mass.gov/info-details/covid-19-response-reporting).

46. California Department of Public Health, COVID-19 Cases - California Open Data, (available at

https://data.ca.gov/dataset/covid-19-cases).

47. California Department of Public Health, COVID-19 Hospital Data - California Open Data, (available at

https://data.ca.gov/dataset/covid-19-hospital-data).

48. Department of Health and Human Hygiene, COVID-19 Daily Counts of Cases, Hospitalizations, and Deaths |

NYC Open Data, (available at https://data.cityofnewyork.us/Health/COVID-19-Daily-Counts-of-Cases-

Hospitalizations-an/rc75-m7u3).

49. Sao Jose do Rio Preto Public Health Office, "COVID-19 Surveillance Data, Sao Jose do Rio Preto" (Sao Jose

do Rio Preto Public Health Office, Sao Jose do Rio Preto, Brazil).

50. New York State Government, Daily Hospitalization Summary by Region. *New York Forward*, (available at

https://forward.ny.gov/daily-hospitalization-summary-region).

51. Massachusetts General Hospital Institute for Technology Assessment, COVID-19 Simulator - Methodology,

(available at https://www.covid19sim.org/images/docs/COVID-

19_simulator_methodology_download_20200507.pdf).

52. CDC, Coronavirus Disease 2019 (COVID-19). *Centers for Disease Control and Prevention* (2020), (available

at https://www.cdc.gov/coronavirus/2019-ncov/cases-updates/commercial-lab-surveys.html).

53. SciPy.org — SciPy.org, (available at https://www.scipy.org/).

54. NumPy, (available at https://numpy.org/).

55. Non-Linear Least-Squares Minimization and Curve-Fitting for Python — Non-Linear Least-Squares

Minimization and Curve-Fitting for Python, (available at https://lmfit.github.io/lmfit-py/).

56. Matplotlib: Python plotting — Matplotlib 3.3.1 documentation, (available at https://matplotlib.org/).

57. Plotly: The front-end for ML and data science models, (available at https://plotly.com/).

Acknowledgments

General: We thank Professor Lee Gehrke for critical reading of the manuscript.

Funding: The study is funded, in part, by a Bill and Melinda Gates Foundation Award

(INV-017872) to E25Bio, Inc. EN is funded by Tufts University DISC Seed Grant. MLN

is supported by a FAPESP grant (#2020/04836-0) and is a CNPq Research Fellow. AFV

is supported by a FAPESP Fellow grant (#18/17647-0). GRFC is supported by a FAPESP

Fellow grant (#20/07419-0). BHGAM is supported by a FAPESP Scholarship

(#19/06572-2). The funders had no role in the design of the study; in the collection,

analyses, or interpretation of data; in the writing of the manuscript, or in the decision to

publish the results.

Author contributions: Conceptualization: BBH. Formal analysis: BN, AB, AR, MB,

NS, ARG, AV, GCDS, TMILDS, BHGAM, MMM, GRFC, FQ, AFNR, MLG, ENN, IB,

BBH. Funding acquisition: IB, BBH. Investigation: BN, AB, AR, MB, NS, ARG, AV,

GCDS, TMILDS, BHGAM, MMM, GRFC, FQ, AFNR, MLG, ENN, IB, BBH.

Methodology: BN, AB, AR, MB, NS, ARG, AV, GCDS, TMILDS, BHGAM, MMM,

GRFC, FQ, AFNR, MLG, ENN, IB, BBH. Project administration: MLN, IB, BBH.

Resources: MLN, IB, BBH. Supervision: MB, MLN, ENN, IB, BBH. Validation: BN,

AB, AR, MB, ENN, BBH. Visualization: BN, AB, AR, MB, AV, ENN, BBH. Writing—

original draft: AR, BBH. Writing—review and editing: BN, AB, AR, MB, NS, ARG,

AV, GCDS, TMILDS, BHGAM, MMM, GRFC, FQ, AFNR, MLG, ENN, IB, BBH.

Competing interests: BN, AB, AR, MB, NS, AG, IB, and BBH are employed by or

affiliated with E25Bio Inc. (www.e25bio.com), a company that develops diagnostics for

epidemic viruses.

821   **Figures and Tables**

823   **TABLES**

825   **Table 1. Clinical validation summary for the direct antigen rapid test (DART) for SARS-**

826   **CoV-2 nucleocapsid protein evaluated using 190 retrospectively collected patient nasal**

827   **swab specimens.**

All Data Summary								
	qRT-PCR (gene average)						95% Confidence Interval	
	+	**-**	**Total**	**Sensitivity**	80.0%	76.1%	83.9%
DART (nucleocapsid protein) **+**	80	8	88	**Specificity**	91.1%	88.2%	94.0%
-	20	82	102	**Positive Predictive Value**	90.9%	87.9%	93.9%
Total	100	90	190	**Negative Predictive Value**	80.4%	76.6%	84.2%
				Prevalence	52.6%	47.8%	57.5%
				Overall Agreement	85.3%	82.8%	87.8%

835 **Table 2. Clinical validation summary for the SARS-CoV-2 direct antigen rapid test**

836 **(DART) for SARS-SoC-2 spike glycoprotein evaluated using 121 retrospectively collected**

837 **patient nasopharyngeal swab specimens.**

All Data Summary						95% Confidence Interval	
	qRT-PCR (gene average)						
	+	-	Total	Sensitivity	84.7%	80.6%	88.9%
DART (spike glycoprotein) +	61	7	68	Specificity	85.7%	80.8%	90.6%
DART (spike glycoprotein) -	11	42	53	Positive Predictive Value	89.7%	86.2%	93.2%
Total	72	49	121	Negative Predictive Value	79.2%	73.6%	84.9%
				Prevalence	59.5%	53.9%	65.1%
				Overall Agreement	85.1%	82.0%	88.3%

Table 3. Details of parameter values used for *SIDHRE-Q* Model.

Parameter	Details & Statistics			
α	α is the probability that an interaction between an undetected infected person and an uninfected person results in a new infection, divided by the average number of uninfected people an undetected infected person comes into contact with on a given day. α is estimated from the data.		Mean	St. Dev.
		MA	0.088	0.051
		LA	0.090	0.034
		NYC	0.067	0.042
		SJRP	0.121	0.042
η	η is the probability that an interaction between an infected person and an uninfected person results in a new infection, divided by the average number of uninfected people a detected infected person comes into contact with on a given day. $\eta = 0.01 \cdot \alpha$ The constant relating η, α accounts for a small but nonzero transmission due to the quarantined (detected) infected population. This value was chosen to be small, assuming a quarantined individual will only infect others with low probability.			
ν	ν is the probability that a symptomatic undetected individual is diagnosed on a given day. ν is estimated from the data. ν is multiplied by sensitivity (assume benchmark sensitivity 100% for PCR, as used when fitting).		Mean	St. Dev.
		MA	0.006	0.005
		LA	0.011	0.006
		NYC	0.0056	0.002
		SJRP	0.015	0.007
ϵ	ϵ is the probability that an asymptomatic undetected infected individual is diagnosed on a given day. $\epsilon = 0$ while fitting (during PCR symptomatic testing). ϵ =(sensitivity/days between tests) when the rapid testing strategy is activated.			
λ	λ is the probability that an undetected infected individual transitions to the recovered state on a given day. $\lambda =$			

			Mean	St. Dev.
	1/14, or the inverse of average recovery time (*51*).			
μ	μ is the probability that an infected individual develops severe symptoms on a given day and transitions into the hospitalized state. The flow from D to H is assumed to be independent of the ratio I/D, but comes only from the detected infected population, hence why it is multiplied by $(I + D)/D$. μ is estimated from the data.		Mean	St. Dev.
		MA	0.0013	9.5e-4
		LA	0.0016	2.4e-4
		NYC	0.0011	6.6e-4
		SJRP	0.0018	8.0e-4
ρ	ρ is the probability that a detected infected individual transitions to the recovered state on a given day. $\rho = 1/14$, or the inverse of the average recovery time (*51*).			
σ	σ is the probability that a hospitalized individual transitions to the recovered state on a given day. $\sigma = 1/11$, or the inverse of the average recovery time for a hospitalized individual (*51*).			
τ	τ is the probability that a hospitalized individual expires on a given day. τ is estimated from the data.		Mean	St. Dev.
		MA	0.034	0.012
		LA	0.016	0.004
		NYC	0.036	0.034
		SJRP	0.032	0.045
γ	γ is the probability of entering either of the quarantine states on a given day from either the Susceptible or Recovered populations. $\gamma = 0$ while fitting (during PCR symptomatic testing). $\gamma = (1 -\text{specificity}) \times (1/\text{days between tests})$ when the rapid testing strategy is activated.			
ψ	ψ is the probability that an individual exits quarantine on a given day. $\psi = 1/14$, or the inverse of the quarantine period for fixed length quarantine.			
Parameter	Details & Statistics			

α	α is the probability that an interaction between an undetected infected person and an uninfected person results in a new infection, divided by the average number of uninfected people an undetected infected person comes into contact with on a given day. α is estimated from the data.		Mean	St. Dev.
		MA	0.088	0.051
		LA	0.090	0.034
		NYC	0.067	0.042
		SJRP	0.121	0.042
η	η is the probability that an interaction between an infected person and an uninfected person results in a new infection, divided by the average number of uninfected people a detected infected person comes into contact with on a given day. $\eta = 0.01 \cdot \alpha$ The constant relating η, α accounts for a small but nonzero transmission due to the quarantined (detected) infected population. This value was chosen to be small, assuming a quarantined individual will only infect others with low probability.			
ν	ν is the probability that a symptomatic undetected individual is diagnosed on a given day. ν is estimated from the data. ν is multiplied by sensitivity (assume benchmark sensitivity 100% for PCR, as used when fitting).		Mean	St. Dev.
		MA	0.006	0.005
		LA	0.011	0.006
		NYC	0.0056	0.002
		SJRP	0.015	0.007
ϵ	ϵ is the probability that an asymptomatic undetected infected individual is diagnosed on a given day. $\epsilon = 0$ while fitting (during PCR symptomatic testing). $\epsilon = $ (sensitivity/days between tests) when the rapid testing strategy is activated.			
λ	λ is the probability that an undetected infected individual transitions to the recovered state on a given day. $\lambda = 1/14$, or the inverse of average recovery time (*51*).			
μ	μ is the probability that an infected individual develops severe symptoms on a		Mean	St. Dev.
		MA	0.0013	9.5e-4

	given day and transitions into the hospitalized state. The flow from D to H is assumed to be independent of the ratio I/D, but comes only from the detected infected population, hence why it is multiplied by $(I + D)/D$. μ is estimated from the data.	LA	0.0016	2.4e-4
		NYC	0.0011	6.6e-4
		SJRP	0.0018	8.0e-4
ρ	ρ is the probability that a detected infected individual transitions to the recovered state on a given day. $\rho = 1/14$, or the inverse of the average recovery time (*51*).			
σ	σ is the probability that a hospitalized individual transitions to the recovered state on a given day. $\sigma = 1/11$, or the inverse of the average recovery time for a hospitalized individual (*51*).			

			Mean	St. Dev.
τ	τ is the probability that a hospitalized individual expires on a given day. τ is estimated from the data.	MA	0.034	0.012
		LA	0.016	0.004
		NYC	0.036	0.034
		SJRP	0.032	0.045

γ	γ is the probability of entering either of the quarantine states on a given day from either the Susceptible or Recovered populations. $\gamma = 0$ while fitting (during PCR symptomatic testing). $\gamma = (1 - \text{specificity}) \times (1/\text{days between tests})$ when the rapid testing strategy is activated.
ψ	ψ is the probability that an individual exits quarantine on a given day. $\psi = 1/14$, or the inverse of the quarantine period for fixed length quarantine.

FIGURES

Figure 1. Graphical scheme displaying the relationships between the stages of quarantine

and infection in *SIDHRE-Q* model. **Q-U**, quarantine uninfected; **S**, susceptible (uninfected); **I**,

infected undetected (pre-testing and infected); **D**, infected detected (infection diagnosis through

testing); **H**, hospitalized (infected with life threatening symptom progression); **R**, recovered

(healed); **E**, extinct (dead); and **Q-R**, quarantine recovered (healed but in quarantine by false

positive testing).

Figure 2. COVID-19 Outcomes in 3 US Regions and Brazil as a result of Frequent Rapid

Testing Protocol using the *SIDHRE-Q* Model. (A) The Cumulative Detected Infected,

Hospitalized, Deceased, Active Infections, Recovered, and Quarantined are modeled over 105

days (top to bottom) using reported data from 4 global regions: Massachusetts, Los Angeles,

New York City, and São José do Rio Preto in Brazil (left to right). The COVID-19 population

spread and outcomes are modeled under a Rapid Testing Protocol (sensitivity 80%, specificity

90%) with variable testing frequencies ranging from 1-21 days between tests. This protocol is

compared to a symptom-based Rapid Testing protocol and a symptom-based PCR protocol. (B)

Effect of Rapid Testing Protocol under variable testing sensitivities (30%-90%) and increasing

frequency under the *SIDHRE-Q* Model. The Cumulative Infections, Maximum Simultaneously

Hospitalized, and Deceased populations are modeled for Massachusetts, Los Angeles, New York

City, and São José do Rio Preto in Brazil with a 90% test specificity.

897 **(A)**

899   **(B)**

909 **Figure 3. Effect of County Based Rapid Test Protocol (A) and Uniform PCR Protocol (B)**

910 **on active infected detected population over time in California (CA).** The legend denotes the

911 thresholds at which testing frequency is determined, the testing frequencies, the percent of CA

912 population under the strategy, and the cost per person per day.