The impact of high frequency rapid viral antigen screening on COVID-19 spread

and outcomes: a validation and modeling study

Authors: Beatrice Nash[1,2*], Anthony Badea[1,3*], Ankita Reddy[1,4*], Miguel Bosch[1,5], Nol

Salcedo[1], Adam R. Gomez[1], Alice Versiani[6], Gislaine Celestino Dutra Silva[6], Thayza

Maria Izabel Lopes dos Santos[6], Bruno H. G. A. Milhim[6], Marilia M. Moraes[6], Guilherme

Rodrigues Fernandes Campos[6], Flávia Quieroz[6], Andreia Francesli Negri Reis[6], Mauricio

L. Nogueira[6], Elena N. Naumova[7], Irene Bosch[1,8], Bobby Brooke Herrera[1,9‡]

*These authors contributed equally to this work.

Affiliations:

[1]E25Bio, Inc., Cambridge, MA, USA

[2]Department of Computer Science, Harvard University School of Engineering and Applied

Sciences, Cambridge, MA, USA

[3]Department of Physics, Harvard University, Cambridge, MA, USA

[4]Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

[5]InfoGeosciences LLC, Houston, TX, USA

[6]Faculdade de Medicina de São José do Rio Preto (FAMERP), São José do Rio Preto,

Brazil

[7]Division of the Nutrition Epidemiology and Data Science, Friedman School of Nutrition

Science and Policy, Tufts University, Boston, MA, USA

[8]Department of Medicine, Mount Sinai School of Medicine, New York, NY, USA

[9]Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of Public

Health, Boston, MA, USA

[‡]Corresponding Author. BBH, email: bbherrera@e25bio.com

Abstract

High frequency screening of populations has been proposed as a strategy in facilitating control of the COVID-19 pandemic. Here we use computational modeling, coupled with clinical data from a rapid antigen test, to predict the impact of frequent rapid testing on COVID-19 spread and outcomes. Using patient nasopharyngeal swab specimens, we demonstrate that the sensitivity and specificity of the rapid antigen test compared to quantitative real-time polymerase chain reaction (qRT-PCR) are 84.7% and 85.7%, respectively; moreover, sensitivity correlates directly with viral load. Based on COVID-19 data from three regions in the United States and São José do Rio Preto, Brazil, we show that high frequency, strategic population-wide rapid testing, even at varied accuracy levels, diminishes COVID-19 infections, hospitalizations, and deaths at a fraction of the cost of nucleic acid detection via qRT-PCR. We propose large-scale antigen-based surveillance as a viable strategy to control SARS-CoV-2 spread and to enable societal re-opening.

INTRODUCTION

The COVID-19 pandemic has taken an unprecedented toll on lives, wellbeing, healthcare systems, and global economies. As of 1 September 2020, there have been more than 25.5 million confirmed cases globally with more than 850,000 confirmed deaths[1]. However, these numbers and the current mapping of disease spread present an incomplete picture of the outbreak largely due to the lack of adequate testing, particularly as undetected infected cases are the main source of disease spread[2–7]. It is estimated that the reported detection rate of actual COVID-19 cases is only 1-2%[5]. As of September 2020, the United States and Brazil remain the top two countries with the highest number of COVID-19 cases and deaths worldwide. As countries begin to re-open their economies, a method for accessible and frequent surveillance of COVID-19, with the necessary rapid quarantine measures, is crucial to prevent the multiple resurgences of the disease.

The current standard of care rightfully places a strong focus on the diagnostic limit of detection, yet frequently at the expense of both cost and turnaround time. This situation has contributed to limited population testing largely due to a dearth of diagnostic resources. Quantitative real-time polymerase chain reaction (qRT-PCR) is the gold-standard method for clinical diagnosis, with high sensitivity and specificity, but these tests are accompanied by the need for trained personnel, expensive reagents and instrumentation, and a significant amount of time to execute. Facilities offering qRT-PCR sometimes require a week or longer to complete and return the results to the patient. During this waiting period the undiagnosed individual may spread the infection and/or receive delayed medical treatment. Moreover, due to the cost and relative inaccessibility of qRT-PCR in both resource-limited and abundant settings, large-scale screening using

qRT-PCR at frequent intervals remains impractical as a way to identify infected but

asymptomatic or mildly symptomatic infections. Numerous studies have reported

asymptomatic COVID-19 cases as well as a variation in viral load within and between

individuals at different time points, suggesting the need for more frequent testing for

informative surveillance.

Technologies alternate to qRT-PCR, such as rapid viral antigen detection,

clustered regularly interspaced short palindromic repeats (CRISPR), and loop-mediated

isothermal amplification (LAMP) of SARS-CoV-2 provide potential large-scale screening

applications, yet their implementation is stymied by requirements for qRT-PCR-like

accuracy before they can reach the market[8]. In countries such as India, where the qRT-

PCR resources would not be sufficient to cover monitoring of the population, the use of

rapid antigen tests is well underway[9,10]. In early May 2020, the United States Food and

Drug Administration (FDA) authorized the first antigen test for the laboratory detection of

COVID-19, citing a need for testing beyond molecular and serological methods. Antigen

testing detects the viral proteins rather than nucleic acids or human antibodies, allowing

for detection of an active infection with relative ease of sample collection and assay.

These rapid assays – like other commercially-available rapid antigen tests - can be mass-

produced at low prices and be administered by the average person without a laboratory

or instrumentation. These tests also take as little as 15 minutes to determine the result,

enabling real-time surveillance and/or diagnosis. Although antigen tests usually perform

with high specificities (true negative rate), their sensitivity (true positive rate) is often lower

when compared to molecular assays. While qRT-PCR can reach a limit of detection as

low as 10^2 genome copies per mL[11], rapid antigen testing detects viral protein that is

assumed to correlate with approximately 10^5 genome copies per mL.

We hypothesize that frequent antigen-based rapid testing even with lower

sensitivities compared to qRT-PCR - along with appropriate quarantine measures - can

be more effective at decreasing COVID-19 spread than less frequent molecular testing of

symptomatic individuals. Keeping in mind the realities of daily testing in resource-limited

regions, we also hypothesize that testing frequency can be adjusted according to the

prevalence of the disease; that is, an uptick in reported cases should be accompanied by

more frequent testing. During the viral incubation period, peak infectivity correlates with

a high viral load that can be detected by either qRT-PCR or rapid antigen testing[12–15].

Rapid tests thus optimize diagnosis for the most infectious individuals. Studies also point

to the relatively small window of time during an individual's incubation period in which the

qRT-PCR assay is more sensitive than rapid tests[12].

In this study we report the development and clinical validation of a direct antigen

rapid test for detection of SARS-CoV-2 spike glycoprotein using 121 retrospectively

collected nasopharyngeal swab specimens. Using the clinical performance data, we

develop a modeling system to evaluate the impact of frequent rapid testing on COVID-19

spread and outcomes using a variation of a SIR model, which has been previously used

to model COVID-19 transmission[16–22]. We build on this model to incorporate quarantine

states and testing protocols to examine the effects of different testing regimes. This model

distinguishes between undetected and detected infections and separates severe cases,

specifically, those requiring hospitalization, from those less so, which is important for

disease response systems such as intensive care unit triaging. We simulate COVID-19

spread with rapid testing and model disease outcomes in three regions in the United

States and São José do Rio Preto, Brazil - the site of the clinical validation study - using

publicly available data. To date, COVID-19 modeling describes the course of disease

spread in response to social distancing and quarantine measures, and a previous

simulation study has shown that frequent testing with accuracies less than qRT-PCR,

coupled with quarantine process and social distancing, are predicted to significantly

decrease infections[12,16,22–26]. This is the first modeling system using publicly-available

data to simulate how potential public health strategies based on testing performance,

frequency, and geography impact the course of COVID-19 spread and outcomes. Our

findings suggest that a rapid test, even with sensitivities lower than molecular tests, when

strategically administered 2-3 times per week, will diminish COVID-19 spread,

hospitalizations, and deaths at a fraction of the cost of nucleic acid testing via qRT-PCR.

RESULTS

Direct Antigen Rapid Test Accuracy Correlates with Viral Load Levels

Rapid antigen tests have recently been considered a viable source for first-line

screening, although concerns about the accuracy of these tests persist. We developed a

direct antigen rapid test in a lateral flow dipstick format for the detection of the spike

glycoprotein from SARS-CoV-2 in nasopharyngeal swab specimens. Of the total number

of nasopharyngeal swab specimens evaluated by qRT-PCR for amplification of SARS-

CoV-2 RNA-dependent RNA polymerase (RdRp), nucleocapsid (N), and envelope (E)

genes, 72 tested positive and 49 tested negative (Table 1).

The overall sensitivity and specificity of the rapid antigen test was 84.7% and

85.7%, respectively (Table 1). Our data demonstrate that the sensitivity of our test is

positively correlated to the viral load level (Fig. 1, Table 2). The rapid test result was

compared to the qRT-PCR cycle threshold (Ct) value measured across RdRp, N, and E

genes, and calculated as sensitivity and specificity.

165
166

Table 1. Clinical validation summary for the SARS-CoV-2 direct antigen rapid test (DART) evaluated using 121 retrospectively collected patient nasopharyngeal swab specimens.

All Data Summary								
		qRT-PCR (gene average)					95% Confidence Interval	
		+	**-**	**Total**	**Sensitivity**	84.7%	80.6%	88.9%
DART	**+**	61	7	68	**Specificity**	85.7%	80.8%	90.6%
	-	11	42	53	**Positive Predictive Value**	89.7%	86.2%	93.2%
	Total	72	49	121	**Negative Predictive Value**	79.2%	73.6%	84.9%
					Prevalence	59.5%	53.9%	65.1%
					Overall Agreement	85.1%	82.0%	88.3%

Table 2. Data summary of direct antigen tapid test (DART) performance in comparison to qRT-PCR results; for each gene being detected and amplified by qRT-PCR (E: envelope, N: nucleocapsid, RdRp: RNA-dependent RNA polymerase, and average), DART performance is organized by qRT-PCR Cycle threshold value increments.

	Cycle threshold (Ct) value	Total Cases	DART Positives	DART Negatives	Percentile Positives in Total Positive Population	Sensitivity	95% Confidence Interval (semi-interval)	Specificity	95% Confidence Interval (semi-interval)	Positive Predictive Value (PPV)	Negative Predictive Value (NPV)	Prevalence	Overall Agreement
RdRp gene	< 10	50	1	49	98.6%	100.0%	30.0%	85.7%	4.9%	12.5%	100.0%	2.0%	86.0%
	< 15	65	16	49	77.8%	100.0%	3.9%	85.7%	4.9%	69.6%	100.0%	24.6%	89.2%
	< 20	90	41	49	43.1%	97.6%	2.8%	85.7%	4.9%	85.1%	97.7%	45.6%	91.1%
	< 25	108	59	49	18.1%	96.6%	2.5%	85.7%	4.9%	89.1%	95.5%	54.6%	91.7%
	< 30	121	72	49	2.8%	84.7%	4.1%	85.7%	4.9%	89.7%	79.2%	59.5%	85.1%
	< 35	121	72	49	0.0%	84.7%	4.1%	85.7%	4.9%	89.7%	79.2%	59.5%	85.1%
N gene	< 10	49	0	49	100.0%	-	-	85.7%	4.9%	0.0%	100.0%	0.0%	85.7%
	< 15	61	12	49	83.3%	100.0%	5.1%	85.7%	4.9%	63.2%	100.0%	19.7%	88.5%
	< 20	90	41	49	43.1%	97.6%	2.8%	85.7%	4.9%	85.1%	97.7%	45.6%	91.1%
	< 25	109	60	49	16.7%	91.7%	3.5%	85.7%	4.9%	88.7%	89.4%	55.0%	89.0%
	< 30	120	71	49	1.4%	85.9%	4.0%	85.7%	4.9%	89.7%	80.8%	59.2%	85.8%
	< 35	121	72	49	0.0%	84.7%	4.1%	85.7%	4.9%	89.7%	79.2%	59.5%	85.1%
E gene	< 10	54	5	49	93.1%	100.0%	10.8%	85.7%	4.9%	41.7%	100.0%	9.3%	87.0%
	< 15	81	32	49	55.6%	100.0%	2.0%	85.7%	4.9%	82.1%	100.0%	39.5%	91.4%
	< 20	91	42	49	41.7%	97.6%	2.7%	85.7%	4.9%	85.4%	97.7%	46.2%	91.2%
	< 25	114	65	49	9.7%	90.8%	3.5%	85.7%	4.9%	89.4%	87.5%	57.0%	88.6%
	< 30	121	72	49	0.0%	84.7%	4.1%	85.7%	4.9%	89.7%	79.2%	59.5%	85.1%
	< 35	121	72	49	0.0%	84.7%	4.1%	85.7%	4.9%	89.7%	79.2%	59.5%	85.1%
Gene Average	< 10	51	2	49	97.2%	100.0%	20.9%	85.7%	4.9%	22.2%	98.9%	3.9%	86.3%
	< 15	72	23	49	68.1%	100.0%	2.8%	85.7%	4.9%	76.7%	98.9%	31.9%	90.3%
	< 20	90	41	49	43.1%	97.6%	2.8%	85.7%	4.9%	85.1%	96.6%	45.6%	91.1%
	< 25	111	62	49	13.9%	91.9%	3.4%	85.7%	4.9%	89.1%	89.4%	55.9%	89.2%
	< 30	121	72	49	0.0%	84.7%	4.1%	85.7%	4.9%	89.7%	79.2%	59.5%	85.1%
	< 35	121	72	49	0.0%	84.7%	4.1%	85.7%	4.9%	89.7%	79.2%	59.5%	85.1%

SARS-CoV-2 qRT-PCR

Figure 1. Performance of direct antigen rapid test (DART) for the detection of SARS-CoV-2 in 121 retrospectively collected patient nasopharyngeal swab specimens. Percentile Positive ranks the samples in order of high Cycle threshold (Ct) value to low Ct value. DART sensitivity is determined by calculating true positive agreement to qRT-PCR; the plot uses an ax^b+c fit and 95% confidence intervals for the sensitivity. Shown are the percentile positive cases of the total positive population conditioned to qRT-PCR Ct for RdRp (RNA-dependent RNA polymerase), N (nucleocapsid), and E (envelope) genes, and the average of these genes.

Ct value cutoffs represent the number of qRT-PCR cycles at which generated

fluorescence crosses a threshold during the linear amplification phase. The x-axis of

Figure 1 describes the percentile positives in the total positive population. Percentile

Positives ranks the samples in order of high qRT-PCR Ct value to low. In other words,

the higher the percentile, the more "positive" because fewer qRT-PCR cycles are required

for gene detection. Because the Ct value is a variable unit based upon qRT-PCR protocol

and instrumentation, we evaluated sensitivity against the percentile of cases across Ct

values.

As the percentile positive increases, the sensitivity between the rapid test and the

gold-standard qRT-PCR increases, reaching 100% sensitivity at 68.1% percentile

positive for the gene average detection. Significantly, even at percentile positives

between 0% and 68.1%, the sensitivity remains above 80%. Taken together, the clinical

data shows that the rapid antigen test performs with increasing accuracy for individuals

with a higher viral load, and are thus the most infectious[13–15].

An Enhanced Epidemiological SIDHRE-Q Model

We propose an enhanced epidemiological modeling system, *SIDHRE-Q,* a variant

of the classical SIR model in order to expand our clinical validation study and to

understand the effects of using frequent rapid tests such as the rapid antigen test on

COVID-19 outbreak dynamics. The changes we make to the basic model to encompass

the unique characteristics of the COVID-19 pandemic are similar to those presented by

Giordano et al.[16] (Fig. 2). The differential equations governing the evolution of the

SIDHRE-Q model and descriptions of the parameter values are provided in the material

and methods section (Equation 2, Table 5).

Figure 2. Graphical scheme displaying the relationships between the stages of quarantine and infection in *SIDHRE-Q* model: **Q-U**, quarantine uninfected; **S**, susceptible (uninfected); **I**, infected undetected (pre-testing and infected); **D**, infected detected (infection diagnosis through testing); **H**, hospitalized (infected with life threatening symptom progression); **R**, recovered (healed); **E**, extinct (dead); and **Q-R**, quarantine recovered (healed but in quarantine by false positive testing).

Figure Supplement 1. Graphical scheme displaying parameters between the stages of quarantine and infection in SIDHRE-Q model.

An individual that begins in **S** may either transition to a Quarantine Uninfected (**Q-**

U) state via a false positive result or to an Infected Undetected (**I**) state via interaction

with an infected individual. Should an individual in **S** move into **Q-U**, they are quarantined

for 14 days before returning to **S**, a time period chosen based on current knowledge of

the infectious period of the disease. One could also conceive of an effective strategy in

which individuals exit quarantine after producing a certain number of negative rapid tests

in the days following their initial positive result or confirm their negative result using qRT-

PCR.

Given that those diagnosed are predominantly quarantined, individuals in **I** interact

more with the **S** population than do those in Infected Detected (**D**). Therefore, the

infectious rate for **I** is assumed to be significantly larger than for **D**. Furthermore, a region's

ability to control an outbreak is directly related to how quickly and effectively people in **I**

test into **D**, reducing their infectiousness through quarantine. This study, in particular,

highlights the critical role frequency of testing, along with strict quarantine, has in

mitigating the spread of the disease and provides specific testing strategies based on

rapid tests we predict to be highly effective.

In this model, we assume that individuals receive a positive diagnosis before

developing severe symptoms and that those with symptoms severe enough to be

potentially fatal will go to the hospital. If an individual develops symptoms, we assume

they are tested daily until receiving a positive result; hence, before severe symptoms

develop, they will be diagnosed with high probability. Those who do not develop

symptoms are tested according to the frequency of tests administered to the general

population. Therefore, there is no modeled connection between **I** and **H** or between **I** and

E. Removing these assumptions would have negligible impact on the results as these

flows are very small.

Should an individual test positive and transition to **D**, they may either develop

serious symptoms requiring care or recover. Those who develop serious symptoms and

transition to state **H** will then transition to either **R** or **E**. The recovered population is

inevitably tested, as infected individuals may recover without being detected. Therefore,

the Quarantined Recovered (**Q-R**) state is introduced with the same connections to **R** as

the connections between **S** and **Q-U**. Though the reinfection rate of SARS-CoV-2 has

been a point of recent debate, it is assumed that the number of re-infected individuals is

small[27–31]. Therefore, individuals cannot transition from **R** to **S**, hence the separately

categorized quarantined populations.

We considered several variations and extensions of the *SIDHRE-Q* model. In

simulations, we tested additional states, such as those in the *SIDARTHE* model, which

include distinctions between symptomatic and asymptomatic cases for both detected and

undetected populations[16]. Incorporating information about the correlations between viral

load and infectivity and sensitivity were also considered. Altogether, our modeling system

has been well tuned to predict the impact of high frequency rapid testing on COVID-19

spread and outcomes.

Frequent Rapid Testing with Actionable Quarantining Dramatically Reduces

Disease Spread

In order to demonstrate how strategies could affect the disease spread in different

geographies and demographics, we used surveillance data obtained from regions of

varying characteristics: the state of Massachusetts (MA), New York City (NYC), Los

Angeles (LA), and São José do Rio Preto (SJRP), Brazil, the site of the rapid antigen test

clinical validation study. These regions are also selected in our study due to the readily

available surveillance data provided by the local governments. We fit the model to the

data from each region starting 1 April 2020. At this time point the disease reportedly is

most advanced in NYC and least advanced in SJRP, Brazil with estimated cumulative

infection rates of 7.11% and 0.12%, respectively.

After calibrating the *SIDHRE-Q* model, the disease spread is observed with varying

validated rapid antigen test performances and frequencies (Fig. 3). Sensitivity (the ratio

of true positives to the total number of positives) and specificity (the ratio of true negatives

to the total number of negatives) compared to gold-standard qRT-PCR were used as

measures of test accuracy.

The rapid test frequency is varied while maintaining an accuracy of 80% sensitivity

and 90% specificity, comparable to our clinical data collected in SJRP, Brazil. These

testing scenarios are then compared to symptomatic testing, in which individuals receive

a rapid test only when presenting symptoms, via either a rapid test or qRT-PCR. Since

the primary testing regiment deployed in MA, LA, NYC and SJRP, Brazil is qRT-PCR-

based and focused on symptomatic individuals, the symptomatic testing protocol via qRT-

PCR is directly estimated from the data to be the rate v (Table 5).

The difference between the qRT-PCR and rapid test simulations (red and orange

lines, respectively) is therefore only sensitivity of testing (Fig. 3). We assumed that test

outcome probability is a function only of whether an individual is infected and independent

of other factors; one can consider this a lower bound on effectiveness of a strategy, as

sensitivity and infectivity are often positively correlated with antigen testing.

Sensitivity: 80.0% Specificity: 90.0%

Figure 3. COVID-19 Outcomes in 3 US Regions and Brazil as a result of Frequent Rapid Testing Protocol using the *SIDHRE-Q* Model. The Cumulative Detected Infected, Hospitalized, Deceased, Active Infections, Recovered, and Quarantined are modeled over 105 days (top to bottom) using reported data from 4 global regions: Massachusetts, Los Angeles, New York City, and São José do Rio Preto in Brazil (left to right). The COVID-19 population spread and outcomes are modeled under a Rapid Testing Protocol (sensitivity 80%, specificity 90%) with variable testing frequencies ranging from 1-21 days between tests. This protocol is compared to a symptom-based Rapid Testing protocol and a symptom-based qRT-PCR protocol.

Figure Supplement 1. COVID-19 Outcomes as a result of Frequent Rapid Testing Protocol with variable test performances using SIDHRE-Q Model.

Figure Supplement 2. Time series of the four fitted parameters α, ν, μ, and τ.

283   To better understand the effect of rapid testing frequency and performance on

284   healthcare capacity and mortality rates, we simulate the testing strategy with 30%-90%

285   sensitivity each with 80% or 90% specificity against the symptomatic testing strategy.

286   Table 3 describes the different test performances implemented in the model matched with

287   corresponding Figure 3 - Figure Supplement 1 plots.

Table 3. Test Performance with corresponding Figure 3 - Figure Supplement 1 designation.

Sensitivity (%)	Specificity (%)	Fig. 3 Supplemental Fig.
90	90	1A
70	90	1B
50	90	1C
30	90	1D
90	80	1E
70	80	1F
50	80	1G
30	80	1H

As per our hypothesis, frequency and symptom-based testing dramatically

reduced infections, simultaneous hospitalizations, and total deaths when compared to the

purely symptom-based testing regiments, and infections, hospitalization, and death were

reduced as frequency increased. Although testing every day was clearly most effective,

even testing every fourteen days with an imperfect test gave an improvement over

symptomatic testing with qRT-PCR. While the strategy works best when implemented at

the very beginning of an outbreak, as demonstrated by the results in SJRP, Brazil, it also

works to curb an outbreak that is already large, as demonstrated by the results in NYC.

The difference between frequencies is more noticeable when the testing strategy is

applied to the outbreak in NYC, leading us to hypothesize that smaller outbreaks require

a lower testing frequency than larger ones; note the difference between the dependence

on frequency to curb a small initial outbreak in SJRP, Brazil versus a large one in NYC

(Fig. 4, Figure 4 - Figure Supplement 1).

For test performance of 80% sensitivity and 90% specificity, the percent of the

population that has been infected in total from the beginning of the outbreak to mid-July

drops from 18% (MA), 11% (LA), 26% (NYC), and 11% (SJRP, Brazil) to 3%, 2%, 12%,

and 0.26%, respectively, using a weekly rapid testing and quarantine strategy (with

regards to predictions of overall infection rates, other studies based on seroprevalence

and epidemiological predictions have reached similar conclusions[32,33]). If testing is

increased to once every three days, these numbers drop further to 1.6% (MA), 1.4% (LA),

9.4% (NYC), and 0.19% (SJRP, Brazil) (Table 4).

Table 4. Summary of results of COVID-19 outcomes in 3 US Regions and Brazil as a result of Frequent Rapid Testing Protocol using SIDHRE-Q Model.

	Massachusetts		Los Angeles		New York City		São José do Rio Preto, Brazil	
	qRT-PCR	1 per 3 days	qRT-PCR	1 per 3 days	qRT-PCR	1 per 3 days	qRT-PCR	1 per 3 days
Total Infected	18.40%	1.60%	11.70%	1.42%	26.40%	9.45%	11.70%	0.186%
Max Hospitalized	0.056%	0.025%	0.028%	0.022%	0.144%	0.130%	0.054%	0.003%
Total Deaths	0.119%	0.029%	0.039%	0.009%	0.226%	0.157%	0.040%	0.003%

To further examine the relationship between frequency and sensitivity, we modeled

the maximum number of individuals in a given state over the 105-day time period for four

geographic regions (Fig. 4). In all four geographic regions, as frequency of testing

increases, the total infections, maximum simultaneous hospitalizations, and total deaths

converge to small percentages regardless of the sensitivity at high frequencies. It is clear

that the difference in frequency required to achieve the same result using tests of differing

sensitivities is very small (Fig. 4) For example, we predict that for the outbreak in LA, a

testing strategy started on 1 April of every 10 days using a test of sensitivity 90% would

have resulted in 2.5% of the population having been infected, while using a test of

sensitivity 30% would require a strategy of every 5 days to achieve the same number.

Thus, we conclude that frequency is more important than sensitivity in curbing the spread,

and a large range of sensitivities prove effective when testing sufficiently often. How

frequently, exactly, depends on the specific outbreak and what stage it is in, which leads

us to the location-based deployment strategy discussed in a later section. Frequency of

testing can be significantly reduced to effectively contain the disease once the initial

outbreak has been controlled; it is clear that this takes only a matter of weeks (Fig. 3).

On the other hand, according to the specificity of the rapid test and the quarantine

duration, larger testing frequency result in a larger percent of the population quarantined

(Fig. 3). Assuming a 90% rapid test specificity and 14-day quarantine duration, for the 1-

, 3- and 7-day frequencies almost 60%, 38% and 20% of the population, respectively,

would be quarantined. This figure may be reduced with additional rules for exiting

quarantine early, such as after complementary testing. An example of such a strategy is

that individuals who test positive are required to either quarantine for two weeks or

produce two consecutive negative rapid tests in the two days following their positive

result. Assuming 80% sensitivity and 90% specificity, those individuals will reenter the

public while still infected with probability 0.04. If uninfected, that individual will exit

quarantine after two days with probability 0.81. However, a compromise between the

reduction of infections and the proportion of the population in quarantine would be part of

the planning for the appropriate testing protocol in each community or region.

Additionally, while high frequency may be necessary to contain a large outbreak

initially, relatively infrequent testing, such as every one or two weeks, is sufficient to keep

controlled outbreaks small, while reducing the number of quarantined individuals to less

than 10% of the population using a two-week mandatory quarantine.

Figure 4. Effect of Rapid Testing Protocol under variable testing sensitivities and increasing frequency under the *SIDHRE-Q* Model. The Cumulative Infections, Maximum Simultaneously Hospitalized, and Deceased populations are modeled for Massachusetts, Los Angeles, New York City, and São José do Rio Preto in Brazil. The effect of increasing frequency of testing is modeled for various testing sensitivities (30%-90%) with a 90% specificity.

Figure 4 - Figure Supplement 1. Effect of Rapid Testing Protocol under variable testing sensitivities and increasing frequency under the *SIDHRE-Q* Model (80% specificity).

A County-Based Testing Strategy Offers a Cost-effective Approach to Large-scale COVID-19 Surveillance

To examine the effects of resource-strategic testing schemes, we modeled the COVID-19 prevalence by varying testing frequency across counties of California. For this analysis, only California was analyzed because of the accessibility of the county level data. In this scheme, the percent of active infected detected individuals in a county determines the frequency of testing. We define thresholds for the number of active detected infections that, when hit, initiate testing protocols of different frequencies depending on the threshold hit. We first tested evenly spaced thresholds for the number of detected active infections up to 1% of the population, but later adopted thresholds that were determined according to Equation 1. In Equation 1, D = population of state **D** at the time of testing. T = number of active infections which, if reached, initiates everyday testing. The days between tests are rounded to the closest integer value.

$$\text{Days between tests} = \max\left(1, 2\log_2(T/D) + 1\right)$$

(1)

The days between tests are chosen such that the detected active infections should remain near to or below T. If the initial detected active infections are greater than T, then the testing frequency of 1 will cause infections to rapidly drop. Both the threshold at which everyday testing begins and the coefficient of $log_2 T/D$ can be modified to produce a strategy that is more or less frequent in testing or resource effective; a range of days between tests from 14 days to 1 day are used (Fig. 5). A scan over different choices of T is shown in the supplements to Figure 5; the threshold we choose in Figure 5 is 0.05% because it is successful in curbing the outbreak within the time period we consider. While

378　these choices work for the epidemic in California at the point we start our simulations, 10

379　April, they do not necessarily reflect the most resource effective choices everywhere. Our

380　analysis could be redone to select the best strategy in other states or in the country as a

381　whole.

Figure 5. Effect of County Based Rapid Test Protocol (A) and Uniform qRT-PCR Protocol
(B) on active infected detected population over time in California (CA). The legend denotes
the thresholds at which testing frequency is determined, the testing frequencies, the
percent of CA population under the strategy, and the cost per person per day.

Figure Supplement 1. Effect of County Based Rapid Testing strategy on COVID-19
outcomes in California.

Figure Supplement 2. Time series of the three fitted pieces of data Cumulative Cases,
Daily Hospitalized, and Cumulative Deaths for each CA county receiving testing.

Using a rapid test with a sensitivity of 80% and specificity of 90%, the county-based

testing with threshold 0.05% reduces the active infections from 0.94% to 0.0005%, while

the uniform strategy with tests administered every 7 days results in double the number of

active infections (Fig. 5). As the threshold is reduced, the total cost increases while the

cumulative infections, maximum percentage hospitalized, and cumulative deaths all

decrease (Figure 5 - Figure Supplement 1).

Strategy B in Figure 5 consists of qRT-PCR testing uniformly applied to the

highlighted population with a frequency of once weekly. The average cost per person per

day is just under $15. Despite this frequency and the accuracy of qRT-PCR, the strategy

does not succeed in curbing the spread as fast as strategy A, which uses a testing

sensitivity and specificity of 80% and 90%, respectively, and testing frequency that vary

between counties depending on the proportion of their population that is currently

infected. The total cost for strategy A is estimated at a fraction of the other at $1.53 per

person per day.

398
399

DISCUSSION

In this study we examine the potential effects of a novel testing strategy to limit the

spread of SARS-CoV-2 utilizing rapid antigen test screening approaches. Our clinical data

and *SIDHRE-Q* modeling system demonstrate that 1) frequent rapid testing even at a

range of accuracies is effective at reducing COVID-19 spread, 2) rapid antigen tests are

a viable source for this strategy and diagnose the most infectious individuals, and 3)

strategic geographic-based testing can optimize disease control with the amount of

available resources. The information from a diagnostic test itself is of tremendous value,

as it can prompt the necessary quarantine measures to prevent spread, guide proper care

and triage, and provide crucial disease-tracking information. Diagnostic testing in the

United States and abroad, however, has been a significant public health hurdle. The

public has witnessed and experienced symptomatic individuals being denied testing due

to shortages, and few testing structures for asymptomatic or mildly symptomatic

individuals - the main source of disease spread. Though several factors contributed to the

stymied early response measures, such as lockdown and quarantine protocols and

adherence, severe testing bottlenecks were a significant culprit[34–36]. Early control

measures have been shown to decrease lives lost by several orders of magnitude[37].

These challenges, though exacerbated during the early months of the pandemic, remain

at the forefront of the public health crises.

Diagnosis of SARS-CoV-2 infection by qRT-PCR is the current standard of care,

yet remains expensive and requires a laboratory and experienced personnel for sample

preparations and experimentation. Significantly, the turnaround time for results can be up

to 10 days[38]. On an individual scale, this leaves the public in limbo, preventing people

from either leaving quarantine if they are negative, or delaying critical care and infecting

others if they are positive. On a societal level, this current testing scheme yields

incomplete surveillance data on which response efforts such as societal reopening and

hospital management depend. Though qRT-PCR is considered the gold-standard

diagnostic method because of its high sensitivity and specificity, the logistical hurdles

render it unrealistic for large-scale screening.

As qRT-PCR remains impractical for this strategy, and rapid tests are facing

regulatory challenges because they do not perform with qRT-PCR-like accuracy, rapid

test screening is either nonexistent in several countries or symptom-based. Even under

best-case assumptions, findings have shown that symptom and risk-based screening

strategies miss more than half of the infected individuals[39]. Some have argued that the

need for widespread testing is overstated due to the variability in test sensitivity and

specificity[40]. Our findings show, however, that test performance, though valuable, is

secondary to widespread test frequency, which is enabled by accessibility and turnaround

time.

Giordano et al. has modeled the evolution of SARS-CoV-2 spread, introducing a

diagnosed state to elucidate the importance of population-wide testing[16]. Mina et al. has

examined how various test sensitivities and frequencies affect the reproductive number[12].

We build upon these findings to show how in affected United States and Brazil regions,

population-wide frequent and rapid testing schemes, with sensitivities ranging from 30%-

90%, can be more effective in curbing the pandemic than a PCR-based scheme.

Integrating real-world surveillance and clinical data into our modeling system has allowed

us to incorporate regional differences - such as variances in healthcare access, state

health policy and adherence, state GDP, and environmental factors - under the same

model. Significantly, our findings hold true across Massachusetts, New York City, Los

Angeles, and São José do Rio Preto, Brazil. We also present the economic

considerations of these testing regimes, showing that widespread rapid testing is more

cost efficient than less frequent qRT-PCR testing. In line with these economic

considerations, our model demonstrates the effectiveness of a geographic-based

frequent testing regime, in which high disease prevalence areas receive more frequent

testing than low disease prevalence areas.

Since COVID-19 is known to affect certain demographics differently, modeling

would benefit from incorporating demographic information correlated with disease

progression and spread to define sub-models and sets of parameters accordingly. Age,

pre-existing conditions, job types, and density of population are examples of possible

categories, each of which influence the risk of contracting and/or dying from COVID-19.

Further studies may benefit from incorporating these ideas should more information

become available.

Our findings also point to low-cost tools for implementation of this testing strategy,

such as a rapid antigen-based test for the detection of SARS-CoV-2 proteins. We show

that the rapid antigen test performs with a range of accuracies under which disease

spread can be dramatically mitigated under our model. Notably, the sensitivity is

correlated to the individual's viral load, effectively diagnosing those who are most

infectious with the highest accuracy. Our findings are significant because these rapid

antigen tests are cheaper than qRT-PCR, can be mass produced to millions per day,

present results within 15 minutes, and can be administered by a nonexpert without a lab

or special equipment.

There are several policy implications for these findings. First, our model supports

that systems of high frequency rapid testing should be implemented as a first-line

screening method. This can be first enabled by a more holistic regulatory evaluation of

rapid diagnostics, such that policy emphasizes accessibility and turnaround time even

under a range of accuracies. One can imagine a less accurate, though rapid method of

first-line screening in schools, public transportation, and airports, or even at home, and a

qRT-PCR-based method for second-line screening (testing those who present severe

symptoms or have been in contact with infected individuals, testing in a clinical setting,

etc). Second, our cost analysis and rapid antigen test data present a viable and potentially

more cost-effective method for screening. Third, our county-based testing scheme

presents a possible method for wide-scale screening while optimizing resources. Future

studies should investigate how this selective testing strategy can be applied to different

location scales to further inform health policy. Moreover, though our models analyze

regions in the United States and Brazil, similar testing strategies can be considered

globally in both resource limited and abundant settings due to the higher accessibility of

rapid tests compared to qRT-PCR.

We emphasize that integral to the effectiveness of diagnostic schemes is 1) the

proper adherence to quarantine measures and 2) the combined use of a variety of

diagnostic methods including nucleic acid, antigen, and antibody tests. According to these

models, rapid antigen tests are an ideal tool for first-line screening. Clinical molecular

tests such as qRT-PCR are vital to the diagnostic landscape, particularly to re-test

suspected cases that were negative on the rapid test. Because rapid tests present a

higher rate of false negatives, methods such as qRT-PCR remain integral to second-line

screening. Antibody tests provide important information for immunity and vaccination

purposes as well as epidemiological surveillance. This model also assumes that

individuals will quarantine themselves before being tested and for 14 days following a

positive diagnostic result.

Our simulations combined with real-world data demonstrate a robust modeling

system and elucidates the significance of this novel testing strategy. However, there are

important limitations to be considered. Differences in disease reporting between the

geographical regions and the incomplete nature of COVID-19 surveillance data, often due

to the lack of testing, are not considered in the model. It is imperative that the testing

results, hospitalization and death statistics, and changes in protocol are reported in real-

time to scientists and policy makers so that models can be accurately tuned as the

pandemic develops. The model also does not take into account infrastructural limitations

such as hospital capacity. Though the rapid antigen test offers several advantages such

as affordability, fast turnaround time, and ease of mass production, we are also assuming

that there are systems in place to implement frequent and safe low-cost screening across

different communities and settings.

Our model underscores the need for a point-of-care or at-home test for frequent

screening, particularly as lockdown restrictions ease. Regulatory agencies such as the

FDA could work towards regulating rapid tests to alternative standards other than

comparison to high sensitivity molecular diagnostics, as our model shows that frequency

and scale of testing may overcome lower sensitivities. Rather, we could refocus policy to

implement first-line screening that optimizes accuracy with efficiency and equitability.

MATERIAL AND METHODS

Development of Direct Antigen Rapid Test for the Detection of SARS-CoV-2

We developed a direct antigen rapid test for the detection of the spike glycoprotein

from SARS-CoV-2 in nasopharyngeal swab specimens as previously described[41]. Briefly,

the rapid antigen test is an immunochromatographic format with a visual readout using

anti-spike mouse monoclonal antibodies (E25Bio, Inc., Cambridge, MA, USA) that are

either coupled to 40 nm gold nanoparticles (Abcam, Cambridge, UK) or adsorbed to

nitrocellulose membranes (Sartorius, Goettingen, Germany). Each rapid antigen test has

a control area adjacent to the paper absorbent pad; the control is an anti-mouse Fc

domain antibody (Leinco Technologies, Fenton, MO, USA) that will capture any of the

antibody-conjugated gold nanoparticles to generate a control visual signal. A visual signal

at the test area reflects SARS-CoV-2 spike glycoprotein that is "sandwiched" between an

anti-spike glycoprotein antibody adsorbed to the nitrocellulose membrane and a second

anti-spike glycoprotein antibody covalently coupled to visible gold nanoparticles.

Validation of Direct Antigen Rapid Test for the Detection of SARS-CoV-2

In a retrospective study of nasopharyngeal swab specimens from human patients,

we compared the accuracy of the rapid antigen test to the viral load of individuals.

Nasopharyngeal swab specimens (n = 121) were tested in Brazil following approved

human subjects use protocols. The age of study participants ranged from 1 to 95 years

with an overall median of 37 years (interquartile range, 27–51 years), and 62% were

female. The demographic summary of the patients are included in Supplementary Table

1. The nasopharyngeal swab specimens were banked refrigerated or frozen samples

from suspected patients submitted to the lab for routine COVID diagnosis. Prior to using

the rapid test, the nasopharyngeal swab samples were validated by qRT-PCR using

GeneFinder^{TM} COVID-19 Plus Real*Amp* Kit (OSANGHealtcare, Anyang-si, Gyeonggi-do,

Republic of Korea I). The primary study under which the samples and data were collected

received ethical clearance from the Faculdade de Medicina de São José do Rio Preto

(FAMERP), protocol number 31588920.0.0000.5415. All excess samples and

corresponding data were banked and de-identified prior to the analyses.

A nasopharyngeal swab specimen (1 mL) was concentrated using a Vivaspin 500

centrifugal concentrator (Sartorius, Goettingen, Germany) at 12,000 x g for 10 minutes.

The concentrated nasopharyngeal swab specimen retentate was transferred to a

collection tube and the rapid antigen test was inserted into the tube with the retentate and

allowed to react for 15 minutes. After processing of the rapid antigen test, the visual

positive or negative signal was documented.

Data for Modeling

As of August 2020, the United States and Brazil have the highest number of

confirmed COVID-19 cases and deaths worldwide, with both countries reporting their first

case on 26 February 2020[1]. Although several affected US regions could have been

modeled, we look at data from Massachusetts, New York, and Los Angeles: these regions

each contained "hotspots", or areas of surging COVID-19 cases, at different points in time

during the pandemic and have publicly available government-provided surveillance data.

Our model is fit using data over 105 days beginning on April 1 for Figures 3 and 4 and

105 days beginning on April 10 for Figure 5 (see "Modeling Parameters" in Methods). In

order to understand the various testing proposals on a global scale, we performed our

clinical study in and expanded the modeling study to Brazil. The specific data we use to

fit our model are cumulative confirmed cases, total deaths, and number of daily

hospitalizations due to COVID-19. This surveillance data was retrieved from government-

provided online databases[42–48].

Modeling Parameters

Equation 2 below provides the exact differential equations governing the model.

$$
\begin{aligned}
d\mathbf{S} &= & -\mathbf{S}\left(\alpha\mathbf{I} + \eta\mathbf{D} + \gamma\right) & & +\psi\mathbf{Q_U} \\
d\mathbf{I} &= & -\mathbf{I}\left(\varepsilon + \lambda + \nu\right) & & +\mathbf{S}\left(\alpha\mathbf{I} + \eta\mathbf{D}\right) \\
d\mathbf{D} &= & -\mathbf{D}\left(\dfrac{\mathbf{D}+\mathbf{I}}{\mathbf{D}}\mu + \rho\right) & & +\mathbf{I}(\nu + \varepsilon) \\
d\mathbf{H} &= & -\mathbf{H}\left(\sigma + \tau\right) & & +\mu\left(\mathbf{D} + \mathbf{I}\right) \\
d\mathbf{E} &= & & & +\tau\mathbf{H} \\
d\mathbf{R} &= & -\gamma\mathbf{R} & & +\rho\mathbf{D} + \lambda\mathbf{I} + \sigma\mathbf{H} + \psi\mathbf{Q_R} \\
d\mathbf{Q_U} &= & -\psi\mathbf{Q_U} & & +\gamma\mathbf{S} \\
d\mathbf{Q_R} &= & -\psi\mathbf{Q_R} & & +\gamma\mathbf{R}
\end{aligned}
$$

570   (2)

In order to determine the values of the parameters defining the flows between states, we

use a least squares regression performed at seven day intervals in the datasets to which

we fit. This allows the model to take into account the time dependent nature of the

parameters, which rely on factors such as social distancing regulations and changes in

testing capacity. We also fit window sizes between 1 and 21 days and find that while the

fit degrades with larger window size, the overall shape of the fits do not change. We

choose seven days assuming policy changes take a week to become effective and that

reasonable parameters can be expected to change within this time period. Also, the seven

day window size accounts for the fact that often data is not reported as diligently over the

weekend. Time series of the values of the parameters for the geographic locations

discussed in this paper can be found in the supplemental materials for Figure 2.

Given the restrictions on data available for the populations of various states,

varying all of the parameters results in an over parameterized system. Therefore, a subset

of the model parameters are fit while the others are either extracted from other sources;

see Table 5.

 Table 5. Details of parameter values used for *SIDHRE-Q* Model.

Parameter	Details & Statistics			
α	α is the probability that an interaction between an undetected infected person and an uninfected person results in a new infection, divided by the average number of uninfected people an undetected infected person comes into contact with on a given day. α is estimated from the data.		Mean	St. Dev.
		MA	0.088	0.051
		LA	0.090	0.034
		NYC	0.067	0.042
		SJRP	0.121	0.042
η	η is the probability that an interaction between an infected person and an uninfected person results in a new infection, divided by the average number of uninfected people a detected infected person comes into contact with on a given day. $\eta = 0.01 \cdot \alpha$ The constant relating η, α accounts for a small but nonzero transmission due to the quarantined (detected) infected population. This value was chosen to be small, assuming a quarantined individual will only infect others with low probability.			
ν	ν is the probability that a symptomatic undetected individual is diagnosed on a given day. ν is estimated from the data. ν is multiplied by sensitivity (assume benchmark sensitivity 100% for PCR, as used when fitting).		Mean	St. Dev.
		MA	0.006	0.005
		LA	0.011	0.006
		NYC	0.0056	0.002
		SJRP	0.015	0.007
ϵ	ϵ is the probability that an asymptomatic undetected infected individual is diagnosed on a given day. $\epsilon = 0$ while fitting (during PCR symptomatic testing). $\epsilon =$ (sensitivity/days between tests) when the rapid testing strategy is activated.			
λ	λ is the probability that an undetected infected individual transitions to the recovered state on a given day. $\lambda = 1/14$, or the inverse of average recovery time.[48]			
μ	μ is the probability that an infected individual develops severe symptoms on a given day and transitions into the hospitalized state. The flow from D to H is assumed to be independent of the ratio I/D, but comes only from the detected infected population, hence why it is multiplied by $(I + D)/D$. μ is estimated from the data.		Mean	St. Dev.
		MA	0.0013	9.5e-4
		LA	0.0016	2.4e-4
		NYC	0.0011	6.6e-4
		SJRP	0.0018	8.0e-4
ρ	ρ is the probability that a detected infected individual transitions to the recovered state on a given day. $\rho = 1/14$, or the inverse of the average recovery time. [48]			

			Mean	St. Dev.
σ	σ is the probability that a hospitalized individual transitions to the recovered state on a given day. $\sigma = 1/11$, or the inverse of the average recovery time for a hospitalized individual. [48]			
τ	τ is the probability that a hospitalized individual expires on a given day. τ is estimated from the data.		Mean	St. Dev.
		MA	0.034	0.012
		LA	0.016	0.004
		NYC	0.036	0.034
		SJRP	0.032	0.045
γ	γ is the probability of entering either of the quarantine states on a given day from either the Susceptible or Recovered populations. $\gamma = 0$ while fitting (during PCR symptomatic testing). $\gamma = (1 - \text{specificity}) \times (1/\text{days between tests})$ when the rapid testing strategy is activated.			
ψ	ψ is the probability that an individual exits quarantine on a given day. $\psi = 1/14$, or the inverse of the quarantine period for fixed length quarantine.			

The fitting procedure minimizes the sum of the squared residuals of the total cases,

current daily hospitalizations, cumulative deaths, and percentage of total infected

individuals currently hospitalized. The first three are present in the data sets while the

latter is derived from the estimates of the ratio between infected undetected to infected

detected individuals from the CDC Laboratory Seroprevalence Survey Data[49]. While this

ratio changes over time, the percentage of infected individuals developing severe

symptoms should remain roughly constant throughout the course of the epidemic in the

different locations studied.

We consider the data sets for outbreaks in MA, NYC, LA, and SJRP, Brazil[42–47].

While each location has testing and fatality information dating back to January,

hospitalization data was not included until late March (for NYC and SJRP) and April (for

MA and LA). Hence we begin our fitting procedure and testing strategy on 1 April for

each of the data sets; by this point, the outbreak is advanced in NYC, substantial in MA,

non-negligible, but far from its peak, in LA, and in early stages in SJRP, Brazil. Starting

simulations at various stages of the outbreak allows one to see the difference in results

between when a testing strategy is administered.

In order to determine the effectiveness of the county-based strategy when applied

to the state of California, we also fit all of the counties in California with a population

greater than 1.5% of that of the entire state and with greater than zero deaths. The results

do not depend on these selections, but instead suggest a practical criteria to administer

limited resources. The fitting is done starting 10 April for these counties, as at this point

the outbreak is sufficiently well-documented in each to successfully model. For the

county-level data we compute a seven day running average of each of the data sets to

which we then fit in order to smooth out fluctuations in the data, likely due to reporting,

which are more significant here than in the other data sets considered, as the county

populations are smaller and hence discrepancies impact the smoothness of the data

more. The fits for each of the counties can be found in the supplementary materials to

Figure 5.

As one can see from Figure 2, these data sets are particularly not smooth, which

indicates inefficiencies in reporting. Additionally, it is difficult to gauge their consistency

within the dates provided or to compare between locations, as reporting mechanisms

changed over time within the same locations. Despite this lack of consistency, our model

and fitting mechanism was successful in reproducing the progress of the outbreak in each

data set studied.

The authors confirm that the data supporting the findings of this study are available within

the article and/or its supplementary materials; any other data will be made available upon

request.   Our   code   can   be   found   on   github:

https://github.com/badeaa3/COVID19_Rapid_Testing. The code is written using python

with the packages scipy, numpy, lmfit, matplotlib and plotly[50–54].

Acknowledgments

We thank Professor Lee Gehrke for critical reading of the manuscript. The study is

funded, in part, by a Bill and Melinda Gates Foundation Award (INV-017872) to E25Bio,

Inc. EN is funded by Tufts University DISC Seed Grant. MLN is supported by a FAPESP

grant (#2020/04836-0) and is a CNPq Research Fellow. AFV is supported by a FAPESP

Fellow grant (#18/17647-0). GRFC is supported by a FAPESP Fellow grant (#20/07419-

0). BHGAM is supported by a FAPESP Scholarship (#19/06572-2). The funders had no

role in the design of the study; in the collection, analyses, or interpretation of data; in the

writing of the manuscript, or in the decision to publish the results.

Competing Interest

BN, AB, AR, MB, NS, AG, IB, and BBH are employed by or affiliated with E25Bio

Inc. (www.e25bio.com), a company that develops diagnostics for epidemic viruses.

References

1. Coronavirus Disease (COVID-19) Situation Reports.

https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports.

2. Menkir, T. F. *et al.* Estimating the number of undetected COVID-19 cases exported

internationally from all of China. *medRxiv* (2020) doi:10.1101/2020.03.23.20038331.

3. Ivorra, B., Ferrández, M. R., Vela-Pérez, M. & Ramos, A. M. Mathematical modeling of the

spread of the coronavirus disease 2019 (COVID-19) taking into account the undetected

infections. The case of China. *Commun Nonlinear Sci Numer Simul* 105303 (2020)

doi:10.1016/j.cnsns.2020.105303.

4. Salathé, M. *et al.* COVID-19 epidemic in Switzerland: on the importance of testing, contact

tracing and isolation. *Swiss Med Wkly* **150**, w20225 (2020).

5. Lau, H. *et al.* Evaluating the massive underreporting and undertesting of COVID-19 cases in

multiple global epicenters. *Pulmonology* (2020) doi:10.1016/j.pulmoe.2020.05.015.

6. Silverman, J. D., Hupert, N. & Washburne, A. D. Using influenza surveillance networks to

estimate state-specific prevalence of SARS-CoV-2 in the United States. *Science Translational*

Medicine (2020) doi:10.1126/scitranslmed.abc1126.

7. Böhning, D., Rocchetti, I., Maruotti, A. & Holling, H. Estimating the undetected infections in

the Covid-19 outbreak by harnessing capture-recapture methods. *Int. J. Infect. Dis.* **97**, 197–

201 (2020).

8. Baek, Y. H. *et al.* Development of a reverse transcription-loop-mediated isothermal

amplification as a rapid early-detection method for novel SARS-CoV-2. *Emerg Microbes Infect*

9, 998–1007 (2020).

9. Leo, L. Mylab gets commercial approval from ICMR for Covid-19 antigen rapid testing kit.

Livemint https://www.livemint.com/news/india/mylab-gets-commercial-approval-from-icmr-

for-covid-19-antigen-rapid-testing-kit-11595434040321.html (2020).

10. Dey, S. Coronavirus testing: Rapid antigen tests now make up nearly half of daily checks |

India News - Times of India. *The Times of India*

https://timesofindia.indiatimes.com/india/rapid-antigen-tests-now-make-up-nearly-half-of-

daily-checks/articleshow/77340459.cms.

11. Vogels, C. B. F. *et al.* Analytical sensitivity and efficiency comparisons of SARS-CoV-2 RT-

qPCR primer-probe sets. *Nat Microbiol* (2020) doi:10.1038/s41564-020-0761-6.

12. Larremore, D. B. *et al.* Test sensitivity is secondary to frequency and turnaround time for

COVID-19 surveillance. *medRxiv* (2020) doi:10.1101/2020.06.22.20136309.

13. Shen, Z. *et al.* Superspreading SARS events, Beijing, 2003. *Emerging Infect. Dis.* **10**, 256–

260 (2004).

14. Peiris, J. S. M. *et al.* Clinical progression and viral load in a community outbreak of

coronavirus-associated SARS pneumonia: a prospective study. *Lancet* **361**, 1767–1772

(2003).

15. He, X. *et al.* Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat. Med.*

26, 672–675 (2020).

16. Giordano, G. *et al.* Modelling the COVID-19 epidemic and implementation of population-wide

interventions in Italy. *Nat. Med.* **26**, 855–860 (2020).

17. Choi, S. & Ki, M. Estimating the reproductive number and the outbreak size of COVID-19 in

Korea. *Epidemiol Health* **42**, e2020011 (2020).

18. Wei, Y. Y. *et al.* [Fitting and forecasting the trend of COVID-19 by SEIR(+CAQ) dynamic

model]. *Zhonghua Liu Xing Bing Xue Za Zhi* **41**, 470–475 (2020).

19. Yang, Z. *et al.* Modified SEIR and AI prediction of the epidemics trend of COVID-19 in China

under public health interventions. *J Thorac Dis* **12**, 165–174 (2020).

20. Cao, S., Feng, P. & Shi, P. [Study on the epidemic development of COVID-19 in Hubei

province by a modified SEIR model]. *Zhejiang Da Xue Xue Bao Yi Xue Ban* **49**, 178–184

(2020).

21. Huang, R., Liu, M. & Ding, Y. Spatial-temporal distribution of COVID-19 in China and its

prediction: A data-driven modeling analysis. *J Infect Dev Ctries* **14**, 246–253 (2020).

22. Godio, A., Pace, F. & Vergnano, A. SEIR Modeling of the Italian Epidemic of SARS-CoV-2

Using Computational Swarm Intelligence. *Int J Environ Res Public Health* **17**, (2020).

23. Gatto, M. *et al.* Spread and dynamics of the COVID-19 epidemic in Italy: Effects of emergency

containment measures. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 10484–10491 (2020).

24. Hou, C. *et al.* The effectiveness of quarantine of Wuhan city against the Corona Virus Disease

2019 (COVID-19): A well-mixed SEIR model analysis. *J. Med. Virol.* **92**, 841–848 (2020).

25. Zhou, T. *et al.* Preliminary prediction of the basic reproduction number of the Wuhan novel
704   coronavirus 2019-nCoV. *J Evid Based Med* **13**, 3–7 (2020).

705   26. Reno, C. *et al.* Forecasting COVID-19-Associated Hospitalizations under Different Levels of
706   Social Distancing in Lombardy and Emilia-Romagna, Northern Italy: Results from an
707   Extended SEIR Compartmental Model. *J Clin Med* **9**, (2020).

708   27. Alizargar, J. Risk of reactivation or reinfection of novel coronavirus (COVID-19). *J. Formos.*
709   *Med. Assoc.* **119**, 1123 (2020).

710   28. Batisse, D. *et al.* Clinical recurrences of COVID-19 symptoms after recovery: viral relapse,
711   reinfection or inflammatory rebound? *J. Infect.* (2020) doi:10.1016/j.jinf.2020.06.073.

712   29. Deng, W. *et al.* Primary exposure to SARS-CoV-2 protects against reinfection in rhesus
713   macaques. *Science* (2020) doi:10.1126/science.abc5343.

714   30. Ota, M. Will we see protection or reinfection in COVID-19? *Nat. Rev. Immunol.* **20**, 351 (2020).

715   31. Victor Okhuese, A. Estimation of the Probability of Reinfection With COVID-19 by the
716   Susceptible-Exposed-Infectious-Removed-Undetectable-Susceptible Model. *JMIR Public*
717   *Health Surveill* **6**, e19097 (2020).

718   32. Gu, Y. COVID-19 Projections Using Machine Learning. *COVID-19 Projections Using Machine*
719   *Learning* https://covid19-projections.com/.

720   33. Stadlbauer, D. *et al.* Seroconversion of a city: Longitudinal monitoring of SARS-CoV-2
721   seroprevalence in New York City. *medRxiv* 2020.06.28.20142190 (2020)
722   doi:10.1101/2020.06.28.20142190.

723   34. Goodnough, A. & Shear, M. D. The U.S.'s Slow Start to Coronavirus Testing: A Timeline. *The*
724   *New York Times* (2020).

725   35. Shear, M. D. *et al.* The Lost Month: How a Failure to Test Blinded the U.S. to Covid-19. *The*
726   *New York Times* (2020).

727   36. Kaplan, S. & Thomas, K. Despite Promises, Testing Delays Leave Americans 'Flying Blind'.
728   *The New York Times* (2020).

37. de Souza, W. M. *et al.* Epidemiological and clinical characteristics of the COVID-19 epidemic

in Brazil. *Nature Human Behaviour* **4**, 856–865 (2020).

38. Mervosh, S. & Fernandez, M. 'It's Like Having No Testing': Coronavirus Test Results Are Still

Delayed. *The New York Times* (2020).

39. Gostic, K., Gomez, A. C., Mummah, R. O., Kucharski, A. J. & Lloyd-Smith, J. O. Estimated

effectiveness of symptom and risk screening to prevent the spread of COVID-19. *eLife* **9**,

e55570 (2020).

40. Zitek, T. The Appropriate Use of Testing for COVID-19. *West J Emerg Med* **21**, 470–472

(2020).

41. Bosch, I. *et al.* Rapid antigen tests for dengue virus serotypes and Zika virus in patient serum.

Sci Transl Med **9**, (2017).

42. Massachusetts Department of Public Health. COVID-19 Response Reporting. *Mass.gov*

https://www.mass.gov/info-details/covid-19-response-reporting.

43. California Department of Public Health. COVID-19 Cases - California Open Data.

https://data.ca.gov/dataset/covid-19-cases.

44. California Department of Public Health. COVID-19 Hospital Data - California Open Data.

https://data.ca.gov/dataset/covid-19-hospital-data.

45. Department of Health and Human Hygiene. COVID-19 Daily Counts of Cases,

Hospitalizations, and Deaths | NYC Open Data. https://data.cityofnewyork.us/Health/COVID-

19-Daily-Counts-of-Cases-Hospitalizations-an/rc75-m7u3.

46. Sao Jose do Rio Preto Public Health Office. *COVID-19 Surveillance Data, Sao Jose do Rio*

Preto.

47. New York State Government. Daily Hospitalization Summary by Region. *New York Forward*

https://forward.ny.gov/daily-hospitalization-summary-region.

48. Massachusetts General Hospital Institute for Technology Assessment. COVID-19 Simulator

-            Methodology.   https://www.covid19sim.org/images/docs/COVID-

19_simulator_methodology_download_20200507.pdf.

49. CDC. Coronavirus Disease 2019 (COVID-19). *Centers for Disease Control and Prevention*

https://www.cdc.gov/coronavirus/2019-ncov/cases-updates/commercial-lab-surveys.html

(2020).

50. SciPy.org — SciPy.org. https://www.scipy.org/.

51. NumPy. https://numpy.org/.

52. Non-Linear Least-Squares Minimization and Curve-Fitting for Python — Non-Linear Least-

Squares Minimization and Curve-Fitting for Python. https://lmfit.github.io/lmfit-py/.

53. Matplotlib: Python plotting — Matplotlib 3.3.1 documentation. https://matplotlib.org/.

54. Plotly: The front-end for ML and data science models. https://plotly.com/.