

# Age-dependence of healthcare interventions for SARS-CoV-2 infection in Ontario, Canada

Irena Papst<sup>\*1</sup>, Michael Li<sup>2,3</sup>, David Champredon<sup>4</sup>, Benjamin M. Bolker<sup>2,5,6</sup>, Jonathan Dushoff<sup>2,5</sup>, and David J.D. Earn<sup>5,6,7</sup>

<sup>1</sup>Center for Applied Mathematics, Cornell University

<sup>2</sup>Department of Biology, McMaster University

<sup>3</sup>South African Centre for Epidemiological Modelling and Analysis, University of Stellenbosch

<sup>4</sup>Department of Pathology and Laboratory Medicine, Western University

<sup>5</sup>Michael G. DeGroote Institute for Infectious Disease Research, McMaster University

<sup>6</sup>Department of Mathematics & Statistics, McMaster University

<sup>7</sup>Department of Mathematics, University of Toronto

\*Corresponding author (email: [ip98@cornell.edu](mailto:ip98@cornell.edu))

Compiled on 2020-09-01 at 23:55:50

## Abstract

**Background:** Patient age is the most salient clinical indicator of risk from COVID-19. Age-specific distributions of known SARS-CoV-2 infections and COVID-19-related deaths are available for most countries. However, relatively little attention has been given to the age distributions of hospitalizations and serious healthcare interventions administered to COVID-19 patients. We examined these distributions in Ontario, Canada, in order to quantify the age-related impacts of COVID-19, and to identify potential risks should the healthcare system become overwhelmed with COVID-19 patients in the future.

**Methods:** We analysed known SARS-CoV-2 infection records from the integrated Public Health Information System (iPHIS) and the Toronto Public Health Coronavirus Rapid Entry System (CORES) between 23 January 2020 and 17 June 2020 (N = 30,546), and estimated the age distributions of hospitalizations, ICU admissions, intubations, and ventilations. We quantified the probability of hospitalization given known SARS-CoV-2 infection, and of survival given COVID-19-related hospitalization.

**Results:** The distribution of COVID-19-related hospitalizations peaks with a wide plateau covering ages 54-90, whereas deaths are sharply concentrated in very old ages, with a maximum at age 90. The estimated probability of hospitalization given known SARS-CoV-2 infection reaches a maximum of 32.0% at age 75 (95% CI 27.5%-36.7%). The probability of survival given COVID-19-related hospitalization is uncertain for children (due to small sample size), and near 100% for adults younger than 40. After age 40, survival of hospitalized COVID-19 patients declines substantially; for example, a hospitalized 50-year-old patient has a 90.4% chance of surviving COVID-19 (95% CI 81.9%-95.7%).

**Interpretation:** Concerted efforts to control the spread of SARS-CoV-2 have kept prevalence of the virus low in the population of Ontario. The healthcare system has not been overstretched, yet the probability of survival given hospitalization for COVID-19 has been lower than is generally recognized for patients over 40. If prevalence of the virus were to increase and healthcare capacities were to be exceeded, survival of individuals in the broad age range requiring acute care would be expected to decrease, potentially expanding the distribution of COVID-19-related deaths toward younger ages.

## 39 Introduction

40 Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) was first confirmed in Ontario, Canada,  
41 on 23 January 2020, in an individual with a recent travel history to Wuhan, Hubei Province, China [1], the  
42 site of the first large-scale outbreak of this novel pathogen [2]. The virus was detected sporadically in  
43 Ontario through February [3], until the number of known infections (KIs) began to rise consistently in March.  
44 The province declared a state of emergency on 17 March 2020 [4], implementing a large-scale economic  
45 shutdown, as well as school closures, to help mitigate the spread of the virus.

46 A major concern associated with the spread of SARS-CoV-2 is the potential for hospitals to become  
47 overwhelmed with COVID-19, the disease caused by the virus, which would compromise care of these  
48 patients, and reduce access to and quality of care for many other illnesses. Severity of COVID-19 presentation  
49 is highly variable among individuals, but is thought to increase with age [5–7]. Deaths attributed to COVID-19  
50 have been found to be strongly concentrated in the elderly [8–10]. Comparatively few studies have explored  
51 the age distribution of serious medical interventions administered to COVID-19 patients [11, 12].

52 In order to understand the potential for the healthcare system to become overwhelmed by SARS-CoV-  
53 2 infections, it is important to identify the demographic groups that are most likely to require significant  
54 medical care. The goal of this paper is to quantify the relationships between COVID-19 patient age and  
55 the administration of serious medical interventions (hospitalizations, Intensive Care Unit (ICU) admissions,  
56 intubations, and ventilations) for the province of Ontario. We compare these age-intervention associations  
57 with the age distributions of KIs and deaths. We also estimate the age-specific probability of hospitalization  
58 given known SARS-CoV-2 infection, and of survival given hospitalization related to COVID-19.

## 59 Methods

60 We use individual-level line lists for known SARS-CoV-2 infections reported up to 17 June 2020 from the  
61 integrated Public Health Information System (iPHIS), maintained by Public Health Ontario, and from the  
62 Toronto Public Health Coronavirus Rapid Entry System (CORES). We include only KIs marked as “resolved”  
63 or “fatal” in the database, to avoid tallying patients whose final outcomes are not yet known. KIs are marked  
64 as “resolved” based on public health unit assessment. In all instances, a record is considered resolved if it is  
65 14 days past the symptom onset date (or specimen collection if symptom onset date is not known), though  
66 public health occasionally performs additional follow-up to update records. For brevity, we use “resolved KIs”  
67 in our analysis to refer to KIs marked as either “resolved” or “fatal” in iPHIS and CORES.

68 These data contain 30,546 records of resolved KIs, which, as of this writing on 1 September 2020, comprises  
69 74% of resolved and fatal KIs accounted for in the publicly-available data [13].

70 These data are also representative of a situation where the healthcare system was not overwhelmed with  
71 COVID-19 patients (due to widespread measures to keep SARS-CoV-2 spread under control). ICU occupancy  
72 for COVID-19 treatment in Ontario never exceeded more than 13% of the total ICU capacity over the period  
73 covered by the data. We calculate this percentage using daily ICU occupancy from publicly-available data  
74 [14] and the number of ICU (critical care) beds available according to the Government of Ontario [15]. Based  
75 on the latter source, we assume the province had a total of 2012 ICU beds before 16 April 2020, and this  
76 capacity increased to 3504 beds on 16 April 2020. We acknowledge that this is a simplification of the ICU  
77 expansion effort and that not all ICU beds are allocated to COVID-19 treatment (although this allocation is  
78 flexible depending on demand).


79 We use 2020 Ontario population projections produced by Statistics Canada [16], specifically projections from  
80 the “M1” medium-growth scenario [17], although all scenarios yield virtually identical projections for the short  
81 time horizon of 2020.

82 We use provincial SARS-CoV-2 testing data from the Ontario Laboratories Information System (OLIS)  
83 database, courtesy of the Institute for Clinical Evaluative Sciences (ICES). This central database records all  
84 tests for active SARS-CoV-2 infection in the province. These data are based on reports up to 9 July 2020  
85 and include 39,598 positive tests and 1,086,892 negative tests.

86 We aggregate age-specific data into two-year age bins (and one wide bin for individuals over 100 years old).  
87 We use a binomial generalized additive model to estimate the age-specific probability of hospitalization  
88 given KI [18], and a binomial generalized linear model to estimate the age-specific survival probability given  
89 hospitalization [19]. Data and source code for all analyses will be provided in an Open Science Framework  
90 repository at a later date.

## 91 Results

92 Figure 1 shows the time course of the SARS-CoV-2 epidemic in Ontario up to 17 June 2020, as represented  
93 by KIs.




**Figure 1:** Known infections (KIs) over time in Ontario, split by whether the KI was resolved (marked as “resolved” or “fatal” in iPHIS and CORES) by 17 June 2020 (see [Methods](#)). Dashed vertical lines mark important dates for the outbreak in the province. The shaded region indicates a 14-day reporting delay from the last record on 17 June 2020.

94 Figure 2 shows the age structure of resolved KIs, population demographics, and SARS-CoV-2 infection  
95 tests. The pattern observed in the raw counts of resolved KIs (panel A) reflects underlying demographics  
96 (panel B) as well as the per capita rate of positive tests, which is in turn related to testing intensity (panel  
97 D). Testing intensity is defined as the number of tests administered per 10,000 population by age group.  
98 Controlling for demography, the number of detected infections (panel C) is relatively low in ages under 20,  
99 increases to a plateau for ages 20-70, and then continually increases after age 70. In ages 15 and under, the  
100 testing intensity is approximately 6.5 times smaller than that of all other ages 75 and under. Testing intensity  
101 increases after age 75.


102 Figure 3 shows the distribution of ages for serious medical interventions (panel A) and deaths (panel B)  
103 related to COVID-19 for resolved KIs. We present raw counts, as opposed to counts normalized by the  
104 age-specific population, because the counts (not per capita rates) determine the pressure on the healthcare  
105 system. Hospitalizations are faceted by the most intensive known intervention (with ventilator use being the  
106 most intensive, followed by intubation, then ICU admission, and hospitalization). Deaths are split by whether  
107 or not the patient has a record of hospitalization for COVID-19 treatment.

108 The distribution of serious medical interventions is much wider than that of deaths, with the latter peaking  
109 at age 90. Hospitalizations are relatively uniformly spread between ages 54-90, while the distribution of  
110 ICU-related interventions (ICU admission, intubation, ventilation) is spread over a slightly younger age range.  
111 The majority of deaths, 60.1% (95% CI 58.2%-62.0%), have occurred in KIs where there is no record of  
112 hospitalization for treatment related to COVID-19.


113 Figure 4 shows the estimated hospitalization probability given known SARS-CoV-2 infection (panel A) and  
114 survival probability given hospitalization for COVID-19 treatment (panel B). There is large uncertainty in both  
115 probability estimates for young and very old age groups due to small numbers of KIs and hospitalizations in



**Figure 2:** Age distribution of known infections (KIs) in Ontario. The distribution of ages for resolved KIs (panel A), Ontario population projections for 2020 (panel B), resolved KIs per 10,000 population (panel C), and SARS-CoV-2 infection tests per 10,000 population (panel D). The y-axes in panels C and D are on a logarithmic scale.



**Figure 3:** COVID-19 outcomes by age in Ontario. The distribution of ages for hospital interventions (panel A), and deaths (panel B). Hospital outcomes are nested and tallied by the most intensive medical intervention used for each patient (ventilator use is the most intensive, followed by intubation, ICU admission, and hospitalization). Deaths are faceted by whether or not the patient also had a record of hospitalization for COVID-19 treatment.



**Figure 4:** Age-dependent COVID-19 hospitalization probability for known SARS-CoV-2 infection (panel A) and survival probability for hospitalized patients (panel B) in Ontario. We estimate the hospitalization probability using a generalized additive model and the survival probability using a generalized linear model (blue curves; see [Methods](#)). Vertical lines give age-specific 95% binomial confidence intervals and point areas are proportional to samples sizes.

116 these ages. The hospitalization probability peaks in the 74-75 age group at 32.0% (95% CI 27.5%-36.7%).  
117 In adults, the survival probability is near 100% until about age 40, where it begins to decline steadily. For  
118 instance, a hospitalized individual in the age group 50-51 has a 90.4% chance of surviving COVID-19 (95%  
119 CI 81.9%-95.7%), implying that roughly 1 in 10 hospitalized COVID-19 patients in this age group die despite  
120 receiving medical care while ICUs and hospitals are below capacity.

## 121 Interpretation

### 122 Summary

123 The age distribution of deaths attributed to COVID-19 in Ontario (Figure 3B) provides limited insight into the  
124 risk that COVID-19 patients could overwhelm Ontario's healthcare system, especially since many deaths to  
125 date have occurred in residential homes, which are independent of the hospital system [20]. The broad age  
126 distributions of hospitalizations, ICU admissions, intubation, and ventilation (Figure 3A), reveal the potential  
127 pressure on the healthcare system from both middle-aged individuals and seniors, if SARS-CoV-2 prevalence  
128 were to increase significantly in the population. Based on known infections (KIs) up to 17 June 2020, the  
129 probability of hospitalization given KI increases with age before peaking at ages 74-75, and then declining.  
130 The probability of survival given hospitalization for COVID-19 treatment declines with age, and decreases  
131 substantially after age 40.

### 132 Explanation of findings

133 Numbers of KIs and testing intensity were lowest in ages younger than 15 (Figure 2), which could reflect  
134 poor detection in these ages, lower age-specific prevalence, or both. We discuss this issue further in the  
135 Appendix. Relatively high numbers of resolved KIs and testing intensity in those over age 75 can be partly  
136 attributed to significant outbreaks in long-term care facilities [20, 21].

137 Early in the outbreak, the province of Ontario expanded coverage for COVID-19-related treatment to include  
138 even individuals who are not usually covered by the Ontario Health Insurance Plan [22]. Access to prompt  
139 and successful medical interventions may have kept a large proportion of COVID-19-related hospitalizations  
140 from resulting in deaths. The majority of COVID-19-related deaths have occurred in patients with no record of  
141 hospitalization (Figure 3B). While we expect that the shape of the age distribution of *need* for hospitalization  
142 (Figure 3A) would remain the same if prevalence were to increase, the distribution of deaths may expand  
143 toward younger ages if hospitals and ICUs reach maximum capacity.

144 The age-dependent hospitalization probabilities that we report (Figure 4A) are likely overestimates, because  
145 they are based on resolved KIs, not all infections; these probabilities depend on how widely SARS-CoV-2  
146 testing has been conducted. Throughout most of the epidemic, testing guidelines selected for sufficiently  
147 symptomatic individuals [23]; testing guidelines were not expanded to include asymptomatic individuals until  
148 29 May 2020 [24]. As testing efforts continue to grow, hospitalization probability estimates can be expected  
149 to decrease, as relatively more asymptomatic and pre-symptomatic individuals are likely to be recorded as  
150 KIs.

151 Our survival probability estimates for hospitalized individuals (Figure 4B) are *not* affected by the same  
152 detection biases present in KI data. Patients admitted to hospital are tested for SARS-CoV-2 as part of  
153 infection control protocols, and thus infection detection in hospitalized individuals is not influenced by testing  
154 guidelines for the general population. Our survival probability estimates do, however, represent an upper  
155 bound with respect to the current standard of care and viral variant. In the absence of significant innovation  
156 in COVID-19 treatment or viral evolution to lower disease severity, we expect survival probabilities would  
157 decrease if ICUs or hospitals were to reach maximum capacity.

### 158 Future directions

159 We have quantified the age distributions of serious medical interventions for SARS-CoV-2 infection in Ontario,  
160 Canada, until 17 June 2020, when the province was first responding to the novel pathogen. The Government  
161 of Canada and the Province of Ontario implemented policies meant to help mitigate SARS-CoV-2 spread  
162 over this period [25]. Future work should consider whether the age dependence of SARS-CoV-2 infection

163 risks is changing over time, as the population adjusts to the pandemic and the testing effort expands. Our  
164 study explores only short-term SARS-CoV-2 infection outcomes; future studies should explore the age  
165 distributions of long-term morbidities from this infection, so that we may better understand the heterogeneous  
166 risks associated with COVID-19. Lastly, all studies relying on known infection counts are subject to bias from  
167 how infections are detected via active infection testing, which may disproportionately affect counts in youth  
168 (see [Appendix](#)). Future work should seek to correct for this bias.

## 169 **Limitations**

170 Our analysis excludes a small proportion of Ontario population because we were unable to access known  
171 infection records from the Ottawa Public Health COVID-19 Ottawa Database and the Middlesex-London  
172 COVID-19 Case and Contact Management tool. In general, the number of KIs underestimates the true  
173 prevalence of SARS-CoV-2 infection in the general population, most notably because provincial testing  
174 guidelines selecting for symptomatic individuals induced bias in these data and the age-specific asymptomatic  
175 proportion is not yet known. The results presented in this study pertaining to children and teenagers are  
176 particularly limited due to undersampling in those age groups, revealed by the testing intensity data, and  
177 should be interpreted with caution. Our study covers a period of extensive economic shutdown; as the  
178 province continues reopening, we expect the age-specific contact structure to change and contact rates to  
179 generally increase, which will affect observed patterns in the age distributions of KIs and deaths.

## 180 **Conclusion**

181 Although the age distribution of COVID-19 deaths in Ontario has been strongly skewed to the elderly  
182 (peaking at 90 years old), the age distribution of COVID-19 hospitalizations is much broader, highlighting  
183 potentially serious risks to public health and the healthcare system in an uncontrolled outbreak of SARS-  
184 CoV-2. Measures that control disease spread and maintain hospital occupancy below capacity will continue  
185 to be important in the absence of improved treatments or vaccination for SARS-CoV-2 infection.

## 186 **Acknowledgments**

187 We thank Sarah Morrison for assistance with literature review. We are grateful to the Ontario COVID-19  
188 Modelling Table for valuable discussions. Data were kindly provided by Public Health Ontario.

189 This study was supported by the Institute for Clinical Evaluative Sciences (ICES) which is funded by the  
190 Ontario Ministry of Health and Long-Term Care (MOHLTC). The opinions, results and conclusions are those  
191 of the authors and are independent from the funding source. No endorsement by ICES or the Ontario  
192 MOHLTC is intended or should be inferred.

193 This study was supported by the Ontario Health Data Platform (OHDP), a Province of Ontario initiative  
194 to support Ontario's ongoing response to COVID-19 and its related impacts. The opinions, results and  
195 conclusions reported in this paper are those of the authors and are independent from the funding sources.  
196 No endorsement by the OHDP, its partners, or the Province of Ontario is intended or should be inferred.

## 197 **Funding statement**

198 DJDE, BMB, and JD were funded by the Natural Sciences and Engineering Research Council of Canada,  
199 the Michael G. DeGroot Institute for Infectious Disease Research at McMaster University, and the Public  
200 Health Agency of Canada.

## 201 **Competing interests**

202 The authors declare no competing interests.



## 203 Appendix

### 204 Detection of and exposure to SARS-CoV-2 in ages under 20

205 For the majority of the period covered by the data in this study, testing guidelines in Ontario selected for  
206 sufficiently symptomatic individuals [23]; it was not until 29 May 2020 that Ontario expanded testing guidelines  
207 to include asymptomatic individuals [24]. Children infected with SARS-CoV-2 may be asymptomatic or have  
208 milder clinical presentations of COVID-19 than adults [26, 27]. Moreover, the symptom profile of pediatric  
209 SARS-CoV-2 infection differs from that of adults [26–29] and may have been poorly represented in screening  
210 guidelines and public awareness campaigns earlier in the pandemic. It is also possible that once the first  
211 KI was established in a household, the rest of the household did not get tested (even in the presence of  
212 symptoms meeting testing criteria) as it was already known that the pathogen was present in close contacts  
213 and testing capacity was initially limited. Since children are not often identified as the index case in a  
214 household SARS-CoV-2 outbreak [30], this effect may disproportionately undercount SARS-CoV-2 infection  
215 in children and teenagers.

216 It is also possible that Ontario youth are simply not being infected with SARS-CoV-2 as frequently compared  
217 to other age groups [6, 30, 31]. However, any studies making this claim by appealing to data based on active  
218 infections in a diagnostic testing scheme are potentially subject to the detection biases outlined above. Well-  
219 timed wide-spread school closures in Ontario may have curtailed SARS-CoV-2 spread in youth [32], though  
220 it is difficult to disentangle the specific effect of school closures from that of all other non-pharmaceutical  
221 interventions undertaken simultaneously [33].

222 Serological studies may help determine the true extent of exposure to SARS-CoV-2 in all ages, but especially  
223 in those under 20, which is particularly important given recent findings of high viral loads in children [34] and  
224 that those between ages 10 and 19 transmit the virus at least as effectively as adults [35]. Public Health  
225 Ontario conducted an age-specific seroprevalence study for the period of 27 Mar 2020 to 30 Jun 2020 [36],  
226 though small sample sizes prevented particularly reliable estimates in youth. A recent, large-scale antibody  
227 study in New York City revealed that the 0-17 age group had the highest antibody test positivity rate [37]  
228 despite having the lowest number of known infections per 100,000 population [38]. While these results may  
229 not be directly transferable to Ontario given the difference in outbreak scale, they suggest that low counts of  
230 KIs in youth may not necessarily imply low age-specific prevalence.

## 231 References

- 232 [1] Ontario confirms first case of Wuhan novel coronavirus, [https://news.ontario.ca/mohltc/en/2020/01/ontario-](https://news.ontario.ca/mohltc/en/2020/01/ontario-confirms-first-case-of-wuhan-novel-coronavirus.html)  
233 [confirms-first-case-of-wuhan-novel-coronavirus.html](https://news.ontario.ca/mohltc/en/2020/01/ontario-confirms-first-case-of-wuhan-novel-coronavirus.html) (2020).
- 234 [2] Novel coronavirus (2019-nCoV) situation report — 1, [https://www.who.int/docs/default-source/](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200121-sitrep-1-2019-ncov.pdf)  
235 [coronaviruse/situation-reports/20200121-sitrep-1-2019-ncov.pdf](https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200121-sitrep-1-2019-ncov.pdf) (2020).
- 236 [3] COVID-19 in Ontario: January 15, 2020 to July 29, 2020 (Daily Epidemiologic Summary),  
237 [https://www.publichealthontario.ca/-/media/documents/ncov/epi/2020/covid-19-daily-epi-summary-](https://www.publichealthontario.ca/-/media/documents/ncov/epi/2020/covid-19-daily-epi-summary-report.pdf?la=en)  
238 [report.pdf?la=en](https://www.publichealthontario.ca/-/media/documents/ncov/epi/2020/covid-19-daily-epi-summary-report.pdf?la=en) (2020).
- 239 [4] Davidson S. Ontario declares state of emergency amid COVID-19 pandemic, [https://toronto.ctvnews.ca/](https://toronto.ctvnews.ca/ontario-declares-state-of-emergency-amid-covid-19-pandemic-1.4856033)  
240 [ontario-declares-state-of-emergency-amid-covid-19-pandemic-1.4856033](https://toronto.ctvnews.ca/ontario-declares-state-of-emergency-amid-covid-19-pandemic-1.4856033) (2020).
- 241 [5] Wu Z, McGoogan J. Characteristics of and important lessons from the coronavirus disease 2019 (COVID-  
242 19) outbreak in China: Summary of a report of 72 314 cases from the Chinese Center for Disease Control  
243 and Prevention. *JAMA*. DOI: [10.1001/jama.2020.2648](https://doi.org/10.1001/jama.2020.2648). (2020).
- 244 [6] Davies NG, Klepac P, Liu Y, et al. Age-dependent effects in the transmission and control of COVID-19  
245 epidemics. *medRxiv*. DOI: [10.1101/2020.03.24.20043018](https://doi.org/10.1101/2020.03.24.20043018). (2020).
- 246 [7] Mizumoto K, Omori R, Nishiura H. Age specificity of cases and attack rate of novel coronavirus disease  
247 (COVID-19). *medRxiv*. DOI: [10.1101/2020.03.09.20033142](https://doi.org/10.1101/2020.03.09.20033142). (2020).

- 248 [8] Onder G, Rezza G, Brusaferro S. Case-fatality rate and characteristics of patients dying in relation to  
249 COVID-19 in Italy. *JAMA*. DOI: [10.1001/jama.2020.4683](https://doi.org/10.1001/jama.2020.4683). (2020).
- 250 [9] Zhang Y, The Novel Coronavirus Pneumonia Emergency Response Epidemiology Team. The epidemio-  
251 logical characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) — China, 2020. *China*  
252 *CDC Weekly* 2020; 2: 113. DOI: [10.46234/ccdcw2020.032](https://doi.org/10.46234/ccdcw2020.032). (2020).
- 253 [10] Medfod A, Trias-Llimos S. Population age structure only partially explains the large number of COVID-19  
254 deaths at the oldest ages. *medRxiv*. DOI: [10.1101/2020.04.09.20056341](https://doi.org/10.1101/2020.04.09.20056341). (2020).
- 255 [11] Myers L, Parodi S, Escobar G, et al. Characteristics of hospitalized adults with COVID-19 in an integrated  
256 health care system in California. *JAMA*. DOI: [10.1001/jama.2020.7202](https://doi.org/10.1001/jama.2020.7202). (2020).
- 257 [12] Zhao M, Wang M, Zhang J, et al. Comparison of clinical characteristics and outcomes of patients with  
258 coronavirus disease 2019 at different ages. *Ageing*; 12. DOI: [10.18632/aging.103298](https://doi.org/10.18632/aging.103298). (2020).
- 259 [13] Status of COVID-19 cases in Ontario (dataset), [https://data.ontario.ca/dataset/status-of-covid-19-cases-](https://data.ontario.ca/dataset/status-of-covid-19-cases-in-ontario)  
260 [in-ontario](https://data.ontario.ca/dataset/status-of-covid-19-cases-in-ontario) (accessed 1 September 2020).
- 261 [14] Li M. COVID19-Canada, <https://wzmli.github.io/COVID19-Canada/> (2020).
- 262 [15] Ontario significantly expands hospital capacity to prepare for any COVID-19 outbreak scenario,  
263 [https://www.news.ontario.ca/opo/en/2020/04/ontario-significantly-expands-hospital-capacity-to-prepare-](https://www.news.ontario.ca/opo/en/2020/04/ontario-significantly-expands-hospital-capacity-to-prepare-for-any-covid-19-outbreak-scenario.html)  
264 [for-any-covid-19-outbreak-scenario.html](https://www.news.ontario.ca/opo/en/2020/04/ontario-significantly-expands-hospital-capacity-to-prepare-for-any-covid-19-outbreak-scenario.html) (2020).
- 265 [16] Table 17-10-0057-01 Projected population, by projection scenario, age and sex, as of July 1 (x 1,000).  
266 DOI: [10.25318/1710005701-eng](https://doi.org/10.25318/1710005701-eng). (2019).
- 267 [17] Population projections for Canada (2018 to 2068), provinces and territories (2018 to 2043), [https:](https://www150.statcan.gc.ca/n1/pub/91-520-x/91-520-x2019001-eng.htm)  
268 [://www150.statcan.gc.ca/n1/pub/91-520-x/91-520-x2019001-eng.htm](https://www150.statcan.gc.ca/n1/pub/91-520-x/91-520-x2019001-eng.htm) (2019).
- 269 [18] Wood S. *Generalized additive models: An introduction with R*. 2nd ed. Chapman; Hall/CRC, 2017.
- 270 [19] R Core Team. *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation  
271 for Statistical Computing, <https://www.R-project.org/> (2019).
- 272 [20] Hsu A, Lane N, Sinha S, et al. Understanding the impact of COVID-19 on residents of Canada's  
273 long-term care homes—ongoing challenges and policy responses, [https://ltccovid.org/wp-content/uploads/](https://ltccovid.org/wp-content/uploads/2020/06/LTCCovid-country-reports_Canada_June-4-2020.pdf)  
274 [2020/06/LTCCovid-country-reports\\_Canada\\_June-4-2020.pdf](https://ltccovid.org/wp-content/uploads/2020/06/LTCCovid-country-reports_Canada_June-4-2020.pdf) (2020).
- 275 [21] Fisman DN, Bogoch I, Lapointe-Shaw L, et al. Risk factors associated with mortality among residents  
276 with Coronavirus Disease 2019 (COVID-19) in Long-term Care Facilities in Ontario, Canada. *JAMA Network*  
277 *Open*; 3. DOI: [10.1001/jamanetworkopen.2020.15957](https://doi.org/10.1001/jamanetworkopen.2020.15957). (2020).
- 278 [22] Ontario expands coverage for care, [https://news.ontario.ca/en/release/56401/ontario-expands-coverage-](https://news.ontario.ca/en/release/56401/ontario-expands-coverage-for-care)  
279 [for-care](https://news.ontario.ca/en/release/56401/ontario-expands-coverage-for-care) (2020).
- 280 [23] COVID-19 provincial testing guidance update, [http://www.health.gov.on.ca/en/pro/programs/publichealth/](http://www.health.gov.on.ca/en/pro/programs/publichealth/coronavirus/docs/2019_covid_testing_guidance.pdf)  
281 [coronavirus/docs/2019\\_covid\\_testing\\_guidance.pdf](http://www.health.gov.on.ca/en/pro/programs/publichealth/coronavirus/docs/2019_covid_testing_guidance.pdf) (2020).
- 282 [24] Ontario opens up COVID-19 testing across the province, [https://news.ontario.ca/opo/en/2020/05/ontario-](https://news.ontario.ca/opo/en/2020/05/ontario-opens-up-covid-19-testing-across-the-province.html)  
283 [opens-up-covid-19-testing-across-the-province.html](https://news.ontario.ca/opo/en/2020/05/ontario-opens-up-covid-19-testing-across-the-province.html) (2020).
- 284 [25] Vogel L. COVID-19: A timeline of Canada's first-wave response, [https://cmajnews.com/2020/06/12/](https://cmajnews.com/2020/06/12/coronavirus-1095847/)  
285 [coronavirus-1095847/](https://cmajnews.com/2020/06/12/coronavirus-1095847/) (2020).
- 286 [26] Hoang A, Chorath K, Moreira A, et al. COVID-19 in 7780 pediatric patients: A systematic review.  
287 *EClinicalMedicine*. DOI: [10.1016/j.eclim.2020.100433](https://doi.org/10.1016/j.eclim.2020.100433). (2020).
- 288 [27] Cui X, Zhao Z, Zhang T, et al. A systematic review and meta-analysis of children with Coronavirus  
289 Disease 2019 (COVID-19). *Journal of Medical Virology*. DOI: [10.1002/jmv.26398](https://doi.org/10.1002/jmv.26398). (2020).

- 290 [28] Xu Y, Li X, Zhu B, et al. Characteristics of pediatric SARS-CoV-2 infection and potential evidence for  
291 persistent fecal viral shedding. *Nature Medicine* 2020; 26: 502–505. DOI: [10.1038/s41591-020-0817-4](https://doi.org/10.1038/s41591-020-0817-4).  
292 (2020).
- 293 [29] Lu X, Zhang L, Du H, et al. SARS-CoV-2 infection in children. *New England Journal of Medicine* 2020;  
294 382: 1663–1665. DOI: [10.1056/NEJMc2005073](https://doi.org/10.1056/NEJMc2005073). (2020).
- 295 [30] Ludvigsson JF. Children are unlikely to be the main drivers of the COVID-19 pandemic—a systematic  
296 review. *Acta Paediatrica*. DOI: [10.1111/apa.15371](https://doi.org/10.1111/apa.15371). (2020).
- 297 [31] Lee P-I, Hu Y-L, Chen P-Y, et al. Are children less susceptible to COVID-19? *Journal of Microbiology,  
298 Immunology, and Infection*. DOI: [10.1016/j.jmii.2020.02.011](https://doi.org/10.1016/j.jmii.2020.02.011). (2020).
- 299 [32] Earn DJD, He D, Loeb MB, et al. Effects of school closure on incidence of pandemic influenza in Alberta,  
300 Canada. *Annals of Internal Medicine* 2012; 156: 173–181. DOI: [10.7326/0003-4819-156-3-201202070-  
301 00005](https://doi.org/10.7326/0003-4819-156-3-201202070-00005). (2012).
- 302 [33] Viner RM, Russell SJ, Croker H, et al. School closure and management practices during coronavirus  
303 outbreaks including COVID-19: A rapid systematic review. *The Lancet Child & Adolescent Health*. DOI:  
304 [10.1016/S2352-4642\(20\)30095-X](https://doi.org/10.1016/S2352-4642(20)30095-X). (2020).
- 305 [34] Yonker LM, Neilan AM, Bartsch Y, et al. Pediatric SARS-CoV-2: Clinical presentation, infectivity, and  
306 immune responses. *The Journal of Pediatrics*. DOI: [10.1016/j.jpeds.2020.08.037](https://doi.org/10.1016/j.jpeds.2020.08.037). (2020).
- 307 [35] Park YJ, Choe YJ, Park O, et al. Contact tracing during coronavirus disease outbreak, South Korea,  
308 2020. *Emerging Infectious Diseases*; 26. DOI: [10.3201/eid2610.201315](https://doi.org/10.3201/eid2610.201315).
- 309 [36] COVID-19 seroprevalence in Ontario: March 27, 2020 to June 30, 2020, [https://www.publichealthontario.ca/  
310 /media/documents/ncov/epi/2020/07/covid-19-epi-seroprevalence-in-ontario.pdf?la=en](https://www.publichealthontario.ca/-/media/documents/ncov/epi/2020/07/covid-19-epi-seroprevalence-in-ontario.pdf?la=en) (2020).
- 311 [37] Goldstein J. 1.5 million antibody tests show what parts of N.Y.C. were hit hardest, [https:  
312 //www.nytimes.com/2020/08/19/nyregion/new-york-city-antibody-test.html](https://www.nytimes.com/2020/08/19/nyregion/new-york-city-antibody-test.html) (2020).
- 313 [38] COVID-19: Data, <https://www1.nyc.gov/site/doh/covid/covid-19-data.page> (accessed 23 August 2020).