Computational Modeling Identifies Embolic Stroke of Undetermined

Source Patients with Potential Arrhythmic Substrate

Savannah F. Bifulco,[1] Griffin D. Scott,[1] Sakher Sarairah,[2] Zeinab Birjandian,[2,5]

Caroline H. Roney,[3] Steven A. Niederer,[3] Christian Mahnkopf,[4] Peter Kuhnlein,[4]

Marcel Mitlacher,[4] David Tirschwell,[5] W. T. Longstreth Jr,[5,6]

Nazem Akoum,[2] Patrick M. Boyle[1,7,8*]*

1: Department of Bioengineering, University of Washington, Seattle, WA, USA.

2: Division of Cardiology, University of Washington, Seattle, WA, USA.

3: School of Biomedical Engineering and Imaging Sciences, King's College London, London, UK.

4: Department of Cardiology, Klinikum Coburg, Coburg, Germany.

5: Department of Neurology, University of Washington, Seattle, WA, USA.

6: Department of Epidemiology, University of Washington, Seattle, WA, USA.

7: Institute for Stem Cell and Regenerative Medicine, University of Washington, Seattle, WA, USA.

8: Center for Cardiovascular Biology, University of Washington, Seattle, WA, USA.

*: Corresponding authors:

PM Boyle; email: pmjboyle@uw.edu; mail: Foege N310H UW Mailbox 355061, Seattle WA 98195, USA;
OR,
N Akoum; email: nakoum@cardiology.washington.edu; mail: 1959 N Pacific St, Seattle WA 98115, USA

Word count (main text): 6404

Abstract

Cardiac magnetic resonance imaging (MRI) has revealed fibrosis in embolic stroke of undetermined source (ESUS) patients comparable to levels seen in atrial fibrillation (AFib). We used computational modeling to understand the absence of arrhythmia in ESUS despite the presence of putatively pro-arrhythmic fibrosis. MRI-based atrial models were reconstructed for 45 ESUS and 45 AFib patients. The fibrotic substrate's arrhythmogenic capacity in each patient was assessed computationally. Reentrant drivers were induced in 24/45 (53%) ESUS and 22/45 (49%) AFib models. Inducible models had more fibrosis (16.7±5.45%) than non-inducible models (11.07±3.61%; P<0.0001); however, inducible subsets of ESUS and AFib models had similar fibrosis levels (P=0.90), meaning the intrinsic pro-arrhythmic substrate properties of fibrosis in ESUS and AFib are indistinguishable. This suggests some ESUS patients have latent pre-clinical fibrotic substrate that could be a future source of arrhythmogenicity. Thus, our work prompts the hypothesis that ESUS patients with fibrotic atria are spared from AFib due to an absence of arrhythmia triggers.

Introduction

Atrial fibrillation (AFib) is the most common cardiac arrhythmia, affecting 1-2% of the world's population and significantly contributing to worldwide morbidity and mortality (1). The primary source of AFib-related mortality is stroke, with around 20% of all ischemic strokes occurring in AFib patients (1). Sub-clinical AFib (i.e., transient, asymptomatic AFib) is implicated as a potential cause of embolic stroke of undetermined source (ESUS), and the current course of clinical care following ESUS is to look for evidence of AFib via an external monitor, an implanted loop recorder, or other forms of wearable monitoring devices. If AFib is diagnosed, treatment with oral anticoagulants is started to mitigate the possibility of recurrent stroke (2). Clinical studies have shown that AFib has been detected in only 30% of patients with long-term rhythm monitoring (3). This creates a frustrating problem for clinicians: in the wake of ESUS events, it is impossible to know which individuals should be treated as high-risk for AFib and therefore monitored accordingly.

Recent evidence from clinical studies suggests that the left atrial fibrosis burden measured

by late gadolinium enhanced (LGE)-MRI is as high in ESUS patients as in AFib patients without

stroke (4). This finding supports the hypothesis that atrial fibrosis is an element of the causal

pathway for stroke, through an atrial cardiopathy, and independent of AFib. The absence of AFib

despite the presence of a fibrotic substrate is intriguing and one potential explanation is that ESUS

patients have pro-arrhythmic fibrotic substrate but lack the triggers needed to initiate arrhythmia.

Another potential explanation is that the fibrosis present in ESUS patients is not pro-arrhythmic.

Patient-derived computational modeling of atrial arrhythmias is uniquely poised to test these

hypotheses. Previously, personalized atrial models have been used to assess arrhythmogenic

propensity of fibrotic substrate and predict AFib ablation targets (5, 6). Applying the same

approach, we can use computational models to predict if, in the presence of appropriate triggers,

fibrotic remodeling in ESUS has the fundamental capacity to harbor reentrant arrhythmic activity.

Thus, we present a large-scale computational study to ascertain whether the fibrotic

substrate with the potential to perpetuate AFib-sustaining reentrant drivers (RDs) exists in ESUS.

Our hypothesis is that a pre-clinical AFib substrate, attributed to a pattern of fibrotic atrial

remodeling that is conducive to RD perpetuation, exists in ESUS patients. By conducting

simulations in models derived from LGE-MRI, we can begin to understand potential pro-

arrhythmic properties of atrial fibrosis in ESUS patients. The study thus provides insights on the

role of atrial fibrosis as a pathophysiological nexus between AFib and stroke.

Results

Patient Characteristics

Ninety patient-derived models were included in our analysis: 45 post-stroke ESUS and 45

pre-ablation AFib patients. Demographic information about both patient groups is provided in

Table 1. There was a significant difference in LA surface area; however, the potential importance

of this feature for interpreting our findings is offset by the lack of difference in LA volume index (a

more commonly used measurement of normalized LA surface area), which suggests higher LA

surface area in those with AFib is a consequence of higher body mass index (BMI). LA fibrosis

burden was not significantly different between ESUS (13.6 ± 6.2%) and AFib patients (14.2 ±

4.5%) (P=0.91), consistent with previous findings (4).

Table 1. *Patient characteristics in ESUS and AFib groups.*

	ESUS (N=45)	AFib (N=45)	P value
Age, y	60±16	62±12	0.504
Female, %	44.0%	32.8%	0.275
BMI, kg/m^2	27.6±4.3	29.5±5.9	0.08
CHA$_2$DVASc score	2.0	1.9	0.345
CHF, n	14.3%	18.4%	0.599
Hypertension, n	68.5%	61.2%	0.468
Diabetes mellitus, n	20.4%	12.2%	0.292
CAD, n	18.4%	18.4%	1.000
Smoking, n	32%	28%	0.679
LA fibrosis, %	13.6±6.2%	14.2±4.5%	0.91
LA surface area, cm^2	109±26	134±40	0.0007
LA volume index, mL/m^2	60±29	57±26	0.607

Induction of arrhythmia and fibrosis quantification in patient-derived models

Personalized LA bilayer models were generated for all ESUS and AFib patients. Examples

of physiological detail incorporated in models can be seen in **Fig. 1**, including patient-specific

patterns of fibrotic remodeling, realistic atrial fiber orientations, and locations of electric pacing

sites; further detail can be found in the Methods section. Rapid electric stimulation caused RD-

sustained arrhythmia in 22 of 45 AFib models (48.8%) and 24 of 45 ESUS models (53.3%). Thus,

the capability of the fibrotic substrate to sustain RDs was not significantly different between the

two groups (P=0.83, **Fig. 2A**). ESUS and AFib models were then sorted by amount of global LA

fibrosis and arranged into quartiles. For five models (21.7%) in the first quartile (fibrosis<9.75%),

six models (28.5%) in the second quartile (fibrosis<12.6%), 15 (62.5%) models in the third quartile

(fibrosis<17. 5%), and 20 (91.0%) models in the top quartile, simulations revealed at least one

pacing site for which stimulation produced an episode of RD-sustained arrhythmia (**Fig. 2B**).

Figure 1: Model generation (A) Reconstruction of LA geometry with anatomical features labelled (RIPV/RSPV/LIPV/LSPV, right/left inferior/superior pulmonary veins; LAA, LA appendage). The LA is modeled as a bilayer comprising nested endocardial and epicardial shells linked in both fibrotic and non-fibrotic regions by 1D linear elements. (B) LA fiber orientations for the endocardium and epicardium, mapped from human atlas geometry as described in Methods. (C) AFib trigger sites as pacing sites (posterior/anterior LIPV, LSPV, RSPV, RIPV, LAA base, mitral valve annulus and posterior wall). (D) Regions of the LA generated as described in methods: (1) atrial floor, (2) anterior wall and LAA, (3) posterior wall, (4) left PVs, (5) right PVs.

Figure 2: Summary of patient-derived model fibrosis with respect to RD inducibility. (A) Histogram of AFib (22/45) and ESUS (24/45) inducible patients. Inducibility was not significantly different by χ^2 test. (B) Patients with ESUS and AFib arranged by percentage of LA fibrosis. Dotted lines indicate the quartiles of fibrosis observed for all 90 patient-derived models. Circles are indicative of stable reentry observed in the model from at least one pacing site after in silico pacing protocol. Triangles indicate no RDs after pacing from all 15 pacing sites independently. Cases that lacked RDs despite high fibrosis (inset I) or were inducible despite low fibrosis (inset II) are highlighted.

To explore potential pro-arrhythmic substrate properties in ESUS and AFib, we analyzed fibrosis burden in the sub-groups of each cohort in which RD-sustained arrhythmias were

inducible and non-inducible (**Fig. 3A**). Fibrosis burden was not significantly different between

inducible ESUS and AFib models with (**Fig. 3A**; P=0.90, confidence interval; CI: [-3.4, 4.1]) or

without induced reentry (**Fig. 3A**; P=1, CI: [-2.1, 2.4]). However, when fibrosis burdens for

inducible and non-inducible models were aggregated across ESUS and AFib groups, a significant

difference was evident (**Fig. 3A**; 16.7 ± 5.46% vs. 11.07 ± 3.61%; P<0.0001, CI: [3.4, 7.6]).

We also investigated potential differences in the number of unique RDs and the number

of AFib triggers that induced reentry in each patient-derived model. In all 46 inducible models, the

median number of unique RDs was 1 for both AFib and ESUS models (AFib range: 1-5; ESUS

range: 1-8). There was no significant difference in the number of unique reentrant morphologies

per model between the two groups (**Fig. 3B**; P=0.83, CI: [-7.7×10^{-5}, 8.8×10^{-5}]). The number of

unique RDs was positively correlated with atrial fibrosis burden (**Fig. 3C**; R=0.63, P<0.0001). For

all RD-inducible cases, the median number of stimulation sites from which rapid pacing led to RD

formation was 2 for AFib and 3 for ESUS models (range for both groups: 1-9). No significant

difference was found in the number of pacing sites that induced reentry per model between AFib

and ESUS (**Fig. 3D**; P=0.79, CI [-2.0,1.0]). The number of RD inducing pacing sites was also

significantly correlated with fibrosis burden (**Fig. 3E**; R=0.62, P<0.0001).

Figure 3: Summary of RD characteristics between ESUS and AFib models. (A) Boxplot of fibrosis
percentage in ESUS and AFib models where reentry was induced (ESUS: N=24, IQR=10.6; AFib: N=22,

IQR=5) and where reentry was not induced (ESUS: N=21, IQR=5; AFib N=23, IQR=3.5). Across ESUS and AFib models, fibrosis burden for RD inducible and RD non-inducible models was significantly different (P<0.0001). (B) Boxplot of number of unique reentrant morphologies elicited by all 15 pacing sites (P=0.83). (C) Correlation plot of fibrosis vs number of RDs (R=0.63, P<0.0001). (D) Boxplot of the number of pacing sites which induced reentry (P=0.79). (E) Correlation plot of fibrosis vs number of pacing sites that induced reentry (R=0.62, P<0.0001).

Arrhythmia Dynamics

Analysis of simulated reentry episodes revealed no qualitative differences in arrhythmia dynamics between AFib and ESUS models. **Figure 4** shows examples of RD-perpetuated *in silico* arrhythmia and instances where stimulation failed to induce reentry for both groups. Of note, this figure highlights two inducible low-fibrosis ESUS models (**Fig. 4A**: 6.9% fibrosis, RD near the LIPV; **Fig. 4B**: 10.0% fibrosis, RD on the atrial floor) and a non-inducible high-fibrosis ESUS model (**Fig. 4C**: 16% fibrosis). In the latter case, dense fibrosis on the posterior wall resulted in conduction block as indicated.

Figure 4: Maps of fibrotic tissue distribution (left) and activation time (right) for ESUS and AFib models in which pacing succeeded (rows 1-2) or failed (row 3) to induce RD-driven arrhythmia. Black arrows indicate

directions of wavefront propagation in RDs. Double lines indicate sites of conduction block. Black-shaded

regions in activation maps indicate locations where activation did not occur during the analysis interval. (A)

ESUS model with 6.9% fibrosis and reentry inferior to LIPV. (B) ESUS model with 10.0% fibrosis and reentry

on the atrial floor. (C) ESUS model with 16% fibrosis with wavefront termination through fibrosis on posterior

wall. (D) AFib model with 9.9% fibrosis and reentry observed adjacent to RIPV on posterior wall. (E) AFib

model with 13.7% fibrosis and reentry observed on the anterior wall. (F) AFib model with 11.6% fibrosis

with wavefront collision on posterior wall.

Figures 4D-E present examples of RD-driven arrhythmia in AFib models (9.9% and 13.7%

fibrosis, respectively), and **Fig. 4F** shows an AFib model (11.6% fibrosis) in which reentry was not

induced due to wavefront collision in the posterior wall region. Overall, this shows that both ESUS

and AFib models exhibited activation patterns consistent with previous definitions of RD-driven

arrhythmia; examples of inducible low-fibrosis and non-inducible high-fibrosis models emphasize

that fibrosis burden alone is an insufficient predictor for a potential arrhythmic substrate.

Properties of RD Localization

As described in Methods, each model was automatically subdivided into five anatomical

regions (see schematic illustrations in **Fig. 1D** and **Supplementary Fig. 1**) and region-wise

inducibility score analysis (IdS, as described in Methods) was used to gauge likelihood of RD

induction in response to rapid electrical stimulation from different locations in the LA. While ESUS

and AFib models had a statistically similar pattern of inducibility rates (P=0.45, by χ^2 test),

stimulation from the posterior wall was ≈2 times more likely to induce RDs in AFib models than

ESUS models (**Fig. 5A**, IdS = 6 vs IdS = 3). In other words, with all other factors held equal, our

simulations suggest that triggered activity in the posterior wall may be more likely to initiate

reentrant arrhythmia in AFib patients compared to ESUS patients. The same IdS values plotted

in **Fig. 5A** were mapped onto representative LA models to facilitate visual comparison of regional

sensitivity to rapid pacing (**Fig. 5B**). Next, we considered the number of unique RD localization

sites in each LA region across the different model groups (i.e., AFib vs. ESUS). The LPV region

was most likely to harbor RDs in the AFib cohort (**Fig. 5C**, N=17). In the ESUS cohort, the atrial

floor was the most likely region to contain an RD (**Fig. 5D**, N=16) The association between the

type of model (ESUS vs AFib) and RD localization was not significant (P=0.13, χ^2 test).

Figure 5: Summary of IdS and RD localization characteristics. (A) Region-wise IdS for both ESUS and
AFib LA models (B) Heat map of the regions in which triggers are most likely to induce arrhythmias depicted
as representative ESUS and AFib models. (C) Histogram of RDs across all AFib and ESUS models binned
by localization to specific LA regions. (D) Heat map of regions in which RDs are most likely to localize
depicted as representative ESUS and AFib models.

In addition to its role as part of the substrate for reentrant arrhythmia, fibrosis may directly

lead to increased AFib trigger incidence via calcium dysregulation leading to localized to regions

of depolarized resting potential (7, 8). To investigate whether cohort-scale differences in this

intrinsic pro-trigger property of fibrosis may explain the lack of arrhythmia in ESUS, we ran

additional simulations to assess the total extent of abnormally depolarized tissue (see Methods

for definition) in ESUS and AFib models; notably, these values are distinct from total fibrosis

burdens, since non-fibrotic tissue can be pulled to a more positive resting potential via electrotonic

coupling. We found no significant difference in these values between ESUS and AFib models

(P=0.32; CI: [–0.007, 0.019]; **Supplemental Fig. 2**); if anything, there was a trend towards *more*

trigger-prone tissue in ESUS. These findings provide preliminary evidence against the notion that

the lack of arrhythmia in ESUS might be due to lower rates of fibrosis-related ectopic pacemaking.

Figure 6: Summary of the region-by-region extent of tissue with a spatial fibrosis pattern (as characterized

by local density and entropy) associated with RD localization (i.e., pro-RD tissue). (A) Maps of pro-RD

tissue for representative AFib and ESUS cases, including boundaries between regions. (B) Region-wise

extent of pro-RD tissue in inducible AFib and ESUS models, depicted as boxplots (**: P < 0.01, Wilcoxon

rank sum test; CI: [–0.149, –0.021]).

Properties of pro-RD Fibrosis

To further examine trends in RD inducibility and localization observed in **Fig. 5**, we also

carried out region-wise analysis of fibrosis spatial pattern. In a previous study (5), we used

machine learning to quantitatively characterize the fibrosis distribution in regions that most

frequently harbored RD organizing centers (i.e., high local fibrosis density *and* entropy). As

described in Methods, the distribution of pro-RD tissue regions was determined for each inducible

model; these values were then subdivided in the five-region schematic (**Fig. 6A**). This analysis

revealed that the extent of pro-RD tissue in the LPV region was significantly higher in AFib

compared to ESUS models (**Fig. 6B**; P=0.01); this was consistent with our findings on regional

inducibility, and RD localization as shown in **Fig. 5**. Likewise, for regions that were associated

with greater inducibility and RD localization in ESUS models (anterior wall + LAA, atrial floor) there

was a trend towards a larger extent of pro-RD tissue compared to AFib models, but these did not

reach the level of significance.

Discussion

This study used a novel computational modeling approach with stimulus locations chosen

based on clinically observed AFib triggers to shed new light on the role of the fibrotic atrial

substrate in the potential for initiation and perpetuation of reentry in ESUS patients. In models

reconstructed from 45 post-stroke ESUS and 45 pre-ablation AFib patients, we showed that the

AFib and ESUS groups did not differ significantly in: (i) the propensity of the fibrotic substrate to

sustain RDs in response to simulated burst pacing; (ii) the LA fibrotic burden of RD-inducible

models or RD-free models; and (iii) the reentrant driver localization or the region-wise inducibility.

One noteworthy difference was that the extent of tissue in the LPV region with a pro-RD fibrosis

spatial pattern was greater in AFib vs. ESUS models. This is the first study to use computational

modeling and simulation to assess potential pro-arrhythmic capacity of LA fibrosis in ESUS

patients. Moreover, to the best of our knowledge, this is the largest cohort ever studied via

computational analysis of atrial electrophysiology in models derived from LGE-MRI, exceeding

the number of patient-specific models (N=50) in the former largest study (9) by a factor of ≈1.8.

Inducibility of reentry and fibrosis quantification in patient-derived atrial models

Experimental findings have shown that atrial fibrosis results in changes that promote reentry (10, 11), but the exact mechanism of this connection is not fully understood. Previous modeling studies have linked RD localization to specific spatial patterns of fibrotic remodeling in AFib (5, 12). Recent clinical findings indicate that atrial fibrosis burden does not differ significantly between AFib and ESUS patients, and yet (by definition) ESUS patients do not demonstrate AFib at the time of stroke or during ambulatory monitoring (4). Given the findings summarized above, a potential explanation is that, notwithstanding the fact that ESUS patients have substantial fibrosis, the particular *spatial distribution* of fibrotic remodeling in their atria is not conducive to arrhythmia perpetuation. Our findings suggest that this is likely not the case. From the standpoint of computational models derived from patient LGE-MRI scans, fibrotic substrate in individuals with ESUS is indistinguishable from that in patients with AFib in terms of the fundamental capacity to sustain RDs. For inducible models from both cohorts, we found that high fibrosis models were more likely to exhibit RDs irrespective of whether they corresponded to ESUS or AFib patients. This is in agreement with prior computational studies, which found the same general association (5, 6), and is consistent with clinical understanding, wherein higher fibrosis burden is associated with poor outcomes in AFib ablation procedures (13). Notably, we did observe several cases in which models defied inducibility expectations based on fibrosis alone. Such models exhibited RDs despite low fibrosis or were non-inducible despite high fibrosis. This observation confirms that, as observed previously in analogous AFib modeling studies (5, 6), assessment of raw fibrosis burden LGE-MRI scans alone is insufficient to fully characterize arrhythmogenic capacity of potentially pro-arrhythmic substrate in ESUS. In the present study, this is confirmed by our region-wise analysis of fibrosis spatial pattern (i.e., density, entropy), which is discussed below.

Consistent with the goal of this research to understand the contribution of fibrotic substrate to potential RD formation in ESUS, we purposefully excluded arrhythmias perpetuated by other

mechanisms from our study design (e.g., self-sustaining activity driven solely by focal sources).

This allows us to study the fibrotic substrate in the absence of all other confounding factors.

Additionally, our analysis of abnormal steady-state depolarization in LA models suggests the

potential for fibrosis itself to serves as a source of arrhythmia triggers is no greater in AFib than

in ESUS models. Potential contributions from the right atrium (RA) were also excluded, since only

the LA was segmented from LGE-MRI as part of the clinical workflow. Either of these factors may

explain the absence of simulated arrhythmia in 23 of 45 AFib models, many of which had very

little LA fibrosis (i.e., AFib in these individuals might have been predominantly focal in nature or

sustained by RDs in the RA). This rate of inducibility is consistent with previous studies (e.g.,13

out of 20 in computational models reconstructed from LGE-MRI scans of persistent AFib patients)

(5), supporting the notion that LA fibrosis is associated with increased arrhythmia inducibility but

fails to tell the whole story. Importantly, neither of these model constraints repudiates the central

finding of our study, which suggests there is no difference between ESUS and AFib patients in

terms of the fundamental capacity of the fibrotic substrate to potentially harbor RDs.

Reentrant driver localization dynamics and morphology in patient-derived atrial models

As discussed above, our qualitative findings suggest that ESUS patients' fibrotic substrate

is no different than that of AFib patients in terms the capacity to sustain RDs per se. We performed

additional analysis to assess whether specific consequences of fibrotic remodeling influenced any

characteristics of RD inducibility in different ways for simulations in models corresponding to

ESUS vs. AFib patients. First, we found that there was no significant difference in global fibrosis

burden between inducible AFib and ESUS models. Thus, our computational modeling suggests

that intrinsic pro-arrhythmic traits of ESUS and AFib fibrotic substrate are indistinguishable. Given

this result, one potential alternative explanation for the lack of arrhythmia in ESUS patients is a

lack of suitable triggers, despite an abundance of fibrotic substrate on par with that observed in

AFib. The plausibility of this explanation is strengthened by the fact that the pacing sites from

which episodes of reentry were induced in our study were based on common AFib trigger sites

as identified in a recent clinical study (14); this is in contrast to previous modeling studies, which

simulated triggered activity from evenly-distributed atrial sites (6). Nevertheless, the premise of

this aspect of our work was not to assess an absence of triggers, but instead ask the following: if

the atria of these ESUS patients were subjected to the same type of triggered activity known to

occur in typical AFib patients, is it possible the result be would sustained arrhythmia? In more

than half of the cohort (24/45), our analysis suggests that the answer is yes. This new hypothesis

could be validated in future studies by designing clinical protocols that systematically monitor

ESUS patients with different levels of fibrosis for potential AFib triggers via electrocardiographic

readouts.

Further analysis was performed to probe potential differences between ESUS and AFib

simulations that went beyond consideration of RD inducibility as a binary variable. Specifically,

we found no difference in the number of unique model-predicted RDs or the number of pacing

sites that induced RDs between ESUS and AFib. Instead, these variables were highly correlated

with LA fibrosis burden, which is consistent with the concept that high fibrosis models are more

susceptible to pacing-induced RDs. This finding further substantiates our principal claim that no

significant differences exist between the detected fibrotic substrate in ESUS and AFib, in that it

holds true for the substrate's capacity to sustain reentry and its susceptibility to triggered activity.

The general implication is that in the presence of simulated triggered activity, both of these

characteristics are closely linked to global fibrosis burden.

RDs identified by non-invasive electrocardiographic imaging (ECGI) and *in silico* phase

singularity identification have been shown to co-localize with fibrosis boundary zones identified

by LGE-MRI (5, 15-17). RD localization dynamics in this study were consistent with these findings,

as illustrated by representative LA fibrotic tissue distributions and corresponding RDs in **Fig. 4**.

RD morphology in this study corroborated previous findings – arrhythmias were perpetuated by

one RD at a time, with activity in the periphery including conduction block, transient reentry, and

wavefront collision (5, 6).

Insights from analysis of RD inducibility and localization by LA region

Further to the macroscopic analysis discussed above, our region-by-region analysis of RD inducibility and localization showed that spatial properties of the fibrotic substrate between AFib and ESUS models are not intrinsically different. However, this analysis yielded several noteworthy findings that suggest subtle distinctions in fibrosis pattern may exist between the two groups. The most striking difference exists for the posterior wall region, where the IdS score was ≈2x higher in AFib compared to ESUS models. Thus, even in cases where posterior LA wall ectopic excitations occur in ESUS patients, our simulations suggest that they could be up to 50% less likely to engage the fibrotic substrate and initiate sustained reentry compared to the same activity in AFib patients. This possibility does not contradict our hypothesis that the lack of arrhythmia in ESUS is due to a dearth of triggers; rather, it is a complementary corollary that can be put to the test in future clinical and computational analysis.

While understanding of RD localization dynamics in AFib remains limited, evidence from ECGI mapping indicates that reentrant activity occurs most frequently in the PV and posterior wall regions (16, 18, 19). Our findings are consistent with these data for AFib but not ESUS models. Many (41.1%) AFib model RDs localized to the LPV region. In contrast, in the ESUS population the RD localization hotspot was the region comprising the anterior wall and LAA (37%). The importance of this finding is unclear, as tendencies toward reentrant activity in particular LA areas have not been meaningfully correlated to clinical arrhythmia properties and are potentially subject to changes in conduction velocity or action potential duration (20). However, it provides a path for future validation studies: if incident AFib in patients who previously presented with ESUS can be characterized by intracardiac mapping, the hypothesis that RDs localize preferentially to the anterior wall can be tested.

Finally, our analysis of fibrosis spatial patterns revealed that the proportion of tissue with the propensity to harbor RDs (as established via machine learning in prior work (5)) was higher in the LPV region for AFib compared to ESUS models. While we acknowledge that a more fulsome

analysis will be required to draw comprehensive conclusions on this subject, we note that our results are consistent with the most prominent observed regional variabilities in RD localization and inducibility between the two model groups. In contrast, ESUS patients trended toward a higher percentage of pro-RD tissue in all anatomical areas except the LPVs. Thus, despite some interesting and potentially consequential differences in regional distribution of potentially pro-arrhythmic fibrosis, our overarching conclusion remains unchanged: our models suggest that if the ESUS substrate were subjected to suitable triggered activity, it could sustain the same types of RDs as those that contribute to AFib perpetuation.

Limitations

In this study, atrial tissue is modeled as a bilayer to drastically reduce computational load. Previous studies have used this modeling framework (15, 21) to represent human atria effectively, but the framework remains a simplification compared to volumetric 3D models. Moreover, clinical-grade MRI resolution limits our ability to detect fine details in anatomical structure and spatial distribution of potentially arrhythmogenic substrate, for instance slow-conducting tracks of fibrotic atrial tissue that could underlie microreentrant circuits (22). While these models are patient-specific in terms of LA anatomy and each individual's unique pattern of fibrotic remodeling, they do not incorporate inter-patient variability in CV and electrophysiological properties such as ion channel expression. Nevertheless, our previous analysis indicates that this representation of atrial architecture with generic "average AFib" electrophysiology is appropriate for use in patient-derived modeling (12, 20).

As in previous studies (23, 24), our models do not differentiate between cell- or tissue-scale properties of atrial electrophysiology between patients with paroxysmal and persistent forms of AFib. Likewise, our approach to characterizing potential arrhythmia propensity in ESUS patients assumes cell- and tissue-scale remodeling based on experimental and clinical data from the AFib milieu. Although this is relevant as a limitation and must be considered when interpreting our results, this aspect of our approach is also one of the major advantages of the modeling and

simulation methodology. Specifically, it allows us to assess whether there are any relevant

differences in the spatial pattern of fibrotic remodeling between ESUS and AFib patients in the

absence of other potentially confounding variables. A related limitation is that patients with stroke

were excluded from the AFib cohort. This was because stroke etiology in the database from which

AFib patients were drawn was not explicitly adjudicated to be cardioembolic, other ischemic such

as atherosclerotic, or hemorrhagic. Therefore, we would not be able to draw reliable conclusions

regarding the role of fibrosis in stroke in this population.

Finally, the mechanism of stroke in ESUS patients may be independent of AFib.

Decreased atrial function due to atrial fibrosis may contribute to reduced hemodynamic efficacy

and thrombus formation in the absence of AFib. Currently, secondary stroke prophylaxis is

dependent on detecting AFib and predicting, through computational modeling, which atria are

more prone to manifest AFib may be of clinical value. Another future research direction that could

prove highly fruitful would be to create multi-scale, multi-physics image-based models of the

fibrotic atria to assess each individual's risk of clot formation in a patient-specific manner (25).

Conclusions

Simulations suggest that the pro-arrhythmic properties of fibrotic substrate in ESUS and

AFib patients are indistinguishable. Our results show that fibrotic remodeling in ESUS patients

has the theoretical capacity to sustain reentry when subjected to common AFib triggers. Thus, we

conclude that fibrotic substrate conducive to perpetuating reentry may exist in up to half of ESUS

patients. As individuals studied in this cohort present with incident AFib over the next few years,

we will be able to put this hypothesis to the test. Our findings also support the notion that the lack

of AFib in this population may be attributable to a lack of suitable arrhythmic triggers, but further

research is needed to fully justify this claim. While the existence of pre-clinical substrate is

correlated with a higher global proportion of fibrotic tissue, many ESUS cases defied these

expectations, suggesting that fibrosis burden alone is insufficient for predicting pre-clinical AFib

substrate. This conclusion justifies the use of computational simulations to probe beyond the

fibrosis as imaged. Overall, these results provide novel insights into the role of atrial fibrotic

remodeling as a critical nexus between the otherwise distinct manifestations of AFib and ESUS.

Methods

Patient Population

Patients were recruited to undergo cardiac LGE-MRI from the University of Washington

(Seattle, WA) and Klinikum Coburg (Coburg, Germany) between July 2016 and June 2019. This

study was approved by the Institutional Review Board (IRB) of the University of Washington (UW)

and the Ethikkommission der Bayerischen Ländesärztekammer München, Bayern, Deutschland;

all participants provided written informed consent. Patients with ESUS met published diagnostic

criteria (26). Patients with paroxysmal (27/45, 60%) or persistent AFib and without stroke were

recruited from the UW Cardiac Arrhythmia Data Repository, an IRB-approved database for

arrhythmia patients. Exclusion criteria for AFib patients included those who had undergone LA

catheter ablation before MRI and those with only atrial flutter. Patients with cardiac implantable

electronic devices, severe claustrophobia, renal dysfunction, and other contraindications to MRI

or gadolinium-based contrast were excluded.

MRI acquisition

Cardiac LGE-MRI was obtained on all participants to quantify the extent of LA fibrosis

using previously described protocols (13). Scans were performed on Philips Ingenia and Siemens

Avanto clinical scanners, 15 to 25 minutes after contrast injection, using a 3-dimensional

inversion-recovery, respiration-navigated, ECG-gated, gradient echo pulse sequence. Acquisition

parameters included transverse imaging volume with a voxel size of 1.25 x 1.25 x 2.5 mm

(reconstructed to 0.625 x 0.625 x 1.25 mm). Scan time was 5-10 minutes dependent on respiration

and heart rate. Fat saturation sequences were used to suppress signal from fatty tissue.

Reconstruction of 3D patient-derived atrial models from LGE-MRI

Geometric models were reconstructed from LGE-MRI and the relative extent of fibrosis in

the LA was quantified via an adaptive histogram thresholding algorithm (27). Clinical-grade

meshes (i.e., coarse discretization) produced by Merisight Inc. (Salt Lake City, UT) were

resampled with a target resolution of 200 μm using an automated process based on gmsh (28).

Each LA model was represented as a bilayer comprising of nested endocardial and epicardial

shells (21), linked at every point by linear connections ($\sigma = 0.8$ S m^{-1}) (**Fig. 1A**). LA bilayer models

were generated by slightly inflating the single-surface mesh to form the epicardial surface (i.e.,

duplicating endocardial points then moving them outward by 100 μm along the surface normal

vector), then connecting the nested shells by attaching linear elements between corresponding

nodes. In each patient-derived model, realistic myocardial fiber orientations were mapped from

an atlas geometry (21) using the universal atrial coordinates (UAC) approach. Briefly, this process

assigned epicardial and endocardial fibers from a previously published bilayer model to the target

atrial geometry (**Fig. 1B**) (29, 30). In all finite element LA meshes, the average element edge

length was ≈188 μm and the number of nodes ranged from ≈600,000 to ≈1.4 million, depending

on LA size. This mesh resolution is consistent with previously established benchmarks for

minimizing numerical error due to spatial discretization in simulations of cardiac wavefront

propagation (31, 32).

Modeling of atrial electrophysiology in fibrotic and non-fibrotic regions

Our methodology for computational modeling at the cell and tissue scale of the fibrotic and

non-fibrotic atrial electrophysiology can be found in previously published papers (5, 16, 33).

Briefly, in non-fibrotic regions, a human atrial action potential model (34) was used to represent

membrane kinetics, including parameter modifications to fit clinical monophasic action potential

recordings from AFib patients (I_{Kur}, I_{to}, I_{CaL} decreased by 50%, 50%, 70%, respectively) (5, 35). At

the tissue scale, conductivity tensor values in non-fibrotic tissue (longitudinal: $\sigma_L = 0.409\,\text{S m}^{-1}$;

transverse: $\sigma_T = 0.0820\,\text{S m}^{-1}$) were calibrated to obtain effective conduction velocity (CV) values

of $71.49\,\text{cm s}^{-1}$ and $37.14\,\text{cm s}^{-1}$ (longitudinal and transverse). These conductivities were chosen

as a compromise between CV values measured in patients induced AFib ($61\pm6\,\text{cm s}^{-1}$) (36),

clinically mapped patients with AFib or atrial flutter (median: $60\,\text{cm s}^{-1}$; inter-quartile range: 22 cm

s^{-1}) (37), and simulations calibrated to match intracardiac mapping data from an individual with

paroxysmal AFib (median: $143\,\text{cm s}^{-1}$ longitudinal, $94\,\text{cm s}^{-1}$ transverse) (38). In fibrotic regions,

modifications to the AFib-like action potential model (I_{CaL}, I_{Na}, and I_{K1} decreased by 50%, 40%,

50%, respectively) were implemented as in prior studies (5, 39), resulting in a 15.4% increase in

action potential duration and a 49.6% decrease in upstroke velocity. These changes represented

the effect of elevated transforming growth factor-β1, a key component of the fibrogenic signaling

pathway. As in previous studies (5, 16, 33), tissue-scale effects of interstitial fibrosis and gap

junction remodeling were represented by reducing overall conductivity and exaggerating the

anisotropy ratio ($\sigma_L{:}\sigma_T$) from 5:1 to 8:1 ($\sigma_L = 0.177\,\text{S m}^{-1}$; $\sigma_T = 0.0221\,\text{S m}^{-1}$).

Simulation of electrical activity and numerical aspects

Electrical propagation in bilayer LA models was simulated by solving the monodomain

equation using the finite element method. This system was coupled with ordinary differential and

algebraic equations representing myocyte membrane dynamics at each node in the mesh, as

described in the prior section. All simulations were executed on the Hyak supercomputer system

at the University of Washington using the openCARP software package (40, 41), which is

available for academic use (see: https://openCarp.org). The compute time required to complete

each unique simulation ranged from 1-10 hours. The total CPU time for all simulations conducted

in all models was 13.4 years.

Induction and analysis of reentrant atrial arrhythmias

Simulations were performed to assess the pro-arrhythmic propensity of the fibrotic

substrate in each patient-derived model. Arrhythmia induction via rapid pacing was attempted

from 15 pacing sites derived from AFib trigger sites (**Fig. 1C**, see caption for detailed anatomical

site descriptions) (14). Clinically relevant AFib trigger sites were chosen over a random pacing

schematic to specifically capture RDs that arise from locations demonstrated to induce AFib. As

in previous publications, a clinically relevant pacing sequence of 12 electrical stimuli was delivered

at each of the 15 locations (5, 39). Individual cell-scale ionic models were paced to limit cycle at

a rate basic cycle length of 500 ms. The electrical stimulus consisted of two initial pulses with a

coupling interval of 300ms, followed by pulses ramping down to 200ms in 20ms intervals. After

the delivery of the final stimulus, simulations were monitored for self-sustaining electrical

wavefront propagation. For all cases in which activity persisted for at least 5,000ms post-pacing,

we applied further analysis to determine whether the cause was an induced RD or macroscopic

re-entry (i.e., continuous repetitive, self-sustaining activation propagating around a non-

conductive obstacle such as the mitral valve or pulmonary vein(s)), which we consider flutter-like

reentry. Instances of macroscopic re-entry were excluded from further analysis.

For each AFib-inducible simulation, we documented whether each pacing site induced

reentry and analyzed patterns of RD localization. Unique RD morphologies in each patient-

derived model were classified as belonging to one of five anatomical regions (**Fig. 1D**), which

were delineated automatically in a two-step process summarized in **Supplemental Fig. 1**. First,

the LA was subdivided into three broad anatomical areas (region 1: LA floor, 2: posterior wall; 3:

anterior wall including LAA) using standardized cutoff values in the UAC space (29). Second, the

left and right PV areas (regions 4 and 5, respectively) were established using a region-growing

approach such that each accounted for 15% of the total LA surface area. We then defined region-

wise inducibility scores (IdS) across all models in a particular group (ESUS or AFib) as the

proportion of pacing sites within a given region from which rapid pacing resulted in initiation of an

RD. For example, since the LPV region contains four pacing sites (anterior/posterior LSPV/LIPV),

the corresponding ESUS IdS value would be derived by summing the number of instances in

which pacing from those locations induced RD across inducible ESUS models then dividing by

four. This ensured our ability to assess spatial heterogeneity of sensitivity to triggered activity in

a manner that was unbiased to the relative abundance of pacing sites in some LA regions.

Quantitative analysis of fibrosis spatial pattern and potentially pro-ectopic effects of fibrosis

Fibrotic tissue areas with both high local fibrosis density (FD) and high local fibrosis

entropy (FE) have increased propensity for RD localization, as shown in computational models

(15) and intracardiac mapping of patients with persistent AFib (17). We used a constraint equation

derived by support vector machine-based classification to delineate such pro-RD regions. As in

prior studies (5, 23), the classification polynomial was: $0.4096\ FD^2 + 3.28(FD)(FE) - 0.1036\ FE^2 -$

$0.7112(FD) - FE + 0.0429$. Maps of pro-RD tissue were subdivided into the five LA regions, as

described above and region-specific burdens of pro-RD fibrosis pattern were calculated.

In some situations, fibrosis itself can develop the capacity to generate ectopic triggers of

arrhythmia (7, 8). Although the cell-scale models used in this study do not undergo early or

delayed afterdepolarizations due to simplified intracellular calcium handling, fibrotic regions in our

models do have a potentially pro-ectopic higher resting membrane voltage (V_m) compared to non-

fibrotic regions due to reduced ion channel expression levels (I_{CaL}, I_{Na}, I_{K1}). Thus, as a surrogate

measure of potential intrinsic pro-trigger capacity in fibrotic tissue, we characterized the extent of

tissue in ESUS and AFib models in which the fibrosis pattern resulted in abnormal depolarization.

To do this, we allowed all 90 models to reach a quasi-equilibrium state (1000 ms in the absence

of pacing). We aggregated resting V_m values across all models and identified the 95[th] percentile

as the threshold for delineation of fibrosis-induced abnormal depolarization (–75.3 mV). Then, we

calculated the proportion of tissue in each model with an equilibrium V_m above that threshold.

Statistical Analysis

LA models for ESUS and AFib patients were divided into quartiles based on the extent of

fibrotic remodeling as measured by LGE-MRI. Continuous variables were compared pairwise

between groups using Wilcoxon rank-sum tests and were reported as mean ± standard deviation.

Confidence intervals were calculated as the interval for the true difference in mean with 95%

certainty. Categorical variables were compared using a χ^2 test. After classifying unique RDs and

number of pacing sites that induced reentry, correlation with fibrosis was assessed with logistic

regression. Statistical significance was established at two-tailed P≤0.05. All statistical analysis

was performed using R (42).

Impact statement: This study uses MRI-based computational modelling to explore why some stroke patients never
have arrhythmias, even though they have cardiac fibrosis that strongly associated with arrhythmia risk in other cohorts.
519
Author contributions: Savannah Bifulco: Conceptualization; Formal analysis; Validation; Investigation; Visualization;
Methodology; Writing - original draft; Writing - review and editing Griffin Scott: Formal analysis; Validation; Writing -
review and editing Sakher Sarairah: Data curation; Formal analysis; Validation; Visualization; Writing - review and
editing Zeinab Birjandian: Data curation; Formal analysis; Validation; Visualization; Writing – review and editing
Caroline Roney: Software; Validation; Methodology; Writing - review and editing Steven Niederer: Software;
Methodology; Writing - review and editing Christian Mahnkopf: Formal analysis; Validation; Investigation; Writing -
review and editing Marcel Mitlacher: Formal analysis; Validation; Investigation; Writing - review and editing Peter
Kuhnlein: Formal analysis; Validation; Investigation; Writing - review and editing David Tirschwell: Formal analysis;
Validation; Investigation; Writing - review and editing Will Longstreth: Formal analysis; Validation; Investigation; Writing
- review and editing Nazem Akoum: Conceptualization; Resources; Data curation; Supervision; Investigation; Writing -
original draft; Project administration; Writing - review and editing Patrick Boyle: Conceptualization; Resources; Data
curation; Software; Formal analysis; Supervision; Funding acquisition; Validation; Investigation; Visualization;
Methodology; Writing - original draft; Project administration; Writing - review and editing.
533
534　**Acknowledgments:** We would like to thank Dr. Colleen Clancy, Dr. Axel Loewe, and our anonymous peer reviewer
535　for extremely helpful and constructive feedback on our manuscript.
536
Funding: ARCS Foundation: Savannah F Bifulco; UKRI | Medical Research Council (MRC): Caroline H Roney,
MR/S015086/1; National Institutes of Health: Steven A Niederer, R01-HL152256; European Research Council: Steven
A Niederer, PREDICT-HF (864055); British Heart Foundation (BHF): Steven A Niederer, RG/20/4/34803; UKRI |
Engineering and Physical Sciences Research Council (EPSRC): Steven A Niederer, EP/P01268X/1; Wellcome Trust:
Steven A Niederer, 203148/Z/16/Z; HHS | National Institutes of Health (NIH): David Tirschwell, Will T Longstreth, NIH
5-U01-NS095869; John Locke Charitable Trust: Nazem Akoum The funders had no role in study design, data collection
and interpretation, or the decision to submit the work for publication.
544
545　**Data availability:** Where possible (**Figs. 2, 3, 5, and 6**), raw numerical data underlying figures are available via figshare
546　[to be linked]. Patient-derived data related to this article will be shared with interested parties for non-commercial reuse
547　on reasonable request to the co-corresponding authors and approval from the Institutional Review Board of the
548　University of Washington.
549
550　**Ethics:** Human Subjects: Yes. Ethics Statement: This study was approved by the Institutional Review Board (IRB) of
551　the University of Washington (UW) and the Ethikkommission der Bayerischen Ländesärztekammer München, Bayern,
552　Deutschland; all participants provided written informed consent. Clinical Trial: No. Animal Subjects: No.
553

References

1.　　　Andrade J, Khairy P, Dobrev D, Nattel S. The clinical profile and pathophysiology of
atrial fibrillation: relationships among clinical features, epidemiology, and mechanisms. Circ
Res. 2014;114(9):1453-68.

2.   Israel C, Kitsiou A, Kalyani M, Deelawar S, Ejangue LE, Rogalewski A, et al. Detection
of atrial fibrillation in patients with embolic stroke of undetermined source by prolonged
monitoring with implantable loop recorders. Thromb Haemost. 2017;117(10):1962-9.

3.   Brachmann J, Morillo CA, Sanna T, Di Lazzaro V, Diener HC, Bernstein RA, et al.
Uncovering Atrial Fibrillation Beyond Short-Term Monitoring in Cryptogenic Stroke Patients:
Three-Year Results From the Cryptogenic Stroke and Underlying Atrial Fibrillation Trial. Circ
Arrhythm Electrophysiol. 2016;9(1):e003333.

4.   Tandon K, Tirschwell D, Longstreth WT, Jr., Smith B, Akoum N. Embolic stroke of
undetermined source correlates to atrial fibrosis without atrial fibrillation. Neurology.
2019;93(4):e381-e7.

5.   Zahid S, Cochet H, Boyle PM, Schwarz EL, Whyte KN, Vigmond EJ, et al. Patient-
derived models link re-entrant driver localization in atrial fibrillation to fibrosis spatial pattern.
Cardiovasc Res. 2016;110(3):443-54.

6.   Boyle PM, Zghaib T, Zahid S, Ali RL, Deng D, Franceschi WH, et al. Computationally
guided personalized targeted ablation of persistent atrial fibrillation. Nat Biomed Eng.
2019;3(11):870-9.

7.   Gouvea de Barros B, Weber dos Santos R, Lobosco M, Alonso S. Simulation of Ectopic
Pacemakers in the Heart: Multiple Ectopic Beats Generated by Reentry inside Fibrotic Regions.
Biomed Res Int. 2015;2015:713058.

8.   Alonso S, Dos Santos RW, Bar M. Reentry and Ectopic Pacemakers Emerge in a Three-
Dimensional Model for a Slab of Cardiac Tissue with Diffuse Microfibrosis near the Percolation
Threshold. PLoS One. 2016;11(11):e0166972.

9.   Roney CH, Beach ML, Mehta A, Sim I, Corrado C, Bendikas R, et al. In silico
comparison of left atrial ablation techniques that target the anatomical, structural and electrical
substrates of atrial fibrillation. Front Physiol. 2020.

10.   Xia Y, Hertervig E, Kongstad O, Ljungstrom E, Platonov P, Holm M, et al. Deterioration
of interatrial conduction in patients with paroxysmal atrial fibrillation: electroanatomic mapping of
the right atrium and coronary sinus. Heart Rhythm. 2004;1(5):548-53.

11.   Litchenberg WH, Norman LW, Holwell AK, Martin KL, Hewett KW, Gourdie RG. The rate
and anisotropy of impulse propagation in the postnatal terminal crest are correlated with
remodeling of Cx43 gap junction pattern. Cardiovasc Res. 2000;45(2):379-87.

12.   Hakim JB, Murphy MJ, Trayanova NA, Boyle PM. Arrhythmia dynamics in computational
models of the atria following virtual ablation of re-entrant drivers. Europace.
2018;20(suppl_3):iii45-iii54.

13.   Marrouche NF, Wilber D, Hindricks G, Jais P, Akoum N, Marchlinski F, et al. Association
of atrial tissue fibrosis identified by delayed enhancement MRI and atrial fibrillation catheter
ablation: the DECAAF study. JAMA. 2014;311(5):498-506.

14.   Santangeli P, Marchlinski FE. Techniques for the provocation, localization, and ablation
of non-pulmonary vein triggers for atrial fibrillation. Heart Rhythm. 2017;14(7):1087-96.

15.   Roney CH, Bayer JD, Zahid S, Meo M, Boyle PM, Trayanova NA, et al. Modelling
methodology of atrial fibrosis affects rotor dynamics and electrograms. Europace. 2016;18(suppl
4):iv146-iv55.

16.   Boyle PM, Hakim JB, Zahid S, Franceschi WH, Murphy MJ, Vigmond EJ, et al.
Comparing Reentrant Drivers Predicted by Image-Based Computational Modeling and Mapped
by Electrocardiographic Imaging in Persistent Atrial Fibrillation. Front Physiol. 2018;9:414.

17.   Cochet H, Dubois R, Yamashita S, Al Jefairi N, Berte B, Sellal JM, et al. Relationship
Between Fibrosis Detected on Late Gadolinium-Enhanced Cardiac Magnetic Resonance and
Re-Entrant Activity Assessed With Electrocardiographic Imaging in Human Persistent Atrial
Fibrillation. JACC Clin Electrophysiol. 2018;4(1):17-29.

18.   Haissaguerre M, Hocini M, Denis A, Shah AJ, Komatsu Y, Yamashita S, et al. Driver
domains in persistent atrial fibrillation. Circulation. 2014;130(7):530-8.

19.   Tanaka K, Zlochiver S, Vikstrom KL, Yamazaki M, Moreno J, Klos M, et al. Spatial
distribution of fibrosis governs fibrillation wave dynamics in the posterior left atrium during heart
failure. Circ Res. 2007;101(8):839-47.

20.   Deng D, Murphy MJ, Hakim JB, Franceschi WH, Zahid S, Pashakhanloo F, et al.
Sensitivity of reentrant driver localization to electrophysiological parameter variability in image-
based computational models of persistent atrial fibrillation sustained by a fibrotic substrate.
Chaos. 2017;27(9):093932.

21.   Labarthe S, Bayer J, Coudiere Y, Henry J, Cochet H, Jais P, et al. A bilayer model of
human atria: mathematical background, construction, and assessment. Europace. 2014;16
Suppl 4:iv21-iv9.

22.   Hansen BJ, Zhao J, Li N, Zolotarev A, Zakharkin S, Wang Y, et al. Human Atrial
Fibrillation Drivers Resolved With Integrated Functional and Structural Imaging to Benefit
Clinical Mapping. JACC Clin Electrophysiol. 2018;4(12):1501-15.

23.   Ali RL, Hakim JB, Boyle PM, Zahid S, Sivasambu B, Marine JE, et al. Arrhythmogenic
propensity of the fibrotic substrate after atrial fibrillation ablation: a longitudinal study using
magnetic resonance imaging-based atrial models. Cardiovasc Res. 2019;115(12):1757-65.

24.   Shade JK, Ali RL, Basile D, Popescu D, Akhtar T, Marine JE, et al. Preprocedure
Application of Machine Learning and Mechanistic Simulations Predicts Likelihood of Paroxysmal
Atrial Fibrillation Recurrence Following Pulmonary Vein Isolation. Circ Arrhythm Electrophysiol.
2020;13(7):e008213.

25.   Boyle PM, Del Alamo JC, Akoum N. Fibrosis, atrial fibrillation and stroke: clinical updates
and emerging mechanistic models. Heart. 2021;107(2):99-105.

26.   Hart RG, Catanese L, Perera KS, Ntaios G, Connolly SJ. Embolic Stroke of
Undetermined Source: A Systematic Review and Clinical Update. Stroke. 2017;48(4):867-72.

27.   Jadidi AS, Cochet H, Shah AJ, Kim SJ, Duncan E, Miyazaki S, et al. Inverse relationship
between fractionated electrograms and atrial fibrosis in persistent atrial fibrillation: combined
magnetic resonance imaging and high-density mapping. J Am Coll Cardiol. 2013;62(9):802-12.

28.   Geuzaine C, Remacle J-F. Gmsh: A 3-D finite element mesh generator with built-in pre-
and post-processing facilities. International Journal for Numerical Methods in Engineering.
2009;79(11):1309-31.

29.   Roney CH, Pashaei A, Meo M, Dubois R, Boyle PM, Trayanova NA, et al. Universal
atrial coordinates applied to visualisation, registration and construction of patient specific
meshes. Med Image Anal. 2019;55:65-75.

30.   Roney CH, Bendikas R, Pashakhanloo F, Corrado C, Vigmond EJ, McVeigh ER, et al.
Constructing a Human Atrial Fibre Atlas. Ann Biomed Eng. 2020.

31.   Niederer SA, Kerfoot E, Benson AP, Bernabeu MO, Bernus O, Bradley C, et al.
Verification of cardiac tissue electrophysiology simulators using an N-version benchmark. Philos
Trans A Math Phys Eng Sci. 2011;369(1954):4331-51.

32.   Boyle PM, Ochs AR, Ali RL, Paliwal N, Trayanova NA. Characterizing the
arrhythmogenic substrate in personalized models of atrial fibrillation: sensitivity to mesh
resolution and pacing protocol in AF models. Europace. 2021;23(suppl_1):i3-i11.

33.   Boyle PM, Hakim JB, Zahid S, Franceschi WH, Murphy MJ, Prakosa A, et al. The
Fibrotic Substrate in Persistent Atrial Fibrillation Patients: Comparison Between Predictions
From Computational Modeling and Measurements From Focal Impulse and Rotor Mapping.
Front Physiol. 2018;9:1151.

34.   Courtemanche M, Ramirez RJ, Nattel S. Ionic mechanisms underlying human atrial
action potential properties: insights from a mathematical model. Am J Physiol.
1998;275(1):H301-21.

35.   Krummen DE, Bayer JD, Ho J, Ho G, Smetak MR, Clopton P, et al. Mechanisms of
human atrial fibrillation initiation: clinical and computational studies of repolarization restitution
and activation latency. Circ Arrhythm Electrophysiol. 2012;5(6):1149-59.

36.   Konings KT, Kirchhof CJ, Smeets JR, Wellens HJ, Penn OC, Allessie MA. High-density
mapping of electrically induced atrial fibrillation in humans. Circulation. 1994;89(4):1665-80.

37.   Verma B, Oesterlein T, Loewe A, Luik A, Schmitt C, Dossel O. Regional conduction
velocity calculation from clinical multichannel electrograms in human atria. Comput Biol Med.
2018;92:188-96.

38.   Roney CH, Whitaker J, Sim I, O'Neill L, Mukherjee RK, Razeghi O, et al. A technique for
measuring anisotropy in atrial conduction to estimate conduction velocity and atrial fibre
direction. Comput Biol Med. 2019;104:278-90.

39.   Zahid S, Whyte KN, Schwarz EL, Blake RC, 3rd, Boyle PM, Chrispin J, et al. Feasibility
of using patient-specific models and the "minimum cut" algorithm to predict optimal ablation
targets for left atrial flutter. Heart Rhythm. 2016;13(8):1687-98.

40.   Vigmond EJ, Hughes M, Plank G, Leon LJ. Computational tools for modeling electrical
activity in cardiac tissue. J Electrocardiol. 2003;36 Suppl:69-74.

41.   Vigmond EJ, Weber dos Santos R, Prassl AJ, Deo M, Plank G. Solvers for the cardiac
bidomain equations. Prog Biophys Mol Biol. 2008;96(1-3):3-18.

42.   R Development Core Team. R: A language and environment for statistical computing.
Vienna, Austria: R Fonudation for Statistical Computing; 2019.
677

Supplemental Figure 1. LA subdivision scheme. (A) Alpha coordinate from universal atrial coordinate

(UAC) system mapped on to a representative LA model. (B) Beta coordinate from UAC mapped on to a

representative LA model. (C) 2D representation of UAC with pulmonary vein locations labeled. Dashed

lines represent edges of the five atrial regions. (D) Segmented LA mapped onto a 3D mesh with numbered

regions corresponding to regions in (C). (E) Final division scheme after edge expansion of the LPVs and

RPVs to generate regions 3 and 4, respectively.

Supplemental Figure 2. Boxplot depicting the percentage of tissue with significantly depolarized tissue

(>95th percentile) between ESUS and AFib models after reaching steady state. P=0.32; CI: [–0.007,

0.019].