

1 **Emergence of the D614A mutation in the spike protein of SARS-CoV-2: Imported cases**  
2 **to the South Korea**

3

4 Ae Kyung Park, Jin Sun No, Eunkyung Shin, Yoon-Seok Chung, Il-Hwan Kim, Byung-Hak

5 Kang, Gi-Eun Rhie, Heui Man Kim, Jeong-Min Kim, Ye Eun Park, Min Joon Kim, Jae Sun

6 Park, Cheon-Kwon Yoo, Junyoung Kim

7

8 Division of High-risk Pathogens, Center for Laboratory Control of Infectious Disease, Korea

9 Centers for Disease Control & Prevention, Cheongju-si, Republic of Korea

10

11

12

13

14

15

16

17

18

19

20 **Abstract**

21 We identified imported cases of a novel D614A mutation in the spike glycoprotein of SARS-  
22 CoV-2 in the South Korea. SARS-CoV-2 harboring the novel mutation was isolated from  
23 returning travelers from Uzbekistan. These results emphasized the possibility that new  
24 mutations have emerged in that area.

25

26 Since a new novel beta-coronavirus, SARS-CoV-2, was first reported in December 2019,  
27 there has been ongoing explosive spread of this virus throughout the world, severely  
28 threatening human health (1). South Korea was one of the first countries to report a  
29 significant Coronavirus Disease-19 (COVID-19) outbreak; however, the epidemic curve has  
30 been quickly flattened due to the outstanding response of the Korean government (2).  
31 Currently, community spread of Covid-19 has slowed in the South Korea. On the other hand,  
32 more than half of confirmed cases were related to travel (2). The Korea Centers for Disease  
33 Control and Prevention (KCDC) has begun to conduct whole-genome sequencing (WGS) of  
34 COVID-19 isolated from travelers to obtain real-time genomic epidemiology data, rapidly  
35 monitor how the virus mutates and investigate sequence variations.

36 On July, 2020, a man who entered Korea from Uzbekistan (Uzbekistan nationality) and had  
37 symptoms of cough, fever, and headache was diagnosed with COVID-19 (3). For WGS, total  
38 RNA was extracted, and cDNA was amplified using primer pools ([https://artic.network/ncov-  
39 2019](https://artic.network/ncov-2019)). Libraries were prepared using the Nextera DNA Flex Library Prep Kit (Illumina,  
40 USA), and sequencing was performed on a MiSeq instrument with 2 x 75 base pairs using a  
41 MiSeq reagent kit V3 (Illumina, USA). The reads were mapped to the reference genome  
42 MN908947.3 using CLC Genomics Workbench version 20.0.3 (CLC Bio, Denmark). A total

43 of 663,666 (98.50%) out of a total of 673,800 reads mapped to the reference genome. We  
44 obtained 99.72% genome coverage at the median genome coverage depth of 1554×  
45 (Supplementary Table 1). The genomic sequences of the SARS-CoV-2 strain were deposited  
46 in GISAID (Accession no. EPI\_ISL\_516783, EPI\_ISL\_516784, EPI\_ISL\_516785). Using the  
47 reference genome, single-nucleotide variants (SNVs) were called using the BioNumerics  
48 version 7.6 SARS-CoV-2 plugin (Applied Maths, Belgium).

49 We identified 14 polymorphic sites in the genome, which corresponded to all SNVs (Table 1).  
50 In total, 13 appeared within the gene-coding regions. Of note, we identified a GAT→GCT  
51 (A23403C) nucleotide change, which leads to a nonsynonymous change (D614A) in the  
52 spike (S) glycoprotein. Based on 77,577 viral genomes downloaded on August 5, 2020, from  
53 the GISAID database (4), the DNA sequences corresponding to the S protein of each genome  
54 were extracted and aligned to the mutated S protein (D614A) (Table 1). The alignment results  
55 indicated that the D614A mutation of the S protein identified in this study was a novel  
56 mutation that has not been identified previously.

57 In addition, two travelers who visited Uzbekistan and returned to Korea were diagnosed with  
58 COVID-19 immediately after entering Korea. They arrived in the South Korea on July after  
59 short-term travel (Supplementary Table 1). The WGS results of the above two isolates also  
60 indicated that they had D614A mutations (Table 1). Due to the lack of WGS data on isolates  
61 from Uzbekistan in GISAID, we were unable to compare our mutation results to those from  
62 Uzbekistan. However, the identification of D614A in the S protein from three individuals  
63 who traveled Uzbekistan in recent days emphasized that new types of mutations might be  
64 emerging and spreading in that area.

65 The significance of the D614G mutation of the S protein for SARS-CoV-2 infection was

66 elucidated in a recent report by Korver *et al.* (5). Korver *et al.* suggested that an association  
67 exists between the G614 variant and higher levels of viral nucleic acid in the upper  
68 respiratory tract in patients, which make it more infectious than D614 and has allowed the  
69 rapid spread of G614 throughout the world. Additionally, this higher level of infection might  
70 require a higher level of antibody, which is correlated with vaccines or antibody-based  
71 therapeutic agents. From a structural perspective, the D614G mutation destabilizes the  
72 interaction between the S1 and S2 units, resulting in the promotion of S1 shedding.  
73 Additionally, this mutation induces the conformational change of the receptor-binding  
74 domain (RBD) of the S protein allosterically to promote the binding of angiotensin-  
75 converting enzyme 2 (ACE2) (6). Additionally, the D614G mutation may modify the  
76 secondary structure of the furin cleavage site to facilitate the cleavage of S1 and S2 (7).  
77 Considering that the D614G mutation is expected to have significant effects on the spread of  
78 SARS-CoV-2, it will be necessary to study and characterize the D614A mutation to evaluate  
79 its phenotypes.

80 In this study, we identified novel D614A mutations in SARS-CoV-2 isolated from travelers  
81 returning to Korea from Uzbekistan in recent days. The results of this study emphasized the  
82 need for enhanced surveillance, which will be essential to combat COVID-19, especially  
83 where genetic epidemiology data are scarce.

84

85 Dr. Park is a senior researcher at the Korea Centers for Diseases Control & Prevention. Her  
86 research interests include the molecular epidemiology of viral and bacterial diseases.

87

88

89 **References**

90 1. Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak  
91 associated with a new coronavirus of probable bat origin. *Nature*. 2020 Mar;579(7798):270-3.

92 2. Korea Centers for Disease Control and Prevention. Press Release. 2020.  
93 [ncov.mohw.go.kr](http://ncov.mohw.go.kr)

94 3. Kim JM, Chung YS, Jo HJ, Lee NJ, Kim MS, Woo SH, et al. Identification of  
95 Coronavirus Isolated from a Patient in Korea with COVID-19. *Osong Public Health Res*  
96 *Perspect*. 2020 Feb;11(1):3-7.

97 4. Shu Y, McCauley J. GISAID: Global initiative on sharing all influenza data - from  
98 vision to reality. *Euro surveillance : bulletin European sur les maladies transmissibles =*  
99 *European communicable disease bulletin*. 2017 Mar 30;22(13).

100 5. Korber B, Fischer WM, Gnanakaran S, Yoon H, Theiler J, Abfalterer W, et al.  
101 Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G Increases Infectivity of the  
102 COVID-19 Virus. *Cell*. 2020 Jul 3.

103 6. Korber B, Fischer WM, Gnanakaran S, Yoon H, Theiler J, Abfalterer W, et al. Spike  
104 mutation pipeline reveals the emergence of a more transmissible form of SARS-CoV-2.  
105 *BioRxiv*. 2020.

106 7. Tang L, Schulkins A, Chen C-N, Deshayes K, Kenney JS. The SARS-CoV-2 Spike  
107 Protein D614G Mutation Shows Increasing Dominance and May Confer a Structural  
108 Advantage to the Furin Cleavage Domain. *Preprints*. 2020.

109

110 Table 1. Single-nucleotide variants (SNVs) of SARS-CoV-2 isolates from travelers returning to Korea from Uzbekistan with respect to the  
 111 reference genome

Gene	Nucleotide position	Reference (MN908947.3)	#1	#2	#3	Mutation
ORF1ab	4579	T	A	-	-	Synonymous
	8782	C	T	T	T	Synonymous
	11230	G	T	T	T	Nonsynonymous (M3655I)
	11970	A	-	G	G	Nonsynonymous (K3902R)
	20580	G	T	T	T	Synonymous
S	23403	A	C	C	C	Nonsynonymous (D614A)
ORF3a	26028	C	T	T	T	Synonymous
	26111	C	T	-	-	Nonsynonymous (P240L)
ORF8	28144	T	C	C	C	Nonsynonymous (L84S)
	28167	G	A	A	A	Nonsynonymous (E91K)
N	28878	G	A	A	A	Nonsynonymous (S202N)
	28887	C	-	T	T	Nonsynonymous (T205I)
	29224	G	-	-	A	Nonsynonymous (M317I)
Noncoding region (3' UTR)	29742	G	A	-	-	Noncoding region

112

113