50 policies, 1 pandemic, 500,000 deaths: Associations between state-level COVID-19

testing recommendations, tests *per capita*, undercounted deaths, vaccination policies,

and doses *per capita* in the United States

Stephanie R. Perniciaro & Daniel M. Weinberger

Department of Epidemiology of Microbial Diseases, Yale School of Public Health, New

Haven, CT, United States.

Email: stephanie.perniciaro@yale.edu

Abstract

Background: State health departments have been responsible for prioritizing and

allocating SARS-CoV-2 tests and vaccines. Testing and vaccination recommendations in

the United States varied by state and over time, as did vaccine rollouts, COVID-19 cases,

and estimates of excess mortality.

Methods: We compiled data about COVID-19 testing, cases, and deaths, and excess

pneumonia + influenza + COVID-19 deaths to assess relationships between testing

recommendations, *per capita* tests performed, epidemic intensity, and excess mortality

during the early months of the COVID-19 pandemic in the United States. We compiled

further data about state-level SARS-CoV-2 vaccination policies and doses administered

during the early months of the vaccine rollout.

Results: As of July 2020, 16 states recommended testing asymptomatic members of the general public. The rate of COVID-19 tests reported in each state correlated with more inclusive testing recommendations and with higher epidemic intensity. Higher *per capita* testing was associated with more complete reporting of COVID-19 deaths, which is a fundamental requirement for analyzing the pandemic. Testing *per capita* during the first three months was associated with vaccination *per capita* in the first three months of rollout. *Per capita* vaccine doses in each state were not associated with adherence to national guidelines.

Conclusions: Reported deaths due to COVID-19 likely represent an undercount of the true burden of the pandemic. States that struggled with testing rollout have also frequently struggled with vaccine rollout. Coordinated, consistent guidelines for COVID-19 testing and vaccine administration should be a high priority for state and national health systems.

Introduction

SARS-CoV-2, the virus that causes COVID-19, was declared a global pandemic by the World Health Organization in March 2020.[1] Worldwide, responses to the pandemic have varied and had varying success. In the United States, each state determined testing guidelines, lockdown procedures, and reopening procedures individually.[2] The Centers for Disease Control and Prevention also maintained testing recommendations and guidelines, and produced guidance for establishing prioritized groups during vaccine rollout, but states and localities were largely responsible for their own policies.[3] The COVID-19 pandemic in the United States is ongoing, costing over half a million lives by the end of February 2021. [4]

Testing for COVID-19 has been a continual hurdle in the United States.[5] Supply chain

issues,[6] doubts about testing accuracy and methodology,[7] and conflicting guidance from

federal, state and local officials about testing have all interfered with testing efforts.[8] State

governments and health departments established testing guidelines in March 2020, which

were sometimes augmented by local governmental health departments.[9] The turnaround

time for test results was strained,[10] and resources for individuals to sustain their household

resources while remaining isolated or quarantined while awaiting the result of a test have

not been widely available.[11]

This situation is now replaying with vaccine rollout and administration. Problems ranging

from misinformation to convoluted registration systems have prevented eligible

individuals from receiving vaccination appointments, echoed in problems including long

wait times[12] and sabotage[13] hampering the actual administration of doses.

Here we describe the changing recommendations for COVID-19 testing to the general

public in the United States in March-July 2020 and the vaccine recommendations in the

December 2020-February 2021.

Methods

The objectives of this analysis were to: 1) establish whether varying the state-level

COVID-19 testing guidelines had any impact on the actual number of tests performed,

and whether the number of tests performed was related to the rate of recognizing COVID-

19 deaths as a cause of excess mortality (reported COVID-19 deaths/excess pneumonia

+ influenza+ COVID-19 (PIC) deaths) and 2) establish whether state-level vaccination

policies had any impact on doses administered. COVID-19 testing guidelines were

documented from the websites of the departments of health in each of the 50 states and

the District of Columbia once monthly from March-July 2020 (Supplemental Table 1). We

ranked recommendations to tests by the general public (as opposed to members of

specific named risk groups) by assigning 0= criterion not specified, 1= accessible to

people with a negative influenza test, 2= accessible to people with symptoms consistent

with COVID-19 illness, 3= accessible to anyone, including asymptomatic people.

Vaccination policies were recorded once monthly for December 2020, January and

February of 2021.[14] Adherence to Advisory Committee on Immunization Practices (ACIP)

guidelines[15] was scored for the first three phases of vaccine rollout as 0= more restrictive

than ACIP guidelines/data not available, 1=identical to ACIP guidelines, 2=more inclusive

than ACIP guidelines.

Correlations between testing recommendation score, tests *per capita*, and percentage of

positive tests were analyzed with Spearman correlation tests, as were correlations

between vaccine recommendation priority match with ACIP guidelines and number of

doses administered total and *per capita*.

We evaluated relationships between the number of COVID-19 tests performed per capita,

testing recommendations score, and epidemic intensity (defined as excess PIC deaths[16]

per capita) using a negative binomial regression model. Total COVID-19 tests performed in each state was the outcome variable, offset by the log of the total population of each state, with the testing recommendations score (categorical) and the log of epidemic intensity as the independent variables.

We assessed the association of adherence to ACIP guidelines for vaccination prioritization and doses administered in a similar fashion, with doses administered as the outcome variable, offset by the log of the population, with adherence to ACIP guidelines and the current stage of vaccination rollout as the independent categorical variables. We investigated possible associations between testing *per capita* in March-May of 2020 and vaccine doses *per capita* in December 2020-February 2021 with monthly and overall Spearman correlation tests and with a negative binomial model as above where vaccine doses administered was the dependent variable, offset by the log of population, and the log of total tests administered/epidemic intensity was the independent variable.

We also used a generalized additive model (negative binomial link) to assess the completeness of death reporting. The outcome variable was the number of reported COVID-19 deaths in each state and month. The number of excess PIC deaths was included as an offset term, so this effectively models the ratio of reported COVID-19 deaths/excess PIC deaths. The *per capita* testing rate in each state and month was included as a penalized spline. Uncertainty in the estimates of excess PIC deaths were ignored in these analyses, and excess PIC deaths were constrained to be greater than

or equal to the reported COVID-19 deaths. The model was fit using the package "mgcv"

in R.

The number of excess PIC deaths per state and the reported number of COVID-19 deaths

were obtained from a repository describing excess deaths during the COVID-19 outbreak

(https://github.com/weinbergerlab/excess_pi_covid).[16] Positive tests and the overall

number of tests in each state were collected from the website of The COVID Tracking

Project (https://covidtracking.com/). Vaccine policy data were gathered from state health

department websites, in a publicly available repository curated by the Kaiser Family

Foundation.[17] Vaccine uptake data were taken from the CDC's COVID-19 Data Tracker

(https://covid.cdc.gov/covid-data-tracker/#vaccinations) Population data were taken from

the United States Census Bureau's 2019 estimates. R (R: A language and environment

for statistical computing, version 4.0.3 R Foundation for Statistical Computing, Vienna,

Austria, https://www.R-project.org/.) was used for statistical analyses.

Results

State Testing Guidelines

From March to July 2020, recommendations of COVID-19 testing for the general public

varied between states and over time (Figure 1), as did testing prioritization for identified

risk groups (Supplemental Figure 1).

As of July, 16 states recommended that asymptomatic members of the general public be

tested for COVID-19. 9 states actively recommended against the testing of asymptomatic

people; 3 of which then reversed this recommendation in July. South Dakota also

recommended against seeking a test if people were experiencing symptoms, citing the

lack of effective COVID-19 therapy as a contraindication for testing. Most states (35/51)

changed their testing recommendations twice between March and July. Only Oklahoma

did not change their testing criteria, though the state health department website specifies

that private testing facilities have their own criteria.

Factors associated with higher rates of testing

Reported COVID-19 tests *per capita* increased steadily, though unevenly, from March to

June, while the percentage of positive tests varied widely between states (Figure 2, 3).

States with more permissive testing recommendations had a lower percentage of positive

tests (rho -0.22, 95% CI -0.34, -0.11), possibly reflecting the inclusion of lower risk

individuals in the testing pool. The minimum percentage of positive tests for a single

month was 0.8% in May in Alaska; the maximum was 47% in April in New Jersey. In a

multivariate model, both more inclusive testing recommendations and higher epidemic

intensity were associated with higher rates of testing.

Testing intensity and completeness of death reporting

The completeness of recording deaths as due to COVID-19 was related to the rate of

testing. (Figure 4). This was particularly true in the first months of the

pandemic when testing levels were low relative to the intensity of the epidemic; in later

months, the level of testing was high and had little influence on the difference between

reported COVID-19 deaths and excess PIC deaths.

State Vaccination Guidelines

48 states included criteria as least as inclusive as ACIP guidelines in their first phase of

vaccination rollout, which dropped 31 to in the second phase, and 30 in the third phase.

Per capita dose administration was highest in Alaska in December (9422 doses per

100,000 population); in Washington D.C. in January (13108 per 100,000); in New Mexico

February (16579 per 100,000) and lowest in Ohio in December (2490 doses per 100,000

population); Idaho in January (2401 per 100,000); in Arkansas in February (7947 per

100,000; Figure 5). State guidelines at least as inclusive as ACIP were not associated

with administered COVID-19 vaccine doses *per capita* each month in correlation tests,

nor overall in the negative binomial model. Rollout of the staged, phased vaccine

administration prioritization schemes varied. All states are still in Phase 1, but have

broken this into three or more stages: as of February 15, 2021, 4 states were on the first

stage of rollout, 43 on the second, 4 on the third (Figure 6). 14 states still did not have

publicly available, fixed criteria prepared for their third stage of rollout, of 13 which were

already actively in the second stage of rollout, however, most of these states have further

subdivided their stages into tiers of prioritized groups, and the timeframe for implementing

each tier, stage, or phase is largely unknown.

The gap between delivered doses (doses received by states) and administered doses

(doses given to patients) varied extensively between states in January and February. In

January, the median percentage of doses *per capita* that were administered was 59%

(minimum: 49% (MO) maximum: 85% (ND)), and by February, the median percentage of

total administered doses *per capita* increased to 85% (minimum: 72% (DC) maximum:

98% (NM and ND)). In February by itself, 48 states administered more than the doses

that had been delivered during that month (presumably doses that were delivered in

previous months were administered in February), showing a ramp-up in vaccine

administration.

Association of per capita vaccination with per capita testing

State-level *per capita* COVID-19 testing was associated with per capita COVID-19

vaccines administered, both overall (rho: 0.71, 95%CI: 0.59-0.81) and for the first month

of testing (March 2020) and vaccine (December 2020) rollout (rho: 0.34, 95%CI: 0.03-

0.59). The negative binomial model showed that *per capita* testing/epidemic intensity was

positively associated with *per capita* COVID-19 vaccine administration in the first three

months of rollout (IRR: 1.17, 95%CI 1.10-1.25). States with less inclusive access to

COVID-19 testing for the general public (score 0 or 1) were not associated with states

with vaccine guidelines less inclusive than ACIP regulations during the first three months

of rollout.

Discussion

In order to obtain accurate statistics on the COVID-19 pandemic, viral testing needs to

scale with epidemic intensity. Particularly in the early months of the pandemic, restrictive

state-level testing recommendations and difficulties in accessing tests resulted in

inadequate testing levels.[5] This was manifest in higher viral positivity rates and greater

differences between reported COVID-19 deaths and excess PIC deaths. From March-

May, testing *per capita* and completeness of mortality data both increased, though

afterwards this apparent relationship became less clear. It is possible that other

characteristics of healthcare systems in each state influenced both the testing rates and

completeness of death reporting (i.e. supply chain or personnel issues). Expanded

access to testing plays a critical role in obtaining an accurate picture of the progression

of the epidemic.

Vaccination prioritization and allocation are, similar to testing recommendations, the

responsibility of the states, and have had a similarly unbalanced and inconsistent rollout.[18]

The distinct difference between the strong associations between inclusive testing policy

and tests performed *per capita* and the lack of association between inclusive vaccination

policy and vaccines administered *per capita* may indicate that other factors, such as

deficiencies in the vaccine administration supply chain, lack of access, or vaccine

hesitancy may have a stronger role in the variation in vaccines administered *per capita*.

The positive association of testing *per capita* and vaccination *per capita* is of interest, as

when both are low, it could be an indication that state-level public health infrastructure is

overtaxed, impeding a rapid rollout of these crucial programs. The nature of this

impediment could include political, economic, logistic, or human factors, each of which

would require a separate course of action to mitigate. Rather than utilizing limited public

health resources and personnel to independently construct 50 separate plans and 50

separate prioritization schemes of varying criteria and complexity, it would likely be more

efficient if the United States would unite behind a common set of guidelines and

standards.[19,20]

This study has certain limitations. Changes in completeness of death reporting could be

influenced by other factors that might also vary over time, such as state-level guidelines

for coding deaths as due to COVID-19. The viral testing criteria established by state health

departments are sometimes superseded by local governments or private clinics.[9] The

testing data counts tests, not individuals, so people tested multiple times will be included

multiple times. The vaccine administration data counts total doses administered, and

since both of the vaccines currently given Emergency Use Authorization in the United

States require two doses, there could be undetected variation in the number of people

receiving their first versus their second dose in each state.

Improving COVID-19 vaccine uptake in the face of not only logistical and administrative

challenges, but also sociobehavioral issues like hesitancy and misinformation, are further

impediments to mass vaccination efforts.[21] United, clear messaging is essential to build

trust and deliver compassionate, competent information to encourage people to get

vaccinated. Developing 50 different state-level uptake improvement plans seems like a

misallocation of resources. Avoiding this requires that centralized guidance be of high

quality and locality-specific adaptability. After a tumultuous pandemic year, resulting in

over 500,000 lives lost, it would be unwise to further rely on overextended state-level

public health infrastructure to be the ultimate source for policy, implementation, and advocacy.

Conclusions

The patchwork of state-level testing guidelines was associated with local variations in testing intensity, and these variations had consequences for the completeness of data on key indicators of the severity and progression of the pandemic. By contrast, the uneven state-level vaccination prioritization policies were not associated with variations in *per capita* doses administered. However, vaccination *per capita* correlated with testing *per capita* in the first three months of onset. More coordinated national policies could strengthen the pandemic tracking and response, and prevent us from making the same mistakes again.

Competing interests: SRP has received travel fees from Pfizer unrelated to this manuscript. DMW has received consulting fees from Pfizer, Merck, GSK, and Affinivax for work unrelated to this manuscript and is Principal Investigator on a grant from Pfizer to Yale University on a project unrelated to this manuscript

Funding: SP was supported by a NIH T32 training grant, award number 2T32AI007210-36A1 (MPI).

Authors' contributions: SP conceived of the study, extracted the data, ran the analyses, and wrote the first draft of the paper. DMW contributed to the design of the study and revisions of the manuscript and plotting.

Acknowledgements: We would like to thank the state health departments in the United

States and the District of Columbia and the COVID Tracking Project for providing data

used for these analyses.

Authors' information: stephanie.perniciaro@yale.edu

References

1. WHO Director-General's opening remarks at the media briefing on COVID-19 - 11

March 2020. https://www.who.int/dg/speeches/detail/who-director-general-s-

opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020.

2. Schuchat, A. Public Health Response to the Initiation and Spread of Pandemic

COVID-19 in the United States, February 24–April 21, 2020. *MMWR Morb. Mortal.*

Wkly. Rep. **69**, (2020).

3. Haffajee, R. L. & Mello, M. M. Thinking Globally, Acting Locally — The U.S.

Response to Covid-19. *N. Engl. J. Med.* **382**, e75 (2020).

4. CDC. COVID Data Tracker. *Centers for Disease Control and Prevention*

https://covid.cdc.gov/covid-data-tracker (2020).

5. Shear, M. D. *et al.* The Lost Month: How a Failure to Test Blinded the U.S. to Covid-

19. *The New York Times* (2020).

6. CNN, P. L. Hogan: Maryland protected coronavirus tests it secured from South

Korea 'like Fort Knox'. *CNN* https://www.cnn.com/2020/04/30/politics/larry-hogan-

coronavirus-masks-national-guard/index.html.

7. Walsh, J. D. How Reliable Are COVID-19 Tests? *Intelligencer*

https://nymag.com/intelligencer/2020/06/how-accurate-are-covid-19-tests.html

(2020).

8. Remarks by President Trump After Tour of the Centers for Disease Control and

Prevention | Atlanta, GA. *The White House* https://www.whitehouse.gov/briefings-

statements/remarks-president-trump-tour-centers-disease-control-prevention-

atlanta-ga/.

9. Georgia Governor And The Mayor Of Atlanta In Turf War Over COVID-19

Restrictions. *NPR.org* https://www.npr.org/sections/coronavirus-live-

updates/2020/07/10/889930319/georgia-governor-and-the-mayor-of-atlanta-in-turf-

war-over-covid-19-restrictions.

10. 'It's Like Having No Testing': Coronavirus Test Results Are Still Delayed - The New

York Times. https://www.nytimes.com/2020/08/04/us/virus-testing-delays.html.

11. HaddenKristie *et al.* Addressing Homelessness and Covid-19 Quarantine: A

Streamlined Assessment and Referral Process. *NEJM Catal. Innov. Care Deliv.*

(2020).

12. KABC. Wait times at Dodger Stadium vaccination site reach 5 hours as multiple

issues plague distribution. *ABC7 Los Angeles* https://abc7.com/9875620/ (2021).

13. Milwaukee, U. Aurora Medical Center – Grafton COVID-19 vaccination update.

Urban Milwaukee https://urbanmilwaukee.com/pressrelease/aurora-medical-center-

grafton-covid-19-vaccination-update/.

14. State COVID-19 Vaccine Priority Populations. *KFF* https://www.kff.org/other/state-

indicator/state-covid-19-vaccine-priority-populations/ (2021).

15. Dooling, K. The Advisory Committee on Immunization Practices' Updated Interim

Recommendation for Allocation of COVID-19 Vaccine — United States, December

2020. *MMWR Morb. Mortal. Wkly. Rep.* **69**, (2021).

16. Weinberger, D. M. *et al.* Estimation of Excess Deaths Associated With the COVID-

19 Pandemic in the United States, March to May 2020. *JAMA Intern. Med.* (2020)

doi:10.1001/jamainternmed.2020.3391.

17. Tolbert, J. & 2021. The COVID-19 "Vaccination Line": An Update on State

Prioritization Plans. *KFF* https://www.kff.org/coronavirus-covid-19/issue-brief/the-

covid-19-vaccination-line-an-update-on-state-prioritization-plans/ (2021).

18. Painter, E. M. Demographic Characteristics of Persons Vaccinated During the First

Month of the COVID-19 Vaccination Program — United States, December 14,

2020–January 14, 2021. *MMWR Morb. Mortal. Wkly. Rep.* **70**, (2021).

19. Weintraub, R. *et al.* COVID-19 vaccine delivery: an opportunity to set up systems for

the future. *Gates Open Res.* **4**, 182 (2020).

20. Maani, N. & Galea, S. COVID-19 and Underinvestment in the Public Health

Infrastructure of the United States. *Milbank Q.* **98**, (2020).

21. The public's role in COVID-19 vaccination: Human-centered recommendations

to enhance pandemic vaccine awareness, access, and acceptance in the United

States | Elsevier Enhanced Reader.

https://reader.elsevier.com/reader/sd/pii/S0264410X20313682?token=F69C9B86CF

A519E9A91CC75BC7680AF790607815EBFA8FC91DE8BD626315D8A40A89248A

BC4EF1A7CB0AA07145B9719D doi:10.1016/j.vaccine.2020.10.059.

Figure 1. SARS-CoV-2 testing guidelines by state. Testing recommendations for the general public was scaled as follows: 0 = criterion not included, 1 = negative influenza test required, 2 = all symptomatic people, 3 = all people, including asymptomatic people. Testing guidelines for special risk groups are shown in Supplemental Figure 1.

	Mar	Apr	May	Jun	Jul			Mar	Apr	May	Jun	Jul
AL	2	0	0	0	0		MO	0	0	0	0	3
AK	2	2	2	2	2		MT	2	0	2	2	2
AZ	0	0	2	0	0		NE	0	0	2	2	3
AR	2	2	2	2	2		NV	0	0	3	3	3
CA	2	2	3	2	2		NH	0	0	3	3	3
CO	2	2	2	2	2		NJ	2	0	2	3	3
CT	0	2	2	2	2		NM	2	2	2	3	3
DE	0	2	2	3	3		NY	1	1	2	3	3
DC	0	2	2	2	2		NC	1	1	2	2	2
FL	0	2	2	2	2		ND	0	2	2	2	2
GA	0	2	3	3	3		OH	0	0	0	0	3
HI	0	0	0	2	2		OK	0	0	0	0	0
ID	2	0	2	2	2		OR	0	2	2	2	2
IL	0	0	0	0	0		PA	0	2	2	2	3
IN	0	0	0	0	3		RI	0	2	2	2	2
IA	0	0	0	0	0		SC	0	1	2	2	2
KS	1	0	0	0	0		SD	0	0	0	0	0
KY	2	0	0	0	0		TN	0	2	3	3	3
LA	1	2	2	2	2		TX	0	2	2	2	2
ME	2	0	0	2	2		UT	0	2	2	2	2
MD	2	0	2	2	3		VT	2	2	3	3	3
MA	0	0	0	2	2		VA	1	2	2	2	2
MI	0	0	2	2	2		WA	0	2	2	2	2
MN	0	0	2	2	2		WV	1	2	3	3	3
MS	2	0	2	2	2		WI	0	0	2	2	2
							WY	1	0	2	2	2

Figure 2. SARS-CoV-2 tests per 1000 population in the United States (A) and the percentage of positive SARS-CoV-2 tests (B), March-July 2020. Each state has a line; colors of lines are determined by geographic region per the inserted map.

370 Figure 3. SARS-CoV-2 testing recommendation scores (A) and SARS-CoV-2 tests per

371 1000 people (B) from March-July, 2020 in the United States.

Figure 4. Association between the rate of testing for SARS-CoV-2 (per capita) and the

completeness of death reporting, as measured by the ratio between reported COVID-19

deaths and excess deaths due to pneumonia/influenza/coronavirus (PIC). The line +/-

95% confidence intervals represents the trend as estimated from a generalized additive

model. The size of the circles is proportional to the number of excess deaths due to PIC

in each state. The color depicts the month March-July) – earlier months are darker colors;

later months are lighter colors.

Figure 5.

COVID-19 vaccine doses administered per 100,000 population in the United States,

December 2020-February 2021. Each state has a line; colors of lines are determined by

geographic region per the inserted map.

Figure 6. Count of states in each phase of COVID-19 vaccine administration rollout, December 2020-February 2021. Darker colors represent states that were at least as inclusive as ACIP with their vaccine prioritization guidelines; pastel colors represent states that were more restrictive than ACIP guidelines.

398   Supplementary Materials

400   Supplemental Table 1. State health department websites, used as the source to

401   determine state-level COVID-19 testing guidelines.

State	Source(s)
AL	https://www.alabamapublichealth.gov/covid19/patients.html
AK	http://dhss.alaska.gov/dph/Epi/id/SiteAssets/Pages/HumanCoV/AKCOVIDTestingGuidance.pdf
AZ	https://www.azdhs.gov/documents/preparedness/epidemiology-disease-control/infectious-diseases-services/coronavirus/arizona-state-public-health-laboratory-testing-matrix.pdf
AR	https://www.healthy.arkansas.gov/programs-services/topics/covid-19-guidance-for-getting-tested
CA	https://www.cdph.ca.gov/Programs/CID/DCDC/Pages/COVID-19/Expanding-Access-to-Testing-Updated-Interim-Guidance-on-Prioritization-for-COVID-19-Laboratory-Testing-0501.aspx
CO	https://covid19.colorado.gov/covid-19-in-colorado/about-covid-19/testing-for-covid-19
CT	https://portal.ct.gov/Coronavirus/Covid-19-Knowledge-Base/COVID-19-Testing
DE	https://coronavirus.delaware.gov/testing/testing-frequently-asked-questions/
DC	https://coronavirus.dc.gov/testing
FL	https://floridahealthcovid19.gov/health-care-providers/
GA	https://dph.georgia.gov/covid-19-testingdirect-patient-lines
HI	https://health.hawaii.gov/coronavirusdisease2019/for-clinicians/evaluating-puis/

ID	https://rebound.idaho.gov/wp-content/uploads/testing-recommendations.pdf	
IL	https://www.dph.illinois.gov/topics-services/diseases-and-conditions/diseases-a-z-list/coronavirus/health-care-providers	
IN	https://www.coronavirus.in.gov/files/IN-COVID-19_FAQforPublic%207.3.20.pdf	https://www.coronavirus.in.gov/files/IN-COVID-19_FAQforPublic%206.13.20.pdf
IA	https://covidtesting.shl.uiowa.edu/	
KS	https://www.coronavirus.kdheks.gov/DocumentCenter/View/393/COVID-19-Testing-Approval-Form-fillable-PDF---4-21-20	
KY	https://chfs.ky.gov/Pages/search.aspx?terms=covid+test&affiliateId=CHFS	https://chfs.ky.gov/agencies/dph/covid19/guidancetohealthcareprovidersregardingCOVID19statelaboratorytesting03102020.pdf
LA	http://ldh.la.gov/assets/oph/Coronavirus/resources/TestingIsolationGuidelines.pdf	
ME	https://www.maine.gov/tools/whatsnew/index.php?topic=DHS+Press+Releases&id=2657874&v=dhhs_article_2020	
MD	https://phpa.health.maryland.gov/Documents/coronavirus_testing_FAQ.pdf	
MA	https://www.mass.gov/info-details/about-covid-19-testing#should-i-be-tested?-	
MI	https://www.michigan.gov/coronavirus/0,9753,7-406-98163-530157--,00.html	
MN	https://www.health.state.mn.us/diseases/coronavirus/symptoms.html#test	
MS	https://msdh.ms.gov/msdhsite/_static/14,21866,420.html	
MO	https://health.mo.gov/emergencies/ert/alertsadvisories/pdf/update42220.pdf	
MT	https://dphhs.mt.gov/publichealth/cdepi/diseases/coronavirusmt	

NE	http://dhhs.ne.gov/Documents/COVID-19%20Guidance%20to%20Public%20and%20Testing.pdf	https://covid19testingne.co1.qualtrics.com/jfe/form/SV_cJa8ezYYRQfZRCl
NV	https://nvhealthresponse.nv.gov/find-covid-19-testing-in-nevada/	
NH	https://business.nh.gov/DOS_COVID19Testing/	
NJ	https://covid19.nj.gov/pages/testing	
NM	https://cv.nmhealth.org/public-health-screening-and-testing/location-details/	
NY	https://coronavirus.health.ny.gov/covid-19-testing	
NC	https://covid19.ncdhhs.gov/about-covid-19/testing	
ND	https://www.health.nd.gov/sites/www/files/documents/Files/MSS/coronavirus/2020_6_16_COVID_PCR_Guidance.pdf	
OH	https://coronavirus.ohio.gov/wps/portal/gov/covid-19/families-and-individuals/MHSF/	
OK	https://coronavirus.health.ok.gov/articles/criteria-guide-evaluation-and-laboratory-testing-covid-19-clinicians-and-laboratories	
OR	https://sharedsystems.dhsoha.state.or.us/DHSForms/Served/le2267.pdf	
PA	https://www.health.pa.gov/topics/disease/coronavirus/Pages/Guidance/Testing-Factsheet.aspx	
RI	https://health.ri.gov/covid/testing/	
SC	https://www.scdhec.gov/sites/default/files/media/document/10466-DHU-06-19-2020-COVID-19.pdf	
SD	https://doh.sd.gov/documents/COVID19/COVID_TestingFacts.pdf	
TN	https://www.tn.gov/governor/covid-19/get-tested.html	
TX	https://dshs.texas.gov/coronavirus/public-health.aspx	

UT	https://coronavirus-download.utah.gov/Health/UDOH%20HAN%20COVID_04142020_Statewide_Guidance%20(1).pdf
VT	https://www.healthvermont.gov/response/coronavirus-covid-19/testing-covid-19
VA	https://www.vdh.virginia.gov/coronavirus/health-professionals/vdh-updated-guidance-on-testing-for-covid-19/

WA	https://www.doh.wa.gov/Emergencies/NovelCoronavirusOutbreak2020COVID19/TestingforCOVID19	https://www.doh.wa.gov/Portals/1/Documents/1600/coronavirus/Interim-2019NovelCoronavirusQuicksheetProviders.pdf

WV	https://dhhr.wv.gov/COVID-19/pages/testing.aspx
WI	https://www.dhs.wisconsin.gov/covid-19/testing.htm
WY	https://health.wyo.gov/department-offers-testing-update-advice-for-wyoming/

Supplemental Figure 1. State-level SARS-CoV-2 testing recommendations for selected

risk groups, March-July 2020. Criteria for (A) Healthcare/Essential Workers, (B) High-

Risk Populations, (C) Long-term Care Facility Residents or Workers (D) Associates of

Known COVID-19 case, (E) Travelers to High-Risk Areas, and (F) Hospitalized Patients

were scaled as follows: 0 = Criterion not listed, 1 = Negative influenza test required, 2 =

Symptomatic individual, 3= Any individual.