

1 **Sleep quality in the chronic stage of concussion is associated with poorer recovery: A**
2 **systematic review**

3 Rebecca Ludwig, OT,¹ Eryen Nelson, MPH,¹ Prasanna Vaduvathiriyana, MLS, AHIP,² Michael
4 A. Rippee MD,³ Catherine Siengasukon, PhD¹

5 ¹Department of Physical Therapy and Rehabilitation Science, University of Kansas Medical
6 Center, 3901 Rainbow Blvd, Mailstop 2002, Kansas City, KS, 66160

7 ² A.R. Dykes Library, University of Kansas Medical Center, 2100 W 39th Ave, Mail Stop 1050,
8 Kansas City, KS 66103

9 ³ Department of Neurology, University of Kansas Medical Center, 3901 Rainbow Blvd, Mailstop
10 2012, Kansas City, KS 66160

11

12 **Corresponding Author:**

13 Catherine Siengasukon, PT, PhD
14 University of Kansas Medical Center
15 Physical Therapy and Rehabilitation Sciences
16 3901 Rainbow Blvd, Mailstop 2002
17 Phone: +1-913-588-6913
18 Fax: +1-913-588-4568
19 E-mail: csiengasukon@kumc.edu

20

21 **Keywords:** sleep, recovery, mild traumatic brain injury, concussion, post-concussion syndrome,
22 recovery, systematic review

23

1 **Abstract**

2 *Objective:* To examine the association between sleep quality during the chronic stage of
3 concussion and post-concussion outcomes.

4 *Literature Survey:* Literature searches were performed during July 1st to August 1st, 2019 in
5 selected databases along with searching grey literature. Out of the 733 results, 702 references
6 were reviewed after duplicate removal.

7 *Methodology:* Three reviewers independently reviewed and consented on abstracts meeting
8 eligibility criteria (n=35). The full-text articles were assessed independently by two reviewers.
9 Consensus was achieved, leaving four articles. Relevant data from each study was extracted
10 using a standard data-extraction table. Quality appraisal was conducted to assess potential bias
11 and the quality of articles.

12 *Synthesis:* One study included children (18-60 months) and three studies included adolescents
13 and/or adults (ranging 12 years to 35). The association between sleep and cognition (two
14 studies), physical activity (one study), and emotion symptoms (one study) was examined. Sleep
15 quality was associated with decreased cognition and emotional symptoms, but not with meeting
16 physical activity guidelines 6 months post-concussion injury.

17 *Conclusions:* The heterogeneity in age of participants and outcomes across studies and limited
18 number of included studies made interpretations difficult. Future studies may consider if
19 addressing sleep quality following concussion will improve outcomes.

20

21

22

23

1 Introduction

2 The prevalence of a concussion is 1.7- 3.8 million people each year in the United States.¹⁻
3 ³ Concussion injuries occur in a variety of settings including motor vehicle accidents, sports,
4 recreation, falls, work environments, and war blast zones^{4,5} and are commonly a result of a high
5 magnitude of force placed to the head, neck, or face.^{2,6} Symptoms caused by a concussion are
6 often categorized into four domains: somatic, cognitive, mood disruption, and sleep
7 dysregulation.^{3,7} *Somatic symptoms* include vertigo, sensitivity to light and sound, nausea,
8 vomiting, headaches, and migraines.⁷ Slower processing speed, decreased focus, disrupted short
9 term memory, and difficulty concentrating are commonly reported impairments in *cognition*
10 following a concussion.⁷ *Mood disruptions* after a concussion consist of an increase or onset of
11 anxiety, depression, or irritability.⁷ *Sleep dysregulation* often manifests in the form of
12 hypersomnia, insomnia, or circadian rhythm disruption.⁷ Acutely following the injury, headache,
13 fatigue, dizziness, and difficulty with information processing are the most endorsed symptoms.^{8,9}
14 Symptoms that are commonly reported at follow-up visits are sleep disturbances, frustration,
15 forgetfulness, and fatigue.⁸

16 Recovery time frames after the concussion injury are different for each individual and
17 commonly are grouped in the following stages: *acute stage* (the first seven days), *sub-acute*
18 *stage* (8-89 days), and *chronic stage* (≥ 90 days) following the concussion injury.¹⁰ Symptoms
19 resolve for 80-90% of individuals 1-3 weeks post- injury.^{1,4,6,7} However, for the remaining 10-
20 20% of individuals, symptoms persist for months to years post injury.¹¹⁻¹⁴

21 While symptom severity and duration are difficult to predict,¹⁵ there are certain factors
22 from prior to the injury, the injury itself, and after the injury that increase the risk of prolonged
23 symptoms. Pre-injury risk factors include age (0-4 years old and 65+ years old), history of

1 learning disabilities, migraines, previous concussions, psychiatric conditions, female gender,
2 lower cognitive reserve, lower socioeconomic status, substance abuse, and premonitory sleep
3 disturbance.¹⁶⁻²⁰ Concussion-specific risk factors for prolonged symptoms consist of loss of
4 consciousness, injuries to other areas of the body, lower (<15) Glasgow Coma score, and a
5 higher velocity mechanism of injury.^{16,17} Post-injury anxiety, depression, post-traumatic stress
6 disorder, migraine, sleep disturbances, fogginess, difficulty concentrating, vestibular
7 dysfunction, and ocular motor dysfunction post-injury have also been associated with prolonged
8 symptom recovery.^{13,14,21} Variability in length of recovery following a concussion is likely due to
9 multiple reasons, including fluctuation of cellular effects, extent of axonal injury, location of
10 impact, and the number and severity of symptoms following a concussion.²²⁻²⁵ Research is still
11 investigating why various risk factors and combinations of risk factors contribute to delayed
12 concussion recovery.

13 Sleep disturbance prior to concussion as well as the development of sleep disturbances
14 following a concussion have also been associated with prolonged concussion symptoms. Poor
15 sleep quality is related to increased pain,²⁶ more frequent headaches,²⁷ and vestibular issues.²⁸
16 Our recently published systematic review found that sleep disturbances in the acute phase have
17 been shown to be predictive of prolonged symptom recovery following a concussion.¹⁷ Short
18 sleep duration and poor quality during the acute and subacute stage of concussion are associated
19 with a range of prolonged post-concussion symptoms including poorer cognitive function,
20 reduced productivity at work or school, decreased social engagement, increased pain, daytime
21 sleepiness, depression, and anxiety.^{17,29,30}

22 The association between sleep and post-concussion symptoms in the acute and sub-acute
23 stages has been well documented.^{19,31-36} There remains a gap in the literature assessing the

1 association between sleep and post-concussion symptoms in the chronic phase of concussion
2 recovery. Therefore, the purpose of this systematic review is to examine the association between
3 sleep disturbance and post-concussive symptoms at >6 months post-injury.

4 Methods

5 To conduct this systematic review, literature searches were performed between July 1st to
6 August 1st, 2019 in the following databases: Ovid MEDLINE(R), Cumulative Index of Nursing
7 & Allied Health Literature (CINAHL), and Web of Science - Science Citation Index Expanded.
8 relevant cited references and grey literature including clinical trials and conference papers were
9 searched. The review was limited to studies published in the English language using the
10 databases' language limit feature. Publication date limits to the search were not applied.

11 A combination of keywords and controlled vocabulary terms were used in Medline and
12 CINAHL to execute the searches in the databases. Medical subject headings including “brain
13 concussion”, “brain injuries”, “traumatic brain injuries”, “sleep”, “pain”, “pain management”,
14 and keywords (text-words) such as “mild traumatic injuries”, “concussion”, “post-concussion”
15 were used in various combinations to increase recall.

16 PRISMA guidelines were followed for the article selection. Studies included in the
17 systematic review, had to meet the following inclusion criteria: (1) included people with
18 concussive or mild traumatic brain injury; (2) assessed sleep using subjective or objective
19 measures > 6 months post-concussion; (3) used a prospective or cross-sectional study design; (4)
20 reported other non-sleep outcomes > 6 months post-concussion; and (5) included statistical
21 analyses to relate sleep outcomes to other reported outcome(s). Studies were excluded if they: (1)
22 included animals; (2) included participants with moderate and/or severe traumatic brain injury;

1 (3) included participants with other neurologic conditions; (4) used a study design other than
2 prospective or cross-sectional; (5) were irrelevant to the objective of the systematic review; (6)
3 were published in a language other than English; or (7) included a duplicate data set.

4 Three reviewers (R.L., K.S., A.P.) assessed titles and abstracts of all potentially eligible
5 articles for inclusion/exclusion criteria to identify which titles/abstracts met those criteria and
6 would be considered for the systematic review. The full-text article of all considered studies was
7 then retrieved and reviewed independently by two reviewers (R.L., E.N.) to establish article
8 relevance. The reviewers came to a consensus on which abstracts met the inclusion/exclusion
9 criteria and which full-text articles were included in the systematic review.

10 *Data Extraction*

11 Relevant data were extracted from each included study by two reviewers (R.L., E.N.)
12 using a data-extraction table. Extracted information included the author(s), year published, study
13 location, study design, sample size, participant demographic characteristics, time since injury,
14 measures used to assess sleep, measures used to assess other outcomes, and results.

15 *Quality Assessment*

16 The risk of potential bias as well as the overall quality of the included studies was
17 assessed using the *Joanna Briggs Institute Critical Appraisal Checklist for Analytical Cross*
18 *Sectional Studies*.³⁷ The checklist is comprised of eight questions regarding defined study
19 inclusion criteria, description of study participants, exposure validity and reliability, standard
20 criteria for measurement conditions, identification of confounding factors, strategies for dealing
21 with confounding factors, valid and reliable outcomes measured, and appropriate statistical
22 analysis. All articles were scored independently by two reviewers (R.L., E.N.) as “yes”, “no”,

1 “unclear”, or “not applicable” for each item on the checklist. The reviewers met to reach
2 consensus.

3 Results

4 The search strategy generated a total of 733 articles. There were 31 duplicates removed,
5 leaving 702 titles/abstracts to review. After the review of title/abstracts for inclusion/exclusion
6 criteria, 662 articles were removed. A total of 40 full-text articles were reviewed. Thirty-six
7 articles were removed due to: reasons related to unclear assessment time since injury (n=22), no
8 statistical analyses conducted to determine the association between sleep and other post-
9 concussion outcomes (n=11), and inability to locate the full-text article (n=3). A total of four
10 articles met the criteria to be included in this review (Figure 1).

11 The studies in this review are summarized in the data extraction table (Table 1) and the
12 risk of bias is reported in the quality appraisal table (Table 2). The number of participants in each
13 study ranged from 49-232 individuals. One article included pediatric patients (18-60 months)³⁸
14 and three articles included adolescents and adults (>12 years old).³⁹⁻⁴¹

15 Landry-Roy et al. (2018)³⁸ evaluated the relationship between sleep and executive
16 function in preschool children. Children aged 18-60 months (n=167) who either had a
17 concussion (n=84) or were typically developing children (TDC) (n=83) participated. The results
18 indicated there was a significant interaction of children’s group (concussion or TDC) and the
19 Children Behavior Checklist (CBCL) sleep scores on the Delay of Gratification assessment
20 (p=.05) at 6-months. Suggesting that children with a concussion and sleep disturbance have
21 disrupted executive function. There was a significant group difference for the Delay of
22 Gratification assessment performance for children with higher levels of sleep problems on the
23 CBCL (p=.03). Children with a concussion had decreased performance compared to TDC on the

1 delay of gratification if they had higher levels of sleep problems on the CBCL. Similarly, in a
2 post hoc analysis there was a significant group difference on Conflict Scale performance for
3 children with shorter sleep duration ($p=.004$). Thus, children with a concussion exhibited poorer
4 conflict scale performance than TDC only if they had shorter nighttime sleep duration. There
5 was a small subgroup of children ($n=68$; 33 concussion 35 TDC) who wore actigraphy but no
6 association between actigraphy and sleep diary and executive function was found. The results of
7 this study indicate that the combination of a concussion and poor sleep result in poorer executive
8 functions in pre-school aged children.

9 The study by Dean and Sterr (2013)³⁹ evaluated working memory and information
10 processing speed in adults ($n=36$; 20-30 years of age) at least one-year post concussion injury
11 compared to a non-head injury control group ($n=36$; 20-30 years of age). Participants were
12 divided into one of four groups: 1. Concussion + Post-Concussion Syndrome (PCS); 2.
13 Concussion – PCS; 3. Control (non-head injury) + PCS; 4. Control (non-head injury) - PCS.
14 Nighttime sleep quality was lower in the Concussion + PCS participants compared to the
15 Concussion - PCS and Control - PCS ($p=.018$). Mean Pittsburg Sleep Quality Index (PSQI)
16 scores for the Concussion + PCS ($8.6 \pm .8$) and the Control + PCS ($6.6 \pm .7$) were indicative of
17 poor nighttime sleep. There was a significant correlation for all individuals with a concussion
18 between lower PSQI scores and poorer performance on the N Back (working memory) and
19 Paced Visual Serial Addition Task (information processing speed) ($Rho=.62$, $p<.001$).

20 Theadom et al. (2018)⁴⁰ evaluated how sleep predicts post-concussion symptoms and
21 community and social engagements in adolescents and adults ($n =232$; age ≥ 16) from the
22 population based BIONIC study^{31,32} four years after the initial concussion injury. The study used
23 the Rivermead Post Concussion Symptoms Questionnaire to assess post-concussion symptoms

1 and the PSQI for sleep quality. The concussion group reported significantly more cognitive
2 symptoms four years post-injury compared to controls. There was no difference between the
3 concussion and control groups on somatic and emotional scores. Regression modeling revealed
4 that sleep quality was a significant predictor for emotional symptoms on the Rivermead Post
5 Concussion symptom questionnaire ($p=.02$) but not for cognitive ($p=.99$) or somatic symptoms
6 ($p=.06$). In particular, poorer sleep quality was a predictor of higher anxiety, depression, and
7 irritable mood symptoms 4 years following concussion.

8 Van Markus-Doornbosch et al. (2019)⁴¹ evaluated the relationship between sleep quality
9 and physical activity level in adolescents and adults who had a concussion. A total of 49
10 participants (12-25 years old) who had a concussion 6-18 months prior were enrolled in the
11 study. If the participant was < 18 years old, a parent completed the questionnaires. Physical
12 activity was assessed by the Activity Questionnaire for Adults and Adolescents, and sleep quality
13 was assessed using the PSQI. Twenty-five out of 49 (51%) participants did not meet the Dutch
14 Health Enhancing Physical Activity (D-HEPA) recommendations. The median PSQI total score
15 was 4.5 for the entire study group with no significant association with meeting or not meeting the
16 D-HEPA recommendations (OR .99, 95% CI .82,1.19).

17 The two contributing factors for bias were not adequately controlling for confounding
18 factors and not using an appropriate assessment tool for measuring sleep outcomes (Table 2). In
19 two out of the four studies, the confounding factors were not adequately described or controlled
20 for in the statistical analysis. One out of the four studies did not use an assessment that was
21 specific to assess sleep outcomes. The sleep portion of the CBCL used in Landry-Roy et al. has
22 good convergent validity, but there is debate on if this assessment is specific to sleep when
23 compared to other pediatric sleep assessments.

1 Discussion

2 This systematic review found that poor sleep quality is associated with concussion
3 symptoms 6 months to 4 years following the injury. Poor sleep quality was associated with
4 impaired executive function in preschoolers at 6 months following concussion,³⁸ and working
5 memory and information processing speed in adults.³⁹ Poor sleep quality was also found to be a
6 predictor of emotional symptoms in adults 4 years post-concussive injury.⁴⁰ However, one study
7 did not find a significant association between sleep quality and meeting physical activity
8 guidelines in adolescents.⁴¹ Only four articles met the criteria for inclusion in this systematic
9 review, thus limiting the scope of interpretation, while simultaneously highlighting the lack of
10 research on how poor sleep quality and sleep disturbances are associated with prolonged
11 symptoms in individuals with chronic concussion.

12 Two articles in this systematic review examined the association between sleep and
13 executive function, working memory, and information processing speed.^{38,39} Cognitive function
14 has been shown to be impaired in the acute and subacute stages of concussion.^{14,16,42} Symptoms
15 such as taking longer to think, decreased focus, and decreased processing speed are commonly
16 reported in the acute and subacute stages.^{14,16,42} What was observed from this systematic review
17 was that executive function, working memory, and processing speed was still impaired.^{38,39}
18 Sleep disturbance and cognitive disturbance are ranked as the most common complaints of
19 patients at follow-up visits.¹⁶ The association between sleep disturbance and cognitive
20 disturbance is not a surprise as sleep disturbance can influence cognition.⁴³ Individuals
21 recovering from a TBI have shown sleep fragmentation and altered sleep architecture on
22 polysomnography.⁴⁴⁻⁴⁶ More specifically, individuals recovering from a TBI spend more time in
23 slow wave sleep (SWS) and less time in rapid eye movement (REM) sleep.⁴⁴ While SWS is

1 necessary for cellular and structural recovery, REM is needed for information processing and
2 memory consolidation. Therefore, a reduction in REM could contribute to impairments of
3 executive functioning, working memory, and information processing that are often experienced
4 by individuals following concussion.⁴⁷⁻⁵¹ Psychological factors such as depression, anxiety, and
5 mood influence cognition as well.⁵²⁻⁵⁴ Sleep disturbances and psychological factors are
6 interrelated, so it is likely a combination of sleep disturbance and psychological factors impact
7 cognitive function following a concussion⁵⁵

8 Sleep quality was also found to be predictive of emotional symptoms including anxiety,
9 depression, and irritability 4 years following a concussion.⁴⁰ This finding supports evidence that
10 chronic poor sleep and insomnia can contribute to the development of depression,
11 anxiety, irritability.⁵⁶⁻⁶⁰ Furthermore, screening and treating insomnia reduces concomitant
12 symptoms of depression, anxiety, and stress.⁶¹⁻⁶³ While sleep quality and insomnia contributes to
13 emotional symptoms, research also shows that emotional symptoms can contribute to the
14 development of sleep issues.⁶⁴ A negative mood (anxious, stressed, tense, sad, or irritable) is
15 more common in individuals with poor sleep.^{65,66} There is a strong relationship between the
16 development of anxiety and the development of poor sleep and insomnia as many core features
17 (intense worry or fear, and avoidance behavior) are the same.⁶⁷ Health care providers should
18 screen for both sleep and emotional issues in people following concussion, and more studies are
19 needed to determine optimal prioritization of treatment for sleep and emotional issues.

20 One study found that meeting physical activity guidelines was not associated with sleep
21 quality six months post- injury.⁴¹ This is surprising as physical activity and sleep have been
22 shown to have a reciprocal relationship.⁶⁸ If an individual does not have a quality night of sleep,
23 there is decreased motivation to be physically active the following day.⁶⁸ Also, physical activity

1 improves sleep duration and quality by increasing sleep drive.⁶⁹ The lack of association in Van
2 Markus-Doornbosch⁴¹ might be due to the mean PSQI score was 4.5, indicating the
3 participations on average had good quality sleep. Also, the authors assessed the relationship
4 between sleep quality and adherence to physical activity guidelines, not physical activity
5 specifically. Van Markus-Doornbosch⁴¹ reported that within their sample 51% of individuals
6 following concussion met physical activity guidelines which is comparable to the national
7 average in the United States. For adults in the United States, 44.8% of individuals meet physical
8 activity guidelines.⁷⁰ Therefore, it is questionable on if guidelines for physical activity matter
9 when assessing sleep in individuals with a concussion.

10 The limitations within the articles for this systematic review is due to not controlling for
11 confounding factors and the specific sleep assessments used. Within the studies reviewed two out
12 of the four articles did not describe confounding factors or address them within their statistical
13 analysis. This made interpretation of the results challenging as it is unclear what effect the
14 confounding factors had on the results described. Additionally, the sleep portion of the CBCL
15 used in Landry-Roy et al., 2018 had good convergent validity, but it is of question on if this
16 assessment is specific enough to assess sleep in comparison to other pediatric sleep
17 questionnaires.

18 This systematic review does not come without limitations. One limitation is there are
19 only four articles included within this review, limiting the breadth and depth for interpretation.
20 Additionally, the studies included in this systematic review includes a variety of outcomes. The
21 heterogeneity limits interpretation as each study addresses different outcomes (cognition,
22 physical activity, and emotional symptoms). There needs to be more studies evaluating these

1 outcomes to strengthen the knowledge base on how sleep affects cognition, physical activity, and
2 emotional symptoms.

3 This systematic review indicates that poor sleep quality is associated with impairment in
4 executive function,³⁸ working memory, information processing,³⁹ and emotional symptoms,⁴⁰
5 but not physical activity in individuals >6 months post-concussion. Additional studies are needed
6 to more thoroughly understand how sleep is associated with various post-concussion symptoms
7 in the chronic stage. Also, investigating how addressing sleep issues may impact symptom
8 resolution and recovery in people with a concussion is warranted.

9

10

11

12

13

14

15

16

17

18

19

20

1 **Acknowledgements:** Kelli Stallbaumer and Alexandra Ptacek assisted with screening abstracts
2 for full text review. This paper is partially funded through the T-32 neurologic rehabilitation
3 training grant through the University of Kansas Medical Center.

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

References

- 1
- 2
- 3 1. Mahooti N. Sports-Related Concussion: Acute Management and Chronic Postconcussive
- 4 Issues. *Child and adolescent psychiatric clinics of North America*. 2018;27(1):93-108.
- 5
- 6 2. Duclos C, Dumont M, Wiseman-Hakes C, et al. Sleep and wake disturbances following
- 7 traumatic brain injury. *Pathologie-biologie*. 2014;62(5):252-261.
- 8
- 9 3. Bramley H, Hong J, Zacko C, Royer C, Silvis M. Mild Traumatic Brain Injury and Post-
- 10 concussion Syndrome: Treatment and Related Sequela for Persistent Symptomatic
- 11 Disease. *Sports Med Arthrosc Rev*. 2016;24(3):123-129.
- 12
- 13 4. Tator CH. Concussions and their consequences: current diagnosis, management and
- 14 prevention. *CMAJ : Canadian Medical Association journal = journal de l'Association*
- 15 *medicale canadienne*. 2013;185(11):975-979.
- 16
- 17 5. Langlois JA, Rutland-Brown W, Wald MM. The epidemiology and impact of traumatic
- 18 brain injury: a brief overview. *The Journal of head trauma rehabilitation*.
- 19 2006;21(5):375-378.
- 20
- 21 6. Chancellor SE, Franz ES, Minaeva OV, Goldstein LE. Pathophysiology of Concussion.
- 22 *Semin Pediatr Neurol*. 2019;30:14-25.
- 23
- 24 7. McCrea MA, Nelson LD, Guskiewicz K. Diagnosis and Management of Acute
- 25 Concussion. *Physical medicine and rehabilitation clinics of North America*.
- 26 2017;28(2):271-286.
- 27
- 28 8. Eisenberg MA, Meehan WP, 3rd, Mannix R. Duration and course of post-concussive
- 29 symptoms. *Pediatrics*. 2014;133(6):999-1006.
- 30
- 31 9. Howell DR, O'Brien MJ, Beasley MA, Mannix RC, Meehan WP, 3rd. Initial somatic
- 32 symptoms are associated with prolonged symptom duration following concussion in
- 33 adolescents. *Acta paediatrica (Oslo, Norway : 1992)*. 2016;105(9):e426-432.
- 34
- 35 10. Krainin BM, Forsten RD, Kotwal RS, Lutz RH, Guskiewicz KM. Mild traumatic brain
- 36 injury literature review and proposed changes to classification. *J Spec Oper Med*.
- 37 2011;11(3):38-47.
- 38
- 39 11. Daneshvar DH, Riley DO, Nowinski CJ, McKee AC, Stern RA, Cantu RC. Long-term
- 40 consequences: effects on normal development profile after concussion. *Physical medicine*
- 41 *and rehabilitation clinics of North America*. 2011;22(4):683-700, ix.
- 42
- 43 12. Iverson GL. Network Analysis and Precision Rehabilitation for the Post-concussion
- 44 Syndrome. *Frontiers in neurology*. 2019;10:489.

- 1 13. McMahon P, Hricik A, Yue JK, et al. Symptomatology and functional outcome in mild
2 traumatic brain injury: results from the prospective TRACK-TBI study. *Journal of*
3 *neurotrauma*. 2014;31(1):26-33.
4
- 5 14. Hiploylee C, Dufort PA, Davis HS, et al. Longitudinal Study of Postconcussion
6 Syndrome: Not Everyone Recovers. *Journal of neurotrauma*. 2017;34(8):1511-1523.
7
- 8 15. Kenzie ES, Parks EL, Bigler ED, Lim MM, Chesnutt JC, Wakeland W. Concussion As a
9 Multi-Scale Complex System: An Interdisciplinary Synthesis of Current Knowledge.
10 *Frontiers in neurology*. 2017;8:513.
11
- 12 16. Tator CH, Davis HS, Dufort PA, et al. Postconcussion syndrome: demographics and
13 predictors in 221 patients. *Journal of neurosurgery*. 2016;125(5):1206-1216.
14
- 15 17. Ludwig R, D'Silva L, Vaduvathiriyam P, Rippee MA, Siengsukon C. Sleep Disturbances
16 in the Acute Stage of Concussion are Associated With Poorer Long-Term Recovery: A
17 Systematic Review. *Pm r*. 2019.
18
- 19 18. Meehan WP, 3rd, Mannix R. Pediatric concussions in United States emergency
20 departments in the years 2002 to 2006. *The Journal of pediatrics*. 2010;157(6):889-893.
21
- 22 19. Schmidt AT, Li X, Hanten GR, McCauley SR, Faber J, Levin HS. A Longitudinal
23 Investigation of Sleep Quality in Adolescents and Young Adults After Mild Traumatic
24 Brain Injury. *Cognitive and behavioral neurology : official journal of the Society for*
25 *Behavioral and Cognitive Neurology*. 2015;28(2):53-62.
26
- 27 20. Papa L, Mendes ME, Braga CF. Mild Traumatic Brain Injury among the Geriatric
28 Population. *Curr Transl Geriatr Exp Gerontol Rep*. 2012;1(3):135-142.
29
- 30 21. Collins MW, Kontos AP, Reynolds E, Murawski CD, Fu FH. A comprehensive, targeted
31 approach to the clinical care of athletes following sport-related concussion. *Knee Surg*
32 *Sports Traumatol Arthrosc*. 2014;22(2):235-246.
33
- 34 22. Giza CC, Hovda DA. The new neurometabolic cascade of concussion. *Neurosurgery*.
35 2014;75 Suppl 4:S24-33.
36
- 37 23. Barkhoudarian G, Hovda DA, Giza CC. The molecular pathophysiology of concussive
38 brain injury. *Clin Sports Med*. 2011;30(1):33-48, vii-iii.
39
- 40 24. Frati A, Cerretani D, Fiaschi AI, et al. Diffuse Axonal Injury and Oxidative Stress: A
41 Comprehensive Review. *Int J Mol Sci*. 2017;18(12).
42
- 43 25. Wiseman-Hakes C, Duclos C, Blais H, et al. Sleep in the Acute Phase of Severe
44 Traumatic Brain Injury: A Snapshot of Polysomnography. *Neurorehabilitation & Neural*
45 *Repair*. 2016;30(8):713-721.

- 1 26. Suzuki Y, Khoury S, El-Khatib H, et al. Individuals with pain need more sleep in the
2 early stage of mild traumatic brain injury. *Sleep medicine*. 2017;33:36-42.
3
- 4 27. Chaput G, Giguere JF, Chauny JM, Denis R, Lavigne G. Relationship among subjective
5 sleep complaints, headaches, and mood alterations following a mild traumatic brain
6 injury. *Sleep medicine*. 2009;10(7):713-716.
7
- 8 28. Sufrinko AM, Howie EK, Elbin RJ, Collins MW, Kontos AP. A Preliminary
9 Investigation of Accelerometer-Derived Sleep and Physical Activity Following Sport-
10 Related Concussion. *The Journal of head trauma rehabilitation*. 2018;33(5):E64-e74.
11
- 12 29. Wickwire EM, Williams SG, Roth T, et al. Sleep, Sleep Disorders, and Mild Traumatic
13 Brain Injury. What We Know and What We Need to Know: Findings from a National
14 Working Group. *Neurotherapeutics*. 2016;13(2):403-417.
15
- 16 30. Raikes AC, Schaefer SY. Sleep Quantity and Quality during Acute Concussion: A Pilot
17 Study. *Sleep*. 2016;39(12):2141-2147.
18
- 19 31. Theadom A, Cropley M, Parmar P, et al. Sleep difficulties one year following mild
20 traumatic brain injury in a population-based study. *Sleep Medicine*. 2015;16(8):926-932.
21
- 22 32. Theadom A, Starkey N, Jones K, et al. Sleep difficulties and their impact on recovery
23 following mild traumatic brain injury in children. *Brain Injury*. 2016;30(10):1243-1248.
24
- 25 33. Mollayeva T, Mollayeva S, Colantonio A. The Risk of Sleep Disorder Among Persons
26 with Mild Traumatic Brain Injury. *Current neurology and neuroscience reports*.
27 2016;16(6):55.
28
- 29 34. Farrell-Carnahan L, Franke L, Graham C, McNamee S. Subjective sleep disturbance in
30 veterans receiving care in the Veterans Affairs Polytrauma System following blast-related
31 mild traumatic brain injury. *Military medicine*. 2013;178(9):951-956.
32
- 33 35. Gosselin N, Lassonde M, Petit D, et al. Sleep following sport-related concussions. *Sleep*
34 *medicine*. 2009;10(1):35-46.
35
- 36 36. Kalmbach DA, Conroy DA, Falk H, et al. Poor sleep is linked to impeded recovery from
37 traumatic brain injury: Characterizing sleep onset insomnia symptoms and short sleep in
38 functional impairment. *Sleep*. 2018;24:24.
39
- 40 37. Moola S MZ, Tufanaru C, Aromataris E, Sears K, Sfetcu R, Currie M, Qureshi R, Mattis
41 P, Lisy K, Mu P-F. Chapter 7: Systematic reviews of etiology and risk. In: Aromataris E
42 MZ, ed. *Joanna Briggs Institute Reviewer's Manual*. 2017.
43
- 44 38. Landry-Roy C, Bernier A, Gravel J, Beauchamp MH. Executive Functions and Their
45 Relation to Sleep Following Mild Traumatic Brain Injury in Preschoolers. *Journal of the*
46 *International Neuropsychological Society*. 2018;24(8):769-780.

- 1 39. Dean PJ, Sterr A. Long-term effects of mild traumatic brain injury on cognitive
2 performance. *Frontiers in Human Neuroscience*. 2013;7:30.
3
- 4 40. Theadom A, Starkey N, Barker-Collo S, et al. Population-based cohort study of the
5 impacts of mild traumatic brain injury in adults four years post-injury. *PLoS ONE*
6 *[Electronic Resource]*. 2018;13(1):e0191655.
7
- 8 41. van Markus-Doornbosch F, Peeters E, van der Pas S, Vlieland TV, Meesters J. Physical
9 activity after mild traumatic brain injury: What are the relationships with fatigue and
10 sleep quality? *European Journal of Paediatric Neurology*. 2019;23(1):53-60.
11
- 12 42. Meares S, Shores EA, Taylor AJ, et al. The prospective course of postconcussion
13 syndrome: the role of mild traumatic brain injury. *Neuropsychology*. 2011;25(4):454-465.
14
- 15 43. Martindale SL, Morissette SB, Rowland JA, Dolan SL. Sleep quality affects cognitive
16 functioning in returning combat veterans beyond combat exposure, PTSD, and mild TBI
17 history. *Neuropsychology*. 2017;31(1):93-104.
18
- 19 44. Parcell DL, Ponsford JL, Redman JR, Rajaratnam SM. Poor sleep quality and changes in
20 objectively recorded sleep after traumatic brain injury: a preliminary study. *Archives of*
21 *physical medicine and rehabilitation*. 2008;89(5):843-850.
22
- 23 45. Schreiber S, Barkai G, Gur-Hartman T, et al. Long-lasting sleep patterns of adult patients
24 with minor traumatic brain injury (mTBI) and non-mTBI subjects. *Sleep medicine*.
25 2008;9(5):481-487.
26
- 27 46. Verma A, Anand V, Verma NP. Sleep disorders in chronic traumatic brain injury.
28 *Journal of clinical sleep medicine : JCSM : official publication of the American Academy*
29 *of Sleep Medicine*. 2007;3(4):357-362.
30
- 31 47. Vyazovskiy VV, Delogu A. NREM and REM Sleep: Complementary Roles in Recovery
32 after Wakefulness. *Neuroscientist*. 2014;20(3):203-219.
33
- 34 48. Perogamvros L, Dang-Vu TT, Desseilles M, Schwartz S. Sleep and dreaming are for
35 important matters. *Front Psychol*. 2013;4:474.
36
- 37 49. Karni A, Tanne D, Rubenstein BS, Askenasy JJ, Sagi D. Dependence on REM sleep of
38 overnight improvement of a perceptual skill. *Science*. 1994;265(5172):679-682.
39
- 40 50. Baran B, Pace-Schott EF, Ericson C, Spencer RM. Processing of emotional reactivity and
41 emotional memory over sleep. *J Neurosci*. 2012;32(3):1035-1042.
42
- 43 51. Rasch B, Born J. About sleep's role in memory. *Physiol Rev*. 2013;93(2):681-766.
44 52. Sleep and sadness: exploring the relation among sleep, cognitive control, and depressive
45 symptoms in young adults. *Sleep Medicine*. 2014;15(1):144 - 149.

- 1 53. Vytal KE, Cornwell BR, Letkiewicz AM, Arkin NE, Grillon C. The complex interaction
2 between anxiety and cognition: insight from spatial and verbal working memory. *Front*
3 *Hum Neurosci.* 2013;7:93.
4
- 5 54. Storbeck J, Clore GL. On the interdependence of cognition and emotion. *Cogn Emot.*
6 2007;21(6):1212-1237.
7
- 8 55. van der Naalt J, Timmerman ME, de Koning ME, et al. Early predictors of outcome after
9 mild traumatic brain injury (UPFRONT): an observational cohort study. *The Lancet*
10 *Neurology.* 2017;16(7):532-540.
11
- 12 56. Zhai L, Zhang H, Zhang D. Sleep duration and depression among adults: a meta-analysis
13 of prospective studies. *Depress Anxiety.* 2015;32(9):664-670.
14
- 15 57. Neckelmann D, Mykletun A, Dahl AA. Chronic insomnia as a risk factor for developing
16 anxiety and depression. *Sleep.* 2007;30(7):873-880.
17
- 18 58. Baglioni C, Battagliese G, Feige B, et al. Insomnia as a predictor of depression: a meta-
19 analytic evaluation of longitudinal epidemiological studies. *J Affect Disord.* 2011;135(1-
20 3):10-19.
21
- 22 59. Li L, Wu C, Gan Y, Qu X, Lu Z. Insomnia and the risk of depression: a meta-analysis of
23 prospective cohort studies. *BMC Psychiatry.* 2016;16(1):375.
24
- 25 60. Ramsawh HJ, Stein MB, Belik SL, Jacobi F, Sareen J. Relationship of anxiety disorders,
26 sleep quality, and functional impairment in a community sample. *J Psychiatr Res.*
27 2009;43(10):926-933.
28
- 29 61. Siengsukon CF, Alshehri M, Aldughmi M. Self-report sleep quality combined with sleep
30 time variability distinguishes differences in fatigue, anxiety, and depression in individuals
31 with multiple sclerosis: A secondary analysis. *Mult Scler J Exp Transl Clin.*
32 2018;4(4):2055217318815924.
33
- 34 62. Siengsukon CF, Alshehri M, Williams C, Drerup M, Lynch S. Feasibility and treatment
35 effect of cognitive behavioral therapy for insomnia in individuals with multiple sclerosis:
36 A pilot randomized controlled trial. *Mult Scler Relat Disord.* 2020;40:101958.
37
- 38 63. Garland SN, Carlson LE, Stephens AJ, Antle MC, Samuels C, Campbell TS.
39 Mindfulness-based stress reduction compared with cognitive behavioral therapy for the
40 treatment of insomnia comorbid with cancer: a randomized, partially blinded,
41 noninferiority trial. *J Clin Oncol.* 2014;32(5):449-457.
42
- 43 64. Alvaro PK, Roberts RM, Harris JK. A Systematic Review Assessing Bidirectionality
44 between Sleep Disturbances, Anxiety, and Depression. *Sleep.* 2013;36(7):1059-1068.

- 1 65. Buysse DJ, Thompson W, Scott J, et al. Daytime symptoms in primary insomnia: a
2 prospective analysis using ecological momentary assessment. *Sleep medicine*.
3 2007;8(3):198-208.
4
- 5 66. McCrae CS, McNamara JP, Rowe MA, et al. Sleep and affect in older adults: using
6 multilevel modeling to examine daily associations. *Journal of sleep research*.
7 2008;17(1):42-53.
8
- 9 67. Jansson-Frojmark M, Lindblom K. A bidirectional relationship between anxiety and
10 depression, and insomnia? A prospective study in the general population. *J Psychosom*
11 *Res*. 2008;64(4):443-449.
12
- 13 68. Chennaoui M, Arnal PJ, Sauvet F, Léger D. Sleep and exercise: a reciprocal issue? *Sleep*
14 *medicine reviews*. 2015;20:59-72.
15
- 16 69. Dolezal BA, Neufeld EV, Boland DM, Martin JL, Cooper CB. Interrelationship between
17 Sleep and Exercise: A Systematic Review. *Adv Prev Med*. 2017;2017:1364387.
18
- 19 70. Zenko Z, Willis EA, White DA. Proportion of Adults Meeting the 2018 Physical Activity
20 Guidelines for Americans According to Accelerometers. *Front Public Health*.
21 2019;7:135.

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

Authors, year, country	Study Design Number of Participants	Participant demographic characteristics	Time since injury	Measures used to assess sleep	Measures used to assess other outcomes	Results Outcomes in favor of concussion
Landry-Roy et al., 2018, Canada	Prospective Cross-sectional 167 participants aged 18-60 months. 84 individuals in the mTBI group 83 in the control typically developing children	Age at injury 36.80 months (11.54) Accidental cause of trauma: car accident, accidental fall, collision related (struck by or against something)	6 months	Child Behavior Checklist- Parent report questionnaire pertaining to children's behavioral difficulties Actigraphy Sleep diary	Executive functions Delay of gratification Conflict Scale Shape Stroop	Demographics between groups The two groups did not differ significantly on age, sex, socioeconomic status, ethnicity, or post-concussive symptoms There was no significant difference between the two groups on executive function measures and sleep measures 6 months post injury Bivariate correlations CBCL sleep scale scores were not correlated with actigraphy-derived variables Significant interaction effect of children's group and CBCL sleep scale scores on delay of gratification (p=.05) There was no significant interaction between children's group and nighttime sleep duration in the prediction of conflict scale (p=.38) Significant group difference for Delay gratification performance for children with higher levels of sleep problems (p=.03) Group differences for children with lower levels of sleep problems was not significant (p=.66) Post hoc analysis Significant group difference on conflict scale performance for children with shorter nighttime sleep duration (p=.004) In children with higher nighttime sleep duration, there was not a significant difference between concussion or TDC on the conflict scale (p=. Pre-injury and 6 month post injury CBCL sleep scale for mTBI children Pre-injury and 6-months post injury CBCL sleep scale were intercorrelated (p=.001) There was no significant difference between pre and post injury CBCL sleep scale scores (p=.66) Both pre-injury and 6-month post injury CBCL sleep scale scores significantly predicted Delay of Gratification performance (p=.03)
Dean et al., 2013, United Kingdom	Cross-sectional 72 participants, 18 per group: mTBI with PCS mTBI with no PCS Controls with PCS Controls with no PCS	Individuals with an mTBI at least one year post injury and controls participants with no history of head injury. (n=72)	one year	Epworth sleepiness scale -daytime sleepiness Pittsburgh Sleep Quality Index - sleep quality Karolinska Sleepiness Scale – Sleepiness in different contexts before and after cognitive testing	N-Back for working memory PVSAT for information processing	Demographics Nocturnal sleep quality was lower in concussion +PCS participants compared to both concussion -PCS (p<.001) and controls-PCS (p=.018) Concussion +PCS participants reported a greater number of PTSD symptoms compared to concussion -PCS participants (p=.009) Mean PSQI scores for concussion +PCS and Control +PCS participants were indicative of poor nocturnal sleep. Correlation Between subjective symptom report and objective cognitive performance Significant correlation between lower sleep quality (PSQI) and poorer performance on PVSAT (p<.001) for concussion participants

<p>Theadom et al., 2018, New Zealand</p>	<p>Prospective Cross-sectional 232 mTBI 232 controls</p>	<p>Mean Age for MTBI: 40.39 (18.5) Male:130 (56%) Female:102 (44%) Prior TBI's in lifetime: None (104:44.8%) 1 to 2 (76 32.8%) 3 or more(43 18.5%) Unknown (9 3.9%)</p>	<p>4 years post injury</p>	<p>Pittsburg sleep quality index – sleep quality</p>	<p>Rivermead Post concussion Symptoms Questionnaire- Post-concussion symptoms Hospital Anxiety and Depression Scale- Anxiety and depression Participation in everyday life- participation/ community integration measure.</p>	<p>Test of difference on mean scores for the RPQ and Community participation There was no statistical difference in total score on the RPQ between mild TBI and controls (p=.009) The concussion group experienced significantly more self-report cognitive symptoms four-years post injury than controls (p=.000) There was no significant difference for somatic and emotional scores between the concussion group and the control group (p=.89; P=.58 res) Concussion group reported significantly poorer participation across all domains of the Community Participation Questionnaire domains: Productivity (P=.000), Social Relations (p=.000), Out and About P=.000 Regression prediction models Sleep quality was not a significant predictor of cognitive symptoms (p=.99) Sleep quality was a significant predictor of emotional symptoms (p=.02) Sleep quality was not a significant predictor of somatic symptoms (p=.06)</p>
<p>Van Markus-Doornbosch et al., 2019, Netherlands</p>	<p>Prospective Cross-sectional 49 Participants with mTBI</p>	<p>Participants were aged 12-25 years Mean Age at injury (16,8 years (3.7) Time since injury mean 1.6 years (.5)</p>	<p>6-18 months post injury</p>	<p>Checklist individual strength- Fatigue Pittsburgh sleep quality index –sleep quality</p>	<p>Activity Questionnaire for Adults and Adolescents- Physical Activity and individual strength</p>	<p>Physical activity minutes Participants performed general physical activity on average 1350 min/week Moderate to vigorous activities were performed on average 793 min/week Sedentary activities were 2670 min/week Meeting guidelines The median PSQI total score was 4.5 for the total group with no significant association with meeting our not meeting the D-HEPA recommen</p>

1


2 Table 1 Results

3

4

Authors (year)	Criteria for inclusion clearly defined	Participants described in detail	Exposure measured in valid and reliable way	Objective standard criteria used for measurement	Confounding factors identified	Strategies to deal with confounding factors	Outcomes measured in a valid and reliable way	Appropriate statistical analysis used
Landry-roy et al 2018	Yes- defined inclusion/ exclusion criteria	Yes- gave specific ages, and criteria for meeting concussion standards	Yes- concussion was evaluated and defined in a valid reliable way	Yes- used standard, valid, and reliable assessment tools	No- did not identify confounding factors	No- did not control for confounding factors in their statistical analysis	Unsure- Use of the CBCL sleep portion may not be specific enough	Unsure- did not describe in detail how the statistical analysis answered their research question
Dean and Sterr, 2013	Yes- defined inclusion/ exclusion criteria	Yes- Gave criteria for meeting concussion standards	Yes- concussion was evaluated and defined in a valid reliable way	Yes- used standard, valid, and reliable assessment tools	No- did not identify confounding factors	Unsure- Did not identify confounding factors and unclear if they statistically controlled for factors.	No- the researchers allowed for rest breaks and potential days between assessment times	Yes- used appropriate statistical analysis
Thea.dom et al., 2018	Yes- defined inclusion/ exclusion criteria	Yes- gave criteria for meeting concussion standard	Yes- concussion was evaluated and defined in a valid reliable way	Yes- used standard, valid, and reliable assessment tools	Yes- identified confound factors	Yes- controlled for confounding factors in statistical analysis	Yes- met in homes or over phone, done by trained research personnel	Yes- used appropriate statistical analysis
Markus- Doornbosch et al., 2018	Yes- defined inclusion/ exclusion criteria	Yes – gave criteria for meeting concussion standard	Yes- concussion was evaluated and defined in a valid reliable way	Yes- used standard, valid, and reliable assessment tools	Yes- identified confounding factors	Not Applicable	Unsure	Yes- used appropriate statistical analysis

1 Table 2 Bias Table



25 Figure 1