Contact tracing efficiency, transmission heterogeneity, and accelerating COVID-

19 epidemics

Billy J. Gardner, A. Marm Kilpatrick*

Department of Ecology and Evolutionary Biology, University of California, Santa Cruz

Correspondence to akilpatr@ucsc.edu

Abstract

Simultaneously controlling COVID-19 epidemics and limiting economic and societal

impacts presents a difficult challenge, especially with limited public health budgets.

Testing, contact tracing, and isolating/quarantining is a key strategy that has been used

to reduce transmission of SARS-CoV-2, the virus that causes COVID-19. However,

manual contact tracing is a time-consuming process and as case numbers increase it

takes longer to reach each cases' contacts, leading to additional virus spread. Delays

between symptom onset and being tested (and receiving results), and a low fraction of

symptomatic cases being tested and traced can also reduce the impact of contact

tracing on transmission. We examined the relationship between cases and delays and

the pathogen reproductive number R_t, and the implications for infection dynamics using

a stochastic compartment model of SARS-CoV-2. We found that R_t increases

sigmoidally with the number of cases due to decreasing contact tracing efficacy. This

relationship results in accelerating epidemics because R_t increases, rather than

declines, as infections increase. Shifting contact tracers from locations with high and

low case burdens relative to capacity to locations with intermediate case burdens

maximizes their impact in reducing R_t (but minimizing total infections is more

complicated). Contact tracing efficacy also decreased with increasing delays between

symptom onset and tracing and with lower fraction of symptomatic infections being

tested. Finally, testing and tracing reductions in R_t can sometimes greatly delay

epidemics due to the highly heterogeneous transmission dynamics of SARS-CoV-2.

These results demonstrate the importance of having an expandable or mobile team of

contact tracers that can be used to control surges in cases, and the value of easy

access, high testing capacity and rapid turn-around of testing results, as well as

outreach efforts to encourage symptomatic infections to be tested immediately after

symptom onset.

Author Summary

A key tool in the control of infectious diseases is contact tracing – the identification of individuals who have contacted the case and may have been infected by a newly detected case. However, to successfully contact and quarantine individuals requires time, and as cases rise, this can result in delays in reaching contacts during which time they may infect other people. Here we examine the quantitative relationships between increasing case numbers, contact tracing efficiency, and the pathogen reproductive number R_t (the number of cases infected by each case) and how these relationships vary with delays and incomplete participation in the testing and tracing process. We built

Introduction

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) emerged in late 2019 spread globally in early 2020 and resulted in rapidly growing local epidemics, large scale mortality, and strains on hospital capacity in many countries (1-4). Initial outbreaks in most countries and US states were arrested only by severe control measures including closing all but essential businesses as well as schools, churches, and other organizations (5), while a few countries were able to limit transmission, at least temporarily, primarily with public health measures (6-8). Severe disease control measures have had devastating impacts on economies and societies (5). Most countries and US states are now attempting to re-open as many business sectors and activities as possible while avoiding a rapid rise in infections.

Although self-isolation, social distancing, and mask wearing have reduced the transmission of SARS-CoV-2, additional interventions, including business closures and working from home, have often been required to keep the pathogen reproductive number R_t below 1 (5, 9, 10), especially in the United States. One public health strategy that has been used effectively to limit transmission in some countries is testing symptomatic individuals, tracing their contacts to people they may have infected, and isolating infected individuals and quarantining people that may have become infected but have yet to show symptoms or test positive for the virus (hereafter abbreviated T-

CT-I/Q) (7, 10, 11). If contacts of cases can be found and quarantined or isolated before

or during their infectious period, this can limit onward spread of the virus.

Numerous studies have examined the effectiveness and limitations of T-CT-I/Q

on transmission of SARS-CoV-2 (11-18). Many studies have shown that T-CT-I/Q can

substantially reduce the pathogen reproductive ratio, R_t, but its efficacy depends on the

importance of pre-symptomatic and asymptomatic transmission, delays between

symptom onset and being tested, and the fraction of infections that are tested and

traced (11, 15, 18). Previous studies have explored various parameter values for

contact tracing efficacy by varying the fraction isolated, the fraction symptomatic, and

the contribution to transmission of undetected infections (11, 15, 18). A key unexplored

challenge in implementing T-CT-I/Q is that tracing contacts and ensuring they can

safely quarantine or isolate is a time-consuming process which results in delays

between detecting a case and successfully quarantining their contacts. This delay

reduces the effectiveness of contact tracing as cases increase. Previous studies have

assumed fixed values for contact tracing parameters and thus have not addressed this

issue.

Our aim was to examine the relationship between increasing cases, contact

tracing efficacy, and the pathogen reproductive ratio, R_t, and to examine the potential

outcomes for disease dynamics. We built a compartment model of SARS-CoV-2

transmission, parameterized it with data from the literature, and examined how R_t varied

with number cases traced, delays between symptom onset and the start of contact

tracing, the numbers of contacts per case, and different fractions of symptomatic cases

being tested and traced. We also simulated a stochastic version of the model with and

without contact tracing to examine how reductions in R_t affected variation in the timing

of epidemics.

Methods

We built a susceptible-exposed-infected-recovered (SEIR) compartment model of

SARS-CoV-2 that included four compartments for infected individuals that reflect

symptom severity (asymptomatic, I_a, pre-symptomatic, I_{ps}, mildly symptomatic, I_{ms}, and

severely symptomatic, I_{ss}) (Fig S1). The equations of the model are:

$dS/dt = -\kappa\beta S/N(\sigma_a I_a + \sigma_{ps} I_{ps} + \sigma_{ms} I_{ms} + \sigma_{ss} I_{ss})$

$dE/dt = \kappa\beta S/N(\sigma_a I_a + \sigma_{ps} I_{ps} + \sigma_{ms} I_{ms} + \sigma_{ss} I_{ss}) - (\varepsilon + q_{E\rightarrow ps} + q_{E\rightarrow a})\ E$

$dI_{ps}/dt = q_{E\rightarrow ps}E - (\varepsilon + q_{ps\rightarrow ms})\ I_{ps}$

$dI_a/dt = q_{E\rightarrow a}E - (\varepsilon + \gamma_a)\ I_a$

$dI_{ms}/dt = q_{ps\rightarrow ms}I_{ps} - (\varepsilon + \tau_{ms} + q_{ms\rightarrow ss} + \gamma_{ms})\ I_{ms}$ [eq. 1]

$dI_{ss}/dt = q_{ms\rightarrow ss}I_{ms} - (\varepsilon + \tau_{ss} + \alpha + \gamma_{ss})\ I_{ss}$

$dQ/dt = \varepsilon I_a + \varepsilon I_{ps} + (\varepsilon + \tau_{ms})\ I_{ms} + (\varepsilon + \tau_{ss})\ I_{ss} - \gamma_Q Q$

$dR/dt = \gamma_a I_a + \gamma_{ms}I_{ms} + \gamma_{ss}I_{ss} + \gamma_Q Q$

where κ is a social distancing factor between 0 and 1 that scales the contact rate β, σ

are relative infectiousness values for each of the I classes, q are transition rates

between classes given by the subscripts separated by the arrow ($q_{E\rightarrow ps}$ is the transition

rate between the E and I_{ps} classes), ε is the rate of removal by contact tracing from the

E or I classes to the quarantined class Q, τ are the removal rate by testing of mildly

symptomatic or severely symptomatic infected individuals, α is the disease-caused

death rate, and γ are the recovery rates to the R class.

The contact tracing removal rate ε is given by:

$\varepsilon = \dfrac{f_{SCT}q_{E\rightarrow ps}}{q_{E\rightarrow ps}+q_{E\rightarrow a}}\ \dfrac{1}{1/\tau_{ms}+0.5\frac{I_{ps}q_{ps\rightarrow ms}N_{cpc}}{N_{CT}N_{CCTD}}}$ [eq. 2]

The first term is the fraction of infections traced; f_{SCT} is the fraction of symptomatic

infections that are traced of those detected with test removal rate τ_{ms} before they

recover or progress to severe symptoms; f_{SCT} is equal to $f_{tmstr}*\tau_{ms}/(\tau_{ms}+q_{ms\rightarrow ss}+\gamma_{ms})$; f_{SCT}

is multiplied by the ratio of transition rates $[q_{E\rightarrow ps}/(q_{E\rightarrow ps}+q_{E\rightarrow a})]$ which is the fraction of

infections that are symptomatic; f_{tmstr} is the fraction of mildly symptomatic cases

detected by testing that were traced. We didn't include tracing from severely

symptomatic cases because, by the time an infection progresses to severe symptoms

5-8 days after symptom onset (19, 20), their contacts will already have finished most of

their infectious period, and quarantining their contacts will have little effect.

The second term is the inverse of the time between symptom onset of the

symptomatic infection and being removed by contact tracing which is the sum of the

delay between symptom onset and receiving a test result, $1/\tau_{ms}$, and the average time

needed to reach the contacts of the newly detected cases (which is half the total time to

call all contacts). $q_{ps \to ms}I_{ps}$ is the rate of new symptomatic infections that need to be

traced, N_{cpc} is the average number of contacts per case, N_{CT} is the number of contact

tracers, and N_{CCTD} is the number of calls each contact tracer can make each day.

We parameterized the model with data from the literature, with all rates in days^{-1}

(Table 1) and used the baseline contact tracing capacity standards suggested by the

US National Association of County and City Health Officials (15 contact tracers per

100,000 people; https://www.naccho.org/uploads/full-width-images/Contact-Tracing-

Statement-4-16-2020.pdf). We note that while notifying individuals that they have had

contact with a case can be done quickly (especially with using a cell-phone tracing app;

(18)), successfully ensuring a contact has their needs met (including food, medicine,

clothing) to quarantine in a safe space where they won't infect their household members

requires substantially more time (https://www.cdc.gov/coronavirus/2019-

ncov/php/notification-of-exposure.html). We assumed the approximate duration required

for a successful contact tracing call was 40 min, resulting in N_{CCTD} = 12

We used the next generation matrix technique to derive an expression for the

pathogen reproductive ratio R_t (21):

$R_t = S/N[\kappa\beta]$ *

$[[(q_{E \to a})(\sigma_a) / (\varepsilon + \gamma_a)] + [(q_{E \to ps})(\sigma_{ps}) / (\varepsilon + q_{ps \to ms})] +$ [eq. 3]

$[(q_{E \to ps})(q_{ps \to ms})(\sigma_{ms}) / ((\varepsilon + q_{ps \to ms})(\varepsilon + \tau_{ms} + q_{ms \to ss} + \gamma_{ms}))] +$

$[(q_{E \to ps})(q_{ps \to ms})(q_{ms \to ss})(\sigma_{ss}) /$

$((\varepsilon + q_{ps \to ms})(\varepsilon + \tau_{ms} + q_{ms \to ss} + \gamma_{ms})(\varepsilon + \tau_{ss} + \alpha + \gamma_{ss}))]] / (\varepsilon + q_{E \to a} + q_{E \to ps})$

This expression can be understood as the fraction of the population that is susceptible,

S/N multiplied by the contact rate β (which is scaled by the social distancing factor κ),

multiplied by the sum of four terms: one for each infected class. Each of the four terms

includes the product of the transition rates q to reach that class from the E class

multiplied by the infectiousness of that class, σ, divided by the recovery, testing and

tracing rates leaving that I class and the I classes before it. Finally, all four terms are

divided by the rates hosts leave the E class which normalizes the rates moving along

each infection pathway.

We examined how R_t and R_0 ($R_t = R_0$ at beginning of epidemic when $S \cong N$)

varied with different numbers of new symptomatic cases detected (a fraction of the

newly symptomatic infection, $I_{ps}\, q_{ps \to ms}$ in eq. 2), delays between symptom onset and the

start of contact tracing ($1/\tau_{ms}$ in eq. 2), and numbers of contacts per case (N_{cpc} in eq. 2).

Note that because these three quantities occur as a ratio (with the number of contact

tracers) in equation (2), any combination of values that produce the same number of

case-contacts per contact tracer calls per day will produce the same value of R_t. We

illustrated the effect of decreasing contact tracing efficiency as infections increased on

disease dynamics and R_t by simulating a deterministic version of the model in eq. 1 and

plotted R_t in real time as an epidemic swept through a population over one year. We

also performed a simple sensitivity analysis by determining how much R_t varied with a

ten percent increase or decrease in each model parameter (Fig S2).

Finally, we explored the implications of stochastic variability and contact tracing

on infection dynamics in a scenario roughly based on a moderate size city with partly

effective social distancing. We simulated a stochastic version of the model given by eq

1 where the number of new infections was drawn from a negative binomial distribution

with mean equal to R_t and dispersion parameter 0.16 which is intermediate between

available estimates for COVID-19 (22-24). We modeled a scenario of a population of

100,000 people where non-pharmaceutical interventions reduced contact rates by 30%

(κ=0.7) resulting in R_0 with/without contact tracing of 1.32/1.67, testing and contact

tracing took place after an average of $1/\tau_{ms}$ = 5 days after symptom onset, and half of

symptomatic cases detected by testing being traced (f_{tmstr} = 0.5). We examined different

initial numbers of latently infected individuals, E, at the start of the epidemic to

understand how stochastic variation could impact the timing of epidemics when initial

infections were low.

R code to reproduce the results is available from:

https://github.com/marmkilpatrick/Contact-Tracing-Efficiency

Results

191   The effectiveness of contact tracing in reducing the pathogen reproductive

192 number, R_t, was highly dependent on three factors: the number of cases being traced

193 (given a fixed number of contact tracers), the delay between symptom onset and the

194 start of tracing, $1/\tau_{ms}$, (including getting tested and receiving result), and the fraction of

195 symptomatic cases that get traced (Figs 1-3).

Figure 1. Pathogen reproductive ratio R_t plotted against the number of new cases

per contact tracer calls per day for 1-5 day delays ($1/\tau_{ms}$) between symptom onset

and the start of contact tracing (including getting tested and receiving result), 10

contacts per case (so the average number of contacts each tracer has to reach

each day is 10x the x-axis values). With no contact tracing R_t increases from 2.03

to 2.37 as the delay $1/\tau_{ms}$ increases from 1 to 5 days, which is evident in the y-axis

difference between curves in the upper right of the graph where new case

burdens are so high contact tracing is ineffective. Delays are indicated by the

small numbers on each curve in the left of the plot.

First, the relationship between R_0 and the number of cases per contact tracer

calls per day was sigmoid (Figs 1-3); at both high and low numbers adding or removing

contact tracers had smaller effects whereas at intermediate case numbers relative to

capacity shifting contact tracers would have a larger impact. When the number of new

daily cases per contact tracer call per day was high (i.e. >1 or 10 contacts per contact

tracer calls per day), contact tracing had relatively little effect in reducing R_0 no matter

how long the delay was between symptom onset and the start of tracing, $1/\tau_{ms}$ (right

side of Figs 1; also evident in Figs 2, 3). This is because contact tracing calls took too

long, on average (>10 days), to reduce the infectious period of contacts. Note that for

the parameter estimates used (Table 1), an average case becomes infectious starting

3.2 days after infection, and is highly or moderately infectious for an average of 7.3

more days (σ_{ps} and σ_{ms}; Table 1). The exponentially distributed durations for the latent

and infectious periods implied by standard compartmental models result in only 37% of

individuals leaving each class after the average duration, which results in a relatively

smaller benefit of contact tracing beyond this case burden, and resultant delay in

reaching contacts of cases.

Figure 2. Pathogen reproductive ratio R_t plotted against the number of Cases per
contact tracer calls per day for four different number of contacts per case (5, 10,
20, 30; indicated by the small numbers on each curve in the middle of the plot),
and an average delay $1/\tau_{ms}$ of 5 days between symptom onset and contact tracing
(including getting tested and receiving result). With no contact tracing R_t = 2.39.

Contact tracers in regions with very high new case numbers relative to contact
tracing capacity (>1 on Fig. 1) would have a larger reduction on R_t if they were tracing
calls with intermediate numbers of cases (0.02 to 1 cases/contact tracer calls per day).
However, it is worth noting that reducing R_t is not the same as preventing new cases
(see Discussion below). Nonetheless, if the goal is to reduce R_t<1 to stop the growth in
cases, the analyses in Figs 1-3 suggest that when new case burdens are high relative
to capacity, population-wide interventions (e.g. social distancing or different levels of
shelter-in-place orders which reduce contact rates, β or κ, and shift the entire curves in
Fig 1-3 downward proportionately; Fig. S2), or orders of magnitude increases in contact
tracing capacity will be needed until R_t can be effectively reduced by contact tracing.
When there are few (<~0.02) cases per contact tracer call per day, contact

243   tracing was as effective as it could be but excess capacity had little impact, especially

244   for realistic delays, $1/\tau_{ms}$, between symptom onset and the start of tracing (e.g. 3-5

245   days). Shifting contact tracers to other areas with more cases per contact tracer (0.02 to

246   0.2) would have minimal impact in increasing R_t but could help substantially reduce R_t in

247   those places with higher new case burdens relative to capacity.

248   Second, increasing delays, $1/\tau_{ms}$, between symptom onset and the start of

tracing greatly influenced the efficacy of contact tracing (Fig. 1). With a 5 day delay,

contact tracing could only reduce R_0 by 33% from 2.6 to 1.6 regardless of contact

tracing capacity per case. In contrast, if all symptomatic people sought care and got

tested within 1 day of after symptom onset and results were returned within the next 24

hours ($1/\tau_{ms}$ = 2 days), contact tracing could reduce R_0 by 73% from 2.2 to 0.6. Note

that the effect of delays in increasing R_0 due to delayed removal by testing alone (τ_{ms}) is

small compared to the effect of delays in reducing contact tracing efficiency (Fig 1:

compare differences among curves in upper right of graph where contact tracing is

having little effect to differences among curves in lower left of graph where it has

maximal effect). The number of contacts per case obviously also influences the time

required to trace these contacts (Fig. 2). If allowable (or illegal) gathering sizes

increase, this leads to more contacts per case and a faster decrease in contact tracing

efficacy.

262   Thirdly, if contacts for a substantial fraction of all symptomatic cases do not get

traced and quarantined, T-CT-I/Q is far less effective. Figures 1 and 2 showed an

optimistic scenario where the fraction of symptomatic infections that are tested and

traced is determined only by the delay between symptom onset and testing results

being returned ($1/\tau_{ms}$) (i.e. all symptomatic infections could be tested and traced). With a

5 day delay ($1/\tau_{ms}$=5) (Fig. 2), this results in 38% of infections being detected by testing

in the mildly symptomatic state (I_{ms}) which is much higher than estimates of case under-

ascertainment based on seroprevalence studies (25). If only half of these cases are

contact traced (f_{tmstr} = 0.5) the maximum impact of contact tracing is smaller: a 21%

reduction in R_0 from 2.39 to 1.89 (Fig 3) rather than a 35% decrease from 2.38 to 1.54

(Fig 2).

Figure 3. Pathogen reproductive ratio R_t plotted against the number of Cases per
274
contact tracer calls per day for four different number of contacts per case (5, 10,
275
20, 30; indicated by the small numbers on each curve in the middle of the plot), 12
276
calls per contact tracer per day, an average delay $1/\tau_{ms}$ of 5 days between
277
symptom onset and contact tracing (including getting tested and receiving
278
result). This Figure is the same Figure 2, except only half as many contacts are
279
traced (f_{tmstr} = 0.5). With no contact tracing, R_t = 2.39.
280

Reduced contact tracing efficiency with increasing cases can result in a transient
282
accelerating epidemic where R_t increases over time (Fig 4). If contact tracing capacity is
283
insufficient to quickly trace contacts, then a decrease in contact tracing efficiency can
284
initially outweigh the depletion of susceptible individuals and lead to an increase in R_t
285
over time until depletion of susceptibles overwhelms this effect (Fig 4: compare
286
rightmost panels with limited contact tracing to leftmost panels where social distancing
287
reduces R_t to the same initial value as contact tracing). With unlimited contact tracing
288
this phenomenon does not arise (Fig 4: compare middle panels with unlimited contact
289
tracing to leftmost panels).
290

Figure 4. Impact of reduced contact tracing efficiency with increasing cases. Left panels show (A) the number of susceptible, exposed, infected, and recovered individuals and (B) reproductive number, R_t, over time with no contact tracing but social distancing ($\kappa = 0.65$) set to give same initial R_0 (1.55) as with contact tracing. Middle panels (C, D) show the same variables with unlimited contact tracing (1500 contact tracers making 12 calls/day; 15 contacts per case; $1/\tau_{ms} =$ 5d; $f_{tmstr} = 1$) but no social distancing ($\kappa = 1$). Right panels (E, F) show the same variables and parameters as (C, D) but with <u>limited</u> contact tracing (5 contact tracers).

The impact of contact tracing in reducing the pathogen reproductive number R_t has two consequences on the temporal timing and establishment of epidemics. First, as is well known, reducing R_t delays and reduces the peak of the epidemic (Fig 5 top vs bottom panels). Second, and less appreciated, stochastic variation in R_t can lead to very different timing of epidemics if the initial number of infected individuals is low (Fig 5 left panels), and variation is larger if R_t is lower (Fig 5A vs Fig 5C). Finally, heterogeneity in individual transmission can result in local fadeout of the pathogen and this is more if contact tracing reduces R_t so that it is closer to 1, and if the number of infected

311   individuals is lower.

**Figure 5. Variability in the timing and outcome of epidemics due to stochastic
variation in individual R_t. Lines show number of latently infected individuals (in
the E class) over time for 1 year with moderate social distancing that reduces
contact rates by 30% (κ = 0.7). Grey lines show a single stochastic simulation and
the black line shows the deterministic outcome. Frac. est. is the fraction of
epidemics that establish (i.e. the maximum number of infected at any time
exceeds the starting number infected). Four scenarios include different starting
numbers of latently infected individuals on day 0, E_0 (A, C: 5; B, D: 50), and with
(A, B) or without (C, D) contact tracing (CT) which lowered R_0 from 1.67 to 1.32.
The delay from symptom onset to testing and tracing was 5d, but only half of
cases were traced (f_{tmstr} = 0.5), as in Fig 3. The modeled population of 100,000
people had 15 tracers making 12 calls/day, and each case had an average of 15
contacts which is intermediate between pre-lockdown and lockdown conditions.**

Discussion

The two main strategies that have been used to control COVID-19 are T-CT-I/Q

and society-wide social distancing interventions (including closing businesses, banning
gatherings, wearing masks, etc.) (9). Closing businesses has had devastating impacts
on employment and economies, as well as cascading impacts on society. T-CT-I/Q has
far smaller economic and societal costs, but its efficacy in controlling epidemics is not
fully understood, and some studies suggest that it is insufficient to keep R_t below 1 in
many settings, especially without digital contact tracing (10, 11, 18). We examined how
the efficacy of contact tracing decreases with increasing case burden. At high case
burdens relative to contact tracing capacity, contact tracing reached most contacts too
late and had little effect on R_t. Conversely when case numbers were very low relative to
contact tracing capacity, there was excess capacity and, all else being equal, contact
tracers could be used more effectively in higher case burden settings with very small
impacts on local transmission. We note that the exact number of contact tracers needed
to reduce R_t depends on the number of contacts per case and the number of calls each
tracer can make each day (Fig 2). However, the key quantity appears as a ratio of case-
contacts per contact tracer calls per day. Thus, each contact tracing team (e.g. a US
County) can use local estimates of contacts per case and the number of calls each
tracer can make each day to determine where they are on the modelled relationships
(Figs 1-3).

A major caveat must be kept in mind in interpreting these results. A smaller
reduction in R_t (e.g. 10%) in one population can prevent more infections (especially over
multiple generations of transmission) than a larger (e.g. 20%) reduction in R_t in a
second population if R_t in the second location is lower (especially when $R_t>1$ in the first
population), or when there is a larger number of infected individuals in the first
population. Thus, transferring contact tracers from a region with a high case burden
relative to contact tracing capacity to maximize their efficacy in reducing R_t should only
be done if other measures (e.g. social distancing) will be put into place to reduce R_t
where case numbers are high. More generally, allocation of contact tracers to maximize
the number of cases prevented given an array of tools would require a complex
dynamic analysis beyond that examined here.

We also found that the efficacy of contact tracing itself, regardless of capacity,
was strongly influenced by delays between the onset of symptoms and the beginning of

tracing, as well as the fraction of symptomatic infections that were traced. Unless delays

were short and the fraction of symptomatic cases that were traced was high, contact

tracing had limited effects in reducing R_t. This finding parallels results from other studies

demonstrating the large effects of delays in reducing efficacy of isolating infections by

testing alone (26). We note that in the model considered here, only symptomatic

individuals were removed by testing (pre-symptomatic and asymptomatic infections

were not tested) which leads to a much smaller impact of testing on R_t. Our results

emphasize the importance of encouraging people to get tested as soon as possible

after mild symptom onset, and having sufficient testing capacity to return their results

quickly. Similarly, the fraction of symptomatic infections that get tested and traced is

poorly known, but if the ratio of infections to cases from seroprevalence studies in some

locations is approximately correct (e.g. 10:1; (25)), then contact tracing will have very

limited effects in reducing transmission.

Allocation of contact tracing resources can be most efficiently deployed in two

ways. First, contact tracing is much more effective when infections are detected soon

after symptom onset. One should prioritize these individual cases for tracing since their

contacts are likely to be earlier in their infections and quarantining them will cut off most

or all of their infectious period. If one knows the date of contact between the case and

the contact, one could also prioritize tracing more recent contacts and those that had

contact with the case during the case's days of peak infectiousness just before and after

symptom onset (27, 28). Second, if one is attempting to limit transmission in multiple

regions (e.g. US counties within a state) one could deploy contact tracers to counties

where they will be able to have the most impact: from places with excess capacity to

those with intermediate numbers of cases per contact tracer calls per day. Conversely,

if contact tracers cannot quarantine the contacts of cases within 10-12 days of the

case's symptom onset, they will be unlikely to effectively reduce transmission from

those contacts.

Our results also offer insight on two phenomena observed in COVID-19

epidemics that are not fully understood. First, epidemic dynamics sometimes differ

enormously between places that seem otherwise similar. This may be due to

differences in social behavior or contact patterns, but our results illustrate that

stochastic chance may also play a role in shifting the timing of epidemics by more than

a month, especially when the initial number of cases is low and R_t is closer to 1 (i.e.

when lockdowns are first lifted) so that populations spend longer periods of time with

few cases where stochastic variation is most important. Second, staged business re-

openings have sometimes led to accelerating or runaway epidemics. These may be due

to sudden changes in social behavior, but decreases in contact tracing efficiency may

also contribute to these accelerating epidemics, if contact tracing was playing a

significant role in limiting transmission. Increasing contact tracing capacity could limit

this epidemic acceleration as cases increase, which suggests that training a reserve

capacity of tracers to be used following case surges or being able to deploy a mobile

tracing force could help limit runaway epidemics.

More broadly, contact tracing could play an important role in limiting transmission

of SARS-CoV-2 and other pathogens. However, we found that its efficacy depends on

participation in seeking testing immediately following symptom onset and quick return of

test results, as well as sufficient contact tracing capacity if case numbers surge.

Shortcomings in each of these factors greatly limit its efficacy, which could lead to

implementation of much more damaging measures to control transmission, including

widespread business and school closures. Investments in public health, including

testing, contact tracing, and public outreach to encourage health seeking when

symptomatic, is likely a much more cost-effective approach to control COVID-19.

Acknowledgements: We thank the dozens of scientists and public health practitioners

who have informed this work through comments and discussions.

417   Literature Cited

1.   J. T. Wu, K. Leung, G. M. Leung, Nowcasting and forecasting the potential domestic and
international spread of the 2019-nCoV outbreak originating in Wuhan, China: a
modelling study. *Lancet* **395**, 689-697 (2020).
2.   A. Remuzzi, G. Remuzzi, COVID-19 and Italy: what next? *Lancet* **395**, 1225-1228
(2020).
3.   P. Zhou *et al.*, A pneumonia outbreak associated with a new coronavirus of probable bat
origin. *Nature* **579**, 270-+ (2020).
4.   N. Zhu *et al.*, A Novel Coronavirus from Patients with Pneumonia in China, 2019. *N.
Engl. J. Med.* **382**, 727-733 (2020).

5.　S. Flaxman *et al.*, Estimating the effects of non-pharmaceutical interventions on COVID-
19 in Europe. *Nature* 10.1038/s41586-020-2405-7, 15.
6.　Y. X. Ng *et al.*, Evaluation of the Effectiveness of Surveillance and Containment
Measures for the First 100 Patients with COVID-19 in Singapore - January 2-February
29, 2020. *MMWR-Morb. Mortal. Wkly. Rep.* **69**, 307-311 (2020).
7.　R. Pung *et al.*, Investigation of three clusters of COVID-19 in Singapore: implications for
surveillance and response measures. *Lancet* **395**, 1039-1046 (2020).
8.　J. R. Koo *et al.*, Interventions to mitigate early spread of SARS-CoV-2 in Singapore: a
modelling study. *Lancet Infect. Dis.* **20**, 678–688 (2020).
9.　B. J. Cowling, A. E. Aiello, Public Health Measures to Slow Community Spread of
Coronavirus Disease 2019. *J. Infect. Dis.* **221**, 1749-1751 (2020).
10.　B. J. Cowling *et al.*, Impact assessment of non-pharmaceutical interventions against
coronavirus disease 2019 and influenza in Hong Kong: an observational study. *Lancet*
Public Health **5**, E279-E288 (2020).
11.　J. Hellewell *et al.*, Feasibility of controlling COVID-19 outbreaks by isolation of cases and
contacts. *Lancet Global Health* **8**, E488-E496 (2020).
12.　J. A. Firth *et al.*, Using a real-world network to model localized COVID-19 control
strategies. *Nat. Med.* 10.1038/s41591-020-1036-8.
13.　G. Giordano *et al.*, Modelling the COVID-19 epidemic and implementation of population-
wide interventions in Italy. *Nat. Med.* 10.1038/s41591-020-0883-7.
14.　S. Marcel *et al.*, COVID-19 epidemic in Switzerland: on the importance of testing,
contact tracing and isolation. *Swiss Medical Weekly* **150** (2020).
15.　S. M. Moghadas *et al.*, The implications of silent transmission for the control of COVID-
19 outbreaks. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 17513-17515 (2020).
16.　C. N. Ngonghala *et al.*, Mathematical assessment of the impact of non-pharmaceutical
interventions on curtailing the 2019 novel Coronavirus. *Math. Biosci.* **325** (2020).
17.　R. P. Walensky, C. del Rio, From Mitigation to Containment of the COVID-19 Pandemic
Putting the SARS-CoV-2 Genie Back in the Bottle. *JAMA-J. Am. Med. Assoc.* **323**, 1889-
1890 (2020).
18.　L. Ferretti *et al.*, Quantifying SARS-CoV-2 transmission suggests epidemic control with
digital contact tracing. *Science* **368**, 619-+ (2020).
19.　F. Zhou *et al.*, Clinical course and risk factors for mortality of adult inpatients with
COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* **395**, 1054-1062
(2020).
20.　J. A. Lewnard *et al.*, Incidence, clinical outcomes, and transmission dynamics of severe
coronavirus disease 2019 in California and Washington: prospective cohort study. *BMJ-*
British Medical Journal **369**, 10 (2020).
21.　O. Diekmann, J. A. P. Heesterbeek, M. G. Roberts, The construction of next-generation
matrices for compartmental epidemic models. *J. R. Soc. Interface* **7**, 873-885 (2010).
22.　B. M. Althouse *et al.*, Stochasticity and heterogeneity in the transmission dynamics of
SARS-CoV-2. *arXiv* arXiv:2005.13689, https://arxiv.org/abs/2005.13689 (2020).
23.　A. Endo, Centre for the Mathematical Modelling of Infectious Diseases COVID-19
Working Group, S. Abbott, A. J. Kucharski, S. Funk, Estimating the overdispersion in
COVID-19 transmission using outbreak sizes outside China. *Wellcome Open Research*
5, https://doi.org/10.12688/wellcomeopenres.15842.12681 (2020).
24.　D. C. Adam *et al.*, Clustering and superspreading potential of SARS-CoV-2 infections in
Hong Kong. *Nat. Med.* 10.1038/s41591-020-1092-0, 10.1038/s41591-41020-41092-
41590 (2020).
25.　E. S. Rosenberg *et al.*, Cumulative incidence and diagnosis of SARS-CoV-2 infection in
New York,. *Ann. Epidemiol.* **48**, 23-29 (2020).

26.   D. B. Larremore *et al.*, Surveillance testing of SARS-CoV-2. *medRxiv*,
478   doi.org/10.1101/2020.1106.1122.20136309 (2020).
479   27.   A. M. Wilson *et al.*, Quantifying SARS-CoV-2 infection risk within the Apple/Google
480   exposure notification framework to inform quarantine recommendations. *medRxiv*
481   10.1101/2020.07.17.20156539 (2020).
482   28.   X. He *et al.*, Temporal dynamics in viral shedding and transmissibility of COVID-19. *Nat.*
483   *Med.* **26**, 672-+ (2020).
484   29.   S. Abbott *et al.*, Estimating the time-varying reproduction number of SARS-CoV-2 using
485   national and subnational case counts [version 1; peer review: awaiting peer review].
486   *Wellcome Open Research* **5**, 112 (2020).
487   30.   R. Verity *et al.*, Estimates of the severity of coronavirus disease 2019: a model-based
488   analysis. *Lancet Infect. Dis.* **20**, 669-677 (2020).
489   31.   T. W. Russell *et al.*, Estimating the infection and case fatality ratio for coronavirus
490   disease (COVID-19) using age-adjusted data from the outbreak on the Diamond
491   Princess cruise ship, February 2020. *Eurosurv.* **25**, 12 (2020).
492   32.   D. C. Buitrago-Garcia *et al.*, Asymptomatic SARS-CoV-2 infections: a living systematic
493   review and meta-analysis. *medRxiv* 10.1101/2020.04.25.20079103 (2020).
494   33.   M. Cevik *et al.*, SARS-CoV-2, SARS-CoV-1 and MERS-CoV viral load dynamics,
495   duration of viral shedding and infectiousness: a living systematic review and meta-
496   analysis. *medRxiv* 10.1101/2020.07.25.20162107 (2020).
497

500 **Tables**

501 **Table 1. Parameter values for the model. All rates are in days^{-1} and many are**

502 **based on the inverse of measured durations.**

Para-meter	Value or range	Description	Reference or Derivation
τ_{ms}	0.2-1	Testing removal rate for I_{ms} (1/delay from onset to testing & tracing)	Scenarios explored
β	0.37	Contact rate	Set to give plausible pre-lockdown $R_0 \cong$ 2-3 (29)
f_{tmstr}	0.5 or 1	Fraction of mildly symptomatic cases detected by testing whose contacts were traced	Scenarios explored
κ	0-1	Social distancing factor	Adjusted to produce $R_t \cong$ 1.2-1.7; consistent with data post-lockdown; (29)
α	0.0025	Disease caused death rate	estimated from IFR* $(\gamma_{ms}+\gamma_{ss}+q_{E\to a})$/(1-IFR) = using infection fatality ratio IFR = 0.0066; (30, 31)
$q_{E\to a}$	0.078	Transition rate exposed to asymptomatically infected	Estimated from $q_{E\to a}$ = $f_{asymp}(q_{E\to ps})$/ (1- f_{asymp}) using fraction asymptomatic (f_{asymp} = 0.2); (32)
$q_{E\to ps}$	1/3.2	1/(duration latent period)	(28)
$q_{ps\to ms}$	1/2.3	1/(duration pre-symptomatic period)	(28)
$q_{ms\to ss}$	1/8	1/duration mild symptoms	(19, 20)
γ_a	1/8	Asymptomatic recovery rate	Based on average ratio of shedding durations compared to symptomatic cases (33)
γ_{ms}	1/5,	Mild infection recovery rate	(28)
γ_{ss}	1/10.7	Severe symptom recovery rate	(20)
γ_Q	1/14	Quarantined recover rate	Does not affect dynamics
σ_a	1	Relative infectiousness	(33)

		asymptomatic:mildly symptomatic	
σ_{ps}	1.81	Relative infectiousness pre-symptomatic:mildly symptomatic	(28)
σ_{ms}	1	Relative infectiousness	(reference level)
σ_{ss}	0.008	Relative infectiousness severe symptoms:mildly symptomatic	(28)

Figure S1. Compartmental model of SARS-CoV-2. See text for equations and

parameter values. Boxes represent Susceptible (S), Exposed (E), Infected (I),

recovered (R), and Quarantined (Q) classes. There are four compartments for

infected individuals that reflect symptom severity (asymptomatic, I_a, pre-

symptomatic, I_{ps}, mildly symptomatic, I_a, and severely symptomatic, I_{ss}) . κ is a

social distancing factor between 0 and 1 that modifies the contact rate β, σ are

infectiousness for each of the I classes, q are transition rates between classes

given by the subscripts separated by the arrow ($q_{E \to ps}$ is the transition rate

between the E and I_{ps} classes), ε is the rate of removal by contact tracing from the

E or I classes to the quarantined class Q, τ are the removal rate by testing of

mildly symptomatic or severely symptomatic infected individuals, α is the

disease-caused death rate, and γ are the recovery rates to the R class. The

dashed line and bracket indicate that all 4 classes of infected individuals

contribute to transmission.

Figure S2. Sensitivity analysis. The plot shows how much R_0 changes from a ten percent increase (red) or ten percent decrease (blue) in that model parameter relative to values in Table 1 (with $\tau_{ms} = 0.2$; $f_{tmstr} = 1$; $\kappa = 1$). R_t scales linearly with β and κ, whereas $q_{ps \to ms}$, τ_{ms}, and σ_{ps} have half as large an effect as β or κ and other parameters are even less influential.