Effectiveness of Non-pharmaceutical Interventions to Contain COVID-19:

A Case Study of the 2020 Spring Pandemic Wave in New York City

Wan Yang,[1*] Jaimie Shaff,[2] Jeffrey Shaman[3]

[1]Department of Epidemiology, Mailman School of Public Health, Columbia University, New

York, NY, USA; [2]New York City Department of Health and Mental Hygiene, New York, NY,

USA; [3]Department of Environmental Health Sciences, Mailman School of Public Health,

Columbia University, New York, NY, USA

Correspondence to:

Wan Yang

Department of Epidemiology

Mailman School of Public Health

Columbia University

722 W 168th Street, Room 514, New York, NY 10032

Phone: (212) 305-0421

Email: wy2202@columbia.edu

Abstract

As COVID-19 continues to pose significant public health threats, quantifying the

effectiveness of different public health interventions is crucial to inform intervention

strategies. Using detailed epidemiological and mobility data available for New York City

and comprehensive modeling accounting for under-detection, we reconstruct the COVID-

19 transmission dynamics therein during the 2020 spring pandemic wave and estimate the

effectiveness of two major non-pharmaceutical interventions—lockdown-like measures

that reducing contact rates and universal masking. Lockdown-like measures were

associated with >50% transmission reduction for all age groups. Universal masking was

associated with a ~7% transmission reduction overall and up to 20% reduction for 65+

year-olds during the first month of implementation. This result suggests that face covering

can substantially reduce transmission when lockdown-like measures are lifted but by itself

may be insufficient to control SARS-CoV-2 transmission. Overall, findings support the need

to implement multiple interventions simultaneously to effectively mitigate COVID-19

spread before the majority of population can be protected through mass-vaccination.

Keywords: COVID-19; transmission dynamics; effectiveness of intervention; social

distancing; lockdown; face covering

Introduction

Since the emergence of SARS-CoV-2 in late 2019, the virus has infected over 79 million

people and killed over 1.75 million worldwide by the end of 2020 (as of 12/27/20).[1] Non-

pharmaceutical interventions such as social distancing and face covering have been the

main strategies to contain COVID-19 during this pandemic in 2020. In late 2020, Phase-III

trials for several SARS-CoV-2 vaccines showed highly promising results and were granted

emergency use in several countries.[2,3] However, before these vaccines become widely

available to the general population (likely in mid- or late 2021), non-pharmaceutical

interventions will need to remain the main strategies to contain COVID-19. In addition,

future (re)emerging infectious disease outbreaks may need to rely on similar non-

pharmaceutical measures. It is thus critical to understand the effectiveness of different

non-pharmaceutical interventions implemented during the COVID-19 pandemic waves in

order to inform effective future planning while balancing economic need. For instance,

with face covering and social distancing by closing businesses as two main interventions,

the more effective face covering is, the more businesses could remain open. As a simplified

calculation, with a basic reproductive number (R_0) of 3 and minimal immunity, a city could

maintain 55% business capacity while curbing epidemic growth, if its residents could

reduce transmission by 40% using face covering [ie, effective reproductive number R_t = 3 ×

55% × (1 – 40%) = 0.99 < 1]; this threshold business capacity would drop to 33% if no

residents used face covering.

However, assessing the effectiveness and impact of a given intervention for COVID-19 has

been challenging due to low infection detection rates (many asymptomatic and mild

infections do not seek care or receive testing),[4] fluctuation of those infection detection

rates, differential disease manifestation by age group,[5,6] and concurrent public health

interventions. As such, while a few studies have assessed the overall effectiveness of

lockdown-like measures, to date, the effectiveness of specific measures including face

covering *under real-world conditions* remains unclear.

To estimate the effectiveness of different non-pharmaceutical public health interventions,

here we thus focus on the 2020 spring COVID-19 pandemic wave in New York City (NYC),

the first COVID-19 epidemic center in the United States, where detailed data are also

available. NYC experienced widespread COVID-19 transmission citywide since early March

and recorded over 200,000 cases and over 21,000 COVID-19 confirmed or probable deaths

during the following three months. To curb this intense transmission, NY State and NYC

implemented multiple intervention measures, including health promotion campaigns in

early March, telecommuting and staggered work schedule recommendations beginning the

week of March 8, public schools closure starting the week of March 15,[7] stay-at-home

orders for non-essential workers starting the week of March 22,[8] and requirements for use

of face covering in public starting the week of April 12.[9] With these overlapping and far

reaching public health interventions, case diagnoses and hospitalizations peaked in April,

and started to decline substantially in late April and May. NYC was able to begin its phased

re-opening of industries starting the week of June 7, 2020.

In this study, we apply a model-inference system[10-12] developed to support the city's

COVID-19 pandemic response to reconstruct the underlying transmission dynamics of

COVID-19 in NYC during March 1 – June 6, 2020 (i.e. prior to the city's reopening). To

address the aforementioned challenges, our model-inference system simultaneously

assimilates three sources of data: 1) confirmed COVID-19 case data, 2) COVID-19

associated death data (both cases and deaths are assimilated by neighborhood and age

group), and 3) neighborhood-level mobility data to constrain the model system. This

enables inference of the overall infection rate (i.e. including those not documented by

surveillance), estimation of key transmission characteristics (e.g., the reproductive

number) through time, and assessment of the effectiveness of different public health

interventions, including social distancing and face covering, implemented over time. We

further incorporate these estimates to project cases and deaths in the weeks beyond our

study period and compare the projections to independent observations in order to evaluate

the accuracy of these estimates. We conclude with a discussion on the implication of our

findings on strategies to safely reopen economies in places COVID-19 continues to pose

substantial public health threats.

Results

Overall Epidemic Trends

Following diagnosis of the first case in NYC, confirmed COVID-19 cases in the entire

population increased nearly exponentially during the first three weeks (Fig 1B) before

slowing down beginning the week of March 22, 2020 when NYC implemented a stay-at-

home order. However, case trajectories differed substantially by age group. Foremost,

reported cases increased with age: the case trajectory for those aged 25-44 years mirrored

the overall epidemic curve, those older than 45 years had higher case rates, and those

under 25 years had the lowest case rates (Fig 1A and Table S1). Infants (i.e. <1 year),

however, had higher case rates than 1-4 and 5-14 year-olds (Fig 1). In addition, the timing

of peak case rate was mixed. Case rates in 25-44, 45-64, and 65-74 year-olds peaked

earliest during the week of March 29, 2020, followed by <1, 1-4, 15-24, and 75+ year-olds

with a 1-week lag; in comparison, the case rate for 5-14 year-olds fluctuated with a less

clear peak during the weeks of March 29 – April 26, 2020.

The epidemic trends based on diagnosed cases, however, were obscured by varying

infection detection rates by age and through time. COVID-19 infections are more likely to

manifest as symptomatic illness and/or more severe disease in individuals with underlying

conditions and in older adults.[5,6] Such differential clinical characteristics by age thus lead to

varying healthcare seeking behaviors and infection detection rates by age. In addition,

testing policies varied over the course of the Spring 2020 COVID-19 pandemic in NYC.

During this time, testing capacity was limited at the federal, state and local levels by

guidelines for who should be tested (due, for example, to limited availability of test kits,

swabbing supplies and reagents), which required prioritizing testing for severely ill

patients and those highly vulnerable to severe disease. Testing capacities expanded during

the week of March 8, 2020;[13] however, by late March, material shortages (including testing

kits and personal protective equipment) again prompted the city to restrict testing to those

severely ill.[14] Using our model-inference system, we estimated that infection detection

rates increased in early March, reaching a peak of around 20% for all ages overall during

the week of March 15, and declined afterwards before increasing again in mid-April.[11,12]

After accounting for infection detection rates to include undiagnosed infections, a different

picture of the NYC spring outbreak emerges (Fig 2 and Fig S3). Estimated infection rates

were highest among 25-44 and 45-64 year-olds (Fig. 2 F and G), followed by 65-74 and 75+

year-olds (Fig. 2 H and I), then 5-14 and 15-24 year-olds (Fig. 2 D and E), and were lowest

among <1 and 1-4 year olds (Fig. 2 B and C; Table S1). Estimated infection rates in the

younger age groups (in particular, 5-14, 15-24, and 25-44 year-olds; Fig. 2 D-F) peaked

during the week of March 22, 2020, followed by the three older age groups (i.e. 45-64, 65-

74, and 75+ year-olds; Fig. 2 G-I) about a week later.

Given the large uncertainties in model estimates, we verified our estimates of infection

rates using available serology data collected during three phases of the pandemic (i.e.

early-phase in March,[15] mid-phase in April,[16] and end of the pandemic wave in June[17]).

Overall, our estimated cumulative infection rates were in line with corresponding

measures from antibody tests, for all three phases of the pandemic wave (for details, see

the Appendix of Yang et al. 2020[11,12])

Overall effectiveness of interventions

The reproductive number at time-t (R_t) measures the average number of persons an

infected individual infects and thus reflects underlying epidemic dynamics. The epidemic

expands in size if R_t is above unity and subsides otherwise. In addition, when the entire

population is susceptible and no interventions are in place, R_t, referred to as the basic

reproductive number (R_0), reflects the transmissibility of an infection in that population.

Here we estimated that R_t was 2.99 [median and interquartile range (IQR): 2.32 – 3.86;

Table S2] during the first week of the pandemic (i.e. the week of March 1) in NYC, similar to

R_0 estimates reported for other places.[18,19] It decreased to around 2.2 during the next two

weeks, when NY State declared a state of emergency and public awareness and voluntary

precautionary measures (e.g. avoiding public transit[20]) increased (Fig 2A). Following the

stay-at-home mandate starting the week of March 22, R_t dropped substantially to 1.37

(IQR: 1.08 – 1.68) during that first week, to 0.93 (IQR: 0.73 – 1.13) a week later, and to a

minimum of 0.56 (IQR: 0.45 – 0.67) during the week of April 12 (Fig 2A). These prompt

decreases in R_t from mid-March to mid-April indicate that implemented public health

messaging and interventions were effective in curtailing COVID-19 transmission.

Similar decreases in R_t occurred among most age groups (Fig 2 B-I). Overall, R_t among

younger age groups (<45 years) decreased one or two weeks earlier than older age groups

(45-64, 65-74, and 75+; Fig 2 C-F vs. Fig 2 G-I). Of note, among the four age groups with

higher contact rates[21] (i.e. 5-14, 15-24, 25-44, and 45-64 year-olds), R_t dropped below 1 the

earliest among 5-14 year-olds (0.99, IQR: 0.74 – 1.30; Table S2) during the week of March

22. This is consistent with the earliest public health interventions to this age group: the

closure of public schools beginning the week of March 15.[7]

Effectiveness of reducing contact via school closure and voluntary or mandated stay-

at-home measures

Several public health interventions were implemented around the same time (Fig. 2), and

some interventions may take longer to produce an effect (e.g., due to slower compliance

with the measure). It is thus challenging to separate the impact of different interventions.

However, a number of measures – including voluntarily working from home during the

early weeks of the pandemic, school closures, and the stay-at-home mandate – in effect

reduce rates of close in-person contact, a key factor for COVID-19 transmission. Thus, here

we focus on estimating the impact of interventions whose primary mechanism of action is

through a reduction in population contact rates. Given the difficulties measuring this

quantity directly, we instead approximated population contact rates using human mobility

data, which record real time population movement based on location changes of individual

mobile devices (see Data). Indeed, the reduction in R_t mirrored the reduction in mobility

(Fig 3). The Pearson correlation (r) between R_t and mobility over the 14-week study period

was 0.96 for all ages overall and ≥0.9 for 1-4, 5-14, 15-24, and 25-44 year-olds (Table S3).

Thus, we focused on mobility as a proxy for contact rates and used this quantity to estimate

the corresponding changes in R_t and segregate the impact of interventions that reduce

population contact rates from other concurrent interventions.

Mobility reduced by 11.6% during the second week of the pandemic in NYC (i.e. the week of

March 8), and by a further 33.5% and 17.3% in the following two weeks, respectively (Fig

3). Using observed mobility data (i.e. our proxy for population contact rates) to estimate

the corresponding changes in R_t, we estimate that, for all ages overall, reductions in

population contact rates were associated with R_t reductions of 10.1% (95% CI: 8.3 –

11.9%) by the second week of the pandemic, and another 29.2% (95% CI: 24.9 – 33.5%)

and 15.0% (95% CI: 14.3 – 15.8%) in the following two weeks, respectively (Fig 3A). By the

week of April 12 when R_t reached its minimum, the reduction in population contact rates

was associated with an R_t reduction of 70.7% (95% CI: 65.0 – 76.4%). In addition, analysis

at the neighborhood level consistently showed large reductions in R_t that were likely due to

reductions in population contact rates (range of median estimates: 66.1 – 90.1% across the

42 neighborhoods in NYC; Fig S4).

In addition, transmission in four age groups (i.e., 5-14, 15-24, 25-44, and 45-64 year-olds)

appeared to be most impacted by changing population contact rates (Fig 3 and Table 1).

The reduction in population contact rates was associated with decreases of the age-specific

R_t by 83.4% (95% CI: 80.1 – 86.7%) for 5-14 year-olds, 65.4% (95% CI: 57.0 – 73.8%) for

15-24 year-olds, 76.5% (95% CI: 68.5 – 84.6%) for 25-44 year-olds, and 68.9% (95% CI:

59.2 – 78.6%) for 45-64 year-olds, by the week of April 12.

Effectiveness of face covering/masking under real-world conditions

Estimated transmission rates (or probability of infection) and the infectious period also

closely tracked changes in mobility (Table S3; $r \geq 0.5$ for most age groups). Thus, it appears

that reducing mobility not only reduces contact rates but also likely reduces 1) the

probability of transmission per contact due to, e.g., increased public spacing and 2) the

effective infectious period per infected individual due to, e.g., more time spent at home and

as a result reduced time for community transmission despite likely unchanged duration of

viral shedding. Given this observation, we hypothesize that the relationship between

mobility and estimated transmission rates (and effective infectious period) can be used to

disentangle the impact of interventions that reduce population contact rates (particularly,

the stay-at-home mandate) and face covering/masking — two major public health

interventions implemented in NYC during the pandemic — on transmission. We make two

predictions if this hypothesis holds. First, predicted transmission rates (infectious period)

using mobility data alone would be higher (longer) than those estimated by the model-

inference system additionally based on case and mortality data *for weeks when face*

covering in public was mandated as it would lead to further reductions in transmission (i.e.,

temporality and direction of the impact). Second, while the efficacy of masking (i.e.,

measured under ideal conditions of mask quality and correct use) likely does not vary by

individual, the effectiveness of masking (i.e., measured under real-world, often imperfect

conditions) and impact of this intervention could vary by subpopulation due, for example,

to different usage rates of masks; as such, we expect the predictive errors to be larger for

age groups with higher compliance of masking (i.e., magnitude of the impact). Our analyses

largely confirmed both predictions. For the first, as shown in Fig 4, transmission rates

predicted using a linear regression model with the observed mobility as the sole predictor

were higher than those estimated by the model-inference system, following the face

covering mandate starting the week of April 12.[9] For the latter, the discrepancies in the two

model estimates (i.e. the gaps between the dashed and solid blue lines; Fig 4) appeared to

increase with age and were largest among the two elderly age groups who have been

reported to more frequently use masks.[22-24] However, infants (<1 year) appeared to have a

larger reduction than other children groups; this could have been due to transmission

reduction related to their sources of infection (e.g. their caretakers and healthcare settings

where they tended to be exposed). Similar patterns held for the effective infectious period

(Fig 4).

Given these observations, we further used the discrepancies in the two model estimates to

approximate the impact of face covering on reducing COVID-19 transmission. Combining

the reduction in the transmission rate and effective infectious period, we estimated that,

for all ages combined, face covering contributed to a 6.6% (95% CI: 0.8 – 12.4%) reduction

during the first month it was implemented and a 3.4% (95% CI: -1.9 – 8.6%) reduction over

the entire 8 weeks prior to the city's reopening (Table 1). As expected, the estimated

impact varied substantially by age group. The effectiveness was 20.8% (95% CI: -0.1 –

41.6%) for 65-74 year-olds and 20.8% (95% CI: -0.9 – 42.5%) for 75+ year-olds during the

first month and remained at similar levels afterwards. For 25-44 and 45-64 year-olds, two

age groups with the highest infection rates (Fig 2), the effectiveness was 4.5% (95% CI: -0.6

– 9.7%) and 8.1% (95% CI: -0.1 – 16.1%) in the first month, respectively; however, it

reduced substantially afterwards, likely due to reversed risk behavior. Of note, in addition

to the likely lower usage rate of face covering in late May – early June, increases in risky

behaviors such as large gatherings at the time[25] may have partially obscured the

effectiveness of masking.

Retrospective projections of cases and deaths

NYC started phased reopening from the week of 6/7/2020, which allows industries to

gradually reopen per a four-phase plan.[26] For instance, manufacturing industries were

allowed to reopen starting the week of 6/7/2020 (dubbed "Phase 1"), whereas real estate

was allowed to reopen starting the week of 6/21/2020 (dubbed "Phase 2"), and personal

care services were allowed starting the week of 7/6/2020 (dubbed "Phase 3"). As such,

population mobility has increased gradually during this time, which could lead to increased

transmission of unknown magnitude. Such changes also offer an opportunity to test the

accuracy of our estimates – should the estimated effectiveness of reducing contact rates

and utilizing face coverings be accurate, these estimates could be used to anticipate

changes in transmission in response to the changing mobility and in turn the epidemic

dynamics after reopening. We thus used these estimates to generate projections of cases

and deaths for the 8 weeks beyond our study period, and compared the projections to

available, independent corresponding observations. Overall, our projections

underestimated the total number of cases (relative error of median projections: -27% over

8 weeks; Fig 5A) but were able to accurately estimate the total number of deaths (relative

error: -2% over 8 weeks; Fig 5B). In addition, examination of age-grouped projections

shows that the underestimation of cases was mostly among younger age groups whose

case rates had increased in June (1-4, 5-14, 15-24 and 25-44 year-olds; Fig S5). These

recent increases in young cases may have resulted from more young adults returning to

work including some in service industries with high contact rates and, relatedly, sending

their children to childcare and/or summer camps due to a lack of caretakers at home

[information from NYC Department of Health and Mental Hygiene (DOHMH) community

investigation; unpublished]. In addition, increased risk behaviors of some young

individuals (e.g., large parties without physical distancing[25]) may have also contributed to

the increased cases among young adults in late June.[27] Consistently, COVID-19 associated

mortality, mostly occurring among older adults continued to decrease and were accurately

predicted for different age groups (Fig S6).

Discussion

The spring 2020 pandemic wave in NYC, the first epidemic center in the US, provides a test

case to study COVID-19 epidemiological characteristics and the effectiveness of public

health interventions. Through comprehensive modeling, we have reconstructed the

transmission dynamics and estimated the effectiveness of two major interventions, social

distancing and mandatory face covering in public. Our results show that reducing contact

rates (mainly via school closures and voluntary or mandated stay-at-home measures) likely

contributed to the largest reduction in transmission in the population overall (~70%) and

for most age groups (>50% for all age groups). Widespread use of face covering likely

contributed to an additional ~7% overall reduction and up to ~20% reduction among 65+

year-olds during the first month face covering was mandated in public places. Our findings

largely consolidate previous model estimates on the impact of lockdown-like

measures[4,28,29] and studies on face covering in reducing COVID-19 transmission. These

findings provide insights that can inform COVID-19 mitigation efforts in the coming months

before the majority of population can be protected through mass-vaccination, as well as

control strategies for other (re)emerging infections in the future.

Lockdown-like measures where confinement at home is encouraged or mandated through

school closures, telework policies, closure of non-essential businesses, and stay-at-home

orders have been a major control measure to curb COVID-19 spread. In effect, such

measures reduce population contact rates and thus transmission. Previous modeling

studies estimated that lockdowns reduced COVID-19 transmission (measured by R_t) by

58% in Wuhan, China,[4] 45% (95% CI: 42-49%) in Italy,[29] and 77% (95% CI: 76-78%) in

France.[28] Our estimate for NYC overall (~70%) is consistent with these previous estimates.

In addition, our estimates show that reducing population contact rates effectively reduced

transmission across all age groups (ranging from a 51% reduction among 1-4 year-olds to

83% among 5-14 year-olds; Table 1). Together, these findings underscore the importance

of reducing contact rates through, for example, physical distancing in places with

continuous community transmission of COVID-19.

The use of surgical masks or cloth face coverings has been another major preventive

measure for COVID-19. Studies overall have shown that surgical masks could substantially

reduce onward transmission albeit with a large range of efficacy estimates across

settings.[30] However, it remains unclear the overall effectiveness of universal face covering

requirements at the population level, especially during a pandemic, due to several factors:

1) The overall effectiveness depends on compliance which may vary across subpopulations

and time; 2) Improper use of face coverings (e.g. without covering the nose and/or mouth

or improper handling[31]) can reduce the effectiveness of face covering; 3) Face coverings

are required and mostly worn in public and thus likely have a lower impact in private

settings, particularly in reducing household transmission; consequently, the relative impact

of face covering depends on the relative contribution of different sources of transmission

(e.g. household vs. community) at a given time and *vice versa*; and 4) Use of face coverings

may lead to complacency and less stringent adherence to social distancing and stay-at-

home behaviors. Here we estimated a ~7% reduction in overall transmission during the

first month of the face covering mandate. However, the estimated effectiveness varied

largely across age groups with much higher effectiveness among older adults (~20% for

both 65-74 and 75+ year-olds vs. <10% for other age groups). This discrepancy was likely

due to the differential compliance and types of face covering used. Observational studies in

Wisconsin and surveys nationwide in April/May reported about 2-fold higher rates of face

covering usage among older adults versus younger adults and minors.[22-24] In addition, due

340 to the shortage of surgical masks during March–May,[32] older adults at higher risk of severe

341 COVID-19 infection were more likely to use surgical masks whereas younger age groups

342 more frequently used non-medical cloth coverings, which are often less effective[33,34] (e.g.

343 measured ultrafine filtration efficiency is ~50% for surgical masks vs. ~10-25% for T-shirt

344 and ~25-35% for cotton covers[34]).

346 When lockdown-like measures are lifted, residents will spend more time outside their

347 homes than during the lockdown. Adjusting for the time spent outside of homes (~8.3

348 hours in April 2020 vs. ~11.5 hours in June-July 2020 and ~13.5 hours pre-pandemic; NYC

349 data[35]), universal face covering would have reduced overall transmission by ~9–11% (i.e.,

350 6.6% multiplied by a factor of 1.4–1.6) during reopening, given the same rates of face

351 covering as in April. However, if the same effectiveness among older adults were achieved

352 among other age groups, universal face covering could reduce overall transmission by up to

353 ~28–32% (i.e., 20% multiplied by a factor of 1.4–1.6). The implication of this latter

354 estimate is two-fold. On the one hand, it suggests that for places with high level of

355 transmission, implementing face covering *alone* is likely insufficient to lower the effective

356 reproductive number R_t to <1 in order to control the epidemic [for instance, for a city with

357 an R_0 = 3, the resulting R_t would be 3 × (1– 30%) = 2.1]. This finding is consistent with the

358 observed resurgence of COVID-19 cases in NYC during fall/winter 2020 despite the

359 concurrent high usage rate of face coverings (~90% of survey respondents in NYC reported

360 always or frequently wearing masks in public in July 2020;[36] and this number was likely

361 higher during fall/winter 2020).   On the other hand, our findings also suggest that

362 improving effective usage rates of face coverings, especially among younger age groups,

could significantly mitigate the risk of resurgence of COVID-19 infections during re-opening

(i.e., ~30% reduction without compromising economic growth). It is thus crucial for future

research to understand reasons for the use/non-use and selection of face coverings by age

group to inform strategies to increase consistent and correct mask use in settings where

social distancing is not possible.

It is important to note, however, that not all individuals have the same opportunities to

physically distance and/or adopt face coverings during a pandemic, despite government

mandates. For instance, over one million frontline workers in NYC (e.g., healthcare

workers, transportation workers, janitors, and grocery clerks, which comprise 25 percent

of the city's workforce) had to continue their essential work during the pandemic.[37] In

addition, data from the U.S. Bureau of Labor Statistics suggest Black and Latino

communities have less opportunities to work from home.[38] Consequently, NYC

neighborhoods with more frontline workers and/or Black and Latino residents tended to

have lower reductions in population mobility during the pandemic.[35] In NYC, these

communities also experience a number of social conditions that are thought to exacerbate

COVID-19, including overcrowded multigenerational households, poverty, and high

prevalence of chronic diseases. These communities, known to also carry a higher burden of

underlying health conditions, suffered greater impacts from COVID-19 and have expressed

fear and experiences of racialized bias when wearing a face covering.[39,40] Further research

is warranted to investigate such health disparities. In addition, future policies should take

into account structural inequities in labor trends, overcrowded housing, and underlying

conditions and adopt additional preventive measures to protect those vulnerable

communities.

We also note there remain large uncertainties in our estimates due to several limitations.

First, we used population mobility as a proxy for contact rates rather than more direct

measures. Similar approximation and uncertainty applied to our estimates of the

effectiveness of face covering. Future studies are thus warranted for further assessment.

For instance, large population scale surveys documenting changes of the intensity and

pattern of contact during the pandemic could provide more accurate measures of contact

rates among different age groups and over time. Second, while we restricted our analysis

on the effectiveness of face covering to a period when masks were mandated, there remain

other residual confounding effects. For instance, increased awareness of COVID-19 and

health risk among key age groups such as the elderly may have contributed to further

reductions of transmission through other precautions in addition to face covering; this may

have led to an overestimation of the effectiveness of face covering for those age groups.

Third, here we focused on estimating the effectiveness of interventions in the general

population without segregating key settings with intense transmission (e.g., long-term care

facilities). Future studies should assess the impact of interventions targeting such high-risk

settings. Lastly, our estimates here were largely based on the first wave of the pandemic

and may not fully capture subsequent changes in awareness and perception of COVID-19

and related behavioral adjustment during later waves. However, we have also used a

similar methodology to estimate the effectiveness of reducing contact rates and face

covering under different city reopening schedules and generated long-term projections for

NYC; the projections generated during June 2020 for the period of June 2020 – May 2021

have been consistent with observations up to the end of 2020 (i.e., at the time of this

writing; see the projected resurgence and second wave in Yang et al.[41] and comparison

with available data in Fig S7). These results thus support the robustness of our estimates

here.

Our study also has several strengths. In particular, our estimates were based on

comprehensive model-inference incorporating multiple data streams and further evaluated

using model projections. Our results thus provide an assessment of two major public heath

interventions (reducing contact rates and face covering) at the population level where the

overall effectiveness depends on multiple factors in addition to the efficacy of a given

intervention. Altogether, our estimates support the need for multiple interventions

(including reducing contact rates by, e.g., restricting occupancy, universal face covering,

and, albeit not studied here, testing, contact tracing, isolation and timely treatment of

cases) in order to effectively mitigate the spread of COVID-19 as it continues to pose

threats to public health.

Methods

Data

COVID-19 cases included all laboratory-confirmed cases by week of diagnosis reported to

the NYC DOHMH. Mortality data by week of death combined confirmed and probable

COVID-19-associated deaths. Confirmed COVID-19-associated deaths were defined as those

occurring in persons with laboratory-confirmed SARS-CoV-2 infection, and probable

COVID-19 deaths were defined as those with COVID-19, SARS-CoV-2, or a similar term

listed on the death certificate as an immediate, underlying, or contributing cause of death,

but did not have laboratory-confirmation of COVID-19.[42] For this study, both weekly case

and mortality data were aggregated by age group (<1, 1-4, 5-14, 15-24, 25-44, 45-64, 65-

74, and 75+ years) for each of the 42 United Hospital Fund (UHF) neighborhoods,[43]

according to the patient's residential address. All data were retrieved on Sep 4, 2020. For a

summary of the spatial variations across the 42 neighborhoods, see Table S3 in the

Appendix of Yang et al.[12]

The mobility data, used to model changes in population contact rates due to public health

interventions implemented during the pandemic (e.g., social distancing), came from

SafeGraph[35,44] and contained counts of visitors to locations in each zip code from the same

zip code and others, separately, based on mobile device locations. The released data were

anonymized and aggregated in weekly intervals. We spatially aggregated these data to the

UHF neighborhood level, for both intra and inter UHF neighborhood mobility. In addition,

SafeGraph also provided an aggregate measure of the length of time spent outside of the

home during each week.

This study was classified as public health surveillance and exempt from ethical review and

informed consent by the Institutional Review Boards of both Columbia University and NYC

DOHMH.

Network transmission model

The epidemic model used in this study was described in Yang et al. 2020[11,12] Briefly, the

model simulated intra- and inter neighborhood transmission of COVID-19 using a

susceptible-exposed-infectious-removed (SEIR) network model:

$$
\begin{cases}
\frac{dS_i}{dt} = -S_i \sum_{j=1}^{j=42} b_j \beta_{city} c_{ij} I_j / N_j \\
\frac{dE_i}{dt} = S_i \sum_{j=1}^{j=42} b_j \beta_{city} c_{ij} I_j / N_j - \frac{E_i}{Z} \\
\frac{dI_i}{dt} = \frac{E_i}{Z} - \frac{I_i}{D} \\
\frac{dR_i}{dt} = \frac{I_i}{D}
\end{cases}
\quad \text{(Eqn 1)}
$$

where S_i, E_i, I_i, R_i, and N_i are the numbers of susceptible, exposed (but not yet infectious),

infectious, and removed (either recovered or deceased) individuals and the total

population, respectively, from a given age group in neighborhood i. Note that due to model

complexity and a lack of information for parameterizing interactions among age groups, we

modeled each age group separately (i.e., combining all sources of transmission to each age

group; see further detail on parameter estimation below); as such, Eqn 1 describes the

spatial transmission across neighborhoods without interactions among age groups. β_{city} is

the citywide transmission rate, which incorporated seasonal variation as observed for

OC43, a beta-coronavirus in humans from the same genus as SARS-CoV-2.[12] To allow

differential transmission in each neighborhood, we included a multiplicative factor, b_i, to

scale neighborhood local transmission rates. Z and D are the latency and infectious

periods, respectively (Table S4).

The contact rates (c_{ij}) in each neighborhood over time and connectivity among

neighborhoods were computed based on mobility data. The model also accounted for

delays from infection to diagnosis using two parameters (gamma distribution with mean T_d

and standard deviation T_{sd} estimated along with other parameters) and death (based on

observed time from diagnosis to death) as well as infection detection rate using a

parameter r (estimated along with other parameters). For further detail, please refer to

Yang et al. 2020.[11,12]

Parameter estimation

To estimate model parameters (e.g., b_i, β_{city}, Z, D, r, and *infection fatality risk*, for $i=1,...,42$)

and state variables (e.g., number of susceptible and infectious individuals in each

neighborhood) for each week, we ran the network-model stochastically with a daily time

step in conjunction with the ensemble adjustment Kalman filter (EAKF)[45] and fit to weekly

case and mortality data from the week starting March 1 to the week ending June 6, 2020.

The posterior distribution of each model parameter/variable was updated for that week at

the same time.[45] This parameter estimation process was done separately for each of the

eight age groups (i.e. <1, 1-4, 5-14, 15-24, 25-44, 45-64, 65-74, and 75+ years). To include

transmission from other age groups, we used measured intra and inter-group contacts

from the POLYMOD study[21] to compute the total number of contacts made with each age

group and adjusted the *prior* range of the transmission rate (β_{city}) for each age group

accordingly. The posterior estimate was computed based on cases and mortality data for

each group, which included all sources of infection. Thus, the estimated transmission rate

for each age group nevertheless included all sources of transmission. To account for

stochasticity in model initiation, we ran the parameter estimation process independently

10 times. Results for each age group were combined from these 10 runs (each with 500

model realizations). We computed age-specific R_t, the effective reproductive number

during week-t, from the posterior estimates of transmission rate (β_{city} and b_i), infectious

period (D), contact matrix (c_{ij}), susceptibility and population size in the neighborhood

using the next generation method.[46] We computed R_t, β_{city}, and D estimates for all ages

overall as a weighted average of the age-specific estimates with weights equal to the

population fraction in each age group.

Estimating the effectiveness of reducing contact rates

The R_t estimates from the model-inference system capture changes in transmission due to

various interventions, i.e., the overall effectiveness of all implemented interventions. To

separately estimate the effectiveness of interventions that reduce contact rates, we used

human mobility as a measure of population contact rate to estimate the changes in R_t in

response to changing population contact rates. Specifically, we regressed the R_t estimates

from the full model-inference system on the mobility data: $R_t = a_0 + a_1 M_{ave,t}$ (Eqn 2),

where $M_{ave,t}$ is the mean of all intra-neighborhood mobility at week-t. We then computed

the effectiveness of reducing contact rate based on $\widehat{R_t}$, the R_t estimate from this regression

model solely based on the observed mobility. That is, the reduction in R_t by week-t, likely

due to reducing contact rate, was computed as $(\widehat{R_t} - \widehat{R_0})/\widehat{R_0}$ (Eqn 3). To test the robustness

of our method, we performed the same analysis for individual UHF neighborhoods (n = 42).

Estimating the effectiveness of face covering/masking

In addition to changes in population contact rates, face covering/masking was another

major control measure implemented beginning the week of April 12, 2020 when NYC

mandated residents wear face coverings in public places. To estimate the effectiveness of

face covering, we first estimated the changes in transmission rate and effective infectious

period, two model parameters determining R_t, due to changes in mobility (as opposed to

masking) using regression models similar to Eqn 2. Specifically, we regressed the estimated

citywide transmission rate (or effective infectious period) from the full model-inference

system on average mobility: $Y_t = a_0 + a_1 M_{ave,t}$ (Eqn 4), where Y_t is β_{city} or D. We then

computed the relative reduction in transmission rate (or effective infectious period) due to

face covering as $\eta_Y = E((Y_t - \widehat{Y}_t)/Y_t)$, where $E(\cdot)$ gives the mean over the relevant

timeframe (here we estimated two timeframes, i.e., 1 month following the mandate and

over 8 weeks up to 6/6/2020). Combining both reductions, we computed the effectiveness

of face covering as $\eta = 1 - (1 - \eta_\beta)(1 - \eta_D)$. Of note, while mechanistically face coverings

act primarily by reducing the probability of transmission (i.e., transmission rate), here we

included both the potential impact on the transmission rate (η_β) and effective infectious

period (η_D), mainly because the multiplicative relationship of the two variables with R_t

makes it challenging to separate the two effects. Nevertheless, reductions in the infectious

period via face covering are possible. A recent study on animals showed that masking could

reduce the severity of infection;[47] if persons with milder infection experience shorter

duration of viral shedding (there is some evidence for this, e.g., from He et al.[48]), milder

symptoms in individuals infected while wearing face covering could lead to shorter

infectious period of these individuals.

Projections of cases and deaths

To evaluate the accuracy of model estimates, in particular, the effectiveness of transmission

reduction by reducing contact rates and use of face covering, we tested if these estimates

along with the model could generate accurate predictions of cases and deaths for 8 weeks

beyond the study period (i.e. from the week of 6/7/2020 to the week of 7/26/2020). We

first projected the citywide transmission rate and infectious period based on observed

mobility using Eqn 4; these estimates thus accounted for changes due to changes in contact

rates. To incorporate the reduction in transmission by face covering, we further reduced

the projected city transmission rate by a factor of $1 - \eta_\beta p_{out}$ and the infectious period by a

factor of $1 - \eta_D p_{out}$, where p_{out} is a factor to adjust for time spent outside of the home

during each week. To reflect longer-term usage rates of face covering, we used η_β and η_D

estimated during the entire 8 weeks face covering was required (i.e. from the week of

4/12/2020 to the week of 5/31/2020). Finally, we used estimates of population

susceptibility and infection rates at the end of the week of 5/31/2020 to model initial

conditions and integrated the SEIR network model forward stochastically for 8 weeks using

the projected transmission rate and infectious period.

Acknowledgments:

This study was supported by the National Institute of Allergy and Infectious Diseases

(AI145883), the National Science Foundation Rapid Response Research Program (RAPID;

2027369), and the NYC DOHMH. We thank the NYC DOHMH Bureau of Vital Statistics team

and EpiData team; in particular, Mary Huynh, Sharon K. Greene, Alice Yeung, Anne Fine,

Miranda S. Moore and Kevin Guerra, for data management and provision. We thank Jennifer

Brite at NYC DOHMH for coordinating discussions on modeling COVID-19 in NYC and Alana

Tornello for amplifying community voices. We also thank Columbia University Mailman

School of Public Health for high performance computing, Safe Graph (safegraph.com) for

providing the mobility data, and Sasikiran Kandula at Columbia University for compiling

the mobility data used in this study. We are also grateful to Sasikiran Kandula, Sharon K.

Greene, Anne Fine, and Hannah Helmy for helpful feedback on this manuscript.

Author contributions

WY designed study, conducted the modeling analyses, and wrote the first draft. J Shaff

provided inputs on COVID-19 transmission dynamics, public health interventions and

community responses in NYC and critically revised the manuscript. J Shaman contributed

to study design, interpreted model results, and critically revised the manuscript.

Conflict of Interest

Shaman and Columbia University disclose partial ownership of SK Analytics. Shaman

discloses consulting for BNI. Other authors have nothing to disclose.

References:

1.   World Health Organization. Coronavirus disease (COVID-2019) situation reports.
10/5/2020 2020. https://www.who.int/emergencies/diseases/novel-coronavirus-
2019/situation-reports/ (accessed 10/6 2020).
2.   US Food and Drug Administration. FDA Takes Additional Action in Fight Against
COVID-19 By Issuing Emergency Use Authorization for Second COVID-19 Vaccine. 2020.
https://www.fda.gov/news-events/press-announcements/fda-takes-additional-action-
fight-against-covid-19-issuing-emergency-use-authorization-second-covid (accessed
12/23/20.

3.   Pharmacy Times. Pfizer, BioNTech Granted Authorization for COVID-19 Vaccine in

the European Union. 2020. https://www.pharmacytimes.com/news/pfizer-biontech-

granted-authorization-for-covid-19-vaccine-in-the-european-union (accessed 12/23/20.

4.   Li R, Pei S, Chen B, et al. Substantial undocumented infection facilitates the rapid

dissemination of novel coronavirus (SARS-CoV-2). *Science* 2020; **368**(6490): 489-93.

5.   The Novel Coronavirus Pneumonia Emergency Response Epidemiology Team. The

epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-

19)—China, 2020. *China CDC Weekly* 2020; **2**(8): 113-22.

6.   CDC COVID-19 Response Team. Preliminary estimates of the prevalence of selected

underlying health conditions among patients with coronavirus disease 2019—United

States, February 12–March 28, 2020. *Morbidity and Mortality Weekly Report* 2020; **69**(13):

382.

7.   New York City Office of the Mayor. New York City to Close All School Buildings and

Transition to Remote Learning. 3/15/2020 2020. https://www1.nyc.gov/office-of-the-

mayor/news/151-20/new-york-city-close-all-school-buildings-transition-remote-learning.

8.   New York State Department of Health. New York State on PAUSE. 2020.

https://coronavirus.health.ny.gov/new-york-state-pause (accessed 5/16/2020 2020).

9.   New York city Office of the Mayor. Transcript: Mayor de Blasio Holds Media

Availability on COVID-19. 4/12/2020 2020. https://www1.nyc.gov/office-of-the-

mayor/news/249-20/transcript-mayor-de-blasio-holds-media-availability-covid-19.

10.   Yang W, Kandula S, Shaman J. Projections of COVID19 Epidemic Outcomes and

Healthcare Demands for New York City (NYC). 2020. https://github.com/wan-

yang/COLUMBIA-COVID19-PROJECTIONS-FOR-NYC (accessed 5/12/2020 2020).

11.   Yang W, Kandula S, Huynh M, et al. Estimating the infection fatality risk of COVID-19

in New York City, March 1-May 16, 2020. *medRxiv* 2020: 2020.06.27.20141689.

12.   Yang W, Kandula S, Huynh M, et al. Estimating the infection-fatality risk of SARS-

CoV-2 in New York City during the spring 2020 pandemic wave: a model-based analysis.

The Lancet Infectious Diseases.

13.   Governor.ny.gov. After Weeks of Demanding Approval, Governor Cuomo Announces

FDA Gives New York State Authority to Conduct All COVID-19 Testing at Public and Private

Labs. March 13, 2020 2020. https://www.governor.ny.gov/news/after-weeks-demanding-

approval-governor-cuomo-announces-fda-gives-new-york-state-authority (accessed

6/22/2020 2020).

14.   Johnson CY, Sun LH, McGinley L. In hard-hit areas, testing restricted to health care

workers, hospital patients. March 21, 2020 2020.

https://www.washingtonpost.com/health/2020/03/21/coronavirus-testing-

strategyshift/ (accessed 6/22/2020 2020).

15.   Havers FP, Reed C, Lim T, et al. Seroprevalence of Antibodies to SARS-CoV-2 in 10

Sites in the United States, March 23-May 12, 2020. *Jama Intern Med* 2020.

16.   www.governor.ny.gov. Amid Ongoing COVID-19 Pandemic, Governor Cuomo

Announces Results of Completed Antibody Testing Study of 15,000 People Showing 12.3

Percent of Population Has COVID-19 Antibodies. MAY 2, 2020 2020.

https://www.governor.ny.gov/news/amid-ongoing-covid-19-pandemic-governor-cuomo-

announces-results-completed-antibody-testing (accessed 5/12/2020 2020).

17.   Government NYS. Video, Audio, Photos & Rush Transcript: Governor Cuomo

Announces U.S. Open to Be Held Without Fans from August 31st to September 13th.

6/16/2020 2020. https://www.governor.ny.gov/news/video-audio-photos-rush-

transcript-governor-cuomo-announces-us-open-be-held-without-fans-august.

18.   Li Q, Guan X, Wu P, et al. Early Transmission Dynamics in Wuhan, China, of Novel

Coronavirus-Infected Pneumonia. *N Engl J Med* 2020; **382**(13): 1199-207.

19.   Wu JT, Leung K, Leung GM. Nowcasting and forecasting the potential domestic and

international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling

study. *Lancet* 2020.

20.   Goldbaum C. Coronavirus in N.Y.: Riders Ditch the Subway as Fear Spreads.

3/13/2020 2020. https://www.nytimes.com/2020/03/13/nyregion/coronavirus-subway-

public-transit.html (accessed 6/22 2020).

21.   Mossong J, Hens N, Jit M, et al. Social contacts and mixing patterns relevant to the

spread of infectious diseases. *PLoS Med* 2008; **5**(3): e74.

22.   Arp NL, Nguyen TH, Graham Linck EJ, et al. Use of face coverings by the public

during the COVID-19 pandemic: an observational study. *medRxiv* 2020:

2020.06.09.20126946.

23.   Haischer MH, Beilfuss R, Hart MR, et al. Who is wearing a mask? Gender-, age-, and

location-related differences during the COVID-19 pandemic. *medRxiv* 2020:

2020.07.13.20152736.

24.   Ritter Z, Brenan M. New April Guidelines Boost Perceived Efficacy of Face Masks.

2020. https://news.gallup.com/poll/310400/new-april-guidelines-boost-perceived-

efficacy-face-masks.aspx (accessed 5/13/2020.

25.   Barone V, Marino J. Hundreds pack St. Marks Place to drink, party despite

coronavirus. 2020. https://nypost.com/2020/06/12/hundreds-pack-st-marks-place-to-

drink-despite-coronavirus/ (accessed 7/14 2020).

26.   New York State Government. Reopening New York City. 2020.

https://forward.ny.gov/reopening-new-york-city (accessed 7/13 2020).

27.   Durkin E. New York City sees uptick in coronavirus cases among young adults. 2020.

https://www.politico.com/states/new-york/albany/story/2020/07/13/new-york-city-

sees-uptick-in-coronavirus-cases-among-young-adults-1300223 (accessed 7/14 2020).

28.   Salje H, Tran Kiem C, Lefrancq N, et al. Estimating the burden of SARS-CoV-2 in

France. *Science* 2020; **369**(6500): 208-11.

29.   Gatto M, Bertuzzo E, Mari L, et al. Spread and dynamics of the COVID-19 epidemic in

Italy: Effects of emergency containment measures. *Proc Natl Acad Sci U S A* 2020; **117**(19):

10484-91.

30.   Liang M, Gao L, Cheng C, et al. Efficacy of face mask in preventing respiratory virus

transmission: A systematic review and meta-analysis. *Travel Med Infect Dis* 2020: 101751.

31.   Parker-Pope T. How NOT to Wear a Mask. 4/8/2020 2020.

https://www.nytimes.com/2020/04/08/well/live/coronavirus-face-mask-mistakes.html.

32.   Bernstein L, Safarpour A. Mask shortage for most health-care workers extended into

May, Post-Ipsos poll shows. 5/20/2020 2020.

https://www.washingtonpost.com/health/mask-shortage-for-most-health-care-workers-

extended-into-may-post-ipsos-poll-shows/2020/05/20/1ddbe588-9a21-11ea-ac72-

3841fcc9b35f_story.html.

33.   Fischer EP, Fischer MC, Grass D, Henrion I, Warren WS, Westman E. Low-cost

measurement of face mask efficacy for filtering expelled droplets during speech. *Science*

Advances 2020; **6**(36): eabd3083.

682 34. O'Kelly E, Pirog S, Ward J, Clarkson PJ. Ability of Fabric Facemasks Materials to Filter

683 Ultrafine Particles at Coughing Velocity - for Home Made and Fabric Face Mask Creation.

684 *medRxiv* 2020: 2020.04.14.20065375.

685 35. SafeGraph. Weekly Patterns: Foot Traffic Data To Understand The COVID-19

686 Pandemic. 2020. https://www.safegraph.com/weekly-foot-traffic-patterns.

687 36. New York Times. Mask-Wearing Survey Data. 2020.

688 https://github.com/nytimes/covid-19-data/tree/master/mask-use (accessed

689 12/22/2020.

690 37. Stringer SM. New York City's Frontline Workers. March 2020 2020.

691 https://comptroller.nyc.gov/reports/new-york-citys-frontline-workers/.

692 38. Gould E, Shierholz H. Not everybody can work from home: Black and Hispanic

693 workers are much less likely to be able to telework. 3/19/2020 2020.

694 https://www.epi.org/blog/black-and-hispanic-workers-are-much-less-likely-to-be-able-

695 to-work-from-

696 home/#:~:text=Less%20than%20one%20in%20five,race%20and%20ethnicity%2C%202

697 017–

698 2018&text=The%20data%20below%20can%20be%20saved%20or%20copied%20directl

699 y%20into%20Excel.&text=The%20next%20figure%20illustrates%20the,who%20can%20

700 telework%20by%20wage. (accessed 8/17 2020).

701 39. Taylor DB. For Black Men, Fear That Masks Will Invite Racial Profiling. 2020.

702 https://www.nytimes.com/2020/04/14/us/coronavirus-masks-racism-african-

703 americans.html.

704 40. Ruiz NG, Horowitz JM, Tamir C. Many Black and Asian Americans Say They Have

705 Experienced Discrimination Amid the COVID-19 Outbreak. 7/1/2020 2020.

706 https://www.pewsocialtrends.org/2020/07/01/many-black-and-asian-americans-say-

707 they-have-experienced-discrimination-amid-the-covid-19-outbreak/.

708 41. Yang W, Kandula S, Shaman J. Simulating Epidemic Outcomes under Different

709 Occupancy and Rollback Policies. Posted online on 6/30/2020. https://github.com/wan-

710 yang/re-opening_analysis/blob/master/report3_occupancy.pdf.

711 42. New York City Department of Health and Mental Hygiene (DOHMH) COVID-19

712 Response Team. Preliminary Estimate of Excess Mortality During the COVID-19 Outbreak

— New York City, March 11–May 2, 2020. *MMWR Morbidity and mortality weekly report*

2020; **69**: 603-5.

43.   NewYork City Department of Health and Mental Hygiene. NYC UHF 42

Neighborhoods. http://a816-dohbesp.nyc.gov/IndicatorPublic/EPHTPDF/uhf42.pdf.

44.   Lasry A, Kidder D, Hast M, et al. Timing of Community Mitigation and Changes in

Reported COVID-19 and Community Mobility - Four U.S. Metropolitan Areas, February 26-

April 1, 2020. *MMWR Morbidity and mortality weekly report* 2020; **69**(15): 451-7.

45.   Anderson JL. An Ensemble Adjustment Kalman Filter for Data Assimilation. *Mon*

Weather Rev 2001; **129**(12): 2884-903.

46.   Heffernan JM, Smith RJ, Wahl LM. Perspectives on the basic reproductive ratio. *J R*

Soc Interface 2005; **2**(4): 281-93.

47.   Chan JF, Yuan S, Zhang AJ, et al. Surgical mask partition reduces the risk of non-

contact transmission in a golden Syrian hamster model for Coronavirus Disease 2019

(COVID-19). *Clinical infectious diseases : an official publication of the Infectious Diseases*

Society of America 2020.

48.   He X, Lau EHY, Wu P, et al. Temporal dynamics in viral shedding and transmissibility

of COVID-19. *Nat Med* 2020; **26**(5): 672-5.

Table

Table 1. Estimated effectiveness of reducing contact rate and face covering. Numbers are

the estimated mean and 95% CIs, in percentage. Note the estimated effectiveness of contact

rate reduction combined all measures that reduce contact rates, including school closures

and voluntary or mandated stay-at-home measures.

	Estimated effectiveness of intervention (%)		
Age	contact rate reduction	Face covering (1st month)	Face covering (2 months)
all	70.7 (65.0, 76.4)	6.6 (0.8, 12.4)	3.4 (-1.9, 8.6)
<1	53.8 (41.6, 66)	9.3 (-4.2, 22.9)	12.8 (0.2, 25.3)
1-4	51.0 (45.8, 56.2)	0.9 (-5.5, 7.4)	6.7 (0.6, 12.8)
5-14	83.4 (80.1, 86.7)	3.0 (-0.5, 6.6)	1.6 (-1.6, 4.8)
15-24	65.4 (57.0, 73.8)	4.3 (-2.8, 11.4)	4.0 (-2.5, 10.6)
25-44	76.5 (68.5, 84.6)	4.5 (-0.6, 9.7)	-1.0 (-5.6, 3.7)
45-64	68.9 (59.2, 78.6)	8.1 (-0.1, 16.1)	4.4 (-2.9, 11.8)
65-74	55.8 (34.5, 77.2)	20.8 (-0.1, 41.6)	18.3 (-0.2, 36.9)
75+	53.8 (32.3, 75.3)	20.8 (-0.9, 42.5)	16.2 (-3.3, 35.7)

Figures

Fig 1. Epidemic dynamics. Reported laboratory confirmed cases (A) and cumulative cases

(B) per 100,000 population by week of diagnosis for all ages overall and by age group.

Fig 2. Estimated changes in the effective reproductive number and infection rates. Blue

lines show the estimated effective reproductive number (R_t) for each week; surrounding

areas show the 50% and 95% CrIs. Superimposed boxes (right y-axis) show estimated

infection rates by week: median (thick vertical lines), 50% CrIs (box edges), and 95% CrIs

(whiskers).

Fig 3. Effectiveness of reducing contact rates. Note the estimated effectiveness combined

all measures that reduce contact rates, including school closures and voluntary or

mandated stay-at-home measures. Dark grey lines show the observed changes in mobility

(right y-axis). Blue lines show R_t estimated using a linear regression model with mobility as

the sole predictor; surrounding areas show the 95% CrIs of the model estimates. The

adjusted r² for the regression model is also shown in each plot. For comparison, dashed

blue lines show R_t estimates from the model-inference system, without accounting for

susceptibility. Percentages attached to the lines show the incremental reductions in either

estimated R_t (in blue) or mobility (in grey).

Fig 4. Effectiveness of face covering in reducing the transmission rate and infectious period. Solid lines show the estimated transmission rate (in blue, left y-axis) and infectious period (in red, right y-axis) using the model-inference system incorporating interventions including face covering. Surrounding areas show the 50% CrIs of model estimates. Dashed lines show corresponding estimates from a linear regression model with mobility as the sole predictor (i.e without accounting for face covering).

766　**Fig 5.** Projections of COVID-19 cases and deaths eight weeks beyond the study period. Blue

767　dots show confirmed cases by week of diagnosis and deaths by week of death, as observed

768　by the surveillance system. Blue lines show model median estimates; surrounding shades

769　show 50% and 90% CrIs. Orange lines show model projected median weekly cases and

770　deaths; surrounding shades show 50% and 90% CIs of the projection.

Supplemental Tables and Figures

Supplemental Tables

Table S1. Total number of reported cases and estimated cumulative infection rate by the week of 5/31/20. Case rate was computed as the number of total cases divided by the population size of the corresponding age group. Estimated infection rate was estimated by model-inference system, normalized to the corresponding population size; numbers are median (and 95% CrIs).

Age	Total number of cases	Case rate (%)	Estimated infection rate (%)
all	205693	2.45	17.6 (13.2, 25.5)
<1	515	0.46	4.4 (2.9, 10.5)
1-4	609	0.14	1.4 (0.9, 3.7)
5-14	2646	0.28	10.5 (6.5, 24.5)
15-24	12562	1.28	12.0 (8.4, 19.5)
25-44	64764	2.45	22.7 (16.8, 31.5)
45-64	74833	3.64	23.1 (18.4, 29.6)
65-74	25469	3.64	15.2 (11.4, 21.7)
75+	24295	4.44	12.7 (9.8, 18.4)

Table S2. Estimated reproductive number by week and age group. Numbers are median (and interquartile range) of the posterior estimates.

date	Age Groups								
	all	<1	1-4	5-14	15-24	25-44	45-64	65-74	75+
3/1/20	2.99 (2.32, 3.86)	1.52 (1.19, 1.96)	1.56 (1.22, 2.02)	3.15 (2.47, 4.09)	2.47 (1.93, 3.22)	4 (3.1, 5.15)	2.97 (2.3, 3.86)	1.77 (1.37, 2.29)	1.75 (1.35, 2.27)
3/8/20	2.25 (1.64, 2.98)	1.43 (1.11, 1.85)	1.28 (1, 1.7)	2.35 (1.74, 3.18)	2.02 (1.53, 2.64)	2.46 (1.57, 3.48)	2.49 (1.97, 3.12)	1.98 (1.55, 2.44)	1.84 (1.48, 2.29)
3/15/20	2.14 (1.62, 2.74)	1.63 (1.32, 2)	1.17 (0.92, 1.48)	1.52 (1.13, 2.06)	2.38 (1.86, 3)	1.45 (0.97, 2.03)	2.99 (2.3, 3.68)	2.84 (2.33, 3.51)	2.87 (2.33, 3.52)
3/22/20	1.37 (1.08, 1.68)	1.59 (1.35, 1.89)	1.05 (0.85, 1.29)	0.99 (0.74, 1.3)	1.29 (1.03, 1.57)	1 (0.76, 1.26)	1.37 (1.06, 1.71)	2.58 (2.15, 3.04)	2.52 (2.06, 3.02)
3/29/20	0.93 (0.73, 1.13)	1.22 (0.97, 1.44)	1.06 (0.82, 1.3)	0.88 (0.66, 1.14)	0.96 (0.75, 1.15)	0.7 (0.52, 0.88)	0.89 (0.71, 1.06)	1.43 (1.17, 1.65)	1.46 (1.27, 1.66)
4/5/20	0.63 (0.5, 0.75)	0.74 (0.6, 0.88)	0.84 (0.65, 1.03)	0.53 (0.41, 0.68)	0.77 (0.59, 0.91)	0.44 (0.33, 0.56)	0.62 (0.5, 0.73)	0.9 (0.79, 1.02)	0.9 (0.78, 1.02)
4/12/20	0.56 (0.45, 0.67)	0.67 (0.54, 0.79)	0.69 (0.53, 0.84)	0.5 (0.38, 0.62)	0.72 (0.57, 0.83)	0.44 (0.34, 0.55)	0.57 (0.47, 0.67)	0.63 (0.55, 0.71)	0.65 (0.57, 0.74)
4/19/20	0.59 (0.48, 0.7)	0.67 (0.54, 0.8)	0.68 (0.54, 0.81)	0.57 (0.44, 0.7)	0.81 (0.66, 0.93)	0.55 (0.42, 0.68)	0.49 (0.41, 0.57)	0.6 (0.52, 0.68)	0.66 (0.58, 0.74)
4/26/20	0.62 (0.5, 0.73)	0.62 (0.5, 0.75)	0.77 (0.62, 0.91)	0.54 (0.42, 0.67)	0.75 (0.61, 0.86)	0.58 (0.45, 0.72)	0.6 (0.5, 0.71)	0.58 (0.5, 0.67)	0.63 (0.56, 0.71)
5/3/20	0.66 (0.54, 0.77)	0.7 (0.56, 0.84)	0.78 (0.62, 0.91)	0.56 (0.44, 0.67)	0.72 (0.61, 0.8)	0.68 (0.54, 0.82)	0.68 (0.56, 0.78)	0.6 (0.52, 0.67)	0.55 (0.48, 0.63)
5/10/20	0.82 (0.67, 0.95)	0.71 (0.56, 0.86)	0.75 (0.6, 0.88)	0.6 (0.47, 0.73)	0.88 (0.73, 0.98)	0.9 (0.72, 1.06)	0.88 (0.74, 0.99)	0.73 (0.64, 0.81)	0.65 (0.57, 0.74)
5/17/20	1.03 (0.83, 1.18)	0.82 (0.65, 0.99)	0.8 (0.65, 0.93)	0.73 (0.57, 0.88)	1.1 (0.89, 1.24)	1.21 (0.95, 1.42)	1.1 (0.91, 1.24)	0.78 (0.69, 0.87)	0.81 (0.72, 0.92)
5/24/20	0.89 (0.73, 1.03)	0.78 (0.61, 0.95)	0.79 (0.63, 0.91)	0.69 (0.53, 0.83)	0.94 (0.78, 1.06)	0.96 (0.76, 1.13)	0.95 (0.78, 1.08)	0.79 (0.7, 0.88)	0.89 (0.78, 0.99)
5/31/20	0.79 (0.66, 0.91)	0.71 (0.55, 0.87)	0.71 (0.56, 0.83)	0.7 (0.55, 0.86)	0.85 (0.72, 0.95)	0.79 (0.64, 0.93)	0.77 (0.65, 0.89)	0.72 (0.64, 0.79)	1.04 (0.93, 1.15)

Table S3. Correlation of key epidemiological parameters with population mobility during the week of March 1 – the week of May 31, 2020.

Age	Rt (ignore susceptibility)	Rt	Transmission rate	Infectious period
all	0.96	0.96	0.73	0.54
<1	0.76	0.77	0.41	0.70
1-4	0.93	0.93	0.05	0.65
5-14	0.99	0.99	0.86	0.92
15-24	0.90	0.91	0.52	-0.05
25-44	0.93	0.96	0.80	0.27
45-64	0.88	0.90	0.60	0.24
65-74	0.57	0.63	0.54	0.65
75+	0.55	0.60	0.50	0.64

Table S4. Prior ranges for main model parameters and variables. The spatial, temporal, and age resolution of each parameter or variable, estimated in the model-inference system, is specified in the column "Resolution". Note posterior parameter estimates can extend outside the specified prior ranges. Note this is the same as Table S1 in Yang et al.(*1*)

Parameter/ variable	Symbol	Resolution	Prior range	Source/rationale
Initial exposed	E(t=0)	neighborhood- and age-group specific, estimated for the beginning of the Week of March 1, 2020	300 – 8000 total citywide, scaled by population size for each age group and neighborhood	Large uncertainties, used very wide range
Initial infectious	I(t=0)	neighborhood- and age-group specific, , estimated for the beginning of the Week of March 1, 2020	150 – 4000 total citywide, scaled by population size for each age group and neighborhood	Assumed to be half the initial exposed
Initial susceptible	S(t=0)	neighborhood- and age-group specific, estimated for the beginning of the Week of March 1, 2020	$N – E – I$	Assumed all were susceptible except for those initially exposed/infectious
Population size in each age group and neighborhood	N	neighborhood- and age-group specific	N/A	NYC intercensal population estimates for 2018(*2*)
Citywide transmission rate	β_{city}	Citywide, age-group specific, estimated for each week	[0.5, 1] per day overall; scaled by contact rate for each age group based on contact data from the POLYMOD study(*3*) (averaged across 8 countries)	Based on R_0 estimates of around 1.5-4 for SARS-CoV-2(*4-6*)
Scaling of neighborhood transmission rate	b_i	neighborhood- and age-group specific, estimated for each week	[0.8, 1.2] for age groups under 65 years; [0.5, 1.5] for age groups 65 or older	Around 1; larger variation for elderly groups based on data

Latency period	Z	Citywide, age-group specific, estimated for each week	[2, 5] days	Incubation period: 5.2 days (95% CI: 4.1, 7)(*4*); latency period is likely shorter than the incubation period
Infectious period	D	Citywide, age-group specific, estimated for each week	[2, 5] days	Time from symptom onset to hospitalization: 3.8 days (95% CI: 0, 12.0) in China,(*7*) plus 1-2 days viral shedding before symptom onset. We did not distinguish symptomatic/asymptomatic infections.
Multiplicative factor for mobility	m_1	Citywide, age-group specific, estimated for each week	[1, 2] for <1 year; [0.5, 1.5] for three age groups 1-24 years; [0.1, 1.5] for age group 25-44; [1, 2.5] for age groups 45 or older	Initial model testing showed transmission rates for younger age groups were more sensitive to changes in mobility whereas the two oldest age groups were not sensitive to mobility. For age groups with contact rates lower than the average (based on the POLYMOD study(*3*)), we raised the diagonal elements in the mobility matrix to the power of the relative contact rate (<1) to account for insensitivity of transmission rate in these age groups to mobility.
Multiplicative factor for neighborhood connectivity	m_2	Citywide, age-group specific, estimated for each week	[0.5, 2]	Likely around 1 but with large uncertainties
Mean of time from viral shedding to diagnosis	T_m	Citywide, age-group specific, estimated for each week	[3, 8] days	From a few days to a week from symptom onset to diagnosis/ reporting,(*7*) plus 1-2 days of viral shedding (being infectious) before symptom onset

Standard deviation (SD) of time from viral shedding to diagnosis	T_{sd}	Citywide, age-group specific, estimated for each week	[1, 3] days	To allow variation in time to diagnosis/reporting
Reporting rate r		Citywide, age-group specific, estimated for each week	Starting from [0.001, 0.05] at time 0 and allowed to increase over time using space re-probing[8]	Large uncertainties
Infection fatality risk (IFR)		Citywide, age-group specific, estimated for each week	$[5, 15] \times 10^{-5}$ for ages under 25; $[5, 15] \times 10^{-4}$ for ages 25-44; $[5, 15] \times 10^{-3}$ for ages 45-64; [0.01, 0.1] for ages 65-74; [0.02, 0.2] for ages 75+;	Based on previous estimates[9] but extend to have wider ranges
Time from diagnosis to death		Citywide	Gamma distribution with mean of 9.36 days and SD of 9.76 days	Based on n=15,686 COVID-19 confirmed deaths in NYC as of May 17, 2020.

Reference:
1. W. Yang, S. Kandula, M. Huynh, S. K. Greene, G. Van Wye, W. Li, H. T. Chan, E. McGibbon, A. Yeung, D. Olson, A. Fine, J. Shaman, Estimating the infection fatality risk of COVID-19 in New York City, March 1-May 16, 2020. *medRxiv*, 2020.2006.2027.20141689 (2020).
2. New York City Department of Health and Mental Hygiene.
3. J. Mossong, N. Hens, M. Jit, P. Beutels, K. Auranen, R. Mikolajczyk, M. Massari, S. Salmaso, G. S. Tomba, J. Wallinga, J. Heijne, M. Sadkowska-Todys, M. Rosinska, W. J. Edmunds, Social contacts and mixing patterns relevant to the spread of infectious diseases. *PLoS Med* **5**, e74 (2008).
4. Q. Li, X. Guan, P. Wu, X. Wang, L. Zhou, Y. Tong, R. Ren, K. S. M. Leung, E. H. Y. Lau, J. Y. Wong, X. Xing, N. Xiang, Y. Wu, C. Li, Q. Chen, D. Li, T. Liu, J. Zhao, M. Liu, W. Tu, C. Chen, L. Jin, R. Yang, Q. Wang, S. Zhou, R. Wang, H. Liu, Y. Luo, Y. Liu, G. Shao, H. Li, Z. Tao, Y. Yang, Z. Deng, B. Liu, Z. Ma, Y. Zhang, G. Shi, T. T. Y. Lam, J. T. Wu, G. F. Gao, B. J. Cowling, B. Yang, G. M. Leung, Z. Feng, Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus–Infected Pneumonia. *New Engl J Med*, (2020).
5. J. T. Wu, K. Leung, G. M. Leung, Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *Lancet*, (2020).
6. R. Li, S. Pei, B. Chen, Y. Song, T. Zhang, W. Yang, J. Shaman, Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV-2). *Science* **368**, 489-493 (2020).

7.   J. Zhang, M. Litvinova, W. Wang, Y. Wang, X. Deng, X. Chen, M. Li, W. Zheng, L. Yi, X. Chen, Q. Wu, Y. Liang, X. Wang, J. Yang, K. Sun, I. M. Longini, Jr., M. E. Halloran, P. Wu, B. J. Cowling, S. Merler, C. Viboud, A. Vespignani, M. Ajelli, H. Yu, Evolving epidemiology and transmission dynamics of coronavirus disease 2019 outside Hubei province, China: a descriptive and modelling study. *The Lancet. Infectious diseases*, (2020).

8.   W. Yang, J. Shaman, A simple modification for improving inference of non-linear dynamical systems. *arXiv*, 1403.6804 (2014).

9.   R. Verity, L. C. Okell, I. Dorigatti, P. Winskill, C. Whittaker, N. Imai, G. Cuomo-Dannenburg, H. Thompson, P. G. T. Walker, H. Fu, A. Dighe, J. T. Griffin, M. Baguelin, S. Bhatia, A. Boonyasiri, A. Cori, Z. Cucunuba, R. FitzJohn, K. Gaythorpe, W. Green, A. Hamlet, W. Hinsley, D. Laydon, G. Nedjati-Gilani, S. Riley, S. van Elsland, E. Volz, H. Wang, Y. Wang, X. Xi, C. A. Donnelly, A. C. Ghani, N. M. Ferguson, Estimates of the severity of coronavirus disease 2019: a model-based analysis. *The Lancet. Infectious diseases*, (2020).

Supplemental Figures

Fig S1. Model fits of reported confirmed COVID-19 cases. Blue dots show reported number of cases by age group and week of diagnosis. Boxes show the model fitted weekly number of cases by age group. Box edges, thick middle lines, and whiskers show the 2.5th, 25th, 50th, 75th, and 97.5th percentiles of model estimates.

Fig S2. Model fits of reported COVID-19 associated deaths. Blue dots show reported weekly number of deaths by age group. Boxes show the model fitted weekly number of deaths by age group. Box edges, thick middle lines, and whiskers show the 2.5th, 25th, 50th, 75th, and 97.5th percentiles of model estimates. Note that deaths among <25 year-olds are combined due to low counts.

Fig S3. Estimated case rates and infection rates by age group. Blue dots show confirmed case rates by age group and week of diagnosis. Blue boxes (left y-axis) show the model fitted weekly case rates and grey boxes (right y-axis) show the model estimated weekly infection rates by age group. Box edges, thick middle lines, and whiskers show the 2.5th, 25th, 50th, 75th, and 97.5th percentiles of model estimates.

Fig S4. Sensitivity analysis on the effectiveness of reducing contact rate by neighborhood. There are 42 United Hospital Fund (UHF) neighborhoods in NYC. (A) shows the changes in human mobility during the pandemic by neighborhood (each colored line). The reductions were substantial in all neighborhoods but to varying degrees. (B) shows the estimated R_t (combining all ages) for each week and neighborhood (each colored line). Note these estimates did not account for changes in susceptibility so as to restrict to changes due to interventions. (C) shows the adjusted r^2 of linear regression model fitting the mobility data (A) to the R_t estimates (B), for each neighborhood, per Eqn 2 in the main text. Adjusted r^2 for most neighborhoods was >0.9. (D) shows the estimated reduction in R_t by the Week of April 12, 2020 based on the Eqn 3 in the main text, for each neighborhood. The histogram is based on the median estimated reduction in R_t.

Fig S5. Projections of COVID-19 cases by age group eight weeks beyond the study period. Blue dots show observed confirmed cases by week of diagnosis (those after the Week of 5/31/2020 were not used in the model). Blue lines show model median estimates; surrounding shades show 50% and 90% CrIs. Orange lines show model projected median weekly cases and deaths; surrounding shades show 50% and 90% CIs of the projection.

Fig S6. Projections of COVID-19 associated deaths by age group eight weeks beyond the study period. Blue dots show observed weekly deaths (those after the Week of 5/31/2020 were not used in the model). Blue lines show model median estimates; surrounding shades show 50% and 90% CrIs. Orange lines show model projected median weekly cases and deaths; surrounding shades show 50% and 90% CIs of the projection.

Fig S7. Comparing long term projection with available observations. The projections on COVID-19 cases, hospitalizations and deaths were generated on June 30, 2020, using case and mortality data up to June 26, 2020. These projections were generated at the time using a simpler model without age grouping but otherwise the same methodology presented in the study. Results were posted online on June 30, 2020 (https://github.com/wan-yang/re-opening_analysis/tree/master/test3_occupancy). Various policy scenarios were tested and here we used the one labeled "sce2fix_asIs" for the "intervention" identifier, which has been the closest to implemented interventions thus far (main settings were capping capacity for all industries including schools at 50%; no shutdown). For comparison with observations thus far, we used data published by the New York City Department of Health and Mental Hygiene, as of Dec 27, 2020 (https://raw.githubusercontent.com/nychealth/coronavirus-data/master/trends/data-by-day.csv). Red line and shaded area show our projections (median and interquartile range) and blue dots show corresponding observations from the New York City Department of Health and Mental Hygiene. Note that estimated infection-detection rates using PCR tests during June – Dec 2020 have been similar to those in June. However, hospitalization rates and estimated infection-fatality-risk in months after June 2020 have been lower than that during the spring wave; these lower hospitalization rates and mortality risks contribute to the lower observed numbers than our projections.