1

COVID-19 Transmission Dynamics and Effectiveness of Public Health Interventions in New York City during the 2020 Spring Pandemic Wave

Wan Yang,[1*] Jaimie Shaff,[2] Jeffrey Shaman[3]

[1]Department of Epidemiology, Mailman School of Public Health, Columbia University, New York, NY, USA; [2]New York City Department of Health and Mental Hygiene, New York, NY, USA; [3]Department of Environmental Health Sciences, Mailman School of Public Health, Columbia University, New York, NY, USA

*Correspondence to: wy2202@cumc.columbia.edu

Abstract

New York City experienced a large COVID-19 pandemic wave during March – May 2020. We model the transmission dynamics of COVID-19 in the city during the pandemic and estimate the effectiveness of public health interventions (overall and for each major intervention separately) for the entire population and by age group. We estimate that the overall effective reproductive number was 2.99 at the beginning of the pandemic wave and reduced to 0.93 one week after the stay-at-home mandate. Most age groups experienced similar reductions in transmission. Interventions reducing contact rates were associated with a 70.7% (95% CI: 65.0 – 76.4%) reduction of transmission overall and >50% for all age groups during the pandemic. Face covering was associated with a 6.6% (95% CI: 0.8 – 12.4%) reduction of transmission overall and up to 20% for 65+ year-olds during the first month of implementation. Accounting for the amount of time masks are in use (i.e. mainly outside homes), these findings indicate universal masking could reduce transmission by up to 28-32% when lockdown-like measures are lifted, if the high effectiveness estimated for older adults were achieved for all ages. These estimates are verified by out-of-fit projections and support the need for implementing multiple interventions simultaneously in order to effectively mitigate the spread of COVID-19.

Keywords: COVID-19; transmission dynamics; effectiveness of intervention; social distancing; lockdown; face covering

Introduction

New York City (NYC) is the largest and most densely populated municipality in the United States with over 60 million travelers each year from places worldwide.(*1*) It is thus perhaps no surprise that NYC became the United States' first epidemic center of the COVID-19 (coronavirus disease 2019) pandemic. NYC experienced widespread COVID-19 transmission citywide since early March and recorded over 200,000 cases and over 21,000 COVID-19 confirmed or probable deaths during the following three months. To curb this intense transmission, NY State and NYC implemented multiple intervention measures, including health promotion campaigns in early March, telecommuting and staggered work schedule recommendations beginning the week of March 8, public schools closure starting the week of March 15,(*2*) stay-at-home orders for non-essential workers starting the week of March 22,(*3*) and requirements for use of face covering in public starting the week of April 12.(*4*) With these overlapping and far reaching public health interventions, case diagnoses and hospitalizations peaked in April, and started to decline substantially in late April and May. NYC was able to begin its phased re-opening of industries starting the week of June 7, 2020.

Despite the overwhelming pandemic wave in Spring 2020, serology surveys indicated that the majority of NYC residents (about 80%; as of June 13, 2020(*5*)) remain susceptible to infection, and thus future epidemics are possible without effective public health interventions. Similar challenges confront populations in other locales in the United States and other countries. It is thus critical to understand the transmission dynamics of COVID-19 in the Spring 2020 pandemic wave and the effectiveness of different public health interventions, in order to more effectively mitigate future COVID-19 epidemics and the devastating morbidity, mortality, and economic burden.

In this study, we apply a model-inference system developed to support the city's COVID-19 pandemic response to reconstruct the underlying transmission dynamics of COVID-19 in NYC during March 1 – June 6, 2020 (i.e. prior to the city's reopening). The inference of COVID-19 transmission dynamics is challenging due to low infection detection rates (many asymptomatic

and mild infections do not seek care or receive testing),(*6*) fluctuation of those infection

detection rates, differential disease manifestation by age group, and concurrent public health

interventions. To address these challenges, our model-inference system simultaneously

assimilates three sources of data: 1) confirmed COVID-19 case data, 2) COVID-19 associated

death data (both cases and deaths are assimilated by neighborhood and age group), and 3)

neighborhood-level mobility data to constrain the model system. This enables inference of the

overall infection rate (i.e. including those not documented by surveillance), estimation of key

transmission characteristics (e.g., the reproductive number) through time, and assessment of

the effectiveness of different public health interventions, including social distancing and face

covering, implemented over time. We further incorporate these estimates to project cases and

deaths in the weeks beyond our study period and compare the projections to independent

observations in order to evaluate the accuracy of these estimates.

Results

Overall Epidemic Trends

Following diagnosis of the first case in NYC, confirmed COVID-19 cases in the entire population

increased nearly exponentially during the first three weeks (Fig 1B) before slowing down

beginning the week of March 22, 2020 when NYC implemented a stay-at-home order.

However, case trajectories differed substantially by age group. Foremost, reported cases

increased with age: the case trajectory for those aged 25-44 years mirrored the overall

epidemic curve, those older than 45 years had higher case rates, and those under 25 years had

the lowest case rates (Fig 1A and Table S1). Infants (i.e. <1 year), however, had higher case

rates than 1-4 and 5-14 year-olds (Fig 1). In addition, the timing of peak case rate was mixed.

Case rates in 25-44, 45-64, and 65-74 year-olds peaked earliest during the week of March 29,

2020, followed by <1, 1-4, 15-24, and 75+ year-olds with a 1-week lag; in comparison, the case

rate for 5-14 year-olds fluctuated with a less clear peak during the weeks of March 29 – April

26, 2020.

The epidemic trends based on diagnosed cases, however, were obscured by varying infection

detection rates by age and through time. COVID-19 infections are more likely to manifest as

symptomatic illness and/or more severe disease in individuals with underlying conditions and in

older adults.(*7, 8*) Such differential clinical characteristics by age thus lead to varying healthcare

seeking behaviors and infection detection rates by age. In addition, testing policies varied over

the course of the Spring 2020 COVID-19 pandemic in NYC. During this time, testing capacity was

limited at the federal, state and local levels by guidelines for who should be tested (due, for

example, to limited availability of test kits, swabbing supplies and reagents), which required

prioritizing testing for severely ill patients and those highly vulnerable to severe disease. Testing

capacities expanded during the week of March 8, 2020;(*9*) however, by late March, material

shortages (including testing kits and personal protective equipment) again prompted the city to

restrict testing to those severely ill.(*10*) Using our model-inference system, we estimated that

infection detection rates increased in early March, reaching a peak of around 20% for all ages

overall during the week of March 15, and declined afterwards before increasing again in mid-

April.(*11*)

After accounting for infection detection rates to include undiagnosed infections, a different

picture of the NYC spring outbreak emerges (Fig 2 and Fig S3). Estimated infection rates were

highest among 25-44 and 45-64 year-olds (Fig. 2 E and F), followed by 65-74 and 75+ year-olds

(Fig. 2 G and H), then 5-14 and 15-24 year-olds (Fig. 2 C and D), and were lowest among <1 and

1-4 year olds (Fig. 2 A and B; Table S1). Estimated infection rates in the younger age groups (in

particular, 5-14, 15-24, and 25-44 year-olds; Fig. 2 C-E) peaked during the week of March 22,

2020, followed by the three older age groups (i.e. 45-64, 65-74, and 75+ year-olds; Fig. 2 F-H)

about a week later.

Overall effectiveness of interventions

The reproductive number at time-*t* (R_t) measures the average number of persons an infected

individual infects and thus reflects underlying epidemic dynamics. The epidemic expands in size

if R_t is above unity and subsides otherwise. In addition, when the entire population is

susceptible and no interventions are in place, R_t, referred to as the basic reproductive number

(R_0), reflects the transmissibility of an infection in that population. Here we estimated that R_t

was 2.99 [median and interquartile range (IQR): 2.32 – 3.86; Table S2] during the first week of

the pandemic (i.e. the week of March 1) in NYC, similar to R_0 estimates reported for other

places.(*12, 13*) It decreased to around 2.2 during the next two weeks, when NY State declared a

state of emergency and public awareness and voluntary precautionary measures (e.g. avoiding

public transit(*14*)) increased (Fig 2A). Following the stay-at-home mandate starting the week of

March 22, R_t dropped substantially to 1.37 (IQR: 1.08 – 1.68) during that first week, to 0.93

(IQR: 0.73 – 1.13) a week later, and to a minimum of 0.56 (IQR: 0.45 – 0.67) during the week of

April 12 (Fig 2A). These prompt decreases in R_t from mid-March to mid-April indicate that

implemented public health messaging and interventions were effective in curtailing COVID-19

transmission.

Similar decreases in R_t occurred among most age groups (Fig 2 B-H). Overall, R_t among younger

age groups (<45 years) decreased one or two weeks earlier than older age groups (45-64, 65-74,

and 75+; Fig 2 C-F vs. Fig 2 G-L). Of note, among the four age groups with higher contact

rates(*15*) (i.e. 5-14, 15-24, 25-44, and 45-64 year-olds), R_t dropped below 1 the earliest among

5-14 year-olds (0.99, IQR: 0.74 – 1.30; Table S2) during the week of March 22. This is consistent

with the earliest public health interventions to this age group: the closure of public schools

beginning the week of March 15.(*2*)

Effectiveness of reducing contact via school closure and voluntary or mandated stay-at-home

measures

Several public health interventions were implemented around the same time (Fig. 2), and some

interventions may take longer to produce an effect (e.g., due to slower compliance with the

measure). It is thus challenging to separate the impact of different interventions. However, a

number of measures – including voluntarily working from home during the early weeks of the

pandemic, school closures, and the stay-at-home mandate – in effect reduce rates of close in-

person contact, a key factor for COVID-19 transmission. Thus, here we focus on estimating the

impact of interventions whose primary mechanism of action is through a reduction in

population contact rates. Given the difficulties measuring this quantity directly, we instead

approximated population contact rates using human mobility data, which record real time

population movement based on location changes of individual mobile devices (see Data).

Indeed, the reduction in R_t mirrored the reduction in mobility (Fig 3). The Pearson correlation

(*r*) between R_t and mobility over the 14-week study period was 0.96 for all ages overall and ≥0.9

for 1-4, 5-14, 15-24, and 25-44 year-olds (Table S3). Thus, we focused on mobility as a proxy for

contact rates and used this quantity to estimate the corresponding changes in R_t and segregate

the impact of interventions that reduce population contact rates from other concurrent

interventions.

Mobility reduced by 11.6% during the second week of the pandemic (i.e. the week of March 8),

and by a further 33.5% and 17.3% in the following two weeks, respectively (Fig 3). Using

observed mobility data (i.e. our proxy for population contact rates) to estimate the

corresponding changes in R_t, we estimate that, for all ages overall, reductions in population

contact rates were associated with R_t reductions of 10.1% (95% CI: 8.3 – 11.9%) by the second

week of the pandemic, and another 29.2% (95% CI: 24.9 – 33.5%) and 15.0% (95% CI: 14.3 –

15.8%) in the following two weeks, respectively (Fig 3A). By the week of April 12 when R_t

reached its minimum, the reduction in population contact rates was associated with an R_t

reduction of 70.7% (95% CI: 65.0 – 76.4%). In addition, analysis at the neighborhood level

consistently showed large reductions in R_t that were likely due to reductions in population

contact rates (range of median estimates: 66.1 – 90.1% across the 42 neighborhoods in NYC; Fig

S4).

In addition, transmission in four age groups (i.e., 5-14, 15-24, 25-44, and 45-64 year-olds)

appeared to be most impacted by changing population contact rates (Fig 3 and Table 1). The

reduction in population contact rates was associated with decreases of the age-specific R_t by

83.4% (95% CI: 80.1 – 86.7%) for 5-14 year-olds, 65.4% (95% CI: 57.0 – 73.8%) for 15-24 year-

olds, 76.5% (95% CI: 68.5 – 84.6%) for 25-44 year-olds, and 68.9% (95% CI: 59.2 – 78.6%) for 45-

64 year-olds, by the week of April 12.

Effectiveness of face covering/masking

Estimated transmission rates (or probability of infection) and the infectious period also closely

tracked changes in mobility (Table S3; $r \geq 0.5$ for most age groups). Thus, it appears that reducing

mobility not only reduces contact rates but also likely reduces 1) the probability of transmission

per contact due to, e.g., increased public spacing and 2) the effective infectious period per

infected individual due to, e.g., more time spent at home and as a result reduced time for

community transmission despite likely unchanged duration of viral shedding. Given this

observation, we hypothesize that the relationship between mobility and estimated

transmission rates (and effective infectious period) can be used to disentangle the impact of

interventions that reduce population contact rates (particularly, the stay-at-home mandate)

and face covering/masking — two major public health interventions implemented in NYC

during the pandemic — on transmission. We make two predictions if this hypothesis holds.

First, predicted transmission rates (infectious period) using mobility data alone would be higher

(longer) than those estimated by the model-inference system additionally based on case and

mortality data *for weeks when face covering in public was mandated* as it would lead to further

reductions in transmission (i.e., temporality and direction of the impact). Second, while the

efficacy of masking (i.e., measured under ideal conditions of mask quality and correct use) likely

does not vary by individual, the effectiveness of masking (i.e., measured under real-world, often

imperfect conditions) and impact of this intervention could vary by subpopulation due, for

example, to different usage rates of masks; as such, we expect the predictive errors to be larger

for age groups with higher compliance of masking (i.e., magnitude of the impact). Our analyses

largely confirmed both predictions. For the first, as shown in Fig 4, transmission rates predicted

using a linear regression model with the observed mobility as the sole predictor were higher

than those estimated by the model-inference system, following the face covering mandate

starting the week of April 12.(*4*) For the latter, the discrepancies in the two model estimates

(i.e. the gaps between the dashed and solid blue lines; Fig 4) appeared to increase with age and

were largest among the two elderly age groups who have been reported to more frequently

use masks.(*16-18*) However, infants (<1 year) appeared to have a larger reduction than other

children groups; this could have been due to transmission reduction related to their sources of

infection (e.g. their caretakers and healthcare settings where they tended to be exposed).

Similar patterns held for the effective infectious period (Fig 4).

Given these observations, we further used the discrepancies in the two model estimates to

approximate the impact of face covering on reducing COVID-19 transmission. Combining the

reduction in the transmission rate and effective infectious period, we estimated that, for all

ages combined, face covering contributed to a 6.6% (95% CI: 0.8 – 12.4%) reduction during the

first month it was implemented and a 3.4% (95% CI: -1.9 – 8.6%) reduction over the entire 8

weeks prior to the city's reopening (Table 1). As expected, the estimated impact varied

substantially by age group. The effectiveness was 20.8% (95% CI: -0.1 – 41.6%) for 65-74 year-

olds and 20.8% (95% CI: -0.9 – 42.5%) for 75+ year-olds during the first month and remained at

similar levels afterwards. For 25-44 and 45-64 year-olds, two age groups with the highest

infection rates (Fig 2), the effectiveness was 4.5% (95% CI: -0.6 – 9.7%) and 8.1% (95% CI: -0.1 –

16.1%) in the first month, respectively; however, it reduced substantially afterwards, likely due

to reversed risk behavior. Of note, in addition to the likely lower usage rate of face covering in

late May – early June, increases in risky behaviors such as large gatherings at the time(*19*) may

have partially obscured the effectiveness of masking.

Retrospective projections of cases and deaths

NYC started phased reopening from the week of 6/7/2020, which allows industries to gradually

reopen per a four-phase plan.(*20*) For instance, manufacturing industries were allowed to

reopen starting the week of 6/7/2020 (dubbed "Phase 1"), whereas real estate was allowed to

reopen starting the week of 6/21/2020 (dubbed "Phase 2"), and personal care services were

allowed starting the week of 7/6/2020 (dubbed "Phase 3"). As such, population mobility has

increased gradually during this time, which could lead to increased transmission of unknown

magnitude. Such changes also offer an opportunity to test the accuracy of our estimates –

should the estimated effectiveness of reducing contact rates and utilizing face coverings be

accurate, these estimates could be used to anticipate changes in transmission in response to

the changing mobility and in turn the epidemic dynamics after reopening. We thus used these

estimates to generate projections of cases and deaths for the 8 weeks beyond our study period,

and compared the projections to available, independent corresponding observations. Overall,

our projections underestimated the total number of cases (relative error of median projections:

-27% over 8 weeks; Fig 5A) but were able to accurately estimate the total number of deaths

(relative error: -2% over 8 weeks; Fig 5B). In addition, examination of age-grouped projections

shows that the underestimation of cases was mostly among younger age groups whose case

rates had increased in June (1-4, 5-14, 15-24 and 25-44 year-olds; Fig S5). These recent

increases in young cases may have resulted from more young adults returning to work including

some in service industries with high contact rates and, relatedly, sending their children to

childcare and/or summer camps due to a lack of caretakers at home [information from NYC

Department of Health and Mental Hygiene (DOHMH) community investigation; unpublished]. In

addition, increased risk behaviors of some young individuals (e.g., large parties without physical

distancing(*19*)) may have also contributed to the increased cases among young adults in late

June.(*21*) Consistently, COVID-19 associated mortality, mostly occurring among older adults

continued to decrease and were accurately predicted for different age groups (Fig S6).

Discussion

The spring 2020 pandemic wave in NYC, the first epidemic center in the US, provides a test case

to study COVID-19 epidemiological characteristics and the effectiveness of public health

interventions. Through comprehensive modeling, we have reconstructed the transmission

dynamics and estimated the effectiveness of two major interventions, social distancing and

mandatory face covering in public. Our results show that reducing contact rates (mainly via

school closures and voluntary or mandated stay-at-home measures) likely contributed to the

largest reduction in transmission in the population overall (~70%) and for most age groups

(>50% for all age groups). Widespread use of face covering likely contributed to an additional

~7% overall reduction and up to ~20% reduction among 65+ year-olds during the first month

face covering was mandated in public places. Our findings largely consolidate previous model

estimates on the impact of lockdown-like measures(*6, 22, 23*) and studies on face covering in

reducing COVID-19 transmission and provide insights to future COVID-19 mitigation efforts.

Lockdown-like measures where confinement at home is encouraged or mandated through

school closures, telework policies, closure of non-essential businesses, and stay-at-home orders

have been a major control measure to curb COVID-19 spread. In effect, such measures reduce

population contact rates and thus transmission. Previous modeling studies estimated that

lockdowns reduced COVID-19 transmission (measured by R_t) by 58% in Wuhan, China,(*6*) 45%

(95% CI: 42-49%) in Italy,(*23*) and 77% (95% CI: 76-78%) in France.(*22*) Our estimate for NYC

overall (~70%) is consistent with these previous estimates. In addition, our estimates show that

reducing population contact rates effectively reduced transmission across all age groups

(ranging from a 51% reduction among 1-4 year-olds to 83% among 5-14 year-olds; Table 1).

Together, these findings underscore the importance of reducing contact rates through, for

example, physical distancing in places with continuous community transmission of COVID-19.

The use of surgical masks or cloth face coverings has been another major preventive measure.

Studies overall have shown that surgical masks could substantially reduce onward transmission

albeit with a large range of efficacy estimates across settings.(*24*) However, it remains unclear

the overall effectiveness of universal face covering requirements at the population level,

especially during a pandemic, due to several factors: 1) The overall effectiveness depends on

compliance which may vary across subpopulations and time; 2) Improper use of face coverings

(e.g. without covering the nose and/or mouth or improper handling(*25*)) can reduce the

effectiveness of face covering; 3) Face coverings are required and mostly worn in public and

thus likely have a lower impact in private settings, particularly in reducing household

transmission; consequently, the relative impact of face covering depends on the relative

contribution of different sources of transmission (e.g. household vs. community) at a given time

and *vice versa*; and 4) Use of face coverings may lead to complacency and less stringent

adherence to social distancing and stay-at-home behaviors. Here we estimated a ~7% reduction

in overall transmission during the first month of the face covering mandate. However, the

estimated effectiveness varied largely across age groups with much higher effectiveness among

older adults (~20% for both 65-74 and 75+ year-olds vs. <10% for other age groups). This

discrepancy was likely due to the differential compliance and types of face covering used.

Observational studies in Wisconsin and surveys nationwide in April/May reported about 2-fold

higher rates of face covering usage among older adults versus younger adults and minors.(*16-*

18) In addition, due to the shortage of surgical masks during March–May,(*26*) older adults at

higher risk of severe COVID-19 infection were more likely to use surgical masks whereas

younger age groups more frequently used non-medical cloth coverings, which are often less

effective(*27, 28*) (e.g. measured ultrafine filtration efficiency is ~50% for surgical masks vs. ~10-

25% for T-shirt and ~25-35% for cotton covers(*28*)).

As lockdown-like measures have been gradually lifted in many places, residents spend more

time outside their homes than during the lockdown. Adjusting for the time spent outside of

homes (~8.3 hours in April 2020 vs. ~11.5 hours in June-July 2020 and ~13.5 hours pre-

pandemic; NYC data(*29*)), universal face covering would have reduced overall transmission by

~9–11% (i.e., 6.6% multiplied by a factor of 1.4–1.6) during reopening, given the same rates of

face covering as in April. However, if the same effectiveness among older adults were achieved

among other age groups, universal face covering could reduce overall transmission by up to

~28–32% (i.e., 20% multiplied by a factor of 1.4–1.6). Thus, our findings suggest that improving

rates of usage of effective face coverings, especially among younger age groups, would

significantly mitigate the risk of resurgence in COVID-19 infections during re-opening. It is thus

crucial for future research to understand reasons for the use/non-use and selection of face

coverings by age group to inform strategies to increase consistent and correct mask use in

settings where social distancing is not possible.

It is important to note, however, that not all individuals have the same opportunities to

physically distance and/or adopt face coverings during a pandemic, despite government

mandates. For instance, over one million frontline workers in NYC (e.g., healthcare workers,

transportation workers, janitors, and grocery clerks, which comprise 25 percent of the city's

workforce) had to continue their essential work during the pandemic.(*30*) In addition, data from

the U.S. Bureau of Labor Statistics suggest Black and Latino communities have less

opportunities to work from home.(*31*) Consequently, NYC neighborhoods with more frontline

workers and/or Black and Latino residents tended to have lower reductions in population

mobility during the pandemic.(*29*) In NYC, these communities also experience a number of

social conditions that are thought to exacerbate COVID-19, including overcrowded

multigenerational households, poverty, and high prevalence of chronic diseases. These

communities, known to also carry a higher burden of underlying health conditions, suffered

greater impacts from COVID-19 and have expressed fear and experiences of racialized bias

when wearing a face covering.(*32, 33*) Further research is warranted to investigate such health

disparities. In addition, future policies should take into account structural inequities in labor

trends, overcrowded housing, and underlying conditions and adopt additional preventive

measures to protect those vulnerable communities.

We also note there remain large uncertainties in our estimates due to several limitations.

Foremost, we used population mobility as a proxy for contact rates rather than more direct

measures. Similar approximation and uncertainty applied to our estimates of the effectiveness

of face covering. Future studies are thus warranted for further assessment. In addition, here we

focused on estimating the effectiveness of interventions in the general population without

segregating key settings with intense transmission (e.g., long-term care facilities). Future

studies should assess the impact of interventions targeting such high-risk settings.

Our study also has several strengths. In particular, our estimates were based on comprehensive

model-inference incorporating multiple data streams and further evaluated using model

projections. Results thus provide an assessment of two major public heath interventions

(reducing contact rates and face covering) at the population level where the overall

effectiveness depends on multiple factors in addition to the efficacy of a given intervention.

Altogether, our estimates support the need for multiple interventions (including reducing

contact rates by, e.g., restricting occupancy, universal face covering, and, albeit not studied

here, testing, contact tracing, isolation and timely treatment of cases) in order to effectively

mitigate the spread of COVID-19 as it continues to pose threats to public health.

Methods

Data

COVID-19 cases included all laboratory-confirmed cases by week of diagnosis reported to the

NYC DOHMH. Mortality data by week of death combined confirmed and probable COVID-19-

associated deaths. Confirmed COVID-19-associated deaths were defined as those occurring in

persons with laboratory-confirmed SARS-CoV-2 infection, and probable COVID-19 deaths were

defined as those with COVID-19, SARS-CoV-2, or a similar term listed on the death certificate as

an immediate, underlying, or contributing cause of death, but did not have laboratory-

confirmation of COVID-19.(*34*) For this study, both weekly case and mortality data were

aggregated by age group (<1, 1-4, 5-14, 15-24, 25-44, 45-64, 65-74, and 75+ years) for each of

the 42 United Hospital Fund (UHF) neighborhoods,(*35*) according to the patient's residential

address. All data were retrieved on Sep 4, 2020.

The mobility data, used to model changes in population contact rates due to public health

interventions implemented during the pandemic (e.g., social distancing), came from

SafeGraph(*29, 36*) and contained counts of visitors to locations in each zip code from the same

zip code and others, separately, based on mobile device locations. The released data were

anonymized and aggregated in weekly intervals. We spatially aggregated these data to the UHF

neighborhood level, for both intra and inter UHF neighborhood mobility. In addition, SafeGraph

also provided an aggregate measure of the length of time spent outside of the home during

each week.

This study was classified as public health surveillance and exempt from ethical review and

informed consent by the Institutional Review Boards of both Columbia University and NYC

DOHMH.

Network transmission model

The epidemic model used in this study was described in Yang et al. 2020(*11*) Briefly, the model

simulated intra- and inter neighborhood transmission of COVID-19 using a susceptible-exposed-

infectious-removed (SEIR) network model:

$$\begin{cases} \frac{dS_i}{dt} = -S_i \sum_{j=1}^{j=42} b_j \beta_{city} c_{ij} I_j / N_j \\ \frac{dE_i}{dt} = S_i \sum_{j=1}^{j=42} b_j \beta_{city} c_{ij} I_j / N_j - \frac{E_i}{Z} \\ \frac{dI_i}{dt} = \frac{E_i}{Z} - \frac{I_i}{D} \\ \frac{dR_i}{dt} = \frac{I_i}{D} \end{cases} \text{(Eqn 1)}$$

where S_i, E_i, I_i, R_i, and N_i are the numbers of susceptible, exposed (but not yet infectious),

infectious, and removed (either recovered or deceased) individuals and the total population,

respectively, from a given age group in neighborhood i. Note that due to model complexity and

a lack of information for parameterizing interactions among age groups, we modeled each age

group separately (i.e., combining all sources of transmission to each age group; see further

detail on parameter estimation below); as such, Eqn 1 describes the spatial transmission across

neighborhoods without interactions among age groups. β_{city} is the citywide transmission rate,

which incorporated seasonal variation as observed for OC43, a beta-coronavirus in humans

from the same genus as SARS-CoV-2.(*37*) To allow differential transmission in each

neighborhood, we included a multiplicative factor, b_i, to scale neighborhood local transmission

rates. Z and D are the latency and infectious periods, respectively (Table S4).

The contact rates (c_{ij}) in each neighborhood over time and connectivity among neighborhoods

were computed based on mobility data. The model also accounted for delays from infection to

diagnosis using two parameters (gamma distribution with mean T_d and standard deviation T_{sd}

estimated along with other parameters) and death (based on observed time from diagnosis to

death) as well as infection detection rate using a parameter r (estimated along with other

parameters). For further detail, please refer to Yang et al. 2020.(*11*)

Parameter estimation

To estimate model parameters (e.g., b_i, β_{city}, Z, D, r, and *infection fatality risk*, for i=1,…,42) and

state variables (e.g., number of susceptible and infectious individuals in each neighborhood) for

each week, we ran the network-model stochastically with a daily time step in conjunction with

the ensemble adjustment Kalman filter (EAKF)(*38*) and fit to weekly case and mortality data

from the week starting March 1 to the week ending June 6, 2020. The posterior distribution of

each model parameter/variable was updated for that week at the same time.(*38*) This

parameter estimation process was done separately for each of the eight age groups (i.e. <1, 1-4,

5-14, 15-24, 25-44, 45-64, 65-74, and 75+ years). To include transmission from other age

groups, we used measured intra and inter-group contacts from the POLYMOD study(*15*) to

compute the total number of contacts made with each age group and adjusted the *prior* range

of the transmission rate (β_{city}) for each age group accordingly. The posterior estimate was

computed based on cases and mortality data for each group, which included all sources of

infection. Thus, the estimated transmission rate for each age group nevertheless included all

sources of transmission. To account for stochasticity in model initiation, we ran the parameter

estimation process independently 20 times. Results for each age group were combined from

these 20 runs (each with 500 model realizations). We computed age-specific R_t, the effective

reproductive number during week-t, from the posterior estimates of transmission rate (β_{city} and

b_i), infectious period (D), contact matrix (c_{ij}), susceptibility and population size in the

neighborhood using the next generation method.(*39*) We computed R_t, β_{city}, and D estimates

for all ages overall as a weighted average of the age-specific estimates with weights equal to

the population fraction in each age group.

Estimating the effectiveness of reducing contact rates

The R_t estimates from the model-inference system capture changes in transmission due to

various interventions, i.e., the overall effectiveness of all implemented interventions. To

separately estimate the effectiveness of interventions that reduce contact rates, we used

human mobility as a measure of population contact rate to estimate the changes in R_t in

response to changing population contact rates. Specifically, we regressed the R_t estimates from

the full model-inference system on the mobility data: $R_t = a_0 + a_1 M_{ave,t}$ (Eqn 2), where $M_{ave,t}$

is the mean of all intra-neighborhood mobility at week-t. We then computed the effectiveness

of reducing contact rate based on $\widehat{R_t}$, the R_t estimate from this regression model solely based

on the observed mobility. That is, the reduction in R_t by week-t, likely due to reducing contact

rate, was computed as $(\widehat{R_t} - \widehat{R_0})/\widehat{R_0}$ (Eqn 3). To test the robustness of our method, we

performed the same analysis for individual UHF neighborhoods (n = 42).

Estimating the effectiveness of face covering/masking

In addition to changes in population contact rates, face covering/masking was another major

control measure implemented beginning the week of April 12, 2020 when NYC mandated

residents wear face coverings in public places. To estimate the effectiveness of face covering,

we first estimated the changes in transmission rate and effective infectious period, two model

parameters determining R_t, due to changes in mobility (as opposed to masking) using

regression models similar to Eqn 2. Specifically, we regressed the estimated citywide

transmission rate (or effective infectious period) from the full model-inference system on

average mobility: $Y_t = a_0 + a_1 M_{ave,t}$ (Eqn 4), where Y_t is β_{city} or D. We then computed the

relative reduction in transmission rate (or effective infectious period) due to face covering as

$\eta_Y = E((Y_t - \widehat{Y_t})/Y_t)$, where E(·) gives the mean over the relevant timeframe (here we

estimated two timeframes, i.e., 1 month following the mandate and over 8 weeks up to

6/6/2020). Combining both reductions, we computed the effectiveness of face covering as $\eta =$

$1 - (1 - \eta_\beta)(1 - \eta_D)$. Of note, while mechanistically face coverings act primarily by reducing

the probability of transmission (i.e., transmission rate), here we included both the potential

impact on the transmission rate (η_β) and effective infectious period (η_D), mainly because the

multiplicative relationship of the two variables with R_t makes it challenging to separate the two

effects. Nevertheless, reductions in the infectious period via face covering are possible. A

recent study on animals showed that masking could reduce the severity of infection;(*40*) if

persons with milder infection experience shorter duration of viral shedding (there is some

evidence for this, e.g., from He et al.(*41*)), milder symptoms in individuals infected while

wearing face covering could lead to shorter infectious period of these individuals.

Projections of cases and deaths

To evaluate the accuracy of model estimates, in particular, the effectiveness of transmission

reduction by reducing contact rates and use of face covering, we tested if these estimates along

with the model could generate accurate predictions of cases and deaths for 8 weeks beyond

the study period (i.e. from the week of 6/7/2020 to the week of 7/26/2020). We first projected

the citywide transmission rate and infectious period based on observed mobility using Eqn 4;

these estimates thus accounted for changes due to changes in contact rates. To incorporate the

reduction in transmission by face covering, we further reduced the projected city transmission

rate by a factor of $1 - \eta_\beta p_{out}$ and the infectious period by a factor of $1 - \eta_D p_{out}$, where p_{out} is

a factor to adjust for time spent outside of the home during each week. To reflect longer-term

usage rates of face covering, we used η_β and η_D estimated during the entire 8 weeks face

covering was required (i.e. from the week of 4/12/2020 to the week of 5/31/2020). Finally, we

used estimates of population susceptibility and infection rates at the end of the week of

5/31/2020 to model initial conditions and integrated the SEIR network model forward

stochastically for 8 weeks using the projected transmission rate and infectious period.

Acknowledgments:

This study was supported by the National Institute of Allergy and Infectious Diseases

(AI145883), the National Science Foundation Rapid Response Research Program (RAPID;

2027369), and the NYC DOHMH. We thank the NYC DOHMH Bureau of Vital Statistics team and

EpiData team; in particular, Mary Huynh, Sharon K. Greene, Alice Yeung, Anne Fine, Miranda S.

Moore and Kevin Guerra, for data management and provision. We thank Jennifer Brite at NYC

DOHMH for coordinating discussions on modeling COVID-19 in NYC and Alana Tornello for

amplifying community voices. We also thank Columbia University Mailman School of Public

Health for high performance computing, Safe Graph (safegraph.com) for providing the mobility

data, and Sasikiran Kandula at Columbia University for compiling the mobility data used in this

study. We are also grateful to Sasikiran Kandula, Sharon K. Greene, Anne Fine, and Hannah

Helmy for helpful feedback on this manuscript.

Conflict of Interest

J Shaman and Columbia University disclose partial ownership of SK Analytics. J Shaman

discloses consulting for BNI. Other authors have nothing to disclose.

References:

1. NYC & Company, 2019 New York City Travel & Tourism Trend Report. 8/17.
   https://adobeindd.com/view/publications/e91e777a-c68b-4db1-a609-
   58664a52cffd/7r7x/publication-web-
   resources/pdf/NYC_Travel&TourismTrendReport_Oct2019.pdf

2. New York City Office of the Mayor, New York City to Close All School Buildings and
   Transition to Remote Learning. https://www1.nyc.gov/office-of-the-mayor/news/151-
   20/new-york-city-close-all-school-buildings-transition-remote-learning

3. New York State Department of Health, New York State on PAUSE.
   https://coronavirus.health.ny.gov/new-york-state-pause

4. New York city Office of the Mayor, Transcript: Mayor de Blasio Holds Media Availability on
   COVID-19. https://www1.nyc.gov/office-of-the-mayor/news/249-20/transcript-mayor-de-
   blasio-holds-media-availability-covid-19

5. N. Y. S. Government, Video, Audio, Photos & Rush Transcript: Governor Cuomo Announces
   U.S. Open to Be Held Without Fans from August 31st to September 13th.
   https://www.governor.ny.gov/news/video-audio-photos-rush-transcript-governor-cuomo-
   announces-us-open-be-held-without-fans-august

6. R. Li, S. Pei, B. Chen, Y. Song, T. Zhang, W. Yang, J. Shaman, Substantial undocumented
   infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV-2). *Science* **368**,
   489-493 (2020).

7. The Novel Coronavirus Pneumonia Emergency Response Epidemiology Team, The
   epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-
   19)—China, 2020. *China CDC Weekly* **2**, 113-122 (2020).

8.   CDC COVID-19 Response Team, Preliminary estimates of the prevalence of selected

underlying health conditions among patients with coronavirus disease 2019—United

States, February 12–March 28, 2020. *Morbidity and Mortality Weekly Report* **69**, 382

(2020).

9.   Governor.ny.gov, After Weeks of Demanding Approval, Governor Cuomo Announces FDA

Gives New York State Authority to Conduct All COVID-19 Testing at Public and Private Labs.

https://www.governor.ny.gov/news/after-weeks-demanding-approval-governor-cuomo-

announces-fda-gives-new-york-state-authority

10. C. Y. Johnson, L. H. Sun, L. McGinley, In hard-hit areas, testing restricted to health care

workers, hospital patients. 6/22/2020.

https://www.washingtonpost.com/health/2020/03/21/coronavirus-testing-strategyshift/

11. W. Yang, S. Kandula, M. Huynh, S. K. Greene, G. Van Wye, W. Li, H. T. Chan, E. McGibbon, A.

Yeung, D. Olson, A. Fine, J. Shaman, Estimating the infection fatality risk of COVID-19 in

New York City, March 1-May 16, 2020. *medRxiv*, 2020.2006.2027.20141689 (2020).

12. Q. Li, X. Guan, P. Wu, X. Wang, L. Zhou, Y. Tong, R. Ren, K. S. M. Leung, E. H. Y. Lau, J. Y.

Wong, X. Xing, N. Xiang, Y. Wu, C. Li, Q. Chen, D. Li, T. Liu, J. Zhao, M. Liu, W. Tu, C. Chen, L.

Jin, R. Yang, Q. Wang, S. Zhou, R. Wang, H. Liu, Y. Luo, Y. Liu, G. Shao, H. Li, Z. Tao, Y. Yang,

Z. Deng, B. Liu, Z. Ma, Y. Zhang, G. Shi, T. T. Y. Lam, J. T. Wu, G. F. Gao, B. J. Cowling, B.

Yang, G. M. Leung, Z. Feng, Early Transmission Dynamics in Wuhan, China, of Novel

Coronavirus-Infected Pneumonia. *N Engl J Med* **382**, 1199-1207 (2020).

13. J. T. Wu, K. Leung, G. M. Leung, Nowcasting and forecasting the potential domestic and

international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling

study. *Lancet*, (2020).

14. C. Goldbaum, Coronavirus in N.Y.: Riders Ditch the Subway as Fear Spreads.

https://www.nytimes.com/2020/03/13/nyregion/coronavirus-subway-public-transit.html

15. J. Mossong, N. Hens, M. Jit, P. Beutels, K. Auranen, R. Mikolajczyk, M. Massari, S. Salmaso,

G. S. Tomba, J. Wallinga, J. Heijne, M. Sadkowska-Todys, M. Rosinska, W. J. Edmunds, Social

contacts and mixing patterns relevant to the spread of infectious diseases. *PLoS Med* **5**, e74

(2008).

16. N. L. Arp, T. H. Nguyen, E. J. Graham Linck, A. K. Feeney, J. H. Schrope, K. L. Ruedinger, A.
Gao, M. Miranda-Katz, A. E. Kates, N. Safdar, Use of face coverings by the public during the
COVID-19 pandemic: an observational study. *medRxiv*, 2020.2006.2009.20126946 (2020).
17. M. H. Haischer, R. Beilfuss, M. R. Hart, L. Opielinski, D. Wrucke, G. Zirgaitis, T. D. Uhrich, S.
K. Hunter, Who is wearing a mask? Gender-, age-, and location-related differences during
the COVID-19 pandemic. *medRxiv*, 2020.2007.2013.20152736 (2020).
18. Z. Ritter, M. Brenan, New April Guidelines Boost Perceived Efficacy of Face Masks.
https://news.gallup.com/poll/310400/new-april-guidelines-boost-perceived-efficacy-face-
masks.aspx
19. V. Barone, J. Marino, Hundreds pack St. Marks Place to drink, party despite coronavirus.
https://nypost.com/2020/06/12/hundreds-pack-st-marks-place-to-drink-despite-
coronavirus/
20. New York State Government, Reopening New York City. https://forward.ny.gov/reopening-
new-york-city
21. E. Durkin, New York City sees uptick in coronavirus cases among young adults.
https://www.politico.com/states/new-york/albany/story/2020/07/13/new-york-city-sees-
uptick-in-coronavirus-cases-among-young-adults-1300223
22. H. Salje, C. Tran Kiem, N. Lefrancq, N. Courtejoie, P. Bosetti, J. Paireau, A. Andronico, N.
Hoze, J. Richet, C. L. Dubost, Y. Le Strat, J. Lessler, D. Levy-Bruhl, A. Fontanet, L. Opatowski,
P. Y. Boelle, S. Cauchemez, Estimating the burden of SARS-CoV-2 in France. *Science* **369**,
208-211 (2020).
23. M. Gatto, E. Bertuzzo, L. Mari, S. Miccoli, L. Carraro, R. Casagrandi, A. Rinaldo, Spread and
dynamics of the COVID-19 epidemic in Italy: Effects of emergency containment measures.
Proc Natl Acad Sci U S A **117**, 10484-10491 (2020).
24. M. Liang, L. Gao, C. Cheng, Q. Zhou, J. P. Uy, K. Heiner, C. Sun, Efficacy of face mask in
preventing respiratory virus transmission: A systematic review and meta-analysis. *Travel*
Med Infect Dis, 101751 (2020).
25. T. Parker-Pope, How NOT to Wear a Mask.
https://www.nytimes.com/2020/04/08/well/live/coronavirus-face-mask-mistakes.html

26. L. Bernstein, A. Safarpour, Mask shortage for most health-care workers extended into May,

Post-Ipsos poll shows. https://www.washingtonpost.com/health/mask-shortage-for-most-

health-care-workers-extended-into-may-post-ipsos-poll-shows/2020/05/20/1ddbe588-

9a21-11ea-ac72-3841fcc9b35f_story.html

27. E. P. Fischer, M. C. Fischer, D. Grass, I. Henrion, W. S. Warren, E. Westman, Low-cost

measurement of face mask efficacy for filtering expelled droplets during speech. *Science*

Advances **6**, eabd3083 (2020).

28. E. O'Kelly, S. Pirog, J. Ward, P. J. Clarkson, Ability of Fabric Facemasks Materials to Filter

Ultrafine Particles at Coughing Velocity - for Home Made and Fabric Face Mask Creation.

medRxiv, 2020.2004.2014.20065375 (2020).

29. SafeGraph, Weekly Patterns: Foot Traffic Data To Understand The COVID-19 Pandemic.

https://www.safegraph.com/weekly-foot-traffic-patterns

30. S. M. Stringer, New York City's Frontline Workers.

https://comptroller.nyc.gov/reports/new-york-citys-frontline-workers/

31. E. Gould, H. Shierholz, Not everybody can work from home: Black and Hispanic workers are

much less likely to be able to telework. https://www.epi.org/blog/black-and-hispanic-

workers-are-much-less-likely-to-be-able-to-work-from-

home/#:~:text=Less%20than%20one%20in%20five,race%20and%20ethnicity%2C%202017–

2018&text=The%20data%20below%20can%20be%20saved%20or%20copied%20directly%2

0into%20Excel.&text=The%20next%20figure%20illustrates%20the,who%20can%20telewor

k%20by%20wage.

32. D. B. Taylor, For Black Men, Fear That Masks Will Invite Racial Profiling.

https://www.nytimes.com/2020/04/14/us/coronavirus-masks-racism-african-

americans.html

33. N. G. Ruiz, J. M. Horowitz, C. Tamir, Many Black and Asian Americans Say They Have

Experienced Discrimination Amid the COVID-19 Outbreak.

https://www.pewsocialtrends.org/2020/07/01/many-black-and-asian-americans-say-they-

have-experienced-discrimination-amid-the-covid-19-outbreak/

34. New York City Department of Health and Mental Hygiene (DOHMH) COVID-19 Response

Team, Preliminary Estimate of Excess Mortality During the COVID-19 Outbreak — New York

City, March 11–May 2, 2020. *MMWR. Morbidity and mortality weekly report* **69**, 603-605

(2020).

35. NewYork City Department of Health and Mental Hygiene, NYC UHF 42 Neighborhoods.

http://a816-dohbesp.nyc.gov/IndicatorPublic/EPHTPDF/uhf42.pdf

36. A. Lasry, D. Kidder, M. Hast, J. Poovey, G. Sunshine, K. Winglee, N. Zviedrite, F. Ahmed, K. A.

Ethier, C. D. C. P. H. L. Program, H. New York City Department of, H. Mental, H. Louisiana

Department of, S. Public Health, C. King, C.-R. T. San Francisco, D. Alameda County Public

Health, D. San Mateo County Health, H. Marin County Division of Public, Timing of

Community Mitigation and Changes in Reported COVID-19 and Community Mobility - Four

U.S. Metropolitan Areas, February 26-April 1, 2020. *MMWR. Morbidity and mortality*

weekly report **69**, 451-457 (2020).

37. W. Yang, S. Kandula, J. Shaman, Eight-week model projections of COVID-19 in New York

City https://github.com/wan-yang/COLUMBIA-COVID19-PROJECTIONS-FOR-

NYC/blob/master/summary_nyc.projection200316_CU_updated.pdf

38. J. L. Anderson, An Ensemble Adjustment Kalman Filter for Data Assimilation. *Mon. Weather*

Rev. **129**, 2884-2903 (2001).

39. J. M. Heffernan, R. J. Smith, L. M. Wahl, Perspectives on the basic reproductive ratio. *J R Soc*

Interface **2**, 281-293 (2005).

40. J. F. Chan, S. Yuan, A. J. Zhang, V. K. Poon, C. C. Chan, A. C. Lee, Z. Fan, C. Li, R. Liang, J. Cao,

K. Tang, C. Luo, V. C. Cheng, J. P. Cai, H. Chu, K. H. Chan, K. K. To, S. Sridhar, K. Y. Yuen,

Surgical mask partition reduces the risk of non-contact transmission in a golden Syrian

hamster model for Coronavirus Disease 2019 (COVID-19). *Clinical infectious diseases : an*

official publication of the Infectious Diseases Society of America, (2020).

41. X. He, E. H. Y. Lau, P. Wu, X. Deng, J. Wang, X. Hao, Y. C. Lau, J. Y. Wong, Y. Guan, X. Tan, X.

Mo, Y. Chen, B. Liao, W. Chen, F. Hu, Q. Zhang, M. Zhong, Y. Wu, L. Zhao, F. Zhang, B. J.

Cowling, F. Li, G. M. Leung, Temporal dynamics in viral shedding and transmissibility of

COVID-19. *Nat Med* **26**, 672-675 (2020).

Table

Table 1. Estimated effectiveness of reducing contact rate and face covering. Numbers are the

estimated mean and 95% CIs, in percentage. Note the estimated effectiveness of contact rate

reduction combined all measures that reduce contact rates, including school closures and

voluntary or mandated stay-at-home measures.

Age	Estimated effectiveness of intervention (%)		
	contact rate reduction	Face covering (1st month)	Face covering (2 months)
all	70.7 (65.0, 76.4)	6.6 (0.8, 12.4)	3.4 (-1.9, 8.6)
<1	53.8 (41.6, 66)	9.3 (-4.2, 22.9)	12.8 (0.2, 25.3)
1-4	51.0 (45.8, 56.2)	0.9 (-5.5, 7.4)	6.7 (0.6, 12.8)
5-14	83.4 (80.1, 86.7)	3.0 (-0.5, 6.6)	1.6 (-1.6, 4.8)
15-24	65.4 (57.0, 73.8)	4.3 (-2.8, 11.4)	4.0 (-2.5, 10.6)
25-44	76.5 (68.5, 84.6)	4.5 (-0.6, 9.7)	-1.0 (-5.6, 3.7)
45-64	68.9 (59.2, 78.6)	8.1 (-0.1, 16.1)	4.4 (-2.9, 11.8)
65-74	55.8 (34.5, 77.2)	20.8 (-0.1, 41.6)	18.3 (-0.2, 36.9)
75+	53.8 (32.3, 75.3)	20.8 (-0.9, 42.5)	16.2 (-3.3, 35.7)

Figures

Fig 1. Epidemic dynamics. Reported laboratory confirmed cases (A) and cumulative cases (B)

per 100,000 population by week of diagnosis for all ages overall and by age group.

 Fig 2. Estimated changes in the effective reproductive number and infection rates. Blue lines
 show the estimated effective reproductive number (R_t) for each week; surrounding areas show
 the 50% and 95% CrIs. Superimposed boxes (right y-axis) show estimated infection rates by
 week: median (thick vertical lines), 50% CrIs (box edges), and 95% CrIs (whiskers).

Fig 3. Effectiveness of reducing contact rates. Note the estimated effectiveness combined all
measures that reduce contact rates, including school closures and voluntary or mandated stay-
at-home measures. Dark grey lines show the observed changes in mobility (right y-axis). Blue
lines show R_t estimated using a linear regression model with mobility as the sole predictor;
surrounding areas show the 95% CrIs of the model estimates. The adjusted r^2 for the regression
model is also shown in each plot. For comparison, dashed blue lines show R_t estimates from the
model-inference system, without accounting for susceptibility. Percentages attached to the
lines show the incremental reductions in either estimated R_t (in blue) or mobility (in grey).

657
658
659
660

Fig 4. Effectiveness of face covering in reducing the transmission rate and infectious period.
Solid lines show the estimated transmission rate (in blue, left y-axis) and infectious period (in
red, right y-axis) using the model-inference system incorporating interventions including face
covering. Surrounding areas show the 50% CrIs of model estimates. Dashed lines show
corresponding estimates from a linear regression model with mobility as the sole predictor (i.e
without accounting for face covering).

Fig 5. Projections of COVID-19 cases and deaths eight weeks beyond the study period. Blue dots
show confirmed cases by week of diagnosis and deaths by week of death, as observed by the
surveillance system. Blue lines show model median estimates; surrounding shades show 50%
and 90% CrIs. Orange lines show model projected median weekly cases and deaths;
surrounding shades show 50% and 90% CIs of the projection.

673
674

Supplemental Tables and Figures

for

COVID-19 Transmission Dynamics and Effectiveness of Public Health Interventions in New York City during the 2020 Spring Pandemic Wave

Wan Yang, Jaimie Shaff, Jeffrey Shaman

Supplemental Tables

Table S1. Total number of reported cases and estimated cumulative infection rate by the week of 5/31/20. Case rate was computed as the number of total cases divided by the population size of the corresponding age group. Estimated infection rate was estimated by model-inference system, normalized to the corresponding population size; numbers are median (and 95% CrIs).

Age	Total number of cases	Case rate (%)	Estimated infection rate (%)
all	205693	2.45	17.6 (13.2, 25.5)
<1	515	0.46	4.4 (2.9, 10.5)
1-4	609	0.14	1.4 (0.9, 3.7)
5-14	2646	0.28	10.5 (6.5, 24.5)
15-24	12562	1.28	12.0 (8.4, 19.5)
25-44	64764	2.45	22.7 (16.8, 31.5)
45-64	74833	3.64	23.1 (18.4, 29.6)
65-74	25469	3.64	15.2 (11.4, 21.7)
75+	24295	4.44	12.7 (9.8, 18.4)

Table S2. Estimated reproductive number by week and age group. Numbers are median (and interquartile range) of the posterior estimates.

date	all	<1	1-4	5-14	15-24	25-44	45-64	65-74	75+
	Age Groups								
3/1/20	2.99 (2.32, 3.86)	1.52 (1.19, 1.96)	1.56 (1.22, 2.02)	3.15 (2.47, 4.09)	2.47 (1.93, 3.22)	4 (3.1, 5.15)	2.97 (2.3, 3.86)	1.77 (1.37, 2.29)	1.75 (1.35, 2.27)
3/8/20	2.25 (1.64, 2.98)	1.43 (1.11, 1.85)	1.28 (1, 1.7)	2.35 (1.74, 3.18)	2.02 (1.53, 2.64)	2.46 (1.57, 3.48)	2.49 (1.97, 3.12)	1.98 (1.55, 2.44)	1.84 (1.48, 2.29)
3/15/20	2.14 (1.62, 2.74)	1.63 (1.32, 2)	1.17 (0.92, 1.48)	1.52 (1.13, 2.06)	2.38 (1.86, 3)	1.45 (0.97, 2.03)	2.99 (2.3, 3.68)	2.84 (2.33, 3.51)	2.87 (2.33, 3.52)
3/22/20	1.37 (1.08, 1.68)	1.59 (1.35, 1.89)	1.05 (0.85, 1.29)	0.99 (0.74, 1.3)	1.29 (1.03, 1.57)	1 (0.76, 1.26)	1.37 (1.06, 1.71)	2.58 (2.15, 3.04)	2.52 (2.06, 3.02)
3/29/20	0.93 (0.73, 1.13)	1.22 (0.97, 1.44)	1.06 (0.82, 1.3)	0.88 (0.66, 1.14)	0.96 (0.75, 1.15)	0.7 (0.52, 0.88)	0.89 (0.71, 1.06)	1.43 (1.17, 1.65)	1.46 (1.27, 1.66)
4/5/20	0.63 (0.5, 0.75)	0.74 (0.6, 0.88)	0.84 (0.65, 1.03)	0.53 (0.41, 0.68)	0.77 (0.59, 0.91)	0.44 (0.33, 0.56)	0.62 (0.5, 0.73)	0.9 (0.79, 1.02)	0.9 (0.78, 1.02)
4/12/20	0.56 (0.45, 0.67)	0.67 (0.54, 0.79)	0.69 (0.53, 0.84)	0.5 (0.38, 0.62)	0.72 (0.57, 0.83)	0.44 (0.34, 0.55)	0.57 (0.47, 0.67)	0.63 (0.55, 0.71)	0.65 (0.57, 0.74)
4/19/20	0.59 (0.48, 0.7)	0.67 (0.54, 0.8)	0.68 (0.54, 0.81)	0.57 (0.44, 0.7)	0.81 (0.66, 0.93)	0.55 (0.42, 0.68)	0.49 (0.41, 0.57)	0.6 (0.52, 0.68)	0.66 (0.58, 0.74)
4/26/20	0.62 (0.5, 0.73)	0.62 (0.5, 0.75)	0.77 (0.62, 0.91)	0.54 (0.42, 0.67)	0.75 (0.61, 0.86)	0.58 (0.45, 0.72)	0.6 (0.5, 0.71)	0.58 (0.5, 0.67)	0.63 (0.56, 0.71)
5/3/20	0.66 (0.54, 0.77)	0.7 (0.56, 0.84)	0.78 (0.62, 0.91)	0.56 (0.44, 0.67)	0.72 (0.61, 0.8)	0.68 (0.54, 0.82)	0.68 (0.56, 0.78)	0.6 (0.52, 0.67)	0.55 (0.48, 0.63)
5/10/20	0.82 (0.67, 0.95)	0.71 (0.56, 0.86)	0.75 (0.6, 0.88)	0.6 (0.47, 0.73)	0.88 (0.73, 0.98)	0.9 (0.72, 1.06)	0.88 (0.74, 0.99)	0.73 (0.64, 0.81)	0.65 (0.57, 0.74)
5/17/20	1.03 (0.83, 1.18)	0.82 (0.65, 0.99)	0.8 (0.65, 0.93)	0.73 (0.57, 0.88)	1.1 (0.89, 1.24)	1.21 (0.95, 1.42)	1.1 (0.91, 1.24)	0.78 (0.69, 0.87)	0.81 (0.72, 0.92)
5/24/20	0.89 (0.73, 1.03)	0.78 (0.61, 0.95)	0.79 (0.63, 0.91)	0.69 (0.53, 0.83)	0.94 (0.78, 1.06)	0.96 (0.76, 1.13)	0.95 (0.78, 1.08)	0.79 (0.7, 0.88)	0.89 (0.78, 0.99)
5/31/20	0.79 (0.66, 0.91)	0.71 (0.55, 0.87)	0.71 (0.56, 0.83)	0.7 (0.55, 0.86)	0.85 (0.72, 0.95)	0.79 (0.64, 0.93)	0.77 (0.65, 0.89)	0.72 (0.64, 0.79)	1.04 (0.93, 1.15)

Table S3. Correlation of key epidemiological parameters with population mobility during the week of March 1 – the week of May 31, 2020.

Age	Rt (ignore susceptibility)	Rt	Transmission rate	Infectious period
all	0.96	0.96	0.73	0.54
<1	0.76	0.77	0.41	0.70
1-4	0.93	0.93	0.05	0.65
5-14	0.99	0.99	0.86	0.92
15-24	0.90	0.91	0.52	-0.05
25-44	0.93	0.96	0.80	0.27
45-64	0.88	0.90	0.60	0.24
65-74	0.57	0.63	0.54	0.65
75+	0.55	0.60	0.50	0.64

Table S4. Prior ranges for main model parameters and variables. The spatial, temporal, and age resolution of each parameter or variable, estimated in the model-inference system, is specified in the column "Resolution". Note posterior parameter estimates can extend outside the specified prior ranges. Note this is the same as Table S1 in Yang et al.(*1*)

Parameter/ variable	Symbol	Resolution	Prior range	Source/rationale
Initial exposed	E(t=0)	neighborhood- and age-group specific, estimated for the beginning of the Week of March 1, 2020	300 – 8000 total citywide, scaled by population size for each age group and neighborhood	Large uncertainties, used very wide range
Initial infectious	I(t=0)	neighborhood- and age-group specific, , estimated for the beginning of the Week of March 1, 2020	150 – 4000 total citywide, scaled by population size for each age group and neighborhood	Assumed to be half the initial exposed
Initial susceptible	S(t=0)	neighborhood- and age-group specific, estimated for the beginning of the Week of March 1, 2020	$N - E - I$	Assumed all were susceptible except for those initially exposed/infectious
Population size in each age group and neighborhood	N	neighborhood- and age-group specific	N/A	NYC intercensal population estimates for 2018(*2*)
Citywide transmission rate	β_{city}	Citywide, age-group specific, estimated for each week	[0.5, 1] per day overall; scaled by contact rate for each age group based on contact data from the POLYMOD study(*3*) (averaged across 8 countries)	Based on R_0 estimates of around 1.5-4 for SARS-CoV-2(*4-6*)
Scaling of neighborhood transmission rate	b_i	neighborhood- and age-group specific, estimated for each week	[0.8, 1.2] for age groups under 65 years; [0.5, 1.5] for age groups 65 or older	Around 1; larger variation for elderly groups based on data

Latency period	Z	Citywide, age-group specific, estimated for each week	[2, 5] days	Incubation period: 5.2 days (95% CI: 4.1, 7)(4); latency period is likely shorter than the incubation period
Infectious period	D	Citywide, age-group specific, estimated for each week	[2, 5] days	Time from symptom onset to hospitalization: 3.8 days (95% CI: 0, 12.0) in China,(7) plus 1-2 days viral shedding before symptom onset. We did not distinguish symptomatic/asymptomatic infections.
Multiplicative factor for mobility	m_1	Citywide, age-group specific, estimated for each week	[1, 2] for <1 year; [0.5, 1.5] for three age groups 1-24 years; [0.1, 1.5] for age group 25-44; [1, 2.5] for age groups 45 or older	Initial model testing showed transmission rates for younger age groups were more sensitive to changes in mobility whereas the two oldest age groups were not sensitive to mobility. For age groups with contact rates lower than the average (based on the POLYMOD study(3)), we raised the diagonal elements in the mobility matrix to the power of the relative contact rate (<1) to account for insensitivity of transmission rate in these age groups to mobility.
Multiplicative factor for neighborhood connectivity	m_2	Citywide, age-group specific, estimated for each week	[0.5, 2]	Likely around 1 but with large uncertainties
Mean of time from viral shedding to diagnosis	T_m	Citywide, age-group specific, estimated for each week	[3, 8] days	From a few days to a week from symptom onset to diagnosis/ reporting,(7) plus 1-2 days of viral shedding (being infectious) before symptom onset

Standard deviation (SD) of time from viral shedding to diagnosis	T_{sd}	Citywide, age-group specific, estimated for each week	[1, 3] days	To allow variation in time to diagnosis/reporting
Reporting rate	r	Citywide, age-group specific, estimated for each week	Starting from [0.001, 0.05] at time 0 and allowed to increase over time using space re-probing(8)	Large uncertainties
Infection fatality risk (IFR)		Citywide, age-group specific, estimated for each week	$[5, 15] \times 10^{-4}$ for ages under 25; $[5, 15] \times 10^{-3}$ for ages 25-44; $[5, 15] \times 10^{-2}$ for ages 45-64; [0.01, 0.1] for ages 65-74; [0.02, 0.2] for ages 75+;	Based on previous estimates(9) but extend to have wider ranges
Time from diagnosis to death		Citywide	Gamma distribution with mean of 9.36 days and SD of 9.76 days	Based on n=15,686 COVID-19 confirmed deaths in NYC as of May 17, 2020.

Reference:

1. W. Yang, S. Kandula, M. Huynh, S. K. Greene, G. Van Wye, W. Li, H. T. Chan, E. McGibbon, A. Yeung, D. Olson, A. Fine, J. Shaman, Estimating the infection fatality risk of COVID-19 in New York City, March 1-May 16, 2020. *medRxiv*, 2020.2006.2027.20141689 (2020).

2. New York City Department of Health and Mental Hygiene.

3. J. Mossong, N. Hens, M. Jit, P. Beutels, K. Auranen, R. Mikolajczyk, M. Massari, S. Salmaso, G. S. Tomba, J. Wallinga, J. Heijne, M. Sadkowska-Todys, M. Rosinska, W. J. Edmunds, Social contacts and mixing patterns relevant to the spread of infectious diseases. *PLoS Med* **5**, e74 (2008).

4. Q. Li, X. Guan, P. Wu, X. Wang, L. Zhou, Y. Tong, R. Ren, K. S. M. Leung, E. H. Y. Lau, J. Y. Wong, X. Xing, N. Xiang, Y. Wu, C. Li, Q. Chen, D. Li, T. Liu, J. Zhao, M. Liu, W. Tu, C. Chen, L. Jin, R. Yang, Q. Wang, S. Zhou, R. Wang, H. Liu, Y. Luo, Y. Liu, G. Shao, H. Li, Z. Tao, Y. Yang, Z. Deng, B. Liu, Z. Ma, Y. Zhang, G. Shi, T. T. Y. Lam, J. T. Wu, G. F. Gao, B. J. Cowling, B. Yang, G. M. Leung, Z. Feng, Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus–Infected Pneumonia. *New Engl J Med*, (2020).

5. J. T. Wu, K. Leung, G. M. Leung, Nowcasting and forecasting the potential domestic and international spread of the 2019-nCoV outbreak originating in Wuhan, China: a modelling study. *Lancet*, (2020).

6.   R. Li, S. Pei, B. Chen, Y. Song, T. Zhang, W. Yang, J. Shaman, Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV-2). *Science* **368**, 489-493 (2020).

7.   J. Zhang, M. Litvinova, W. Wang, Y. Wang, X. Deng, X. Chen, M. Li, W. Zheng, L. Yi, X. Chen, Q. Wu, Y. Liang, X. Wang, J. Yang, K. Sun, I. M. Longini, Jr., M. E. Halloran, P. Wu, B. J. Cowling, S. Merler, C. Viboud, A. Vespignani, M. Ajelli, H. Yu, Evolving epidemiology and transmission dynamics of coronavirus disease 2019 outside Hubei province, China: a descriptive and modelling study. *The Lancet. Infectious diseases*, (2020).

8.   W. Yang, J. Shaman, A simple modification for improving inference of non-linear dynamical systems. *arXiv*, 1403.6804 (2014).

9.   R. Verity, L. C. Okell, I. Dorigatti, P. Winskill, C. Whittaker, N. Imai, G. Cuomo-Dannenburg, H. Thompson, P. G. T. Walker, H. Fu, A. Dighe, J. T. Griffin, M. Baguelin, S. Bhatia, A. Boonyasiri, A. Cori, Z. Cucunuba, R. FitzJohn, K. Gaythorpe, W. Green, A. Hamlet, W. Hinsley, D. Laydon, G. Nedjati-Gilani, S. Riley, S. van Elsland, E. Volz, H. Wang, Y. Wang, X. Xi, C. A. Donnelly, A. C. Ghani, N. M. Ferguson, Estimates of the severity of coronavirus disease 2019: a model-based analysis. *The Lancet. Infectious diseases*, (2020).

Supplemental Figures

Fig S1. Model fits of reported confirmed COVID-19 cases. Blue dots show reported number of cases by age group and week of diagnosis. Boxes show the model fitted weekly number of cases by age group. Box edges, thick middle lines, and whiskers show the 2.5th, 25th, 50th, 75th, and 97.5th percentiles of model estimates.

Fig S2. Model fits of reported COVID-19 associated deaths. Blue dots show reported weekly number of deaths by age group. Boxes show the model fitted weekly number of deaths by age group. Box edges, thick middle lines, and whiskers show the 2.5th, 25th, 50th, 75th, and 97.5th percentiles of model estimates.

Fig S3. Estimated case rates and infection rates by age group. Blue dots show confirmed case rates by age group and week of diagnosis. Blue boxes (left y-axis) show the model fitted weekly case rates and grey boxes (right y-axis) show the model estimated weekly infection rates by age group. Box edges, thick middle lines, and whiskers show the 2.5th, 25th, 50th, 75th, and 97.5th percentiles of model estimates.

Fig S4. Sensitivity analysis on the effectiveness of reducing contact rate by neighborhood. There are 42 United Hospital Fund (UHF) neighborhoods in NYC. (A) shows the changes in human mobility during the pandemic by neighborhood (each colored line). The reductions were substantial in all neighborhoods but to varying degrees. (B) shows the estimated R_t (combining all ages) for each week and neighborhood (each colored line). Note these estimates did not account for changes in susceptibility so as to restrict to changes due to interventions. (C) shows the adjusted r^2 of linear regression model fitting the mobility data (A) to the R_t estimates (B), for each neighborhood, per Eqn 2 in the main text. Adjusted r^2 for most neighborhoods was >0.9. (D) shows the estimated reduction in R_t by the Week of April 12, 2020 based on the Eqn 3 in the main text, for each neighborhood. The histogram is based on the median estimated reduction in R_t.

Fig S5. Projections of COVID-19 cases by age group eight weeks beyond the study period. Blue dots show observed confirmed cases by week of diagnosis (those after the Week of 5/31/2020 were not used in the model). Blue lines show model median estimates; surrounding shades show 50% and 90% CrIs. Orange lines show model projected median weekly cases and deaths; surrounding shades show 50% and 90% CIs of the projection.

Fig S6. Projections of COVID-19 associated deaths by age group eight weeks beyond the study period. Blue dots show observed weekly deaths (those after the Week of 5/31/2020 were not used in the model). Blue lines show model median estimates; surrounding shades show 50% and 90% CrIs. Orange lines show model projected median weekly cases and deaths; surrounding shades show 50% and 90% CIs of the projection.