

1 **Vitamin D and COVID-19 susceptibility and severity in the COVID-19 Host Genetics**
2 **Initiative: A Mendelian randomization study**

3
4 **Authors:**

5 *Guillaume Butler-Laporte^{1,2}, *Tomoko Nakanishi^{1,3,4,5}, Vincent Mooser^{3,6}, David R.
6 Morrison¹, Tala Abdullah¹, Olumide Adeleye¹, Noor Mamlouk¹, Nofar Kimchi^{1,7}, Zaman
7 Afrasiabi¹, Nardin Rezk¹, Annarita Giliberti⁸, Alessandra Renieri^{8,9}, Yiheng Chen¹, Sirui
8 Zhou^{1,2}, Vincenzo Forgetta¹, J Brent Richards^{1,2,3,10}

9
10 *These authors made equal contributions to this manuscript.

11
12 **Affiliations:**

- 13 1) Lady Davis Institute, Jewish General Hospital, McGill University, Montréal,
14 Québec, Canada
15 2) Department of Epidemiology, Biostatistics and Occupational Health, McGill
16 University, Montréal, Québec, Canada
17 3) Department of Human Genetics, McGill University Montréal, Québec, Canada
18 4) Kyoto-McGill International Collaborative School in Genomic Medicine, Graduate
19 School of Medicine, Kyoto University, Kyoto, Japan
20 5) Japan Society for the Promotion of Science
21 6) Canada Excellence Research Chair in Genomic Medicine, McGill University,
22 Montréal, Québec, Canada
23 7) Ruth and Bruce Rappaport Faculty of Medicine, Technion Israel Institute of
24 Technology, Israel
25 8) Medical Genetics, University of Siena, Siena, Italy
26 9) Genetica Medica, Azienda Ospedaliera Universitaria Senese, Siena, Italy
27 10) Department of Twin Research, King's College London, London, United Kingdom
28

29 **Corresponding author:**

30 Brent Richards, Professor of Medicine
31 McGill University
32 Senior Lecturer, King's College London (Honorary)

33 Contact:

34 Pavilion H-413, Jewish General Hospital
35 3755 Côte-Ste-Catherine
36 Montréal, Québec, Canada, H3T 1E2
37 T: +1 514 340 8222 x24362 F: +1 514 340 7529

38 E: brent.richards@mcgill.ca

39 www.mcgill.ca/genepi

40
41 **Funding:**

42 The Richards research group is supported by the Canadian Institutes of Health
43 Research (CIHR: 365825; 409511), the Lady Davis Institute of the Jewish General
44 Hospital, the Canadian Foundation for Innovation, the NIH Foundation, Cancer
45 Research UK, Genome Québec, the Public Health Agency of Canada and the Fonds de
46 Recherche Québec Santé (FRQS). GBL is supported by the CIHR, and a joint
47 scholarship from the FRQS and Québec's Ministry of Health and Social Services. TN is
48 supported by Research Fellowships of Japan Society for the Promotion of Science
49 (JSPS) for Young Scientists and JSPS Overseas Challenge Program for Young
50 Researchers. JBR is supported by a FRQS Clinical Research Scholarship. Support from
51 Calcul Québec and Compute Canada is acknowledged. TwinsUK is funded by the

52 Welcome Trust, Medical Research Council, European Union, the National Institute for
53 Health Research (NIHR)-funded BioResource, Clinical Research Facility and Biomedical
54 Research Centre based at Guy's and St Thomas' NHS Foundation Trust in partnership
55 with King's College London. VM is supported by the Canada Excellence Research Chair
56 Program. The funders had no role in study design, data collection and analysis, decision
57 to publish, or preparation of the manuscript.

58

59 **Competing interests:**

60 JBR has served as an advisor to GlaxoSmithKline and Deerfield Capital. These
61 agencies had no role in the design, implementation or interpretation of this study.

62

63 **Article type:** Research

64 **Word count:** 3480 (Introduction, Methods, Results, and Discussion)

65 **Key words:** Vitamin D, COVID-19, SARS-CoV-2, Mendelian Randomization, Outcome

66

67

68 **Abstract**

69 *Background*

70 Increased vitamin D levels, as reflected by 25OHD measurements, have been proposed
71 to protect against COVID-19 disease based on *in-vitro*, observational, and ecological
72 studies. However, vitamin D levels are associated with many confounding variables and
73 thus associations described to date may not be causal. Vitamin D Mendelian
74 randomization (MR) studies have provided results that are concordant with large-scale
75 vitamin D randomized trials. Here, we used two-sample MR to assess evidence
76 supporting a causal effect of circulating 25OHD levels on COVID-19 susceptibility and
77 severity.

78
79 *Methods and findings*

80 Genetic variants strongly associated with 25OHD levels in a genome-wide association
81 study (GWAS) of 443,734 participants of European ancestry (including 401,460 from the
82 UK Biobank) were used as instrumental variables. GWASs of COVID-19 susceptibility,
83 hospitalization, and severe disease from the COVID-19 Host Genetics Initiative were
84 used as outcome GWASs. These included up to 14,134 individuals with COVID-19, and
85 1,284,876 without COVID-19, from 11 countries. SARS-CoV-2 positivity was determined
86 by laboratory testing or medical chart review. Population controls without COVID-19
87 were also included in the control groups for all outcomes, including hospitalization and
88 severe disease. Analyses were restricted to individuals of European descent when
89 possible. Using inverse-weighted MR, genetically increased 25OHD levels by one
90 standard deviation on the logarithmic scale had no significant association with COVID-19
91 susceptibility (OR = 0.97; 95% CI: 0.95, 1.10; P=0.61), hospitalization (OR = 1.11; 95%
92 CI: 0.91, 1.35; P=0.30), and severe disease (OR = 0.93; 95% CI: 0.73, 1.17; P=0.53).
93 We used an additional 6 meta-analytic methods, as well as sensitivity analyses after
94 removal of variants at risk of horizontal pleiotropy and obtained similar results. These
95 results may be limited by weak instrument bias in some analyses. Further, our results do
96 not apply to individuals with vitamin D deficiency.

97
98 *Conclusion*

99 In this two-sample MR study, we did not observe evidence to support an association
100 between 25OHD levels and COVID-19 susceptibility, severity, or hospitalization. Hence,
101 vitamin D supplementation as a means of protecting against worsened COVID-19
102 outcomes is not supported by genetic evidence. Other therapeutic or preventative
103 avenues should be given higher priority for COVID-19 randomized controlled trials.

104

105

106 **Author Summary**

107

108 • *Why was this study done?*

109 - Vitamin D levels have been associated with COVID-19 outcomes in
110 multiple observational studies, though confounders are likely to bias these
111 associations.

112 - By using genetic instruments which limit such confounding, Mendelian
113 randomization studies have consistently obtained results concordant with
114 vitamin D supplementation randomized trials. This provides rationale to
115 undertake vitamin D Mendelian randomization studies for COVID-19
116 outcomes.

117 • *What did the researchers do and find?*

118 - We used the genetic variants obtained from the largest consortium of
119 COVID-19 cases and controls, and the largest study on genetic
120 determinants of vitamin D levels.

121 - We used Mendelian randomization to estimate the effect of increased
122 vitamin D on COVID-19 outcomes, while limiting confounding.

123 - In multiple analyses, our results consistently showed no evidence for an
124 association between genetically predicted vitamin D levels and COVID-19
125 susceptibility, hospitalization, or severe disease.

126 • *What do these findings mean?*

127 - Using Mendelian randomization to reduce confounding that has
128 traditionally biased vitamin D observational studies, we did not find
129 evidence that vitamin D supplementation in the general population would
130 improve COVID-19 outcomes

131 - These findings, together with recent randomized controlled trial data,
132 suggest that other therapies should be prioritized for COVID-19 trials.

133 **Introduction**

134 SARS-CoV-2 infection has killed millions of individuals and has led to the largest
135 economic contraction since the Great Depression [1]. Therefore, therapies are required
136 to treat severe COVID-19 disease and to prevent its complications. Therapeutic
137 development, in turn, requires well-validated drug targets to lessen COVID-19 severity.

138
139 Recently, vitamin D status, as reflected by 25-hydroxy-vitamin D (25OHD) level has
140 been identified as a potentially actionable drug target in the prevention and treatment of
141 COVID-19 [2]. As the pre-hormone to the biologically active calcitriol, 25OHD has been
142 epidemiologically linked to many health outcomes [3,4]. Given calcitriol's recognized *in-*
143 *vitro* immunomodulatory role [5], as well as observational and ecological studies
144 associating measured 25OHD blood levels with COVID-19 [6,7], the vitamin D pathway
145 might be a biologically plausible target in COVID-19. This could be of public health
146 importance, given that the prevalence of vitamin D insufficiency is high in most countries,
147 and that more than 37% of elderly adults in the USA take vitamin D supplements [8].
148 Further, 25OHD supplementation is inexpensive and reasonably safe—thus providing a
149 potential avenue to lessen the burden of the SARS-CoV-2 pandemic.

150
151 However, observational studies on 25OHD are prone to confounding and reverse
152 causation bias. Confounding happens when the relationship between exposure (25OHD)
153 and the outcome (COVID-19) is influenced by unobserved, or improperly controlled
154 common causes. Reverse causation happens when the outcome itself is a cause of the
155 exposure. Likewise, conclusions drawn from *in-vitro* may not be applicable *in-vivo*.
156 Accordingly, randomized controlled trials (RCTs) on 25OHD supplementation have been
157 undertaken to test their effect on disease outcomes where observational studies have
158 supported a role for 25OHD level. However, across endocrinology, respiratory,
159 cardiology, and other specialties, these trials have most often not demonstrated
160 statistically significant benefits [9–11]. Some RCTs have even shown detriment to
161 25OHD supplementation [12]. In the field of infectious diseases, an individual patient
162 data meta-analysis of randomized controlled trial of 25OHD supplementation [13]
163 showed some benefit to prevent respiratory tract infections (OR 0.80, 95% CI: 0.69 to
164 0.93). However, this effect was driven by generally benign upper respiratory tract
165 infections, was not observed in lower respiratory tract disease (OR: 0.96, 95% CI: 0.83
166 to 1.10) and even showed numerically worse all-cause mortality (OR: 1.39, 95% CI: 0.85
167 to 2.27). Likewise, a recent trial on sepsis obtained a numerically higher mortality rate in
168 patients who received 25OHD supplementation [14]. At present, we are aware of two
169 RCTs testing the role of vitamin D supplementation on COVID-19 outcomes, both using
170 high-dose vitamin D given at time of hospital admission for COVID-19. The first [15] was
171 a small trial (n=75) showing less intensive care unit admissions in the vitamin D treated
172 arm. However, the follow-up time for mortality varied, and the open-label design put it at
173 high risk of bias. The second [16] was a larger study (n=240) using a double-blind
174 design, and showed no effect on mortality, risk of mechanical ventilation, and length of
175 stay. Nevertheless, questions remain on the use of pre-illness vitamin D
176 supplementation and its effect on disease susceptibility. While RCTs can control for
177 confounding and provide unbiased estimates of the effect of 25OHD supplementation in
178 COVID-19, large well-designed RCTs require considerable resources and time.

179
180 Mendelian randomization (MR) is a genetic epidemiology method that uses genetic
181 variants as instrumental variables to infer the causal effect of an exposure (in this case
182 25OHD level) on an outcome (in this case, COVID-19 susceptibility and severity) [17].
183 MR overcomes confounding bias since genetic alleles are randomized to the individual

184 at conception, thereby breaking associations with most confounders. Similarly, since
185 genetic alleles are always assigned prior to disease onset, they are not influenced by
186 reverse causation. MR has been used in conjunction with proteomics and metabolomics
187 to prioritize drug development and repurposing, and support investment in RCTs which
188 have a higher probability of success [18,19]. In the case of vitamin D, MR has been able
189 to provide causal effect estimates consistently in line with those obtained from RCTs
190 [9,20–24], or support the use of vitamin D supplementation in preventing diseases in at
191 risk individuals (most notably multiple sclerosis [25]). Hence, MR may support
192 investments in 25OHD supplementation trials in COVID-19, if a benefit was shown.
193 Further, since MR results can be generated rapidly, such evidence may provide interim
194 findings while awaiting RCT results.

195
196 However, MR relies on several core assumptions [26]. First, genetic variants must be
197 associated with the exposure of interest. Second, they should not affect the outcome
198 except through effects on the exposure (also known as lack of horizontal pleiotropy).
199 Specifically, MR also assumes that the relationship between the exposure and the
200 outcome is linear. However, this assumption still provides a valid test of the null
201 hypothesis when studying population-level effects [27], as MR then measures the
202 population-averaged effect on the outcome of a shift in the distribution of the exposure.
203 Third, genetic variants should not associate with the confounders of the exposure-
204 outcome relationship. Of these, the most problematic is the second assumption. Yet, in
205 the case of 25OHD, many of its genetic determinants reside at loci that harbour genes
206 whose roles in 25OHD production, metabolism and transport are well known [25].
207 Leveraging this known physiology can help to prevent the incorporation of genetic
208 variants that could lead to horizontal pleiotropy.

209
210 Here, we used genetic determinants of serum 25OHD from a recent genome-wide
211 association study (GWAS) and meta-analysis of more than 443,734 participants of
212 European ancestry [28] in an MR study to test the relationship between increased
213 25OHD level and COVID-19 susceptibility and severity.

214 215 **Methods**

216 We used a two-sample MR approach to estimate the effect of 25OHD levels on COVID-
217 19 susceptibility and severity. In two-sample MR [29], the effect of genetic variants on
218 25OHD and on COVID-19 outcomes are estimated in separate GWASs from different
219 populations. This allows for increased statistical power by increasing the sample size in
220 both the exposure and outcome cohorts. This study is reported as per the Strengthening
221 the Reporting of Observational Studies in Epidemiology (STROBE) guideline [30]
222 (**Supplement 1**).

223
224 Our study did not employ a prospective protocol. Analyses were first planned and
225 performed in July 2020 and updated following peer-review in December 2020. Three
226 major changes were made during the update. First, we used the most up to date COVID-
227 19 Host Genetics Initiative (COVID-19 HGI) GWAS summary statistics. These were
228 made available during the peer-review process. Second, to alleviate potential selection
229 and collider bias, we modified the outcome phenotypes to include population controls.
230 We also performed additional MR sensitivity analyses to check for the robustness of our
231 results. The latter two modifications were made at the request of peer-reviewers. Finally,
232 minor changes to the results' interpretations were made following further peer-review in
233 February 2021.

234

235 *Choice of 25OHD genetic instruments*

236 To find genetic variants explaining 25OHD levels [28], we used a GWAS from our group,
237 which is the largest published GWAS of 25OHD levels, to the best of our knowledge.
238 Importantly, this meta-analysis controlled for season of vitamin D measurement to obtain
239 genetic variants significantly associated with 25OHD levels. From the list of conditionally
240 independent variants provided, we further selected SNPs whose effect on 25OHD level
241 was genome-wide significant ($P < 5 \times 10^{-8}$), minor allele frequency was more than 1%, and
242 with linkage disequilibrium coefficients (r^2) of less than 5% (using the LDlink [31] tool and
243 the European 1000 Genomes dataset, excluding Finnish populations). For SNPs that
244 were not available in the outcome GWAS or with palindromic alleles of intermediate
245 frequency (between 42% and 58%), we again used the LDlink [31] tool to find genetic
246 proxies in the European 1000 Genomes dataset (excluding Finnish populations) using
247 linkage disequilibrium r^2 of 90% or more.

248
249 *COVID-19 outcome definitions and GWASs*

250 We used the COVID-19 HGI outcome definitions and GWAS summary statistics for
251 COVID-19 susceptibility, hospitalization, and severe disease outcomes [32]. For all
252 outcomes, a COVID-19 infection was defined as a positive SARS-CoV-2 laboratory test
253 (e.g. RNA RT-PCR or serology tests) or electronic health record evidence of SARS-CoV-
254 2 infection (using International Classification of Diseases or physician notes). The
255 susceptibility phenotype compared COVID-19 cases, with controls which were defined
256 as any individuals without a history of COVID-19. The hospitalized outcome compared
257 cases defined as hospitalized patients with COVID-19, and controls as any individuals
258 not experiencing a hospitalization for COVID-19, which includes those without COVID-
259 19. The severe disease outcome cases were defined as hospitalized individuals with
260 COVID-19 who required respiratory support. Respiratory support was defined as
261 intubation, CPAP, BiPAP, continuous external negative pressure, or high flow nasal
262 cannula. Controls for the severe COVID-19 outcome were defined as individuals without
263 severe COVID-19 (including those without COVID-19). The inclusion of COVID-19
264 negative participants as controls in each outcome decreases the possibility of collider
265 bias [33] and allows for better population level comparisons. These three outcome
266 phenotypes are referred to as C2, B2, and A2, respectively, in the COVID-19 HGI
267 documentation.

268
269 For our study, we used the October 20th 2020 (v4) COVID-19 HGI fixed effect meta-
270 analysis of GWAS from up to 22 cohorts, performed in up to 11 countries. Every
271 participating cohort was asked to provide summary statistics from a GWAS on the above
272 three outcomes, and including the following non-genetic covariates: age, sex, age*age,
273 age*sex, 20 genetic principal components, as well as any locally relevant covariates at
274 the discretion of participating studies (e.g. hospital, genotype panel, etc.). Cohorts were
275 asked to follow common sample and variant quality control, and only performed analysis
276 if they enrolled 100 cases or more. Analyses were done separately for each major
277 ancestry group to further control for population stratification. For the purposes of our
278 study, we used the meta-analysis results from European ancestry cohorts, except for the
279 severe COVID-19 outcome, for which this meta-analysis was not available. Further
280 details on the three phenotypes and participating cohorts are found in [Table 1](#) and
281 [Supplement 2](#).

282
283 *Primary MR analysis*

284 The effect of 25OHD level on COVID-19 outcomes was obtained for each SNP by using
285 the Wald ratio method. The effect of each SNP was given in standardized log-

286 transformed 25OHD level. Each estimate was meta-analyzed using the inverse-variance
287 weighted (IVW) method, and we performed variant heterogeneity tests to check
288 robustness of IVW results. Allele harmonization and computations were performed using
289 the TwoSampleMR package [34].

290

291 *Horizontal pleiotropy sensitivity analysis*

292 We undertook multiple analyses to assess the risk of horizontal pleiotropy (a violation of
293 the second MR assumption). First, we used MR Egger method, which allows for an
294 additional intercept (alpha) term which also provides an estimate of directional horizontal
295 pleiotropy. This method relies upon the assumption that the size of the direct effects of
296 the genetic variants on the outcome that do not operate through the exposure are
297 independent of the variant's effect on the exposure. Given possible instability in MR
298 Egger estimates [35], we also used the bootstrap MR Egger method to meta-analyze the
299 causal effect estimates from each SNP instrument. Further, we used four additional
300 meta-analysis methods known to be more robust to presence of horizontal pleiotropy (at
301 the expense of statistical power): penalised weighted median, simple mode, weighted
302 median, and weighted mode [36].

303

304 Second, we restricted our choices of SNPs to those whose closest gene is directly
305 involved in the vitamin D pathway. These genes have an established role in vitamin D
306 regulation through its synthesis (*DHCR7/NADSYN1* and *CYP2R1*), transportation (*GC*),
307 and degradation (*CYP24A1*) (**Supplement 3**). This decreases the risk of selecting a
308 genetic variant that affects COVID-19 outcomes independent of its effect on 25OHD
309 levels.

310

311 Third, we used the Phenoscanner tool [37,38] on the remaining SNPs to check for
312 variants associated (at a genome-wide significant threshold of $p=5 \times 10^{-8}$) with
313 phenotypes at risk of affecting COVID-19 outcomes independent of 25OHD, making
314 them at higher risk of horizontal or vertical pleiotropy. Note that vertical pleiotropy, which
315 happens when the COVID-19 outcome is influenced by a phenotype directly in the
316 causal pathway between 25OHD level and COVID-19 outcome, does not violate MR
317 assumptions.

318

319 *Research Ethics*

320 Each cohort included in this study received their respective institutional research ethics
321 board approval to enroll patients. All information used for this study are publicly available
322 as deidentified GWAS summary statistics.

323

324 **Results**

325 *Choice of 25OHD genetic instruments*

326 We obtained our 25OHD genetic instruments from our previously published GWAS on
327 circulating 25OHD levels in 401,460 white British participants in the UK Biobank (UKB)
328 [39], which was meta-analyzed with a GWAS on 25OHD levels of 42,274 participants of
329 European ancestry [40]. Of the 138 reported conditionally independent SNPs (explaining
330 4.9% of the 25OHD variance), 100 had a minor allele frequency of more than 1%, of
331 which 78 were directly available in the COVID-19 HGI GWAS summary statistic and had
332 linkage disequilibrium coefficient of less than 5%. Additionally, 3 more variants had good
333 genetic proxies ($r^2 > 90\%$) and were therefore added to our instrument lists, for a total of
334 81 variants. These explained 4.3% of the variance in 25OHD serum levels. The full list of
335 SNPs used can be found in **Supplement 4**.

336

337 *COVID-19 outcome definitions and GWASs*

338 Using the COVID-19 HGI results restricted to cohorts of European ancestry, we used
339 a total of 14,134 cases and 1,284,876 controls to define COVID-19 susceptibility, 6,406
340 cases and 902,088 controls to define COVID-19 hospitalization, and 4,336 cases and
341 623,902 controls to define COVID-19 severe disease. **Table 1** summarizes the definition
342 and sample size of both the exposure and outcome GWASs. Since the UKB was used in
343 the two phases of the MR study, some overlap between the exposure and the outcome
344 GWASs was unavoidable (**Supplement 2**).

345
346 *Primary MR analysis*

347 We first used IVW meta-analysis to combine effect estimates from each genetic
348 instrument. For a standard deviation increase in log-transformed 25OHD level, we
349 observed no statistically significant effect upon odds of susceptibility (OR = 0.97; 95%
350 CI: 0.85, 1.10; P = 0.61). Of note, in the UKB, the distribution of 25OHD levels has a
351 mean of 48.6 nmol/L and a standard deviation of 21.1 nmol/L. This standard deviation is
352 comparable to what can be achieved with vitamin D supplementation, especially over
353 short therapeutic courses [41]. Similarly, we observed no significant difference in risk of
354 hospitalization (OR = 1.11; 95% CI: 0.91, 1.35; P = 0.30) or risk of severe disease (OR =
355 0.93; 95% CI: 0.73, 1.17; P = 0.53) associated with a standard deviation increase in log-
356 transformed 25OHD level (**Table 2** and **Figure 1**).

357
358 *Horizontal pleiotropy assessment and sensitivity analysis*

359 Using the MR Egger intercept terms, we did not observe evidence of horizontal
360 pleiotropy. While they have less statistical power than IVW meta-analysis, the 6
361 sensitivity meta-analyses we used also showed no evidence of an association between
362 25OHD levels and COVID-19 susceptibility, hospitalization, and severe disease, with
363 each confidence interval crossing the null in the primary analysis using all SNPs (**Figure**
364 **1** and **Supplement 5**). Our results are therefore unlikely to be strongly biased by
365 horizontal pleiotropy.

366
367 Second, we restricted our analysis to SNPs which reside close to the four genes directly
368 involved in 25OHD metabolism. This left 12 SNPs, explaining 3.2% of 25OHD variation.
369 Using IVW, each standard deviation increase in log-transformed 25OHD was again not
370 associated with COVID-19 susceptibility (OR = 0.96; 95% CI: 0.83, 1.11; P = 0.59),
371 hospitalization (OR = 1.07 [95% CI: 0.78, 1.47]; P = 0.67) and severe disease (OR =
372 0.87; 95% CI: 0.63, 1.19; P = 0.38). For the three phenotypes, the MR Egger intercept
373 term did not support bias from directional horizontal pleiotropy.

374
375 Lastly, we used the Phenoscanner [37,38] tool to check if the SNPs used in the MR
376 study were associated with other phenotypes. Using Phenoscanner, rs11723621 was
377 associated with white blood cell level, and rs6127099 was associated with glomerular
378 filtration rate [42,43]. In both cases, the association with each phenotype was mild
379 compared to their effect on 25OHD level, as rs11723621 explained less than 0.03% of
380 the variance in white blood cell counts, and rs6127099 explained less than 0.001% of
381 the glomerular filtration rate variance. Removing these SNPs from the 12 SNPs above
382 further decreased the proportion of 25OHD variance explained to 1.7%. While
383 confidence intervals widened, effect estimates when restricting our analysis to these
384 SNPs remained null for susceptibility (0.80; 95% CI: 0.77, 1.23; P=0.80), hospitalization
385 (1.09; 95% CI: 0.68, 1.75; P=0.71), and severe disease (0.91; 95% CI: 0.54, 1.55;
386 P=0.73).

387

388 *Genetic instruments heterogeneity*

389 Overall, our results showed little evidence of heterogeneity of effect between our genetic
390 instruments (**Table 2**). We nonetheless observed that for at least one of the three
391 analyses, we would have rejected the null hypothesis of homogeneous genetic effects in
392 the COVID-19 hospitalization phenotype. However, given the large number of
393 hypotheses tested, this may be due to chance.

394

395 **Discussion**

396 In this large-scale MR study, we did not find evidence to support increasing 25OHD
397 levels in order to protect against COVID-19 susceptibility, hospitalization, or severity.
398 This lack of evidence was consistent across phenotypes, sensitivity analyses, and
399 choice of genetic instruments. Differences between our findings and those reported in
400 observational studies [6] may reflect the fact that associations between vitamin D and
401 COVID-19 may be confounded due to factors difficult to control for even with advanced
402 statistical adjustments, such as socio-economic status, institutionalization or medical
403 comorbidities associated with lower vitamin D levels. While our study assessed the
404 association between genetically determined levels of 25OHD and COVID-19, these
405 results can still inform us on the role of vitamin D supplementation. Specifically, in
406 contrast to observational studies, our findings do not support an association between
407 higher 25OHD level and better COVID-19 outcome, and therefore do not support the use
408 of vitamin D supplementation to prevent COVID-19 outcomes. Further, while a
409 randomized trial [15] showed benefit of vitamin D supplementation using an endpoint at
410 risk of bias due to the unblinded intervention (admission to the critical care unit) and a
411 small sample size (n=75), a larger randomized trial [16] of 240 patients showed no effect
412 of a single high dose of vitamin D3 on mortality, length of stay, or risk of mechanical
413 ventilation. Thus, findings from the largest randomized trial to date are thus concordant
414 with our MR results.

415

416 Our study's main strength is MR's track record of predicting RCT outcomes for multiple
417 medical conditions [9–11,21–24,44,45]. Our study also leverages the largest
418 cohort of COVID-19 cases and controls currently available (even outside of genetic
419 studies) and the largest study on genetic determinants of 25OHD levels to date. Using
420 these data sources, we were able to obtain results robust to multiple sensitivity analysis.

421

422 Our study still has limitations. First, our results do not apply to individuals with vitamin D
423 deficiency, and it remains possible that truly deficient patients may benefit from
424 supplementation for COVID-19 related protection and outcomes. However, individuals
425 who are found to have frank vitamin D deficiency, should undergo replacement for bone
426 protection. Second, our study may suffer from weak instrument bias, especially within
427 sensitivity analyses that restricted to smaller sets of genetic instruments. In two-sample
428 MR, this bias would tend to make estimates closer to the null. Nonetheless, similar
429 studies have been able to use MR to establish an association between 25OHD levels
430 and other diseases (most notably multiple sclerosis [25]), suggesting that these
431 instruments are strong enough to find such associations. Further, given the large
432 percentage of shared individuals from the UKB between the vitamin D exposure GWAS
433 [28] and the severe COVID-19 phenotype, this analysis is close to a one-sample MR,
434 which would show bias towards the observational study association. Given that this
435 analysis also shows largely null effects, we do not suspect that weak instruments bias is
436 a significant issue in our results. Third, given that vitamin D levels are affected by
437 season (with higher levels after sunlight exposure), even if our SNP-instruments were
438 obtained from a GWAS that controlled for season of blood draw, effect attenuation by

439 averaging the effect of 25OHD levels on COVID-19 over all seasons may influence
440 results. Nevertheless, a recent study in a Finnish cohort (where sun exposure greatly
441 varies by season) showed that genetic determinants of 25OHD level were able to
442 discriminate between individuals with predisposition to varying levels of 25OHD,
443 regardless of the season [46]. Therefore, while the cyclical nature of 25OHD level is not
444 completely modelled by MR, the size of this bias is likely small. Fourth, our MR analyses
445 assume a linear exposure-outcome relationship. While this may slightly bias our results,
446 simulation studies have previously shown that this assumption provides adequate results
447 when looking at a population effect [27]. Therefore, for the purpose of vitamin D
448 supplementation in the general population, our conclusions should still be valid.
449 However, as pointed out above, we are not able to test the effect of vitamin D deficiency
450 on COVID-19 outcomes. Lastly, as we only studied the effect of 25OHD and COVID-19
451 in individuals of European ancestry, it remains possible that 25OHD levels might have
452 different effects on COVID-19 outcomes in other populations. However, previous RCTs
453 on vitamin D supplementation have given similar results in populations of various
454 ancestries [44,45].

455

456 In conclusion, using a method that has consistently replicated RCT results from vitamin
457 D supplementation studies in large sample sizes, we find no evidence to support a
458 protective role for higher 25OHD on COVID-19 outcomes. Specifically, vitamin D
459 supplementation as a public health measure to improve COVID-19 outcomes is not
460 supported by this MR study. Most importantly, our results suggest that investment in
461 other therapeutic or preventative avenues should be prioritized for COVID-19 RCTs.

462

463 **Acknowledgements**

464 We thank the patients and investigators who contributed to the COVID-19 HGI
465 (**Supplement 6**) and the Vitamin D GWAS consortium. Members of the GEN-COVID
466 study are acknowledged in **Supplement 7**. This research has been conducted using the
467 UK Biobank Resource (project number: 27449).

468

469 **Contributions**

470 Conception and design: GBL, TN, JBR. Data acquisition and standardization: AR, AG,
471 DRM, TA, OA, NM, NK, ZA. Data analyses: GBL and TN. Interpretation: GBL, TN, VM,
472 DRM, TA, OA, NM, NK, ZA, AR, AG, SZ, YC, VF, JBR. Computational resources and
473 support: VF, JBR. Writing original draft: GBL, TN, JBR. All authors were involved in
474 reviewing the manuscript and critically reviewed its content. All authors gave final
475 approval of the version to be published. The corresponding author attests that all listed
476 authors meet authorship criteria and that no others meeting the criteria have been
477 omitted.

478

479 **Supplementary files captions**

480 **Supplement 1:** STROBE case-control study checklist

481 **Supplement 2:** Cohorts used for each outcome phenotype for the COVID-19 Host
482 Genetics Initiative.

483 **Supplement 3:** Vitamin D metabolism pathway and genes involved.

484 **Supplement 4:** Genetic instruments summary statistics.

485 **Supplement 5:** results from Mendelian randomization sensitivity analyses.

486 **Supplement 6:** Acknowledgement to data contributors and the COVID-19 Host Genetics
487 Initiative.

488 **Supplement 7:** GEN-COVID Multicenter Study.

489 **References**

490

- 491 1. McKee M, Stuckler D. If the world fails to protect the economy, COVID-19 will
492 damage health not just now but also in the future. *Nat Med*. 2020;26: 640–642.
493 doi:10.1038/s41591-020-0863-y
- 494 2. Mansur JL, Tajer C, Mariani J, Inserra F, Ferder L, Manucha W. Vitamin D high
495 doses supplementation could represent a promising alternative to prevent or treat
496 COVID-19 infection. *Clin e Investig en Arterioscler Publ Of la Soc Esp*
497 *Arterioscler*. 2020. doi:10.1016/j.arteri.2020.05.003
- 498 3. Charoengam N, Holick MF. Immunologic Effects of Vitamin D on Human Health
499 and Disease. *Nutrients*. 2020;12. doi:10.3390/nu12072097
- 500 4. Marcos-Pérez D, Sánchez-Flores M, Proietti S, Bonassi S, Costa S, Teixeira JP,
501 et al. Low Vitamin D Levels and Frailty Status in Older Adults: A Systematic
502 Review and Meta-Analysis. *Nutrients*. 2020;12. doi:10.3390/nu12082286
- 503 5. Skrobot A, Demkow U, Wachowska M. Immunomodulatory Role of Vitamin D: A
504 Review. *Adv Exp Med Biol*. 2018;1108: 13–23. doi:10.1007/5584_2018_246
- 505 6. Martineau AR, Forouhi NG. Vitamin D for COVID-19: a case to answer? *Lancet*
506 *Diabetes Endocrinol*. 2020. doi:10.1016/S2213-8587(20)30268-0
- 507 7. Munshi R, Hussein MH, Toraih EA, Elshazli RM, Jardak C, Sultana N, et al.
508 Vitamin D insufficiency as a potential culprit in critical COVID-19 patients. *J Med*
509 *Virol*. 2020. doi:10.1002/jmv.26360
- 510 8. Rooney MR, Harnack L, Michos ED, Ogilvie RP, Sempos CT, Lutsey PL. Trends
511 in Use of High-Dose Vitamin D Supplements Exceeding 1000 or 4000
512 International Units Daily, 1999-2014. *JAMA*. 2017;317: 2448–2450.
513 doi:10.1001/jama.2017.4392
- 514 9. Manson JE, Cook NR, Lee I-M, Christen W, Bassuk SS, Mora S, et al. Vitamin D
515 Supplements and Prevention of Cancer and Cardiovascular Disease. *N Engl J*
516 *Med*. 2019;380: 33–44. doi:10.1056/NEJMoa1809944
- 517 10. Pittas AG, Dawson-Hughes B, Sheehan P, Ware JH, Knowler WC, Aroda VR, et
518 al. Vitamin D Supplementation and Prevention of Type 2 Diabetes. *N Engl J Med*.
519 2019;381: 520–530. doi:10.1056/NEJMoa1900906
- 520 11. Wactawski-Wende J, Kotchen JM, Anderson GL, Assaf AR, Brunner RL,
521 O’Sullivan MJ, et al. Calcium plus vitamin D supplementation and the risk of
522 colorectal cancer. *N Engl J Med*. 2006;354: 684–696.
523 doi:10.1056/NEJMoa055222
- 524 12. Sanders KM, Stuart AL, Williamson EJ, Simpson JA, Kotowicz MA, Young D, et
525 al. Annual High-Dose Oral Vitamin D and Falls and Fractures in Older Women: A
526 Randomized Controlled Trial. *JAMA*. 2010;303: 1815–1822.
527 doi:10.1001/jama.2010.594
- 528 13. Martineau AR, Jolliffe DA, Hooper RL, Greenberg L, Aloia JF, Bergman P, et al.
529 Vitamin D supplementation to prevent acute respiratory tract infections:
530 systematic review and meta-analysis of individual participant data. *BMJ*.
531 2017;356: i6583. doi:10.1136/bmj.i6583
- 532 14. Ginde AA, Brower RG, Caterino JM, Finck L, Banner-Goodspeed VM, Grissom
533 CK, et al. Early High-Dose Vitamin D(3) for Critically Ill, Vitamin D-Deficient
534 Patients. *N Engl J Med*. 2019;381: 2529–2540. doi:10.1056/NEJMoa1911124
- 535 15. Entrenas Castillo M, Entrenas Costa LM, Vaquero Barrios JM, Alcalá Díaz JF,
536 López Miranda J, Bouillon R, et al. “Effect of calcifediol treatment and best
537 available therapy versus best available therapy on intensive care unit admission
538 and mortality among patients hospitalized for COVID-19: A pilot randomized
539 clinical study.” *J Steroid Biochem Mol Biol*. 2020/08/29. 2020;203: 105751.

- 540 doi:10.1016/j.jsbmb.2020.105751
- 541 16. Murai IH, Fernandes AL, Sales LP, Pinto AJ, Goessler KF, Duran CSC, et al.
- 542 Effect of a Single High Dose of Vitamin D3 on Hospital Length of Stay in Patients
- 543 With Moderate to Severe COVID-19: A Randomized Clinical Trial. *JAMA*. 2021.
- 544 doi:10.1001/jama.2020.26848
- 545 17. Davies NM, Holmes M V, Davey Smith G. Reading Mendelian randomisation
- 546 studies: a guide, glossary, and checklist for clinicians. *BMJ*. 2018;362: k601.
- 547 doi:10.1136/bmj.k601
- 548 18. Zheng J, Haberland V, Baird D, Walker V, Haycock PC, Hurle MR, et al.
- 549 Phenome-wide Mendelian randomization mapping the influence of the plasma
- 550 proteome on complex diseases. *Nat Genet*. 2020. doi:10.1038/s41588-020-0682-
- 551 6
- 552 19. Suhre K, Arnold M, Bhagwat AM, Cotton RJ, Engelke R, Raffler J, et al.
- 553 Connecting genetic risk to disease end points through the human blood plasma
- 554 proteome. *Nat Commun*. 2017;8: 14357. doi:10.1038/ncomms14357
- 555 20. Hysinger EB, Roizen JD, Mentch FD, Vazquez L, Connolly JJ, Bradfield JP, et al.
- 556 Mendelian randomization analysis demonstrates that low vitamin D is unlikely
- 557 causative for pediatric asthma. *J Allergy Clin Immunol*. 2016/08/20. 2016;138:
- 558 1747-1749.e4. doi:10.1016/j.jaci.2016.06.056
- 559 21. Ye Z, Sharp SJ, Burgess S, Scott RA, Imamura F, Langenberg C, et al.
- 560 Association between circulating 25-hydroxyvitamin D and incident type 2 diabetes:
- 561 a mendelian randomisation study. *Lancet Diabetes Endocrinol*. 2015;3: 35–42.
- 562 doi:10.1016/S2213-8587(14)70184-6
- 563 22. He Y, Timofeeva M, Farrington SM, Vaughan-Shaw P, Svinti V, Walker M, et al.
- 564 Exploring causality in the association between circulating 25-hydroxyvitamin D
- 565 and colorectal cancer risk: a large Mendelian randomisation study. *BMC Med*.
- 566 2018;16: 142. doi:10.1186/s12916-018-1119-2
- 567 23. Trajanoska K, Morris JA, Oei L, Zheng H-F, Evans DM, Kiel DP, et al.
- 568 Assessment of the genetic and clinical determinants of fracture risk: genome wide
- 569 association and mendelian randomisation study. *BMJ*. 2018;362: k3225.
- 570 doi:10.1136/bmj.k3225
- 571 24. Manousaki D, Mokry LE, Ross S, Goltzman D, Brent Richards J. Mendelian
- 572 Randomization Studies Do Not Support a Role for Vitamin D in Coronary Artery
- 573 Disease. *Circ Cardiovasc Genet*. 2016;9: 349–356.
- 574 doi:10.1161/CIRCGENETICS.116.001396
- 575 25. Mokry LE, Ross S, Ahmad OS, Forgetta V, Smith GD, Leong A, et al. Vitamin D
- 576 and Risk of Multiple Sclerosis: A Mendelian Randomization Study. *PLoS Med*.
- 577 2015;12: e1001866. Available: <https://doi.org/10.1371/journal.pmed.1001866>
- 578 26. Davey Smith G, Ebrahim S. ‘Mendelian randomization’: can genetic epidemiology
- 579 contribute to understanding environmental determinants of disease?*. *Int J*
- 580 *Epidemiol*. 2003;32: 1–22. doi:10.1093/ije/dyg070
- 581 27. Burgess S, Davies NM, Thompson SG. Instrumental variable analysis with a
- 582 nonlinear exposure-outcome relationship. *Epidemiology*. 2014;25: 877–885.
- 583 doi:10.1097/EDE.000000000000161
- 584 28. Manousaki D, Mitchell R, Dudding T, Haworth S, Harroud A, Forgetta V, et al.
- 585 Genome-wide Association Study for Vitamin D Levels Reveals 69 Independent
- 586 Loci. *Am J Hum Genet*. 2020;106: 327–337.
- 587 doi:<https://doi.org/10.1016/j.ajhg.2020.01.017>
- 588 29. Lawlor DA. Commentary: Two-sample Mendelian randomization: opportunities
- 589 and challenges. *Int J Epidemiol*. 2016/07/17. 2016;45: 908–915.
- 590 doi:10.1093/ije/dyw127

- 591 30. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP.
592 The Strengthening the Reporting of Observational Studies in Epidemiology
593 (STROBE) statement: guidelines for reporting observational studies. *PLoS Med*.
594 2007;4: e296. doi:10.1371/journal.pmed.0040296
- 595 31. Machiela MJ, Chanock SJ. LDlink: a web-based application for exploring
596 population-specific haplotype structure and linking correlated alleles of possible
597 functional variants. *Bioinformatics*. 2015;31: 3555–3557.
598 doi:10.1093/bioinformatics/btv402
- 599 32. The COVID-19 Host Genetics Initiative, a global initiative to elucidate the role of
600 host genetic factors in susceptibility and severity of the SARS-CoV-2 virus
601 pandemic. *Eur J Hum Genet*. 2020;28: 715–718. doi:10.1038/s41431-020-0636-6
- 602 33. Griffith GJ, Morris TT, Tudball MJ, Herbert A, Mancano G, Pike L, et al. Collider
603 bias undermines our understanding of COVID-19 disease risk and severity. *Nat*
604 *Commun*. 2020;11: 5749. doi:10.1038/s41467-020-19478-2
- 605 34. Walker VM, Davies NM, Hemani G, Zheng J, Haycock PC, Gaunt TR, et al. Using
606 the MR-Base platform to investigate risk factors and drug targets for thousands of
607 phenotypes. *Wellcome open Res*. 2019;4: 113.
608 doi:10.12688/wellcomeopenres.15334.2
- 609 35. Burgess S, Davies NM, Thompson SG. Bias due to participant overlap in two-
610 sample Mendelian randomization. *Genet Epidemiol*. 2016; 1–12.
611 doi:10.1002/gepi.21998
- 612 36. Slob EAW, Burgess S. A comparison of robust Mendelian randomization methods
613 using summary data. *Genet Epidemiol*. 2020;44: 313–329.
614 doi:<https://doi.org/10.1002/gepi.22295>
- 615 37. Kamat MA, Blackshaw JA, Young R, Surendran P, Burgess S, Danesh J, et al.
616 PhenoScanner V2: an expanded tool for searching human genotype-phenotype
617 associations. *Bioinformatics*. 2019;35: 4851–4853.
618 doi:10.1093/bioinformatics/btz469
- 619 38. Staley JR, Blackshaw J, Kamat MA, Ellis S, Surendran P, Sun BB, et al.
620 PhenoScanner: a database of human genotype-phenotype associations.
621 *Bioinformatics*. 2016;32: 3207–3209. doi:10.1093/bioinformatics/btw373
- 622 39. Bycroft C, Freeman C, Petkova D, Band G, Elliott LT, Sharp K, et al. The UK
623 Biobank resource with deep phenotyping and genomic data. *Nature*. 2018;562:
624 203–209. doi:10.1038/s41586-018-0579-z
- 625 40. Manousaki D, Dudding T, Haworth S, Hsu Y-H, Liu C-T, Medina-Gómez C, et al.
626 Low-Frequency Synonymous Coding Variation in CYP2R1 Has Large Effects on
627 Vitamin D Levels and Risk of Multiple Sclerosis. *Am J Hum Genet*. 2017;101:
628 227–238. doi:<https://doi.org/10.1016/j.ajhg.2017.06.014>
- 629 41. Żebrowska A, Sadowska-Krępa E, Stanula A, Waśkiewicz Z, Łakomy O, Bezuglov
630 E, et al. The effect of vitamin D supplementation on serum total 25(OH) levels and
631 biochemical markers of skeletal muscles in runners. *J Int Soc Sports Nutr*.
632 2020;17: 18. doi:10.1186/s12970-020-00347-8
- 633 42. Astle WJ, Elding H, Jiang T, Allen D, Ruklisa D, Mann AL, et al. The Allelic
634 Landscape of Human Blood Cell Trait Variation and Links to Common Complex
635 Disease. *Cell*. 2016;167: 1415-1429.e19. doi:10.1016/j.cell.2016.10.042
- 636 43. Wuttke M, Li Y, Li M, Sieber KB, Feitosa MF, Gorski M, et al. A catalog of genetic
637 loci associated with kidney function from analyses of a million individuals. *Nat*
638 *Genet*. 2019;51: 957–972. doi:10.1038/s41588-019-0407-x
- 639 44. Aloia JF, Talwar SA, Pollack S, Yeh J. A randomized controlled trial of vitamin D3
640 supplementation in African American women. *Arch Intern Med*. 2005;165: 1618–
641 1623. doi:10.1001/archinte.165.14.1618

- 642 45. Aspray TJ, Chadwick T, Francis RM, McColl E, Stamp E, Prentice A, et al.
643 Randomized controlled trial of vitamin D supplementation in older people to
644 optimize bone health. *Am J Clin Nutr.* 2019;109: 207–217.
645 doi:10.1093/ajcn/nqy280
646 46. Sallinen RJ, Dethlefsen O, Ruotsalainen S, Mills RD, Miettinen TA, Jääskeläinen
647 TE, et al. Genetic Risk Score for Serum 25-Hydroxyvitamin D Concentration
648 Helps to Guide Personalized Vitamin D Supplementation in Healthy Finnish
649 Adults. *J Nutr.* 2020. doi:10.1093/jn/nxaa391
650
651

Phenotype	Source of genetic variants	
	Cohort	Participants
25OHD circulating levels	Manousaki <i>et al</i>	Meta-analysis of two 25OHD GWAS: <ul style="list-style-type: none"> - 401,460 adult white British participants from the UKB - 42,274 from an international consortium of adult individuals of European ancestry
COVID-19 susceptibility	Susceptibility	Meta-analysis of 22 GWAS performed in individuals of European ancestry from 11 countries: <ul style="list-style-type: none"> - Cases: 14,134 individuals with COVID-19 by laboratory confirmation, chart review, or self-report - Controls: 1,284,876 individuals without confirmation or history of COVID-19
COVID-19 severity	Hospitalized	Meta-analysis of 13 GWAS performed in individuals of European ancestry from 11 countries: <ul style="list-style-type: none"> - Cases: 6,406 hospitalized individuals with COVID-19 - Controls: 902,088 without hospitalization with COVID-19
	Severe Disease	Meta-analysis of 12 GWAS performed in individuals of European ancestry from 9 countries: <ul style="list-style-type: none"> - Cases: 4,336 SARS-CoV-2 infected hospitalized individuals who died or required respiratory support (intubation, CPAP, BiPAP, continuous external negative pressure, high flow nasal cannula). - Controls: 623,902 without severe COVID-19

Table 1: Sources of data for the analysis. COVID-19 susceptibility and severity outcomes are taken from the COVID-19 HGI. See **Supplement 2** for details on cohorts of COVID-19 susceptibility and severity phenotypes. 25OHD: 25-hydroxy vitamin D. GWAS: genome-wide association study. UKB: UK Biobank. CPAP: continuous positive airway pressure ventilation. BiPAP: bilevel positive airway pressure ventilation.

Outcome	nSNPs	IVW OR (95% CI)	IVW p-value	IVW SNP Heterogeneity p-value	Egger alpha	Alpha p-value
25OHD primary analysis with all SNPs						
Susceptibility	81	0.97 (0.85, 1.10)	0.61	0.008	0.003 (-0.004, 0.009)	0.43
Hospitalization	81	1.11 (0.91, 1.35)	0.30	0.066	0.0002 (-0.010, 0.011)	0.97
Severe disease	81	0.93 (0.73, 1.17)	0.53	0.126	0.010 (-0.002, 0.022)	0.11
25OHD sensitivity analysis restricted to genes in the vitamin D pathway						
Susceptibility	12	0.96 (0.83, 1.11)	0.59	0.160	0.005 (-0.021, 0.032)	0.70
Hospitalization	12	1.07 (0.78, 1.47)	0.67	0.004	0.031 (-0.027, 0.088)	0.32
Severe disease	12	0.87 (0.64, 1.19)	0.38	0.105	0.056 (0.006, 0.106)	0.05
25OHD sensitivity analysis after removal of SNPs identified by Phenoscanner						
Susceptibility	10	0.97 (0.77, 1.23)	0.80	0.083	0.004 (-0.032, 0.040)	0.83
Hospitalization	10	1.09 (0.68, 1.75)	0.71	0.010	0.014 (-0.059, 0.088)	0.71
Severe disease	10	0.91 (0.54, 1.55)	0.73	0.095	0.072 (-0.003, 0.146)	0.01

Table 2: MR results. SNP: single nucleotide polymorphism. nSNPs: number of SNPs retained for this analysis. IVW: inverse-variance weighted method. CI: confidence interval. Confidence intervals were obtained using Normal approximations, explaining minor discrepancies with p-values close to the alpha=5% statistical significance threshold.




Figure 1: Odds ratio point estimates and 95% confidence intervals for a one standard deviation increase in 25OHD levels (on the log scale) on COVID-19 susceptibility and severity. Restricted to 25-OHD Genes: analysis restricted to SNPs near the 4 genes involved in known vitamin D metabolic pathways. Phenoscaner Filtered: analysis restricted to the 4 genes above, and with removal of SNPs identified to have other associations in Phenoscaner. Full results including odds ratios, confidence intervals, and p-values are available in [Supplement 5](#).