1 # Neural markers of suppression in impaired binocular vision
2

3   Freya A. Lygo[1,2], Bruno Richard[3], Alex R. Wade[1,4], Antony B. Morland[1,4] & Daniel H. Baker[1,4,5]
4
1. Department of Psychology, University of York, Heslington, York, UK
2. Institute of Cognitive Neuroscience, University College London, London, UK
3. Department of Mathematics and Computer Science, Rutgers University–Newark, Newark,
New Jersey, USA
4. York Biomedical Research Institute, University of York, Heslington, York, UK
5. Corresponding author: daniel.baker@york.ac.uk

11
12

13 **Precis** (35 words)
14 Evidence of atypical interocular suppression in individuals with impaired binocular vision was
15 obtained using EEG and fMRI, with a single experimental paradigm. This may underlie the deficits
16 in stereopsis experienced by treated amblyopes.

18 *Keywords*: interocular suppression; dichoptic; fMRI; SSVEP; V1
19
20 **Financial support:** This work was supported in part by a Wellcome Trust (ref: 105624) grant,
21 through the Centre for Chronic Diseases and Disorders (C2D2) at the University of York. FL was
22 funded by a PhD student from the University of York. The sponsor or funding organization had no
23 role in the design or conduct of this research.

24
25 **Conflict of interest**: No conflicting relationship exists for any author
26
27 **Running head**: Suppression in impaired binocular vision
28
29 This article contains additional online-only material. The following should appear online-only:
30 Figure S1, Figure S2.
31
32

Abstract

Objective/Purpose: Even after conventional patching treatment, individuals with a history of
amblyopia typically lack good stereo vision. This is often attributed to atypical suppression
between the eyes, yet the specific mechanism is still unclear. Guided by computational models of
binocular vision, we tested explicit predictions about how neural responses to contrast might differ
in individuals with impaired binocular vision.

Design: A 5×5 factorial repeated measures design was used, in which all participants completed
a set of 25 conditions (stimuli of different contrasts shown to the left and right eyes).

Participants: 25 individuals with a history of amblyopia, and 19 control participants with typical
visual development, participated in the study.

Methods: Neural responses to different combinations of contrast in the left and right eyes, were
measured using both electroencephalography (EEG) and functional magnetic resonance imaging
(fMRI). Stimuli were sinusoidal gratings with a spatial frequency of 3c/deg, flickering at 4Hz. In
the fMRI experiment, we also ran population receptive field and retinotopic mapping sequences,
and a phase-encoded localiser stimulus, to identify voxels in primary visual cortex (V1) sensitive
to the main stimulus.

Main outcome measures: The main outcome measures were the signal-to-noise ratio of the
steady state visual evoked potential, and the fMRI β weights from a general linear model.

Results: Neural responses generally increased monotonically with stimulus contrast. When
measured with EEG, responses were attenuated in the weaker eye, consistent with a fixed tonic
suppression of that eye. When measured with fMRI, a low contrast stimulus in the weaker eye
substantially reduced the response to a high contrast stimulus in the stronger eye. This effect was
stronger than when the stimulus-eye pairings were reversed, consistent with unbalanced dynamic
suppression between the eyes.

Conclusions: Measuring neural responses using different methods leads to different conclusions
about visual differences in individuals with impaired binocular vision. Both of the atypical
suppression effects may relate to binocular perceptual deficits, e.g. in stereopsis, and we
anticipate that these measures could be informative for monitoring the progress of treatments
aimed at recovering binocular vision.

Introduction

The binocular visual system is exquisitely sensitive, and has the ability to detect differences
(disparities) between the eyes of well below one minute of arc[1]. This results in a vivid perception
of depth from stereopsis[2] that benefits everyday tasks such as fine motor control (e.g. threading
a needle) and the judgement of relative object distance (e.g. during driving). But in a substantial
minority of individuals (around 1.4%[3]), an optical (e.g. anisometropia) or muscular (e.g.
strabismus) asymmetry between the eyes during childhood disrupts the development of binocular
vision. This can lead to amblyopia, in which vision through the affected eye is significantly
impaired[4]. Such problems can be treated to some extent by orthoptic or surgical interventions,
which recover sensitivity in the weaker eye in a proportion of cases[5]. But even if treatment is
successful in improving vision in the amblyopic eye, binocular vision may not be restored and
stereopsis rarely reaches normal levels[6]. An enduring mystery is the identity of the neural
mechanism that disrupts binocular vision, even in situations where the eyes have similar acuity
and sensitivity.

In clinical practice, binocular visual disturbances in amblyopia are typically attributed to a process
of suppression, whereby the fellow eye suppresses signals from the amblyopic eye[7,8,9]. This
suppression could take several different forms. For example, 'tonic' suppression should persist
even when there is no input to the fellow eye (e.g. if it is closed, patched or pressure blinded, or
simply shown a blank display). This amounts to a fixed attenuation of the signal in the amblyopic
eye that is invariant to signals from the fellow eye[10]. Alternatively, a more 'dynamic' form of
suppression would depend on the current stimulation of the two eyes, such that higher contrasts
in one eye produce greater suppression of the other eye. Interocular suppression has been widely
studied in intact binocular vision, and has several perceptual consequences, such as ocularity
invariance (the observation that our general perception of the world is unchanged whether one or
both eyes are open[11]) and binocular rivalry (the alternation in perception between conflicting
images shown to the two eyes[12]). Impaired binocular vision might result from an imbalance of
these existing processes of interocular suppression.

Distinguishing between these, and other, explanations for binocular impairments has proved
challenging. In some psychophysical paradigms, such as dichoptic contrast discrimination, similar
performance can result even over a wide range of relative amounts of suppression between the
eyes[10]. In other paradigms, tonic and dynamic suppression are equally able to account for the
results[13]. Isolating a direct neural measure of suppression would allow us to distinguish between
different models, and potentially provide an objective index of binocular impairment that could be
used to track improvements during treatment. In the present study we measured visual responses
to stimuli of different contrasts directly with two methods: functional magnetic resonance imaging
(fMRI) and electroencephalography (EEG). These methods have complementary strengths and
weaknesses: fMRI has excellent spatial precision, but poor temporal resolution, whereas EEG
has poor spatial precision, but good temporal resolution. Previous work measuring visual
responses in amblyopia with one or other of these methods has reported generally weaker
responses to stimuli in the amblyopic eye[14,15]. However, the two techniques have not previously
been directly compared using common stimuli.

Here we use both methods to measure contrast-response functions (Figure 1a) for factorial
combinations of contrast shown to the left and right eyes (Figure 2g; see also[11]). We plot the
results as a series of functions, where stimuli of increasing contrast are shown to one eye (the
'target' stimuli) in the presence of a fixed-contrast (but otherwise identical) stimulus in the other
eye (the 'mask' stimulus). The general character of these functions can be predicted by
contemporary models of binocular signal combination[16,17], and correspond well to previous
measurements in intact binocular visual systems using both fMRI[17] and EEG[18]. When the mask
is absent (0% contrast), the model produces a monotonically increasing monocular contrast
response function (left-most curve in Figure 1a). As mask contrast increases, the overall response
becomes larger because it combines the target and mask signals together. However, interocular
suppression causes a surprising reduction in the response to a high contrast mask when a target
of intermediate contrast is added[11,19]. This produces the u-shaped function shown in the final
curve of Figure 1a - the response increase caused by excitation is outweighed by the response
reduction caused by suppression. Responses therefore go down before they go up, giving a direct
measure of interocular suppression.

131
Figure 1: Model predictions and diagrams. Panel (a) shows predictions of the two-stage model (Meese et
al., 2006) for different combinations of mask and target contrasts shown to the left and right eyes. The
model diagram features multiple stages of gain control (boxes), exponentiation (to powers m, p and q),
inhibition (orange arrows), and binocular summation (denoted by Σ). Panel (b) shows a variant of the same
model[10], where the input to the left eye is attenuated prior to any other processing (orange box in the
diagram). This affects the model's behaviour by reducing the response to the affected (e.g. amblyopic) eye
(orange dashed curves). Panel (c) shows a further variant in which there is stronger inhibition from one eye
onto the other (red arrow in the diagram). This has no effect for monocular stimulation (leftmost function),
but increases suppression with high mask contrasts (rightmost functions). Further model details and
equations are given in Appendix 1.

We can disrupt the model shown in Figure 1a in two key ways. First, we can implement tonic
suppression by attenuating the signal in one eye by a constant factor[10]. This reduces the response
in the affected eye, and also weakens its impact on the fellow eye (Figure 1b). Second, we can
implement dynamic suppression by increasing the weight of suppression from one eye onto the
other (Figure 1c). This has no effect on monocular presentations (as there is no signal in the
opposite eye to cause suppression), but with high-contrast masks there is a much greater
reduction in response for intermediate signal contrasts (the u-shaped functions in the right-most
plot become deeper). This experimental paradigm therefore has the potential to distinguish
between these two types of suppression.

152
Our aim in this study was to empirically test specific predictions of these competing models by
measuring neural responses with fMRI and EEG to a common set of visual stimuli. In addition to
testing control participants with typical binocular vision, we also recruited individuals with a history
of binocular disturbance. Although these participants do not all currently meet the diagnostic
criteria for amblyopia (owing to successful treatment), they would very likely have done so in
childhood and/or had they not been treated. Given the widespread incidence of treatment in
countries with developed healthcare systems, understanding the residual binocular deficits in
treated amblyopes is of substantial clinical importance. To summarise our results, we find
attenuated responses to stimuli in the amblyopic eye when measured using EEG, and increased
and asymmetrical interocular suppression in individuals with impaired binocular vision when
measured using fMRI. Surprisingly this takes the form of stronger suppression of the dominant
eye by the weaker eye.

165
166 **Methods**

167
168 *Participants*

169
A total of 44 adult participants completed the EEG experiment, 19 of whom were control
participants with no history of binocular visual abnormalities, and clinically normal vision. The
remaining 25 participants had been diagnosed with amblyopia, or treated for strabismus during
childhood (see Table 1 for further details). Their ages ranged from 17 to 49, with a mean age of
23.8 years (SD of 7.7 years). Approximately half of these participants (12/25, highlighted in bold)
still met the diagnostic criteria for amblyopia at the time of testing, based on a corrected visual
acuity difference of two lines or more between the eyes. The MRI experiments were completed
by 10 of the control participants, and 12 of the patients (A1 - A12 in Table 1). Participants gave
written informed consent, and received financial compensation for their time (£20 per experiment).
The study protocols were approved by the research governance committee of the York
Neuroimaging Centre, and were consistent with the original wording of the Declaration of Helsinki.

181
182 *Apparatus and stimuli*

183
In the MRI scanner, stimuli were displayed using a ProPixx DLP projector (VPixx Ltd., Quebec,
Canada) with a refresh rate of 120Hz, and a resolution of 1920 × 1080 pixels. Viewed from a
distance of 57cm, there were 36 pixels per degree of visual angle. The projector was driven by a

high performance PC. A circular polariser interleaved images intended for the left and right eyes
(effective refresh rate of 60Hz per eye). Images were projected onto a custom acrylic display
panel that maintained the polarisation, and viewed through a front-silvered mirror and passive
stereo polarizer glasses. The maximum luminance was 356cd/m^2 when viewed through the
glasses. In the EEG lab, stimuli were displayed using a gamma-corrected ViewPixx 3D LCD
display (VPixx Ltd.) with a refresh rate of 120Hz, and a resolution of 1920 × 1080 pixels. Viewed
from a distance of 57cm, there were 36 pixels per degree of visual angle. The display was driven
by a Mac Pro computer. Active stereo shutter goggles (NVidia 3D Vision), synchronised by an
infrared signal, allowed segregation of images to the left and right eyes. Through the goggles, the
maximum luminance was 26cd/m^2. Both display systems had low levels of crosstalk, as measured
using a photometer[20]. All experiments were programmed in Matlab, using the Psychophysics
Toolbox extensions[21–23].
199
200 Table 1: summary of patient demographics, clinical history and acuity measurements. Those highlighted in
201 bold currently meet the clinical criteria for amblyopia (acuity difference of two lines or more on a Snellen
202 chart). None of the participants had a residual strabismus.

Participant	Sex	Amblyopic Eye	R acuity	L acuity	Stereo acuity	Detected	Patching	Surgery	Correction
A1	F	Right	20/32	20/25	None	18 mo	1 - 6 y	<18mo	None
A2	M	Right	20/50	20/20	80s	9 y	9 y	None	None
A3	M	Left	20/20	20/50	140s	4 y	4 y	None	None
A4	M	Left	20/20	20/32	140s	18 mo	5 y	18 mo	None
A5	F	Left	20/20	20/40	40s	5 y	5 - 6 y	None	LE: -2.5; RE: + 0.75
A6	M	Left	20/20	20/40	80s	4 y	4 y	None	None
A7	F	Right	20/40	20/20	60s	~10 y	None	None	None
A8	F	Left	20/40	20/80	None	11 y	11 - 12 y	None	None
A9	F	Left	20/16	20/16	400s	6 y	None	None	None
A10	F	Left	20/16	20/20	None	18 mo	4 y	None	LE: +4; RE:+3.75
A11	M	Right	20/20	20/20	140s	5 y	5 y	None	None
A12	F	Right	20/20	20/20	400s	3 y	None	3 & 4 y	None
A13	F	Left	20/20	20/50	None	5 y	5 - 6 y	None	None
A14	F	Left	20/25	20/32	40s	4 y	4 - 6 y	None	None
A15	M	Left	20/16	20/32	None	20 mo	4 y	1 & 7 y	Not known
A16	M	Right	20/120	20/25	400s	9 y	None	None	None
A17	M	Right	20/100	20/16	None	4 y	4 y	None	None
A18	F	Left	20/16	20/16	None	7 y	7 - 10 y	None	None
A19	F	Left	20/20	20/200	None	18 mo	18 mo	None	None
A20	M	Left	20/50	20/50	200s	<6 y	6 y	None	None
A21	F	Right	20/20	20/16	40s	5 y	5 y	None	LE: -1.25; RE: -1.75
A22	F	Left	20/20	20/32	None	~6 y	6 - 7 y	~6 y	LE: +10.5; RE: +9.5
A23	M	Right	20/32	20/20	400s	5 y	5 y	16 y	None
A24	F	Left	20/20	20/40	40s	4 y	4 - 5 y	None	None
A25	M	Left	20/16	20/20	140s	2 y	2 - 3 y	None	None

204
Figure 2: Example stimuli and methodological details. Panels (a,b,c,d,f) show stimuli used in different parts
of the study. Panels (a,b) show ring and wedge plaid stimuli used in retinotopic mapping, with black arrows
(not presented) indicating the direction of motion. Panel (c) shows the noise bar used in population receptive
field (pRF) mapping, which followed the temporal sequence illustrated in panel (e). Panel (d) shows the
plaid localiser stimulus, which followed the positional sequence indicated by the arrows (only one plaid was
visible in a given 6-second window, and the coloured rings were not shown). Panel (f) shows the sine-wave
grating stimuli used to measure contrast response functions. These were presented to the left and right
eyes in different contrast combinations, as illustrated in panel (g). The gratings flickered on and off for 12
seconds according to a 4Hz sine-wave, as shown in panel (h).
214

Retinotopic mapping involved binocularly presented ring and wedge stimuli constructed from a
radial square wave plaid, as illustrated in Figure 2a,b. The plaid had an angular wavelength of 45
degrees (i.e. 8 complete cycles in 360 degrees), a radial frequency of 0.8 cycles per degree, and
flickered in counterphase at 4Hz. Expanding rings had a width of 1 plaid cycle, and a period of 12
seconds per sequence (4 ring positions). Rotating wedges were 45 degrees (1 cycle) wide, with
a period of 24 seconds per rotation (8 positions in 45 degree clockwise steps). Population
receptive field (pRF) mapping used a drifting bar (0.5×10 degrees) of dynamic 1/f noise, with an
RMS contrast of 0.2 (see Figure 2c). The bar drifted at a speed of 0.4 deg/sec, and followed the
sequence illustrated in Figure 2e. The pRF stimulus was presented to either the left or right eye
in different blocks, with the other eye viewing mean luminance. We used a phase-encoded
localiser stimulus, constructed from a radial plaid with a width of 4 degrees (see Figure 2d),
presented binocularly. The localiser stimulus counterphase flickered at 4Hz, and changed position
every 6 seconds, according to the sequence illustrated in Figure 2d. Each stimulus location had
an x-y offset of ±2.34 degrees from fixation. Stimuli for the main experiments were four horizontal
sine-wave gratings with a spatial frequency of 3c/deg, a cosine-blurred spatial window, and a
width of 4 degrees (see Figure 2f). Five different Michelson contrast values (defined as $100*(L_{max}-$
$L_{min})/(L_{max}+L_{min})$, where L is luminance) were presented in different combinations (see Figure 2g).
The stimuli flickered sinusoidally between 0 and their nominal contrast (on/off flicker) at a
frequency of 4Hz (see Figure 2h). The grating stimuli had x-y offsets of ±2.34 degrees from
fixation. In all experiments, a static binocular texture was presented to aid fusion. This was
constructed from low spatial frequency bandpass filtered noise, and filled the display beyond the
central 12 degree stimulus aperture (see Figure 2 for examples).

MRI acquisition

All MRI data were acquired using a GE 3T HDx Excite MRI scanner. We collected two high
resolution T1-weighted structural scans (TR 7.8 ms; TE 3 ms; voxel size $1 \times 1 \times 1$ mm; 12° flip
angle; matrix size 256×256; FOV 256 mm), and two T2*-weighted fast gradient recalled echo
scans (TR 400 ms; TE 4.2 ms; voxel size $1 \times 1 \times 2$ mm; 25° flip angle; matrix size 128×128;
FOV 260×260 mm), using an 8-channel surface coil (Nova Medical, Wilmington, MA, USA). We
acquired functional images using an EPI sequence with a 16-channel posterior surface coil (Nova
Medical, Wilmington, MA, USA), to optimize signal-to-noise ratio at the occipital pole. The slice
prescription covered the region containing the calcarine sulcus and occipital pole with 39 axial
slices (TR 3000 ms; TE 30 ms; voxel size $2 \times 2 \times 2$ mm; 90° flip angle; matrix size 96×96; FOV
192×192 mm). We also acquired an in-plane proton density scan using the same slice
prescription to aid alignment with the structural scans.

Participants completed the MRI experiments in two sessions. In the first session, we collected the
structural scans, and the retinotopic mapping and pRF data. Each pRF sequence lasted 396
seconds (132 TRs), and either the left or right eye was stimulated. Two repetitions for each eye
were completed. The retinotopic mapping (ring and wedge) sequences were collected as a single
scan lasting 204 seconds (68 TRs). In the second session, the phase-encoded localiser and
contrast response function data were collected. The localiser scan lasted 156 seconds (52 TRs),

and consisted of a blank period (12 seconds), followed by 6 repetitions of the localiser sequence.
The contrast response function sequence lasted 612 seconds (204 TRs), and tested each of the
25 conditions (see Figure 2g) once, with 12-second blank periods between each 12-second trial.
This was repeated four times for each participant. During all functional scans, participants
performed a fixation task, in which they monitored a grid of 9 squares (3x3, each 0.14 degrees
wide) with random luminances in the centre of the screen. They were instructed to press a button
whenever the fixation marker was changed by re-randomising the luminances. This occurred at
randomly determined times, on average once every 48 seconds (i.e. once every two trials). The
task was intended to maintain attention and fixation, and we did not record the responses.

MRI analysis

Primary MRI analysis was conducted in Matlab using the mrVista toolbox
(https://github.com/vistalab/vistasoft). Functional data were motion-corrected within and between
scans, and aligned to the in-plane (proton density) scan, and subsequently to the participant's
anatomical space. The first 12 seconds (4 TRs) of each functional scan were discarded to account
for magnetic saturation effects. Structural scans were processed using Freesurfer[24,25] to generate
a 3D model of the cortex. We created flat patches of unfolded cortex for each hemisphere (120
mm in diameter), centred on the occipital pole (see Figure 3) to facilitate data visualisation and
the creation of Regions of Interest (ROIs). The ring and wedge retinotopy and localiser scans
were summarised by a coherence (travelling wave) analysis[26] to calculate the phase of the BOLD
response at the repetition frequency of the stimulus for each voxel. The pRF data were fit by
estimating (at each voxel) the parameters of a 2D Gaussian function that best predicted the BOLD
timecourse, given the position of the bar stimulus[27]. This was done independently for the left and
right eye scans. The contrast response function data were combined across repetition and
analysed using a general linear model (GLM), with regressors (β weights) for each of the 25
conditions. We used a combination of retinotopy and pRF results to define a V1 ROI on the
flattened cortex using the location of the calcarine sulcus and reversals of phase angle. ROIs
were further restricted using the localiser data, by retaining voxels with a coherence exceeding
0.3. Results were saved as Matlab files, and imported into *R* for statistical analysis and
visualisation. We also converted GLM β weights to MNI space (using tools from FSL[28]), and
averaged them across participants for visualisation on an inflated cortex in the Connectome
Workbench software[29].

EEG acquisition

All EEG data were acquired using a 64-channel ANT Neuroscan system, with electrodes
positioned in a Waveguard cap according to the 10-20 system. Signals were recorded at 1000Hz,
and referenced to the whole-head average. Low-latency digital triggers were sent from the
stimulus computer to the EEG amplifier using a parallel cable, and recorded stimulus onset and
condition codes to the EEG trace. Participants completed 8 repetitions of the contrast response
function experiment. On each repetition, stimuli were presented for trials of 12 seconds, with an
intertrial interval of 3 seconds. All 25 conditions (see Figure 2g) were presented once per
repetition in a random order, taking 375 seconds per block. Participants were given breaks

between blocks. The same fixation task as described for the MRI experiments was performed
throughout the experiment to maintain attention.
304
305   *EEG analysis*
306
Raw data were converted to a compressed *csv* format using functions from EEGlab[30], and were
then imported into *R* for analysis. We took the Fourier transform of the EEG waveform at each
electrode, for a ten-second window beginning one second after stimulus onset (to avoid onset
transients). Fourier spectra were averaged across four occipital electrodes (*Oz*, *POz*, *O1* and *O2*),
and across repetition, using coherent averaging (i.e. retaining the phase information). We then
calculated signal-to-noise ratios (SNRs) by dividing the absolute amplitude in the signal bin (4Hz)
by the mean of the ten adjacent bins (±0.5Hz in steps of 0.1Hz). These SNRs were averaged
across participants, and standard errors were calculated using bootstrapping. To plot the
timecourse of SSVEP activity, we repeated the Fourier transform using a sliding one-second
window (in steps of 10ms), and scaled by the two adjacent bins (±1 Hz) to calculate the SNR.
317
318   *Data and script availability*
319
320   Data, experiment code (in Matlab) and analysis code (in Matlab and R) is available at:
321   http://dx.doi.org/10.17605/OSF.IO/X9ZR8
322
323   **Results**
324
We used the results from retinotopic mapping scans with ring and wedge stimuli (Figure 2a,b),
and population receptive field (pRF) sequences (Figure 2c,e), to identify primary visual cortex
(V1) on flattened discs of occipital cortex for each hemisphere. Example flat maps are shown in
Figure 3 for one control participant (see the project repository for equivalent plots for all
participants: https://osf.io/x9zr8/). The phase angle of the BOLD response to the ring stimuli, and
the pRF eccentricity values, showed a typical central-to-peripheral gradient[27], and were highly
consistent (e.g. across left and right eye pRF sequences). The phase angle across the wedge
and pRF scans showed strong hemisphere/hemi-field segregation (i.e. the right hemisphere
responded to stimuli in the left hemi-field and vice versa), as well as the expected phase
reversals[31] that were used to determine boundaries between V1, V2 and V3 (shown by the white
triangles in each map - the middle triangle is V1). The V1 region-of-interest (ROI) was further
restricted using the responses to a phase-encoded localiser stimulus (see Figure 2d). These
responses also showed strong hemisphere/hemi-field and dorsal/ventral segregation, and we
retained voxels in the V1 ROI that produced responses with a coherence of 0.3 or higher (right-
most flat maps in Figure 3). This resulted in a mean total V1 ROI size (combined across
hemispheres) of 601 voxels, and no significant difference in ROI size between patients and
controls ($t(20)=0.54$, $p=0.60$).
342

Figure 3: Summary of retinotopic mapping and functional localizer results. The icons in the upper right corner indicate the region of cortex in each hemisphere used to make the flat maps (each 120 mm in diameter). The left-most columns show eccentricity estimates from a plaid ring localiser (top row), and pRF models for stimuli shown to the left (middle row) and right (lower row) eyes. The middle columns show the polar angle parameter from a rotating plaid wedge localiser (top row) and the two pRF models (middle and lower rows). The flat maps in the right-most columns show the responses to the phase-encoded plaid localiser. Colour maps for each measure are shown in the lower right corner. In each flat map, dark grey regions indicate sulci, and white triangles show the locations of visual areas V1 (middle triangle) and V2/3v/d (outer triangles). The phase encoded retinotopy and localiser results were thresholded at a coherence value of 0.3, and the pRF results were thresholded at 10% of explained variance.

The localiser-restricted V1 ROIs for each participant were then used to estimate neural responses to stimuli of different contrasts presented to the two eyes (Figure 2f). The BOLD response in this ROI had a typical timecourse[32], which was modulated by stimulus contrast (see Figure 4a). We fitted a general linear model (GLM) to the full timecourse to estimate a β coefficient for each stimulus condition at each voxel. The β weights were strongly modulated by stimulus contrast at the occipital pole (red shading in Figure 4b). In a separate experiment using identical stimuli, we also recorded steady-state visual evoked potentials (SSVEPs) using EEG. These showed clear modulation of signal-to-noise ratio (SNR) with stimulus contrast at the flicker frequency of the stimuli (4Hz, Figure 2g), but with a less sluggish timecourse than the BOLD response (see Figure 4c). The responses were well-isolated in the Fourier amplitude spectrum (see Figure 4d), and localised to occipital electrodes (Figure 4d inset). We therefore used the fMRI β weights averaged across the ROI, and the SSVEP SNRs averaged across four occipital electrodes (black points in the Figure 4d inset) to calculate contrast response functions for each experiment. Exploratory analyses in V2 and V3, and using different localiser thresholds to define the V1 ROI, produced very similar functions (not shown), as did using raw amplitude values for the EEG experiment (rather than SNR values).

Figure 4: Timecourses and topographies of visual responses measured using fMRI and EEG. Panel (a) shows the BOLD timecourse in the localiser-restricted V1 ROI, for binocular presentation at five stimulus contrasts (see legend). The grey rectangle adjacent to the x-axis indicates the period when the stimulus was presented, and error bars indicate ±1SE across participants (N=22). Panel (b) shows averaged beta weights (unthresholded) from the general linear model, projected on a posterior view of each hemisphere, for the non-zero stimulus contrasts averaged across all participants (subtracting the 0% condition as a baseline). Panel (c) shows the SSVEP timecourse as a signal-to-noise ratio (SNR) at the target flicker frequency (4Hz), calculated using a 1000ms sliding window (centred at the time indicated on the x-axis). Shaded regions indicate bootstrapped 95% confidence intervals of the median, calculated across participants (N=44). The grey shaded rectangle represents the period when the stimulus was presented. Panel (d) shows the Fourier spectrum for a 96% contrast binocular target, averaged across all participants (N=44), for 10-second windows of stimulation. Clear peaks in SNR are apparent at the stimulus flicker frequency (4Hz) and its harmonics (especially 8Hz). The grey shaded region indicates 95% confidence intervals of the median. The inset scalp topography shows that responses were strongest at posterior electrode sites over early visual areas. For panels (c,d), signals were averaged across the electrodes indicated in black on the scalp plot (*Oz*, *POz*, *O1* and *O2*).

Figure 5 shows contrast response functions from both experiments, split by participant group (control participants in Figure 5a,c, patients in Figure 5b,d). The first contrast response function in each row (circle symbols) is for monocular stimulus presentation, and shows the expected monotonic increase for each data set. As predicted by our computational models (see Figure 1), as mask contrast increased, the functions rose from baseline across the plot. Statistically, there were significant main effects of both target and mask contrast, and significant interactions, for all data sets (see Table 2). For the patients there was a very slight reduction of response in the amblyopic eye (orange circles) compared with the fellow eye (blue circles) at the highest target contrast in the fMRI data (Figure 5b), though this was not significant ($t(11)=1.20$, $p=0.26$, $d=0.35$). The difference was more pronounced in the SSVEP data (Figure 5d), and was significant at 24% target contrast ($t(24)=2.91$, $p<0.01$, $d=0.58$), though not at 96% contrast ($t(24)=1.96$, $p=0.06$, $d=0.39$).

For the control participants, there was evidence of interocular suppression (a u-shaped function)
for the highest mask contrast when measured using fMRI (final function in Figure 5a), though this
was not statistically significant (paired samples t-test comparing 0% and 6% target contrast
conditions when a 96% contrast mask was present, $t(9)=1.8$ $p=0.10$, $d=0.57$, mean difference of
0.13 β units). For the control EEG data interocular suppression occurred at a higher target
contrast (96%, see final function in Figure 5c), but was also not significant ($t(18)=0.96$, $p=0.35$,
$d=0.22$, mean difference of 0.44 SNR units). The u-shaped function was more pronounced in the
patients (Figure 5b,d). For the fMRI data, suppression was more substantial when the mask was
shown to the fellow eye (orange inverted triangles comparing 0% and 6% target levels with a 96%
mask; $t(11)=3.25$, $p=0.008$, $d=0.94$, mean difference of 0.46 β units) than the amblyopic eye (blue
inverted triangles; $t(11)=2.51$, $p=0.029$, $d=0.72$, mean difference of 0.22 β units). A very subtle
suppression effect was qualitatively apparent for the SSVEP data (Figure 5d), but this did not
reach statistical significance for either eye (both $p>0.05$). SSVEP responses at the second
harmonic frequency (8Hz) were broadly similar to those at the fundamental (see Supplementary
Figure S1).
418
419   Table 2: ANOVA results for contrast response functions.

Data set	Effect	F-ratio (df)	p-value	Effect size (ω^2)
Control fMRI	Left eye contrast	18.82 (4,36)	<0.001	0.27
Control fMRI	Right eye contrast	19.56 (4,36)	<0.001	0.22
Control fMRI	Interaction	2.91 (16,144)	<0.001	0.18
Patient fMRI	Fellow eye contrast	12.80 (4,44)	<0.001	0.14
Patient fMRI	Amblyopic eye contrast	8.47 (4,44)	<0.001	0.14
Patient fMRI	Interaction	2.92 (16,176)	<0.001	0.15
Control SSVEP	Left eye contrast	29.93 (4,72)	<0.001	0.31
Control SSVEP	Right eye contrast	27.21 (4,72)	<0.001	0.28
Control SSVEP	Interaction	6.43 (16,288)	<0.001	0.14
Patient SSVEP	Fellow eye contrast	34.88 (4,96)	<0.001	0.38
Patient SSVEP	Amblyopic eye contrast	19.26 (4,96)	<0.001	0.17
Patient SSVEP	Interaction	2.96 (16,384)	<0.001	0.04

420
421

422
423   Figure 5: Contrast response functions measured using fMRI and EEG. Data for control participants (panels
424   a,c) are averaged across complementary conditions for the left and right eyes. Data for patients (panels
425   b,d) are plotted considering the fellow eye as the 'target' eye (blue) and also considering the amblyopic eye
426   as the 'target' eye (orange). These data are identical, but are re-ordered to aid interpretation. Error bars in
427   each plot indicate ±1SE across participants.
428

To further investigate the interocular suppression effect, we plotted full BOLD timecourses for the
condition where the mask only was shown (96% contrast mask to one eye, 0% contrast target to
the other), and the condition where the same mask was paired with a 6% contrast target in the
other eye (see Figure 6). Interocular suppression is clear in each data set, as the white points
appear below the black points over much of the function. Our computational model (see right-
most functions in Figure 1a,c) predicts that this happens because the excitatory impact of the 6%
contrast stimulus is outweighed by its suppression of the response to the 96% contrast stimulus
in the other eye: overall activity goes down instead of up. However this effect is much more
substantial for the fellow eye of the patients (Figure 6b) than for the amblyopic eye (Figure 6c),
consistent with our finding from the β weights (Figure 5b). The cortical meshes along the upper
row of Figure 6 show the difference in β weights between conditions, with blue shading indicating
stronger suppression. Suppression is apparent at the occipital pole, and is again strongest for the
fellow eye of the patients (Figure 6b).
442

443
Figure 6: Timecourse and spatial distribution of interocular suppression measured using fMRI, and polar
plots of SSVEP responses to monocular stimuli. Upper plots show the difference of β weights at each voxel
(unthresholded), between a monocular condition where one eye saw 96% contrast, and a dichoptic
condition where the eyes saw 96% and 6% contrast. Blue shading indicates a suppressive effect (and red
shading a facilitatory effect) of the 6% component. Lower panels show the timecourse of % signal change
for the same two conditions. Data are shown for (a) control participants, (b) the 96% contrast mask in the
fellow eye of the patients, and (c) the 96% contrast mask in the amblyopic eye of the patients. Error bars
indicate ±1SE across participants, and grey shaded rectangles show the duration of stimulus presentation.
Panels (d,e) show SSVEP responses for monocularly presented stimuli at 24% (d) and 96% (e) contrast.
This representation shows a phase lag (i.e. angular difference) between the amblyopic (orange) and fellow
(blue) eyes. Shaded regions indicate ±1SE, calculated independently for amplitude and phase values.

The lower two panels of Figure 6 show polar plots comparing monocular SSVEP responses to
stimuli of 24% contrast (Fig 6d), and 96% contrast (Fig 6e). There is a phase lag between the
fellow and amblyopic eye in both panels. For the 24% contrast target, this is approximately 112
degrees, which corresponds to a lag of around 78 ms at the 4Hz flicker frequency used here. For
the 96% contrast, the lag is around 28 degrees (20 ms). This latter estimate corresponds well with
previously reported phase lags using a similar paradigm in magnetoencephalography (MEG)[33].
461
462   **Discussion**
463
We measured neural responses to different combinations of contrast in the left and right eyes,
using both EEG and fMRI. In participants with atypical binocular vision, we found reduced
responses in the amblyopic eye using EEG, and increased suppression between the eyes
(compared with controls) in V1 using fMRI. These different effects are consistent with greater
tonic and dynamic suppression (respectively) in individuals with impaired binocular vision, and
may be responsible for the deficits in stereopsis experienced by the majority of these participants
(see Table 1). We now discuss why the results differ across measurement methods, what these
findings tell us about amblyopic suppression, and how treatments might be targeted towards the
development of functioning binocular vision.
473
474   *Comparison of EEG and fMRI measures*
475
This is the first study to use both EEG and fMRI to investigate contrast processing in impaired
binocular vision. Previous studies using either EEG or MEG[14,33] have typically found larger
amblyopic deficits in the monocular response than those using fMRI[34,35] (though some work has
shown substantial fMRI deficits[36]), mirroring our results here (left-most functions of Figure 5b,d).
Considering only this previous work, heterogeneity of stimuli and participants across studies might
well have explained the differences. However in the present experiments we used identical
stimuli, and the same participants completed both experiments (Supplementary Figure S2 shows
the EEG data for only the participants who completed the MRI experiments). The reduction in
response amplitude to stimuli in the weaker eye is clearer using SSVEP than using fMRI.
However, differences in dynamic interocular suppression are only apparent when measured with
fMRI (right-most functions in Figure 5a,b; Figure 6a-c).
487
Analogous differences between these two methods have recently been reported in the study of
attention. Itthipuripat et al.[37] found that spatial attention produces a change in baseline response
when measured using fMRI, but not when measured using EEG (including evoked potentials and
SSVEPs). Instead fMRI, which measures global neural activity indirectly via oxygen consumption,
appears to be more closely related to late ERP components and alpha band oscillations recorded
using EEG[37], rather than the stimulus onset transients detected by SSVEP and early ERP
components. Although our contrast response functions do not involve baseline shifts, the other
differences between our fMRI and EEG results are consistent with different features of neural
activity being probed by these two methods.
497

One possible explanation for the differences in monocular response between methods is that the
firing of visual neurons responsive to the amblyopic eye might be desynchronised. This would
have a greater effect on phase-locked SSVEP responses – which depend upon synchronised
firing – than on fMRI BOLD responses, which are a proxy for overall neural activity. Instead,
asynchronous activity in higher frequency bands (50 – 200Hz) shows a closer correspondence
with the BOLD response[38], and these signals can be detected with extracranial techniques such
as MEG[39]. In terms of the differences in suppression, this could reflect processing in different
layers of cortex. Evoked responses measured using EEG correspond most closely to activity in
the more superficial (supragranular) layers of cortex, whereas much inhibitory processing involves
deeper (granular) layers[40]. Future work using laminar fMRI at higher field strength (7T or above[41]),
or multi-unit electrophysiology[42,43], could allow dissociation of excitatory and inhibitory responses
in amblyopia. In addition, these methods can resolve ocular dominance columns in V1, allowing
eye-specific inputs to be measured directly.

What is amblyopic suppression?

Suppression in amblyopia is measured very differently in clinical practice compared with the
methods used in lab-based neuroscience research. The most common clinical measures are the
Worth 4-dot test, and Bagolini striated lenses, both of which give a qualitative indication of whether
one eye's image is substantively suppressed by the other eye during binocular viewing. Lab-
based measures involve a variety of paradigms, including psychophysical approaches such as
dichoptic masking[10,44–46], or assessing binocular fusion of edges[47] or gratings[48], and more direct
neurophysiological estimates of suppression in animal models[49,50]. However, because both eyes
are being stimulated during testing, clinical suppression could in principle be explained by either
tonic or dynamic suppression, and most psychophysical work also cannot distinguish between
these possibilities. Many of our patients (around half) no longer met the clinical criterion for
amblyopia, yet as a group they still exhibited greater dynamic interocular suppression than our
control participants, as well as tonic suppression of one eye. The finding that both types of
suppression are apparent, even in individuals who have received patching or surgical
interventions, strongly suggests that both will also be present in untreated amblyopes. Amblyopic
suppression might therefore involve a combination of these two processes.

One feature that remains to be determined is whether one type of suppression is a primary cause
of amblyopia, and the other a later consequence. Cortical suppression, characterised as a
process of gain control[51], is a dynamic, adaptive process that acts to optimise the sensory
response[52]. It is clear that binocular vision is particularly plastic, as the relative weighting of the
two eyes can be altered following a brief period of occlusion[53], and this may be the mechanism
by which clinical patching treatment improves vision. Our finding of increased suppression of the
fellow eye by the amblyopic eye could be an attempt to rebalance an asymmetrical system. If so,
this might influence the development of novel treatments geared towards modulating interocular
suppression.

Perceptual consequences of suppression, and potential for targeted treatment

It is generally assumed that clinical suppression results in viewing the world 'through' the fellow
eye, with signals from the amblyopic eye being completely suppressed. However, other work has
shown that when signals in the amblyopic eye are boosted by an appropriate amount, information
is still summed binocularly[54]. Indeed, the principle of ocularity invariance (that the world does not
change when one eye is closed) makes it very difficult to distinguish between suppression and
fusion outside of the laboratory or clinic. However, even if the amblyopic eye still contributes to
perception, stereopsis is extremely sensitive to imbalances between the eyes, and breaks down
when one eye receives a stronger input[55]. The two types of suppression we identify here would
likely unbalance signals in exactly this way, which may contribute to the poorer stereopsis for
most of our patients (see Table 1).
552
Consistent with this idea, several treatments have recently been proposed that aim to reduce
suppression between the eyes. For example, antisuppression therapy[56] involves presenting dot
motion stimuli, in which signals are shown to one eye, and noise distractors to the other. To
perform the task, patients must favour information from the signal (amblyopic) eye. Performance
improves over time, implying that suppression is reduced. This approach also improves acuity
and stereopsis, even in amblyopic adults far beyond the critical period (in childhood) for traditional
treatment. Related treatments that involve playing dichoptic video games[57–59], or watching
dichoptic movies[60], may work in a similar way. Measuring the two types of suppression identified
here throughout treatment would reveal causal relationships between suppression and visual
function, potentially allowing clinicians to optimise treatment schedules and monitor progress.
563
564   **Appendix 1 – details of computational models**
565
566   The two stage model of Meese et al.[16] comprises an intial stage of monocular gain control,
567   followed by binocular summation, defined as:
568

$$binsum = \frac{C_L^m}{S + C_L + \omega_R C_R} + \frac{C_R^m}{S + C_R + \omega_L C_L}$$

569
570
571   where m = 1.3, S = 1, and $\omega_R = \omega_L$ = 1, C_L and C_R are the input contrasts to the left and right eyes.
572   The output of the model follows a further nonlinearity:
573

$$resp = \frac{binsum^p}{Z + binsum^q}$$

574
575
576   where p = 8, q = 6.5, and Z = 0.1. The predictions shown in Figure 1a are the output of the model
577   for different combinations of C_L and C_R. To predict the effects of attenuating one eye's input
578   (Figure 1b), an attenuator parameter[10] is added to one eye's input:
579

$$binsum = \frac{aC_L^m}{S + aC_L + \omega_R C_R} + \frac{C_R^m}{S + C_R + \omega_L aC_L}$$

580
581

582   where a = 0.5. Finally, to assess the impact of unbalanced interocular suppression (Figure 1c),
583   we set ω_R = 2, but left ω_L = 1 (and set a = 1). Precise parameter values, e.g. of exponents, are
584   not essential to produce the overall model behaviour.
585
586   **Acknowledgements**
587

588   We are grateful to all of our participants for their involvement in this work, and to Robert Hess for
589   helpful discussions on suppression in amblyopia.

590
591   **References**
592

1. Coutant BE, Westheimer G. Population distribution of stereoscopic ability. Ophthalmic and
Physiological Optics 1993;13:3–7.
2. Julesz B. Stereoscopic vision. Vision Res 1986;26:1601–1612.
3. Fu Z, Hong H, Su Z, et al. Global prevalence of amblyopia and disease burden projections through
2040: a systematic review and meta-analysis. British Journal of Ophthalmology 2020;104:1164–1170.
4. Hess RF. Contrast sensitivity assessment of functional amblyopia in humans. Trans Ophthalmol Soc U
K 1979;99:391–397.
5. Levi DM. Rethinking amblyopia 2020. Vision Research 2020;176:118–129.
6. Levi DM, Knill DC, Bavelier D. Stereopsis and amblyopia: A mini-review. Vision Research
2015;114:17–30.
7. Jampolsky A. Characteristics of Suppression in Strabismus. Archives of Ophthalmology 1955;54:683–
696.
8. Pratt-Johnson JA, Tillson G. Suppression in strabismus--an update. British Journal of Ophthalmology
1984;68:174–178.
9. Travers T aB. Suppression of vision in squint and its association with retinal correspondence and
amblyopia. British Journal of Ophthalmology 1938;22:577–604.
10. Baker DH, Meese TS, Hess RF. Contrast masking in strabismic amblyopia: attenuation, noise,
interocular suppression and binocular summation. Vision Res 2008;48:1625–1640.
11. Baker DH, Meese TS, Georgeson MA. Binocular interaction: contrast matching and contrast
discrimination are predicted by the same model. Spatial Vision 2007;20:397–413.
12. Levelt WJM. The alternation process in binocular rivalry. British Journal of Psychology 1966;57:225–
238.
13. Ding J, Levi DM. Rebalancing binocular vision in amblyopia. Ophthalmic and Physiological Optics
2014;34:199–213.
14. Baker DH, Simard M, Saint-Amour D, Hess RF. Steady-State Contrast Response Functions Provide
a Sensitive and Objective Index of Amblyopic Deficits. Investigative Ophthalmology & Visual Science
2015;56:1208–1216.
15. Barnes GR, Hess RF, Dumoulin SO, et al. The cortical deficit in humans with strabismic amblyopia.
The Journal of Physiology 2001;533:281–297.
16. Meese TS, Georgeson MA, Baker DH. Binocular contrast vision at and above threshold. J Vis
2006;6:1224–1243.
17. Moradi F, Heeger DJ. Inter-ocular contrast normalization in human visual cortex. J Vis 2009;9:13.1-
22.
18. Baker DH, Wade AR. Evidence for an Optimal Algorithm Underlying Signal Combination in Human
Visual Cortex. Cerebral Cortex 2017;27:254–264.
19. Baker DH, Meese TS, Georgeson MA. Paradoxical Psychometric Functions ("Swan Functions") are
Explained by Dilution Masking in Four Stimulus Dimensions. i-Perception 2013;4:17–35.
20. Baker DH, Kaestner M, Gouws AD. Measurement of crosstalk in stereoscopic display systems used
for vision research. Journal of Vision 2016;16:14.
21. Brainard DH. The Psychophysics Toolbox. Spat Vis 1997;10:433–436.
22. Kleiner M, Brainard DH, Pelli DG. What's new in Psychtoolbox-3? Perception 2007;36(S):14.

23. Pelli DG. The VideoToolbox software for visual psychophysics: transforming numbers into movies.
Spat Vis 1997;10:437–442.
24. Dale AM, Fischl B, Sereno MI. Cortical surface-based analysis. I. Segmentation and surface
reconstruction. Neuroimage 1999;9:179–194.
25. Fischl B. FreeSurfer. Neuroimage 2012;62:774–781.
26. Engel SA, Rumelhart DE, Wandell BA, et al. fMRI of human visual cortex. Nature 1994;369:525.
27. Dumoulin SO, Wandell BA. Population receptive field estimates in human visual cortex. NeuroImage
2008;39:647–660.
28. Jenkinson M, Beckmann CF, Behrens TEJ, et al. FSL. Neuroimage 2012;62:782–790.
29. Marcus DS, Harwell J, Olsen T, et al. Informatics and data mining tools and strategies for the human
connectome project. Front Neuroinform 2011;5:4.
30. Delorme A, Makeig S. EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics
including independent component analysis. J Neurosci Methods 2004;134:9–21.
31. Sereno MI, Dale AM, Reppas JB, et al. Borders of Multiple Visual Areas in Humans Revealed by
Functional Magnetic Resonance Imaging. Science 1995;268:889–893.
32. Boynton GM, Engel SA, Glover GH, Heeger DJ. Linear systems analysis of functional magnetic
resonance imaging in human V1. J Neurosci 1996;16:4207–4221.
33. Chadnova E, Reynaud A, Clavagnier S, Hess RF. Latent binocular function in amblyopia. Vision
Research 2017;140:73–80.
34. Conner IP, Odom JV, Schwartz TL, Mendola JD. Monocular activation of V1 and V2 in amblyopic
adults measured with functional magnetic resonance imaging. J AAPOS 2007;11:341–350.
35. Li X, Dumoulin SO, Mansouri B, Hess RF. Cortical Deficits in Human Amblyopia: Their Regional
Distribution and Their Relationship to the Contrast Detection Deficit. Investigative Opthalmology &
Visual Science 2007;48:1575.
36. Hess RF, Li X, Lu G, et al. The contrast dependence of the cortical fMRI deficit in amblyopia; a
selective loss at higher contrasts. Human Brain Mapping 2010;31:1233–1248.
37. Itthipuripat S, Sprague TC, Serences JT. Functional MRI and EEG Index Complementary Attentional
Modulations. The Journal of Neuroscience 2019;39:6162–6179.
38. Hermes D, Nguyen M, Winawer J. Neuronal synchrony and the relation between the blood-oxygen-
level dependent response and the local field potential. PLOS Biology 2017;15:e2001461.
39. Kupers ER, Wang HX, Amano K, et al. A non-invasive, quantitative study of broadband spectral
responses in human visual cortex. PLOS ONE 2018;13:e0193107.
40. Bruyns-Haylett M, Luo J, Kennerley AJ, et al. The neurogenesis of P1 and N1: A concurrent
EEG/LFP study. NeuroImage 2017;146:575–588.
41. de Hollander G, van der Zwaag W, Qian C, et al. Ultra-high resolution fMRI reveals origins of
feedforward and feedback activity within laminae of human ocular dominance columns. BioRxiv
preprint; 2020. Available at: http://biorxiv.org/lookup/doi/10.1101/2020.05.19.102186.
42. Hirsch JA, Martinez LM, Pillai C, et al. Functionally distinct inhibitory neurons at the first stage of
visual cortical processing. Nature Neuroscience 2003;6:1300–1308.
43. Martinez LM, Wang Q, Reid RC, et al. Receptive field structure varies with layer in the primary visual
cortex. Nature Neuroscience 2005;8:372–379.
44. Harrad RA, Hess RF. Binocular integration of contrast information in amblyopia. Vision Res
1992;32:2135–2150.
45. Huang P-C, Baker DH, Hess RF. Interocular suppression in normal and amblyopic vision: Spatio-
temporal properties. Journal of Vision 2012;12:29–29.
46. Zhou J, Reynaud A, Yao Z, et al. Amblyopic Suppression: Passive Attenuation, Enhanced Dichoptic
Masking by the Fellow Eye or Reduced Dichoptic Masking by the Amblyopic Eye? Investigative
Opthalmology & Visual Science 2018;59:4190.
47. Spiegel DP, Baldwin AS, Hess RF. The Relationship Between Fusion, Suppression, and Diplopia in
Normal and Amblyopic Vision. Investigative Opthalmology & Visual Science 2016;57:5810.
48. Ding J, Klein SA, Levi DM. Binocular combination in abnormal binocular vision. Journal of Vision
2013;13:14.
49. Sengpiel F, Blakemore C. Interocular control of neuronal responsiveness in cat visual cortex. Nature
1994;368:847–850.
50. Shooner C, Hallum LE, Kumbhani RD, et al. Asymmetric Dichoptic Masking in Visual Cortex of
Amblyopic Macaque Monkeys. The Journal of Neuroscience 2017;37:8734–8741.

51. Carandini M, Heeger DJ. Normalization as a canonical neural computation. Nat Rev Neurosci
691   2012;13:51–62.
692   52. Westrick ZM, Heeger DJ, Landy MS. Pattern Adaptation and Normalization Reweighting. J Neurosci
693   2016;36:9805–9816.
694   53. Lunghi C, Burr DC, Morrone C. Brief periods of monocular deprivation disrupt ocular balance in
695   human adult visual cortex. Current Biology 2011;21:R538–R539.
696   54. Baker DH, Meese TS, Mansouri B, Hess RF. Binocular Summation of Contrast Remains Intact in
697   Strabismic Amblyopia. Investigative Opthalmology & Visual Science 2007;48:5332.
698   55. Legge GE, Gu YC. Stereopsis and contrast. Vision Res 1989;29:989–1004.
699   56. Hess RF, Mansouri B, Thompson B. A binocular approach to treating amblyopia: antisuppression
700   therapy. Optom Vis Sci 2010;87:697–704.
701   57. Backus BT, Dornbos BD, Tran TA, et al. Use of virtual reality to assess and treat weakness in human
702   stereoscopic vision. Electronic Imaging 2018;2018:109-1-109–6.
703   58. Li J, Thompson B, Deng D, et al. Dichoptic training enables the adult amblyopic brain to learn.
704   Current Biology 2013;23:R308–R309.
705   59. Vedamurthy I, Nahum M, Bavelier D, Levi DM. Mechanisms of recovery of visual function in adult
706   amblyopia through a tailored action video game. Scientific Reports 2015;5. Available at:
707   http://www.nature.com/articles/srep08482.
708   60. Bossi M, Tailor VK, Anderson EJ, et al. Binocular Therapy for Childhood Amblyopia Improves Vision
709   Without Breaking Interocular Suppression. Invest Ophthalmol Vis Sci 2017;58:3031–3043.
710

Online Supplemental materials

Figure S1: SSVEP data for the second harmonic (2F) response. The main trends are the same
as in Figure 5c,d, though the SNRs are overall lower (note y-axis scaling). Panel (a) shows data
from control participants (N=19), and panel (b) shows data from patients (N=25).

Figure S2: SSVEP data for only the participants who completed the fMRI experiment. The main
trends are the same as in Figure 5c,d. Panel (a) shows data from control participants (N=10), and
panel (b) shows data from patients (N=12).