5

Slight reduction in SARS-CoV-2 exposure viral load due to masking results in a

10   **significant reduction in transmission with widespread implementation**

Ashish Goyal[1], Daniel B. Reeves[1], E. Fabian Cardozo-Ojeda[1], Bryan T. Mayer [1]†, Joshua T.

15   Schiffer[1,2,3]*†

[1] Vaccine and Infectious Diseases Division, Fred Hutchinson Cancer Research Center

[2] Department of Medicine, University of Washington, Seattle

[3] Clinical Research Division, Fred Hutchinson Cancer Research Center

20   † These authors contributed equally to the work.

Corresponding author: Joshua T. Schiffer, jschiffe@fredhutch.org

Masks are a vital tool for limiting SARS-CoV-2 spread in the population. Here we utilize a mathematical model to assess the impact of masking on transmission within individual transmission pairs and at the population level. Our model quantitatively links mask efficacy to reductions in viral load and subsequent transmission risk. Our results reinforce that the use of masks by both a potential transmitter and exposed person substantially reduces the probability of successful transmission, even if masks only lower exposure viral load by ~50%. Slight increases in mask adherence and/or efficacy above current levels would reduce the effective reproductive number (R_e) substantially below 1, particularly if implemented comprehensively in potential super-spreader environments. Our model predicts that moderately efficacious masks will lower exposure viral load 10-fold among people who get infected despite masking, potentially limiting infection severity. Because peak viral load tends to occur pre-symptomatically, we also identify that antiviral therapy targeting symptomatic individuals is unlikely to impact transmission risk. Instead, antiviral therapy would only lower R_e if dosed as post-exposure prophylaxis and if given to ~50% of newly infected people within 3 days of an exposure. These results highlight the primacy of masking relative to other biomedical interventions under consideration for limiting the extent of the COVID-19 pandemic prior to widespread implementation of a vaccine.

Introduction

Masks are a barrier method to prevent the spread of respiratory viral infections. A mask essentially serves as a filter that prevents passage of some portion of viruses from the airway of the transmitter to the airway of exposed contacts. Mask efficacy is therefore mediated by a reduction in exposure viral load (*1, 2*). If a potential transmitter as well as an exposed contact are masked, then this filtering process occurs twice potentially amplifying protection.

Mask efficacy is inferior to that of other, less permeable barrier methods to prevent infections such as condoms (*1, 3, 4*). The most commonly used cloth and hospital masks do not provide a perfect facial seal and mask fabric does not block emission or inhalation of all aerosolized viral particles (*5, 6*). N95 masks may bypass these shortcomings but are in short supply and difficult to wear for long periods of time (*7, 8*). As a result of these imperfections, recommendations for mask use have varied over the course of the COVID-19 pandemic. Nevertheless, widespread mask use is recognized as a critical component of any viable public health strategy against COVID-19 (*9-11*). Recent models demonstrate that slight increases in mask utilization could be the single most important factor that prevents exponential growth in incident cases (*12-14*).

Quantifying mask efficacy in real-world settings remains challenging. Elegant experimental work demonstrated the efficacy of masks in animal models (*15*). Many studies have been performed in hospital settings where mask compliance is uniform and other complementary infection prevention methods are more commonly employed than in other public gathering or work locations (*16, 17*). To the best of our knowledge, no study has captured the impact of masking on the likelihood of super-spreader events, with specific consideration of intermittent compliance.

Here we develop a mathematical model capturing viral load-mediated effects of mask use on transmission probability within transmission pairs and at the population level. We use this approach to estimate the efficacy of masks in real world settings, and to characterize effects on super-spreader events as well as exposure viral loads of those who get infected despite masking. Finally, we compare the preventative impact of masking to the use of antiviral therapies given early during symptomatic infection, or when used as post-exposure prophylaxis (PEP).

Results

Baseline mathematical model of SARS-CoV-2 viral load dependent transmission. We
employed a previously developed mathematical model which links viral load shedding at the
individual level with population level epidemic spread (*18, 19*), to determine the impact of masks
on epidemics. Briefly, the model is built upon the assumptions that each transmitter has a
specific number of exposure contacts per day and that each exposure contact has a certain
probability of successful transmission based on the viral load of the transmitter. This probability
is based on a transmission dose (TD) response curve. We fit the model to frequency histograms
describing heterogeneity of individual R0, or the number of secondary infections attributed to
each infected person (*20-24*). The individual R0 distribution for SARS-CoV-2 transmission is
highly over-dispersed, meaning that most infected people do not spread infection while a
minority infect a large number of people. We also fit the model to distributions of individual
serial intervals, the time from the onset of symptoms in the transmitter to symptom onset in the
secondarily infected person (*25*). The overall qualitative conclusions of this model were that the
period of contagiousness for SARS-CoV-2 is quite short, typically less than a day, and that
super-spreader events are largely attributable to high variability in the number of exposure
contacts per day among infected people.

***Predicted impact of transmitter or exposure contact masking on transmission probability
within transmission pairs.*** We added masking to this model by assuming that a mask decreases
the exposure viral load in a transmission pair by a value that we refer to as the combination mask
efficacy (ε_C). This efficacy represents the proportion of viruses filtered by masks worn by both
the transmitter and exposed person. If the transmitter is wearing mask with efficacy ε_T and the
exposed person is wearing a mask with efficacy ε_E, then the exposure viral load V_E can be related
to the transmitter viral load V_T by: $V_E = V_T(1-\varepsilon_T)(1-\varepsilon_E)$. The combination mask efficacy is then
$\varepsilon_C=1-(1-\varepsilon_T)(1-\varepsilon_E)$, which takes on a value of zero when both parties are not wearing a mask or
wearing masks that are totally ineffective **(Fig 1).**

As in our prior model, the exposure viral load impacts contagiousness, which is the
probability that virus is passaged to the exposed person's airway, as well as infectiousness, the
probability of cellular infection given the presence of virus in the airway. Each of these

properties is associated with a dose response curve (contagiousness dose (CD) response curve and infectiousness dose (ID) response curve), the product of which is the TD response curve.

105   In this context, we first establish a baseline probability of transmission given no use of masks on both sides of a potential transmission pair **(Fig 2a)**. The absolute & relative reductions in transmission probability with more effective masks vary as V_T increases. At lower viral loads ($<10^{7.5}$ viral RNA copies), a moderate to highly effective mask worn by either a transmitter ($0.9 > \varepsilon_T \geq 0.5$) or an exposed contact ($0.9 > \varepsilon_E \geq 0.5$) is sufficient to partially lower the absolute

110   probability of transmission **(Fig 2b-c, Sup fig 1a).** The relative reduction in transmission probability increases linearly with increasing mask efficacy at 10^7 viral RNA copies with an increasingly concave, curvilinear relationship at higher viral loads **(Sup fig 1c)**.

        At higher transmitter viral load ($10^{7.5}$–10^9 viral RNA copies), a moderate to highly effective mask worn by either a transmitter ($0.9 > \varepsilon_T \geq 0.5$) or an exposed contact ($0.9 > \varepsilon_E \geq 0.5$)

115   insignificantly lowers the absolute probability of transmission **(Fig 2b-c, Sup fig 1a).** At high viral loads, the relative reduction in transmission probability increases dramatically with extremely effective masks of efficacy ≥ 0.9, when the mask is worn by either a transmitter or an exposed contact **(Sup fig 1c)**.

120   ***Predicted impact of dual masking on transmission probability within transmission pairs.*** If both the transmitter and exposed person wear masks (dual masking), then lower mask efficacies are sufficient to significantly lower transmission risk at a wider range of exposure viral loads **(Fig 2d)**. At viral loads $<10^8$ viral RNA copies, masks worn by both transmitter and exposed contact of more than moderate efficacy ($\varepsilon_T \geq 0.5$, $\varepsilon_E \geq 0.5$ resulting in $\varepsilon_C \geq 0.75$), is sufficient to

125   partially lower the absolute probability of transmission **(Fig 2d, Fig 3, Sup fig 1b).** The relative reduction in transmission probability according to mask efficacy increases more rapidly with dual **(Sup fig 1d)** compared to single **(Sup fig 1c)** masking. If both transmitter and exposed contacts wear masks with $\varepsilon_T = 0.9$ and $\varepsilon_E = 0.9$ ($\varepsilon_C = 0.99$), then transmission probability is reduced to $<5\%$ for viral loads $<10^{8.5}$ viral RNA copies and to ~20% for transmitter viral load of

130   10^9 viral RNA copies **(Fig 2d, 3d & Sup fig 1c)**.

        Predicted impact of transmitter and exposure contact masking on effective reproduction number (R_e) at different levels of implementation. We next explore the impact of general

masking adherence rates on population level metrics of infection by simulating 3000 potential

135　transmitters assuming heterogeneity in viral load trajectories and exposure contact networks among individuals. Reduction in the effective reproductive number (R_e) depends on both the mask efficacy levels (ε_T and ε_E) and the level of adherence to masking **(Fig 4)**. If we assume 25% of people wear masks 25% of the time, then most variability in results occur due to the stochastic nature of the model **(Fig 4a)**. If we assume that 50% of people wear masks 50% of the

140　time, then the use of masks with high efficacy of ~0.9 results in a drop of R_e from ~1.8 to ~1.0 **(Fig 4b)**. With 75% of people wearing masks 75% of the time, a mask efficacy of ~0.5 allows for a reduction of R_e from ~1.8 to ~1.0 **(Fig 4c)**. With 100% of people wearing masks 100% of the time, then a mask efficacy of ~0.3 is sufficient to achieve R_e ~1.0, and efficacy of 0.5 in both transmitter and exposed contacts lowers R_e to less than 0.6 **(Fig 4d)**.

145　　　　Current estimates of daily mask use over the last 2 months vary between states in the United States between 40 and 60% (http://www.healthdata.org/acting-data/maps-mask-use), and R_e has varied between 0.8 and 1.2 (https://covid19-projections.com/us). These results suggest that panel **Fig 4c** is likely the closest to current U.S. epidemic conditions and that ε_T and ε_E likely fall roughly between 0.4 and 0.6 in a real-world setting, if mask efficacy is equivalent between

150　transmitter and exposed contacts. While a combined efficacy ε_C of 0.65-0.85 can be roughly estimated from the model, the possibility of superior efficacy of masks in transmitters versus exposed, or vice versa, cannot be excluded.

Proportions of infections attributable to masked and unmasked transmission pairs. We next

155　project the proportion of transmission events attributable to different masking profiles among transmission pairs assuming equally effective mask (ε) used by both transmitters and exposed contacts. In circumstances with low mask utilization (25% of people wearing masks 25% of the time), nearly all transmissions occur from an unmasked person to an unmasked person **(Fig 5a)**. A similar trend is noted for moderate mask utilization (50%), particularly as mask efficacy

160　increases **(Fig 5b)**.

　　　　For high (75%) and extremely high (90%) mask utilization scenarios, if mask efficacy is moderate (ε~0.6) as is currently believed, then a higher proportion of transmissions occur to or from a person wearing a mask, despite the fact that the total number of transmissions dramatically decreases **(Fig 5c, d)**.

165   We project that at current epidemic conditions in the United States, including ~50% mask use (similar to 75% of people wearing masks 75% of the time) and mask efficacy of 0.3-0.6, that fewer than half of ongoing transmissions occur within unmasked pairs and that transmission between masked transmitters or masked exposed contacts likely contributes significantly to R_e **(Fig 5c).**

Predicted impact of transmitter and exposure contact masking on super-spreader events. We next identified that increased mask compliance and efficacy dramatically decreases the proportion of infected people who successfully transmit to another person **(Fig 6a-d).** If 50% of people were to wear 50% effective masks half of the time, then the likelihood of an individual

175   transmitting decreases from 30% to 20% **(Fig 6b).** If mask compliance is increased to 75% of people 75% of the time, then the likelihood of a person transmitting decreases to ~15% **(Fig 6c)**.

       When masking is applied homogeneously across the population, the proportion of infectors (transmitters who infect at least one person) who pass the infection to 5 or more people decreases, as mask utilization and efficacy increase **(Sup fig 2a-d).** Increased mask utilization

180   and increased mask efficacy leads to an even reduction of all types of transmission events, including transmissions to small numbers (1-3) of people, or super-spreader events to >5, >10, >20 or >50 people **(Fig 7a-d)**. Improvements in mask efficacy have a larger impact as utilization of mask use increases, including against super-spreader events **(Fig 7c, d)**. Under all simulations, super-spreader events with transmission to >5 people persist and make a nearly equivalent

185   contribution proportionally to overall R_e, though their absolute impact is considerably lessened with higher mask compliance and efficacy.

       Our results suggest that with current levels of masking in the United States **(Fig 7c, $\varepsilon=$ 0.3-0.6)**, most of the contribution to R_e still comes from super-spreader events involving >5 secondary infections. We therefore simulated masking applied to 100% of people with >10

190   exposure contacts per day **(Sup fig 3a-d)** and found that even modest uptake of moderately effective masks (~0.5) in the remainder of the population appeared to be sufficient to maintain R_e <1.

Predicted impact of transmitter and exposure contact masking on viral inoculum at time of

195   ***infection.*** Another theoretical benefit of masks is reduction in exposure viral load which in

animal models of SARS-CoV-1 and MERS, leads to less severe infection (*26-28*). Simulating under the assumption that 75% of people wear masks 75% of the time (i.e., a situation representing current levels of masking in the United States), we identified that viral load of a transmitter required to generate secondary infections increases slightly with higher

200   implementation of more efficacious masks, particularly with dual masking of transmitters and exposed contacts **(Fig 8a)**. Viral load exposure at the time of a successful transmission decreased according to efficacy of mask, particularly if both transmitters and exposed contacts are masked **(Fig 8b)**. With dual masking in place with efficacies of 0.6, exposure viral load decreased by ~1 log **(Fig 8b)**. With dual masking in place with efficacies of 0.9, exposure viral load decreased by

205   ~2 logs **(Fig 8b)**.

Predicted impact of antiviral therapy during early symptomatic infection on R_e. Treatment as prevention is a highly effective means for reducing person-to-person transmission of HIV (*29, 30*). Our models predict that initiation of potent antiviral therapy early after symptom onset is

210   likely to have therapeutic benefit (*18*)**.** We therefore tested whether early symptomatic therapy which rapidly eliminates shedding might also decrease secondary transmissions. Owing to the fact that symptomatic therapy would almost invariably occur after peak viral shedding **(Sup Fig 4a)**, our simulations suggest that even widespread implementation of early symptomatic therapy would not lower R_e **(Fig 9a),** percentage of infected people who transmit to at least one other

215   person **(Fig 9c)** or percentage of infectors who transmit to at least 5 other people **(Fig 9e)**.

Predicted impact of PEP on R_e. PEP is also a potential method for lowering SARS-CoV-2 transmissions. Because PEP is given in the pre-symptomatic phase and would usually fall before or near peak viral shedding **(Sup Fig 4b)**, our simulations suggest an inverse linear relationship

220   between uptake of PEP and R_e **(Fig 9b),** percentage of infected people who transmit to at least one other person **(Fig 9d)** or percentage of infectors who transmit to at least 5 other people **(Fig 9f)**. To achieve R_e <1 would require PEP efficacy of 50% and ~75% uptake in the population, which would in turn require 75% of SARS-CoV-2 cases to be contact traced. Increases in PEP efficacy beyond 0.5 would provide minimal to no enhancement of this benefit **(Fig 9b).**

225   We further determine that the timing of PEP is critical. If PEP is initiated within 2 days of exposure, then the percent of people receiving effective treatment is highly predictive of R_e **(Fig 10)**. However, from day 3 onwards, the impact of effective PEP diminishes **(Fig 10)**.

Overall, these results highlight the fact that masking is likely to have more of an impact on R_e than any form of licensed antiviral therapy that emerges during the course of the pandemic.

Discussion

Relative to other barrier methods for preventing the spread of infectious diseases such as condoms, masks are imperfect (*31*). Surgical and cloth masks, which are now used commonly by members of the public, do not completely eliminate droplet and airborne emission of viral particles by a transmitter (*1*). Nor do they prevent viral exposure to airway cells among exposed contacts. Their effectiveness in real world settings is further limited by intermittent compliance and improper masking technique. As a result, masks only prevent a proportion of person-to-person transmissions.

Nevertheless, our results demonstrate that in the absence of a licensed vaccine, based on moderate efficacy, low cost, high availability, and ease of use, masks are the most effective currently available biomedical intervention. If implemented widely and strategically, on top of baseline levels of physical distancing, masking would be sufficient to suppress ongoing spread of SARS-CoV-2 until widespread deployment of a vaccine is possible. More specifically, our model suggests that increased masking would lower the effective reproduction number (R_e), lower the percentage of infected people who transmit the virus, decrease the total number of super-spreader events, and lower the exposure viral loads among infected people, possibly leading to less severe infections overall (*26*).

Importantly, there appears to be a critical threshold of compliance. We predict massive additional benefits accrued with an increase in masking compliance from 75% of people masking 75% of the time to 90% of people masking 90% of the time. Masking also highlights the critical nature of suppressing super-spreader events. If nearly 100% compliance could be achieved among persons with 10 or more exposure contacts per day, then this would be sufficient for maintaining R_e less than one. This result highlights that policies mandating the proper use of masks at all times by all persons at sites of known super-spreader events including high risk work environments, locker rooms, weddings, social gatherings, and schools should be considered.

Slight increases in mask efficacy could also drive R_e to much lower levels. We believe that different types of masks should be comparatively tested with the same scientific rigor applied to clinical trials of small molecular agents and vaccines.

An important artifact of widespread masking is that while the total number of incident cases is expected to decrease dramatically, the proportion of transmissions in which at least one

member of the transmission pair is masked will be higher. Therefore, anecdotal documentation of successful infections between masked individuals, or even super-spreader events in which many infected people were masked, should not be misinterpreted as failure of masking policy. The counterfactual, that masks limited the severity of these events, is likely to be true. Only longitudinal incidence data, along with shifts in level of masking in a given region, are appropriate for inferring the effectiveness of masks.

It is unlikely that antiviral therapy will prove nearly as useful as widespread masking for preventing transmissions. We previously demonstrated that antiviral therapy given during the early symptomatic phase of infection has the potential to limit duration of shedding and infection associated inflammation and is likely to be more efficacious than therapy given later during COVID-19 to hospitalized patients (*18*). Unfortunately, early symptomatic therapy would likely occur after peak viral load in a majority of cases. Our simulations suggest that even 100% penetrance of extremely potent antiviral therapies would have a negligible impact on population level spread of the virus. Therefore, while treatment as a prevention is a vital piece of HIV public health policy (*30*), it is unlikely to impact the COVID-19 pandemic.

On the other hand, treatment in the earliest pre-symptomatic phase of infection, which could only realistically occur in the setting of PEP, happens prior to peak viral load and therefore could limit secondary transmissions. However, the gains from this approach diminish with each day following exposure. In order to meaningfully impact R_e, over 50% of exposed contacts would need to receive fully effective therapy within 3 days of an exposure. Given that no available agent yet achieves this level of efficacy and that identifying 50% of post-exposure contacts is unrealistic in most countries, it is clear that relative to masking, PEP will only have an adjunctive role in managing the pandemic. Potential areas of implementation are among high risk populations such as skilled nursing facility residents or cancer center patients and among populations where masking is difficult or impossible.

Our prior work strongly suggests the presence of a transmission dose response curve in which exposure viral load is a key determinant of transmission risk (*32, 33*). Our current analysis is built upon this assumption. We project that masks will lead to a lower exposure viral load among newly infected people, particularly if both the transmitter and exposed individual are successfully masked. Animal models of SARS-CoV-1 and MERS (*27, 28, 34*), as well as challenge studies with influenza H1N1 in humans (*35*), all demonstrate that lower exposure dose

is associated with less severe disease, and human data from the Hong Kong SARS-CoV-1

295   outbreak in Hong Kong suggest a similar trend (*36*). While data for SARS-CoV-2 in humans is

lacking, it is notable that age-adjusted hospitalization and death rates may be decreasing since

more widespread utilization of masks. A mask related reduction in exposure viral load is a

plausible but unproven reason for this observation.

Our work has key limitations. First, based on available data, it is impossible to know the

300   true average efficacy of a mask worn by a transmitter or an exposed contact. Many epidemics at

the state level have demonstrated a reduction in the effective reproductive number ranging from

0.2-0.5 when more widespread masking was implemented, even as physical distancing levels

waxed and waned. Our model suggests that if 50-75% of people wear masks 50-75% of the time,

which is roughly in accordance with state level observations of mask compliance, then a broad

305   estimate for real world mask efficacy is ~0.4-0.6, assuming that efficacy is equal between

transmitters and exposed contact, and that masks are properly used to optimize their efficacy. If

transmitter masking is more efficacious, while exposed contact masking is proportionally less

efficacious, then similar results can be expected. The real-world estimate is inclusive of multiple

factors including variability in mask type and masking technique. Regardless of the precise

310   estimate, it is clear that wider implementation would yield significant reductions in spread of

SARS-CoV-2 at the population level.

Second, our generalized model is not region-specific for the current pandemic. The

relative impact of super-spreader events, intensity of transmission and proportion of symptomatic

cases may vary from region to region based on contact network structure and age demographics.

315   Nevertheless, the general qualitative conclusions about masking are insensitive to these

differences and are likely to be generalizable across the globe.

Finally, our model does not include a standard SIR format and therefore does not capture

other dynamic features that might alter the force of infection such as herd immunity or time-

variant shifts in degree of physical distancing. Another missing feature that could be captured

320   with an SIR modeling framework is the possibility of an assortative mixing pattern, in which

individuals with lower adherence to masking might preferentially interact with others who have

low adherence to masking (*12*). Such an effect could allow persistence of SARS-CoV-2 within

this sub-population, even if masking is sufficient for containment in the rest of the population.

In conclusion, we developed a mechanistic model to demonstrate how masks reduce
325   exposure SARS-CoV-2 viral load and transmission probability. Widespread use of even
modestly effective masks is predicted to severely limit epidemic spread and represents the key
available intervention along with physical distancing, to mitigate the number of infections, and
perhaps the proportion of infections that are severe, while the world awaits a widely available
and effective vaccine.

Methods

SARS-CoV-2 within-host model. We used the within-host model describing the SARS-CoV-2 infection from our previous study (*18*). This model assumes that the contact of SARS-CoV-2 (V) with susceptible cells (S) produces infected cells at rate βVS which then generates new virus at a per-capita rate π. The model also incorporates the death of infected cells mediated by (1) the innate responses (δI^k) and (2) the acquired immune responses ($\frac{mE^r}{E^r+\phi^r}$) by SARS-CoV-2-specific effector cells (E). The magnitude of the innate immunity is dependent on the infected cell density and the exponent k. The nonlinearity of the acquired responses is captured by the Hill coefficient r that allows for rapid saturation of the killing. Finally, the parameter ϕ defines level of SARS-CoV-2-specific effector cells at which the killing of infected cells becomes half maximal. In the model, the rise of SARS-CoV-2-specific effector cells rise is described in a two-stage manner. The first stage defines the proliferation of the first precursor cell compartment (M_1) at rate $\omega I M_1$ and differentiation into a second precursor cell compartment (M_2) at a per capita rate q. Finally, second precursor cells differentiate into effector cells at the same per capita rate q and are cleared at rate δ_E.

The model is expressed as a system of ordinary differential equations:

$$\frac{dS}{dt} = -\beta VS$$
$$\frac{dI}{dt} = \beta VS - \delta I^k I - m\frac{E^r}{E^r+\phi^r}I$$
$$\frac{dV}{dt} = \pi I - \gamma V$$
$$\frac{dM_1}{dt} = \omega I M_1 - qM_1 \qquad\qquad (1)$$
$$\frac{dM_2}{dt} = q(M_1 - M_2)$$
$$\frac{dE}{dt} = qM_2 - \delta_E E$$

The initial conditions for the model were assumed as $S(0) = 10^7$ cells/mL, $I(0) = 1$ cells/mL, $V(0) = \frac{\pi I(0)}{c}$ copies/mL, $M_1(0) = 1$, $M_2(0) = 0$ and $E_0 = 0$. For simulations we sampled parameter values from a nonlinear mixed-effect model as described in (*19*), with the following fixed effects and standard deviation of the random effects (in parenthesis): $Log_{10}\beta$: -7.23 (0.2) virions^{-1} day^{-1}; δ: 3.13 (0.02) day^{-1} cells^{-k}; k: 0.08 (0.02); $Log10(\pi)$: 2.59 (0.05) day^{-1};

355   m: 3.21 (0.33) days^{-1}cells^{-1}; Log10(ω): -4.55 (0.01) days^{-1}cells^{-1}. We also assumed $r = 10$;

$\delta_E = 1$ day^{-1}; $q = 2.4 \times 10^{-5}$ day^{-1} and $c = 15$ day^{-1}.

Dose-response model. We employed our previously developed dose-response model to estimate

the probability of virus entering the airway given a transmitter viral load (i.e., contagiousness)

360   and the probability of cellular infection given a transmitter viral load, (i.e., infectiousness)

$P_t[V(t)]$ (response) based on viral loads $V(t)$ (dose) (*19*). The relation between the response

and the dose follows, $P_t[V(t); \lambda, \alpha] = \frac{V(t)^\alpha}{\lambda^\alpha + V(t)^\alpha}$, being λ the viral load that corresponds to 50%

infectiousness and 50% contagiousness and α the Hill coefficient that controls the sharpness in

the dose-response curve. We assumed that the viral load-dependent contagiousness (i.e., the

365   probability that virus is passaged to the exposed person's airway) is the same as infectiousness.

We estimate the transmission risk as the product of the infectiousness and contagiousness (*19*).

Transmission model and reproduction number. As in our previous model (*19*), we determined

the total exposed contacts of a transmitter within a time step (Δ_t) using a gamma distribution, i.e.

370    $\eta_{\Delta_t} \sim \Gamma\left(\frac{\theta}{\rho}, \rho\right) \Delta_t$, where θ and ρ represent the average daily contact rate and the dispersion

parameter, respectively. The true number of exposure contacts (with viral airway exposure) was

then obtained by multiplying the total exposed contacts and the contagiousness of the transmitter

(i.e., $\zeta_t = \eta_{\Delta_t} P_t$). We modelled infectiousness as a Bernoulli event with mean P_t, yielding the

number of secondary infections within a time step as $T_{\Delta_t} = Ber(P_t) P_t \eta_{\Delta_t}$. Finally, we summed

375   up the number of secondary infections over 30 days since the time of exposure to obtain the

individual reproduction number, i.e. $R_0 = \sum_{\Delta_t} T_{\Delta_t}$. For each successful transmission, we further

assumed that it takes τ days for the first infected cell to produce virus.

        In simple steps, we followed the procedure below to estimate R$_e$,

1. Simulate viral load $V(t)$ of a simulated infected individual using **the within-host**

380   **model.**

2. For a given combination of $(\lambda, \tau, \alpha, \theta, \rho)$

        a. For each time step Δ_t

                i. Compute $P_t[V(t); \lambda, \alpha]$

      ii. Draw $\eta_{\Delta_t} \sim \Gamma\left(\frac{\theta}{\rho}, \rho\right)\Delta_t$

385   **iii.** Calculate $T_{\Delta_t} = Ber(P_t)P_t\eta_{\Delta_t}$

    b. Calculate $R_e = \sum_{\Delta_t} T_{\Delta_t}$

  3. Repeat Steps 1 and 2 to estimate R_e for 3,000 infected individuals. The population level R_0 can then be calculating by taking the mean of 3,000 individual R_e values.

390   ***Parameter values for the transmission model.*** For simulations, we used the parameter set [α, λ, τ, θ, ρ] = [0.8, 10^7, 0.5, 4, 40]) as they most closely reproduces empirically observed individual R_0 and serial interval histograms as well as mean R_0 across individuals ($R_0 \in [1.4, 2.5]$) and mean serial interval across individuals (SI $\in [4.0, 4.5]$) early during the pandemic (*20-22, 24, 25*).

Modeling mask use. To evaluate the impact of the use of mask on epidemics, we first assumed that a mask decreases the exposure viral load by a fraction $(1 - \epsilon)$, being ϵ the mask efficacy or the proportion of viruses filtered by the mask of transmitter or exposed individuals. If the transmitter is wearing mask with efficacy ϵ_T and the exposed person is wearing a mask with

400 efficacy ϵ_E, then the combined mask efficacy ϵ_C, is given by $1 - (1 - \epsilon_T)(1 - \epsilon_E)$. Infectiousness or contagiousness reduction by the use of mask can be computed as:

$$\xi = 1 - \frac{P_t(\epsilon > 0)}{P_t(\epsilon = 0)} = \frac{1 - (1-\epsilon_T)^\alpha(1-\epsilon_E)^\alpha}{1 + (1-\epsilon_T)^\alpha(1-\epsilon_E)^\alpha\frac{V_t^\alpha}{\lambda^\alpha}} \qquad (2)$$

Similarly, the transmission risk reduction using mask can be computed as:

$$\xi' = 1 - \frac{P_t(\epsilon>0)P_t(\epsilon>0)}{P_t(\epsilon=0)P_t(\epsilon=0)} = \xi^2 + 2\xi\frac{(1-\epsilon_T)^\alpha(1-\epsilon_E)^\alpha(V_t^\alpha+\lambda^\alpha)}{(1-\epsilon_T)^\alpha(1-\epsilon_E)^\alpha V_t^\alpha+\lambda^\alpha}. \qquad (3)$$

405   Finally, we modeled the *compliance* of an individual wearing mask in the population as a Bernoulli event with mean σ and the *adherence* of wearing it at time step Δ_t as a Bernoulli event with mean ϑ. Compliance is defined as whether the person ever wears a mask. Adherence is the percentage of time that a mask wearer wears a mask.

410 ***Simulating secondary transmissions with mask use.*** For a specific scenario with selected σ and ϑ, we followed the procedure below to estimate the population level R_0:

  1. Simulate $V(t)$ for a transmitter using the within-host model in **eq. 1.**

2. Simulate transmitter mask compliance using $Ber(p = \sigma)$.

3. Discretize the time-space of 30 days over time steps Δ_t. For each time step,

415
    a. If the transmitter is *compliant* in using a mask:

       i. Simulate transmitter mask *adherence* at time step Δ_t using $Ber(p = \vartheta)$

       ii. If transmitter is wearing a mask at time step Δ_t

         1) Draw $\eta_{\Delta_t} \sim \Gamma\left(\frac{\theta}{\rho}, \rho\right)\Delta_t$.

         2) Determine masking adherence among exposed contacts at time step Δ_t

420
            using $\kappa \sim Ber(p = \sigma\vartheta)$. If $\kappa = 1$, then there is 100% adherence among exposed contacts and if $\kappa = 0$, then there is 0% adherence among exposed contacts.

         3) Determine the number of exposed contact wearing a mask (i.e.,

$\eta_{\Delta_{t\_Mask}} = \eta_{\Delta_t}\kappa$) and the number of exposed contact not wearing a

425
mask (i.e., $\eta_{\Delta_{t\_Unmask}} = \eta_{\Delta_t}(1 - \kappa)$.

      A. Compute $P_{t_{MasktoMask}}[(1 - \epsilon_T)(1 - \epsilon_E)V(t); \lambda, \alpha]$

      B. Calculate $T_{\Delta_{t_{MasktoMask}}} =$

$Ber\left(P_{t_{MasktoMask}}\right)P_{t_{MasktoMask}}\eta_{\Delta_{t_{Mask}}}$

      C. Compute $P_{t_{MasktoUnmask}}[(1 - \epsilon_T)V(t); \lambda, \alpha]$

430
      D. Calculate $T_{\Delta_{t_{MasktoUnmask}}} =$

$Ber\left(P_{t_{MasktoUnmask}}\right)P_{t_{MasktoUnmask}}\eta_{\Delta_{t\_Unmask}}$

       iii. If the transmitter is not adhering with the use of a mask at time step Δ_t, which is determined at step (i).

         1) Draw $\eta_{\Delta_t} \sim \Gamma\left(\frac{\theta}{\rho}, \rho\right)\Delta_t$.

435
         2) Determine masking adherence among exposed contacts at time step Δ_t using $\kappa \sim Ber(p = \sigma\vartheta)$.

         3) Determine the number of exposed contact wearing a mask (i.e.,

$\eta_{\Delta_{t\_Mask}} = \eta_{\Delta_t}\kappa$) and the number of exposed contact not wearing a

mask (i.e., $\eta_{\Delta_{t\_Unmask}} = \eta_{\Delta_t}(1 - \kappa)$

440
      A. Compute $P_{t_{UnmasktoMask}}[(1 - \epsilon_E)V(t); \lambda, \alpha]$

B. Calculate $T_{\Delta t_{UnmasktoMask}} =$

$Ber\left(P_{t_{UnmasktoMask}}\right)P_{t_{UnmasktoMask}}\eta_{\Delta t_{Mask}}$

C. Compute $P_{t_{UnmasktoUnmask}}[V(t); \lambda, \alpha]$

D. Calculate $T_{\Delta t_{UnmasktoUnmask}} =$

$Ber\left(P_{t_{UnmasktoUnmask}}\right)P_{t_{UnmasktoUnmask}}\eta_{\Delta t\_Unmask}$

    b. If the transmitter is not compliant in wearing a mask

        1) Draw $\eta_{\Delta t} \sim \Gamma\left(\frac{\theta}{\rho}, \rho\right)\Delta_t$.

        2) Determine masking adherence among exposed contacts at time step Δ_t using $\kappa \sim Ber(p = \sigma\vartheta)$.

        3) Determine the number of exposed contact wearing a mask (i.e., $\eta_{\Delta t\_Mask} = \eta_{\Delta t}\kappa$) and the number of exposed contact not wearing a mask (i.e., $\eta_{\Delta t\_Unmask} = \eta_{\Delta t}(1 - \kappa)$

        A. Compute $P_{t_{UnmasktoMask}}[(1 - \epsilon_E)V(t); \lambda, \alpha]$

        B. Calculate $T_{\Delta t_{UnmasktoMask}} =$

$Ber\left(P_{t_{UnmasktoMask}}\right)P_{t_{UnmasktoMask}}\eta_{\Delta t_{Mask}}$

        C. Compute $P_{t_{UnmasktoUnmask}}[V(t); \lambda, \alpha]$

        D. Calculate $T_{\Delta t_{UnmasktoUnmask}} =$

$Ber\left(P_{t_{UnmasktoUnmask}}\right)P_{t_{UnmasktoUnmask}}\eta_{\Delta t\_Unmask}$

    c. Calculate $T_{\Delta t} = T_{\Delta t_{UnmasktoUnmask}} + T_{\Delta t_{UnmasktoMask}} + T_{\Delta t_{MasktoUnmask}} +$ $T_{\Delta t_{MasktoMask}}$.

4. Calculate $R_e = \sum_{\Delta t} T_{\Delta t}$.

5. Repeat Steps **1** to **4** to estimate R_0 for 3,000 infected individuals (transmitters). R_e can then be calculating by taking the mean of 3,000 individual R_0 values.

Modeling antiviral treatment. We simulate the antiviral treatment by assuming that the antiviral treatment reduces the viral production (π) by $(1 - \epsilon_{treat})$, where ϵ_{treat} is the efficacy of

treatment. Here, $\epsilon_{treat} = 0$ and $\epsilon_{treat} = 1$ represent the case of completely ineffective and

470   100% effective treatment, respectively. For the transmission simulations we model *coverage* to

treatment with as a Bernoulli event with mean ψ.

In the presence of treatment with mean coverage ψ and efficacy ϵ_{treat}, we follow the

procedure below to estimate the population level R_e:

1. Determine adherence to treatment using $Ber(p = \psi)$

475   2. Determine the time of start of antiviral treatment (T_{treat}) for treatment in symptomatic phase

   and pre-symptomatic phase by randomly drawing a number from Uniform distributions

   $(U(0.5 + I_I, 5\_I_I))$ and $(U(0.5,5))$, respectively, where I_I is the incubation period of the

   infected individual.

3. Simulate viral load $V(t)$ of a simulated infected individual using the within-host model in

480   **eq. 1** with $\epsilon_{treat} = 0$ for $t \leq T_{treat}$ and $0 < \epsilon_{treat} \leq 1$ for $t > T_{treat}$.

4. For a given combination of $(\lambda, \tau, \alpha, \theta, \rho)$

   a. For each time step Δ_t

      i. Compute $P_t[V(t); \lambda, \alpha]$

      ii. Draw $\eta_{\Delta_t} \sim \Gamma\left(\frac{\theta}{\rho}, \rho\right) \Delta_t$

485   iii. Calculate $T_{\Delta_t} = Ber(P_t)P_t\eta_{\Delta_t}$

   b. Calculate $R_e = \sum_{\Delta_t} T_{\Delta_t}$

5. Repeat Steps 1 and 4 to estimate R_e for 3,000 infected individuals. The population level R_e

   can then be calculating by taking the mean of 3,000 individual R_e values.

Figure 1. Mathematical model of mask impact on SARS-CoV-2 exposure viral load a. The viral load emitted by a potential transmitter (V_t) can be filtered, resulting in lower exposure viral loads due to a single mask worn by a transmitter or exposed individual with efficacy e_T or e_E respectively. Dual masking lowers exposure viral load further by filtering virus twice. **b.** Dual masking may prevent super-spreader events to a greater extent than a masked transmitter or masked exposed individual.

Figure 2. Impact of masking of the transmitter alone, the exposed contact alone or both members of the transmission pair, on transmission risk given a single exposure contact. a-d. Each panel is based on simulations of 1000 transmission pairs. **a.** No masking, **b.** Transmitter is masked, **c.** Exposed contact is masked, **d**. Both members are masked.

Figure 3. Impact of masking of both members of a transmission pair on transmission risk according to transmitter viral load. a-d. Each panel is based on simulations of 1000 transmission pairs. **a.** Low viral load, **b.** Moderate viral load, **c.** High viral load, **d.** Extremely high viral load.

Figure 4. Effect of mask utilization and efficacy on mean SARS-CoV-2 R0. Each heat map is based on simulations of 3000 transmitters with varying daily exposure contacts. **a.** Low mask utilization, **b.** Moderate mask utilization, **c.** High mask utilization, **d.** Extremely high mask utilization.

Figure 5. Effect of mask utilization and efficacy on proportion of masked transmissions contributing to total R0. Each histogram bar in each panel is based on simulations of 3000 transmitters with varying daily exposure contacts. **a.** Low mask utilization, **b.** Moderate mask utilization, **c.** High mask utilization, **d.** Extremely high mask utilization.

Figure 6. Effect of mask utilization and efficacy on percent of infected people who transmit to others. Each heat map is based on simulations of 3000 transmitters with varying daily exposure contacts. **a.** Low mask utilization, **b.** Moderate mask utilization, **c.** High mask utilization, **d.** Extremely high mask utilization.

Figure 7. Effect of mask utilization and efficacy on proportion of super-spreader transmissions. Each histogram bar in each panel is based on simulations of 3000 transmitters with varying daily exposure contacts. **a.** Low mask utilization, **b.** Moderate mask utilization, **c.** High mask utilization, **d.** Extremely high mask utilization.

Figure 8. Effect of mask efficacy on exposure viral load during successful transmissions. Each boxplot in each panel is based on simulations of 3000 transmitters with varying daily exposure contacts. Boxplots are median and interquartile range (IQR) and lines are 1.5 the IQR. **a.** Viral load of transmitters, **b.** Viral load exposure in secondarily infected persons.

Figure 9. Projected impact of antiviral therapy on SARS-CoV-2 R0. Each line in each panel is based on simulations of 3000 transmitters with varying daily exposure contacts. Panels a, c and e are for early symptomatic therapy. Each simulation assumes 100% efficacy. Panels b, d and f are for post-exposure prophylaxis given during pre-symptomatic infection. Colored lines assume different antiviral efficacies (inset). **a, b.** Projected R0 given different amounts of uptake in the population. **c, d.** Projected percent of transmitters who infect at least one person given different amounts of uptake in the population. **e, f.** Projected percent of infectors who infect at least five people given different amounts of uptake in the population.

Figure 10. Projected impact of post-exposure prophylaxis on SARS-CoV-2 according to the day of implementation. Each line in is based on simulations of 3000 transmitters with varying daily exposure contacts and are colored by percent of transmitters receiving therapy.

Supplementary figure 1. Impact of masking of the transmitter or exposed contact alone or both members of the transmission pair, on transmission risk given an exposure viral load. Each panel is based on simulations of 1000 transmission pairs. **a-b.** Absolute reduction in transmission risk. **c-d.** Relative reduction in transmission risk. **a,c.** Transmitter is masked only **b, d.** Both transmitter and exposed contact are masked

Supplementary figure 2. Effect of mask utilization and efficacy on percent of infectors who supers-spread to others. Each heat map is based on simulations of 3000 transmitters with varying daily exposure contacts. **a.** Low mask utilization, **b.** Moderate mask utilization, **c.** High mask utilization, **d.** Extremely high mask utilization.

Supplementary figure 3. Effect of mask utilization and efficacy on mean SARS-CoV-2 R0 assuming all persons with >10 exposure contacts per day are masked. Each heat map is based on simulations of 3000 transmitters with varying daily exposure contacts. **a.** Low mask utilization, **b.** Moderate mask utilization, **c.** High mask utilization, **d.** Extremely high mask utilization.

Supplementary figure 4. Viral load simulations on antiviral therapy. a-b. Solid lines trajectories are assumed with therapy, whereas dashed lines are counterfactuals with no therapy. **a.** Simulated early symptomatic therapy. **b.** Simulated PEP.

References

1. M. Liang *et al.*, Efficacy of face mask in preventing respiratory virus transmission: A systematic review and meta-analysis. *Travel Med Infect Dis*, 101751 (2020).

2. E. P. Fischer *et al.*, Low-cost measurement of face mask efficacy for filtering expelled droplets during speech. *Science Advances* **6**, eabd3083 (2020).

3. R. Chou, T. Dana, R. Jungbauer, C. Weeks, M. S. McDonagh, Masks for Prevention of Respiratory Virus Infections, Including SARS-CoV-2, in Health Care and Community Settings: A Living Rapid Review. *Ann Intern Med*, (2020).

4. A. S. Magaret *et al.*, Effect of Condom Use on Per-act HSV-2 Transmission Risk in HIV-1, HSV-2-discordant Couples. *Clin Infect Dis* **62**, 456-461 (2016).

5. A. Konda *et al.*, Aerosol Filtration Efficiency of Common Fabrics Used in Respiratory Cloth Masks. *ACS Nano* **14**, 6339-6347 (2020).

6. E. E. Sickbert-Bennett *et al.*, Filtration Efficiency of Hospital Face Mask Alternatives Available for Use During the COVID-19 Pandemic. *JAMA Intern Med*, (2020).

7. N. H. L. Leung *et al.*, Respiratory virus shedding in exhaled breath and efficacy of face masks. *Nat Med* **26**, 676-680 (2020).

8. J. Howard *et al.*, Face Masks Against COVID-19: An Evidence Review. *Preprints* **2020040203 (doi: 10.20944/preprints202004.0203.v3).** , (2020).

9. R. Zhang, Y. Li, A. L. Zhang, Y. Wang, M. J. Molina, Identifying airborne transmission as the dominant route for the spread of COVID-19. *Proc Natl Acad Sci U S A* **117**, 14857-14863 (2020).

10. Y. Wang *et al.*, Reduction of secondary transmission of SARS-CoV-2 in households by face mask use, disinfection and social distancing: a cohort study in Beijing, China. *BMJ Glob Health* **5**, (2020).

11. W. Lyu, G. L. Wehby, Community Use Of Face Masks And COVID-19: Evidence From A Natural Experiment Of State Mandates In The US. *Health Aff (Millwood)* **39**, 1419-1425 (2020).

12. D. N. Fisman, A. L. Greer, A. R. Tuite, Bidirectional impact of imperfect mask use on reproduction number of COVID-19: A next generation matrix approach. *Infect Dis Model* **5**, 405-408 (2020).

13. C. J. Worby, H. H. Chang, Face mask use in the general population and optimal resource allocation during the COVID-19 pandemic. *Nat Commun* **11**, 4049 (2020).

525   14.   R. O. J. H. Stutt, R. Retkute, M. Bradley, C. A. Gilligan, J. Colvin, A modelling framework to assess the likely effectiveness of facemasks in combination with 'lock-down' in managing the COVID-19 pandemic. *Proc Math Phys Eng Sci* **476**, 20200376 (2020).

      15.   J. F. Chan *et al.*, Surgical mask partition reduces the risk of non-contact transmission in a
530   golden Syrian hamster model for Coronavirus Disease 2019 (COVID-19). *Clin Infect Dis*, (2020).

      16.   X. Wang, E. G. Ferro, G. Zhou, D. Hashimoto, D. L. Bhatt, Association Between Universal Masking in a Health Care System and SARS-CoV-2 Positivity Among Health Care Workers. *JAMA*, (2020).

535   17.   J. J. Bartoszko, M. A. M. Farooqi, W. Alhazzani, M. Loeb, Medical masks vs N95 respirators for preventing COVID-19 in healthcare workers: A systematic review and meta-analysis of randomized trials. *Influenza Other Respir Viruses* **14**, 365-373 (2020).

      18.   A. Goyal, E. F. Cardozo-Ojeda, J. T. Schiffer, Potency and timing of antiviral therapy as determinants of duration of SARS CoV-2 shedding and intensity of inflammatory
540   response. *medRxiv*, 2020.2004.2010.20061325 (2020).

      19.   A. Goyal, D. B. Reeves, E. F. Cardozo-Ojeda, J. T. Schiffer, B. T. Mayer, Wrong person, place and time: viral load and contact network structure predict SARS-CoV-2 transmission and super-spreading events. *medRxiv*, 2020.2008.2007.20169920 (2020).

      20.   A. Endo, S. Abbott, A. J. Kucharski, S. Funk, C. f. t. M. M. o. I. D. C.-W. Group,
545   Estimating the overdispersion in COVID-19 transmission using outbreak sizes outside China. *Wellcome Open Res* **5**, 67 (2020).

      21.   Q. Bi *et al.*, Epidemiology and transmission of COVID-19 in 391 cases and 1286 of their close contacts in Shenzhen, China: a retrospective cohort study. *Lancet Infect Dis* **20**, 911-919 (2020).

550   22.   Y. Zhang, Y. Li, L. Wang, M. Li, X. Zhou, Evaluating Transmission Heterogeneity and Super-Spreading Event of COVID-19 in a Metropolis of China. *Int J Environ Res Public Health* **17**, (2020).

      23.   D. Miller *et al.*, Full genome viral sequences inform patterns of SARS-CoV-2 spread into and within Israel. *medRxiv*, 2020.2005.2021.20104521 (2020).

555   24.   D. Adam *et al.*, Clustering and superspreading potential of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infections in Hong Kong. *Europe PMC* **10.21203/rs.3.rs-29548/v1**, (2020).

      25.   Z. Du *et al.*, Serial Interval of COVID-19 among Publicly Reported Confirmed Cases. *Emerg Infect Dis* **26**, (2020).

560   26.   M. Gandhi, C. Beyrer, E. Goosby, Masks Do More Than Protect Others During COVID-19: Reducing the Inoculum of SARS-CoV-2 to Protect the Wearer. *J Gen Intern Med*, (2020).

27.   T. Watanabe, T. A. Bartrand, M. H. Weir, T. Omura, C. N. Haas, Development of a dose-response model for SARS coronavirus. *Risk Anal* **30**, 1129-1138 (2010).

565   28.   A. Totura *et al.*, Small Particle Aerosol Exposure of African Green Monkeys to MERS-CoV as a Model for Highly Pathogenic Coronavirus Infection. *Emerg Infect Dis* **26**, (2020).

29.   M. Das *et al.*, Decreases in community viral load are accompanied by reductions in new HIV infections in San Francisco. *PLoS One* **5**, e11068 (2010).

570   30.   M. S. Cohen *et al.*, Antiretroviral Therapy for the Prevention of HIV-1 Transmission. *N Engl J Med* **375**, 830-839 (2016).

31.   S. Weller, K. Davis, Condom effectiveness in reducing heterosexual HIV transmission. *Cochrane Database Syst Rev*, CD003255 (2002).

32.   B. T. Mayer *et al.*, Estimating the Risk of Human Herpesvirus 6 and Cytomegalovirus
575   Transmission to Ugandan Infants from Viral Shedding in Saliva by Household Contacts. *Viruses* **12**, (2020).

33.   I. Boucoiran *et al.*, Nonprimary Maternal Cytomegalovirus Infection After Viral Shedding in Infants. *Pediatr Infect Dis J* **37**, 627-631 (2018).

34.   L. M. Gretebeck, K. Subbarao, Animal models for SARS and MERS coronaviruses. *Curr*
580   *Opin Virol* **13**, 123-129 (2015).

35.   M. J. Memoli *et al.*, Validation of the wild-type influenza A human challenge model H1N1pdMIST: an A(H1N1)pdm09 dose-finding investigational new drug study. *Clin Infect Dis* **60**, 693-702 (2015).

36.   C. M. Chu *et al.*, Viral load distribution in SARS outbreak. *Emerg Infect Dis* **11**, 1882-
585   1886 (2005).