Screening electrocardiogram in young athletes and military members: a systematic review and meta-analysis

Aaron Lear MD, MSc; Cleveland Clinic Akron General, Akron, OH USA
Niraj Patel, DO HeyDoctor by GoodRx, San Francisco, CA USA
Chanda Mullen, PhD; Cleveland Clinic Akron General, Akron, OH USA
Marian Simonson, MS; Cleveland Clinic Foundation, Cleveland, OH USA
Vince Leone, MD; Northeast Ohio Medical University (NEOMED), Akron, OH USA
Constantinos Koshiaris, PhD; University of Oxford, Oxford, UK
David Nunan, PhD; University of Oxford, Oxford, UK

Word Count: 2985

Contact author:
Aaron Lear
1 Akron General Way
Ctr for Fam Medicine
Akron, OH 44307 USA
ORCID iD: 0000-0002-1353-7785
leara@ccf.org
office: 330.344.6047 (Angela Watkins assistant)

Abstract

Objective: To determine the effect of electrocardiogram (ECG) screening on prevention of sudden cardiac arrest and death (SCA/D) in young athletes and military members.

Design: Systematic review and meta-analysis

Data Sources: MEDLINE, Embase, Cochrane CENTRAL, Web of Science, BIOSIS, Scopus, SPORT discus, PEDro, were searched from inception to date searched between 2/21/19 and 3/1/19, and Clinicaltrials.gov on 7/29/19. All databases were searched without limits on language.

Eligibility criteria for selected studies: Randomized and non-randomized controlled trials, where pre-participation examination including ECG was the primary intervention used to screen athletes or military 40 years of age or younger. Accepted controls were no screening, usual care, or pre-participation examination without ECG

Results: 4 studies (11,689,172 participants) were included, all at high risk of bias. Pooled data from two studies (n= 3,869,274; very low quality) showed a 42% relative decrease in sudden cardiac death, equating to an absolute risk reduction of .0016%. Uncertainty was high, with a potential 67% relative decrease to a 45% relative increase in those screened with ECG based on 95% confidence intervals (RR 0.58; 95%CI 0.23, 1.45). Heterogeneity was found to be high as measured with I^2 statistic (71%).

Conclusion: There is very low quality evidence ECG screening decreases risk of sudden cardiac death in young athletes and military members. Decisions need to consider evidence that ECG screening could also increase risk of sudden cardiac death based on the findings of meta-analysis.

PROSPERO Registration: CRD42019125560

Key Words: athlete, military, electrocardiogram screening, sudden cardiac arrest, sudden cardiac death

Introduction

Efforts to reduce the incidence of sudden cardiac arrest and death (SCA/D) in young athletes

has led the European Society of Cardiology (1) to recommend electrocardiogram (ECG)

screening as part of a pre-participation examination (PPE) of young competitive athletes prior

to participation in 2005. Professional bodies around the world have followed this

recommendation with statements of their own. With some in agreement including the

International Olympic Committee (2); some agreeing but with limitations, such as the

Australasian Society for Sports Physicians (3); while organizations in the United States have

resisted calls for blanket screening (4,5). The evidence-base to support inclusion of ECG

screening for reducing incidence of SCA and SCD in young athletes has not undergone

systematic review. A previous systematic review assessed the effectiveness of ECG screening to

detect potentially lethal cardiac disorders, but did not address the impact on SCA, SCD and the

potential negative effects of ECG screening (6).

The provision of a systematic summary of existing data on the outcomes of ECG screening will

provide both the public, health care practitioners, and policy makers with vital information

about the health effects of ECG screening in these populations when compared with history

and physical examination alone.

The aim of this study was to review all available evidence assessing the effect of the addition of

ECG screening as part of PPE in young athletic, and military populations on incidence of SCA/D,

and to synthesize available research to evaluate the effect of the addition of ECG on the occurrence of SCA/D.

Methods

This review is part of a project with two objectives: identifying the global incidence of SCA/D in athletes and military members, and evaluating the effect of screening ECG on SCA/D in the same population. It was performed according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)(7) guidelines, and was registered at PROSPERO March 18th 2019 under CRD42019125560 (8).

Data Sources and Searches

The search strategy was designed in conjunction with a medical librarian experienced in systematic reviews (MS), and the search strategy used combined the dual objectives into a single search. The search strategy is included in the supplementary appendix. We searched MEDLINE, Embase, Cochrane CENTRAL, Web of Science, BIOSIS, Scopus, SPORT discus, PEDro, between 2/22/19 and 3/1/19 and Clinicaltrials.gov on 7/29/19. Review articles and position statements were reviewed for eligible articles (9–11).   There was no limitation on language or date of publication.

Study Selection

Studies eligible for inclusion were randomized and non-randomized controlled trials, where pre-participation examination including ECG was the primary intervention used to screen

athletes or military 40 years of age or younger. Accepted controls were no screening, usual

care, or pre-participation examination without ECG. Age 40 was selected due to the increased

incidence of coronary artery disease as cause of SCA/D with increasing age, and the desire to

focus on etiologies other than coronary artery disease (10,12).

The pre-specified primary outcome was the difference in SCA/D in athletes and military

populations screened with ECG compared to control groups not screened with ECG. Secondary

outcomes to assess the impact of ECG screening included the rate of screened participants

removed from sport/activity for cardiac conditions; rate of participants with abnormal cardiac

findings and treated as a result of screening; the proportion of those removed from sport for

cardiac conditions, treated and returned to sport. Pre-planned subgroup analysis were

considered based on age, gender, race, and by type and level (i.e. elite, scholastic, recreational)

of sport played or military role.

Data Extraction and Quality Assessment

Independent dual-investigator article screening, selection, risk of bias (ROB) assessments, and

extraction was performed with Covidence (Covidence.org, Veritas Health Innovation,

Melbourne, Australia 2018). The primary author (AL) screened all titles, and the second

reviewer was from a team of three (CM, NP, VL). Google Translate (translate.google.com,

Google LLC, Mountain View, CA) was used for articles in foreign languages when technically

possible. When not technically possible, physician translators were used. The Cochrane risk of

bias tool (13), native to Covidence was used for ROB assessment.   Disagreements were resolved by discussion with primary author (AL), and the second reviewer.

Data Synthesis and Analysis

Meta-analysis was performed with the statistical package native to Review Manager (Cochrane Collaboration, London, UK) using the random-effects Mantel-Haenszel method based on the clinical heterogeneity within the included studies(14). Data are presented as relative risk with 95% confidence interval in those screened with ECG compared with those not screened with ECG where possible. Heterogeneity is reported with summary statistics I^2 and Chi^2, with pre-specified values of <30% considered low; 30-70% considered moderate; and >70% considered high. A p-value of 0.10 or lower for Chi^2 statistic was indicated statistical heterogeneity. Sensitivity analysis and publication bias assessment (e.g. funnel plot asymmetry) were planned but not performed due the small number of included studies.

Role of Funding Source

No funding was received for the completion of this project.

Results

After removal of 10,780 duplicates, and addition of a further 11 titles after hand search, 20,059 titles and abstracts were screened from database searches, and additions from review of recent review articles. Full text screening was carried out on 322 articles, four of these met criteria for inclusion (Figure 1).

[INSERT FIGURE ONE HERE]

Included studies

All included studies were non-randomized controlled studies. Three were on athletes(15–17)

and included a total of 6,431,380 subjects, and 1 was in military males only(18) including

5,257,792 subjects. Three studies included a historical control group, comparing the rate of

SCD in a control cohort prior to the implementation of ECG screening, to an intervention cohort

after ECG screening began (15,17,18). No studies were identified which reported an outcome

of SCA. Two studies were included in meta-analysis, both on athletes , and in both it was

unclear if the historical control group received a screening PPE, or no examination(15,16). In

the two studies not included in meta-analysis, one including only military males(18), and one in

athletes(16) compared the rate of SCD in a cohort who had been screened with PPE including

ECG, with the rate in a cohort that received a PPE without ECG. Full descriptions and

characteristics of these four studies are presented in the supplementary appendix.

We found no studies reporting data on our secondary objectives. Corrado, et. al., (2006) do

report observational data on a subgroup of their intervention cohort which received further

testing after initially abnormal ECG screening. Subgrouping by level of athlete, age, sex, race

was not possible based on the reported data in any of the included studies.

All included studies were found to have either unknown, or high risk of bias in the large majority of categories evaluated (Table 1). No included studies reported funding which was determined to increase their risk of bias.   Further details on the risk of bias determination is included in the characteristics of included studies table in the supplementary appendix.

Included Studies	Sequence generation	Allocation concealment	Subject/participant blinding	Blinding outcome assessors	Incomplete outcome data	Selective outcome reporting	Other risk of bias
Corrado 2006	red	yellow	yellow	red	green	red	red
Maron, Haas 2009a	red	yellow	yellow	red	green	red	red
Steinvil 2011	red	yellow	yellow	red	red	green	red
Abacherli 2014	red	yellow	yellow	yellow	yellow	yellow	yellow

Table 1: Risk of bias in included non-randomized trials in objective 2; based on Cochrane risk of bias tool
● Signifies low risk of bias; ●signifies unclear risk of bias; ●signifies high risk of bias

Effectiveness of ECG screening

Corrado, et. al. (2006) reports an analysis on SCD in athletes in the Veneto region of Italy. The authors report a primary outcome of an 89% decrease in incidence of SCD after the implementation of ECG screening program for athletes in 1982. This is done by comparing a two-year time period prior to the initiation of screening (1979-80), to the final two years of the screening period (2003-04). In our analysis, data from 23 years of athletes screened with ECG compared with the three years of athletes not screened with ECG showed a 63% decreased risk in the ECG screened group (RR 0.37; 95%CI 0.20 to 0.69).

Steinvil, et. al. (2011) reports on SCD events in athletes reported in two Israeli newspapers covering 90% of the country, before and after the initiation of a cardiac screening program. The authors showed a 5% decrease in risk in those athletes undergoing cardiac screening with ECG and stress test, to those which did not receive this screening (RR: 0.95; 95%CI 0.43 to 2.13). The findings were highly uncertain, with a potential 57% decrease in risk of SCD to 113% increased risk.

A conference abstract by Abacherli, et. al. (2014) details a comparison of SCD in Swiss male military conscripts separated into age groups 16-19, 20-14, 25-29. The authors compare episodes of SCD after the initiation of ECG screening, compared to historical controls prior to ECG screening. A statistically significant reduction in the ECG screened 20-24 age group with a point estimate of 0.56 (CI95% 0.35 to 0.91) was reported. The same comparison in men aged 16-19 was found to be 0.89, and 25-29 was found to 1.04. These were described as non-significant, with only the point estimates, and no confidence intervals reported. The abstract is unclear as to whether the statistical method used was relative risk or odds ratio, making interpretation of the findings difficult. No extractable data were present. We were unable to find a published article based on the abstract by Abacherli, et. al. and attempts to obtain additional data from the lead author were unsuccessful.

Maron, et. al. (2009), in comparing a cohort of U.S. athletes in the state of Minnesota who have undergone PPE without ECG screening, with a proportion of ECG screened athletes from the

Corrado, et. al. (2006) Italian cohort over a similar time period, report a 6% decrease in risk of

SCD but estimates included considerable benefit and harm (RR 0.94; 95%CI 0.41, 2.12). To

avoid double counting the Corrado, et. al. (2006) intervention group the Maron, et. al. (2009)

article was not included in our meta-analysis.

Two studies (15,17) including athlete participants were included in meta-analysis (Figure 2).

The results show a relative decrease risk of 42% for SCD in athletes screened by ECG but

uncertainty was high, with a potential 77% relative decrease to a 45% relative increase in those

screened with ECG (RR 0.58, 95%CI 0.23 to 1.45; I^2 = 71%; Chi^2 3.41, p=0.06).

[INSERT FIGURE TWO HERE]

Corrado, et. al. (2006) reported data beyond the outcome of SCD, and this was done only in a

portion of the intervention cohort, not allowing comparison with the control group. The

authors report that 9% (3,914 of 42,386) of athletes were referred for further testing after their

initial ECG screening and a further 2 % (879 of 42,386) were ultimately removed from sport.

The authors did not describe the treatment rendered to athletes with positive screening ECG

beyond the further diagnostic studies used and did not report on athletes returning to sport

after treatment.

Quality of evidence

For the primary outcome of SCD we judged the evidence to be of very low certainty due mainly

to high risk of bias, serious inconsistency, and serious imprecision.

GRADE Summary of Findings Table

Does screening ECG in athletes/military members prevent sudden cardiac arrest and death?

Population: Competitive athletes participating in an organized sport, and active duty military members age 40 or younger.
Intervention: Performance of pre-participation examination (PPE) with screening electrocardiogram (ECG) with or without echocardiogram to find conditions which are known to lead to sudden cardiac arrest associated with sports/athletic activity.
Comparison: No PPE, or PPE without screening ECG
Outcome: 1) SCA/D in screened vs. those not screened with ECG 2) Rate of athletes/military members removed from sport/activity 3) Rate of athletes/military members with abnormal findings, who underwent treatment as a result of screening 4) The number of treated athletes/military members who returned to sport/activity

Outcome	Intervention	Number of Studies	Absolute Risk No ECG or Exam with no ECG	Absolute Risk ECG screened	Relative Effect (95% CI)	Number of Participants	Quality of the Evidence*
Sudden cardiac death in athletes	ECG in 1 study, ECG + exercise stress test in 1 study	2 studies evaluating prevention of sudden cardiac death	.00328%	.00171%	0.58 (0.23, 1.45)	3,869,274 (2 historically controlled trials)	Very Low

*Summary of decisions on quality of the evidence:
Study design: Non-randomized historically controlled trials at high risk of bias
Risk of bias: Very serious concern. Downgraded for significant risk of bias in these historically controlled trials, including in categories of randomization, allocation, blinding and sequence generation, as well as risk of significant confounding due to the historical nature of the studies.
Inconsistency: Serious concern. Based on the differences in the estimate of effect and the high heterogeneity with I^2 of 71%, and significant Chi^2 p value of 0.06.
Indirectness: No concern. The studies included evaluate the event of interest, in the population of interest.
Imprecision: Serious concern. Downgraded for wide confidence intervals despite meeting the optimal information size. With the calculated event rates in these combined studies, a power analysis with p=0.05, and Beta=0.80, a total of 3,177,808 [1,548,556 per group] is suggested for a randomized trial.
Publication Bias: not evaluated due to small number of studies.
Other factors: There is likely significant confounding which has occurred in these trials. The confounding present is likely to have increased the effect estimate.

Discussion

Our systematic, pre-registered, comprehensive, and up-to-date review found very low-quality

evidence that ECG screening decreases risk of sudden cardiac death in young athletes and

military members. However, caution is needed when considering this finding. Firstly, we were able to perform meta-analysis only with two studies. The absolute risk reduction from pooling these studies was 0.00157%. Using a single year assessment(19), this would result in a number needed to screen to prevent one death in one year of 63,694. Secondly, only one of the four studies included reached statistical significance when evaluated the effect of ECG screening on SCD, and the remaining studies report confidence intervals which include both considerable decreased and increased risk with ECG screening. Thirdly, the findings of one study are based on unpublished data(18). And fourth, due to high risk of bias, high heterogeneity and poor precision of effect estimates, the overall certainty of evidence on the effectiveness of ECG screening was judged to be very low. Taken together, we have very low confidence that our findings would not change substantially with further high-quality research.

The existing evidence base to support the use of ECG screening to prevent SCD in athletes is largely confined to the data presented in the Corrado, et. al. (2006) article included in the review. There are significant methodologic concerns about this article, including the inherent bias, and likely confounding present, when comparing a small historical control group to a much larger intervention group some 20 years later. There have also been concerns raised about the transparency of the data reported, and further follow up data on the Italian screening program(20). While there have been no recent controlled studies published, two recently published cohort studies may call into question the ability of screening athletes to prevent SCA/D. Both report on cohorts of mostly male, professional soccer players who underwent ECG screening as part of pre-participation examinations(21,22). Both reported results with

relatively high rates of SCA/D at 6.8, and 63 per 100,000 athlete years compared to accepted

estimates of SCA/D in athletic populations. When considering the ability of ECG screening to

prevent episodes of SCA/D, it is notable that published data on events of SCA/D suggest that

approximately 60% of cardiac conditions which cause SCA/D in athletes may be identifiable

with ECG screening(23,24).

While there remains disagreement, and a general lack of empiric data to support the use of ECG

to prevent SCA/D, multiple authors have advocated it's use as an effective addition to the pre-

participation examination due to the ability to better identify conditions putting athletes at risk

of SCA/D(25). A systematic review in 2015(6) compared the likelihood of history (hx), physical

exam (PE), and ECG to identify potentially lethal cardiac disorders reporting the superiority of

ECG in sensitivity (ECG=94%; hx 20%; PE 9%), positive likelihood ratio (ECG=14.8; hx=3.22;

PE=2.93), and false positive rate (ECG=6%; hx=8%; PE=10%). Results in a recent cohort study

focusing on collegiate athletes in the U.S. comparing the results history and physical exam to

additional ECG screening in the same cohort of patients(26) show false positive rates of 33.3%

for history alone, 2% with physical exam alone, and 3.4% with ECG alone. Sensitivity with ECG

was reported as 100% compared to 15.4% for history, and 7.7% for physical exam.

As demonstrated in this review, there remains little trial data to compare information on how

ECG screening affects athlete's removal from sport, follow up treatment, and potential return

to sport. There is cohort data on some of these outcomes, the most notable again being from

the included Corrado, et. al. (2006) article which reports on a subgroup of the screened athletes

included, with 9% found to be abnormal and receiving further testing, ultimately resulting in 2%

excluded from sport. A 2014 scientific statement from the American Heart Association(27)

details a selection of published articles on ECG screening cohort studies, with no control groups

and community ECG screening projects in athletes 12-25 years old. Articles reviewed report

abnormal ECG rates from 2.5-25%, further testing of 2.5-24%, and disqualification from sport

rates 0f 0.2-2% of those screening. More modern cohort studies referenced above (26,28)

comparing history and physical exam to additional ECG screening in the same cohort of patients

have shown lower levels of initially positive results in ECG screened athletes to those

completing the American Heart Association(29) history questionnaire, and physical exam alone.

It is difficult to compare how these findings affected further testing and treatment, as all

subjects received history, physical, and ECG screening interventions. Drezner, et. al. (2016)

report the identification of 0.25% of athletes with a serious cardiac condition after full

evaluation in collegiate athletes in the United States (26). Drezner, et. al. (2016) also report an

average of 2.6 days out of sport to evaluate those with ECG abnormalities on screening. It is

also notable, that within studies using ECG screening and reporting the cardiac abnormalities

identified, that Wolff-Parkinson-White syndrome is often far and away the most frequent

finding, often making up the majority of identifiable cardiac conditions considered serious

(26,28,30), which is of questionable significance when attempting to prevent SCA/D(31).

Over the past decade, refinement of the ECG criteria for diagnosing these potentially lethal

cardiac conditions has continued to improve the sensitivity, and decrease the false positive

rates(32). While these advancements in the diagnostic capability of ECG screening have

occurred, there have been no controlled trials published on independent cohorts of patients comparing the ability of ECG to PPE with history and physical alone to prevent SCA/D. There is great need for a prospective study which tests the utility of screening ECG to prevent SCA/D in athletes. Carrying out a prospective study on this topic would be a daunting task, and with the rarity of the condition may not be possible. To undertake such a project, one could consider randomizing clusters of high school and collegiate athletes to ECG screening with PPE compared to PPE. A model such as this may be even more feasible in the military, where large numbers of recruits enter into service every year. It would be possible to randomize subjects in this setting, which could lead to the data needed. Short of these two trials, it should be possible to compare the rates of SCA/D in a cohort study in the U.S. Many universities have transitioned to testing their athletes, and comparing the rates of SCA/D in these universities, to comparable universities who do not screen, would be technically feasible.

We believe the strength of this review lies in the breadth of the search for controlled trials of any kind which report on the ability of screening ECG in athletes or military members to prevent SCA/D. The primary limitation of our review is the low quality of evidence provided by the included studies leading to uncertainty for decision making. The limitations lie in both the paucity of, and the poor quality of the identified research reporting outcomes on SCA/D in our population.

Conclusion

There is very low-quality evidence ECG screening decreases risk of sudden cardiac death in young athletes and military members. Decisions regarding the use of screening ECG need to consider evidence here that ECG screening could also increase risk of sudden cardiac death. We have very low confidence that our findings would not change substantially with further high-quality research.

Acknowledgements: This work was previously presented in abstract form at the AMSSM annual

meeting 2020, and as a master's dissertation at Oxford University.

Competing Interests: No authors report competing interests with this work

Funding: No funding was received for this work

What is already known:
- Multiple cardiology and sports medicine professional organizations have made differing recommendations on the use of the electrocardiogram (ECG) in screening in young athletes.
- ECG screening of athletes has been shown to be more effective than history and physical examination alone to diagnose conditions which put the athlete at risk for sudden cardiac arrest or death (SCA/D).
- Few data are available to answer the question of the effectiveness of ECG screening in preventing SCA/D in young athletes.

What are the new findings:
- We identified only four published accounts (3 full papers and one conference abstract) of non-randomized trials reporting on the effectiveness of ECG screening to prevent SCA/D in young athletes and military members.
- The quality of the published evidence is judged to be of very low quality to answer the question of whether ECG screening prevents episodes of SCA/D.
- No difference was identified between screened and non-screened athletes in data synthesis of two of the published articles eligible for meta-analysis (RR 0.58; 95%CI 0.23, 1.45).

References

1. Corrado D, Pelliccia A, Bjørnstad HH, Vanhees L, Biffi A, Borjesson M, et al. Cardiovascular pre-participation screening of young competitive athletes for prevention of sudden death: Proposal for a common European protocol - Consensus Statement of the Study Group of Sport Cardiology of the Working Group of Cardiac Rehabilitation an. Eur Heart J. 2005;26(5):516–24.

2. Ingersoll CD. The periodic health evaluation of elite athletes: A consensus statement from the international olympic committee. J Athl Train. 2009;44(5):453.

3. Exeter D, Kuah D, Carbon R, Shawdon A, Bolzonello D. Australasian College of Sport and Exercise Physicians (ACSEP) Position Statement on Pre- Participation Cardiac Evaluation in Young Athletes [Internet]. 2018. p. 1–4. Available from: https://www.acsep.org.au

4. Hainline B, Drezner J, Baggish A, Harmon KG, Emery MS, Myerburg RJ, et al. Interassociation consensus statement on cardiovascular care of college student-athletes. J Athl Train. 2016;51(4):344–57.

5. Maron BJ, Douglas PS, Graham TP, Nishimura RA, Thompson PD. Task Force 1: Preparticipation screening and diagnosis of cardiovascular disease in athletes. J Am Coll Cardiol. 2005;45(8):1322–6.

6. Harmon KG, Zigman M, Drezner JA. The effectiveness of screening history, physical exam, and ECG to detect potentially lethal cardiac disorders in athletes: A systematic review/meta-analysis. J Electrocardiol [Internet]. 2015;48(3):329–38. Available from: http://dx.doi.org/10.1016/j.jelectrocard.2015.02.001

7.	PRISMA [Internet]. 2009 [cited 2019 Jul 16]. p. Flow Diagram. Available from: http://prisma-statement.org

8.	Lear A. A systematic review to determine the rate of sudden cardiac arrest and death in athletes, and to determine the benefits and harms of the electrocardiogram screening of athletes [Internet]. PROSPERO. 2019. Available from: http://www.crd.york.ac.uk/PROSPERO/display_record.php?ID=CRD42019125560

9.	Drezner JA, O'Connor FG, Harmon KG, Fields KB, Asplund CA, Asif IM, et al. AMSSM Position Statement on Cardiovascular Preparticipation Screening in Athletes: Current evidence, knowledge gaps, recommendations and future directions.[Erratum appears in Br J Sports Med. 2018 Mar 6;:; PMID: 29510974]. Br J Sports Med. 2017;51(3):153–67.

10.	Harmon KG, Drezner JA, Wilson MG, Sharma S. Incidence of sudden cardiac death in athletes: A state-of-the-art review. Br J Sports Med. 2014;48(15):1185–92.

11.	Mohananey D, Masri A, Desai RM, Dalal S, Phelan D, Kanj M, et al. Global Incidence of Sports-Related Sudden Cardiac Death. J Am Coll Cardiol. 2017;69(21):2672–3.

12.	Drezner JA, Sharma S, Baggish A, Papadakis M, Wilson MG, Prutkin JM, et al. International criteria for electrocardiographic interpretation in athletes: Consensus statement. Br J Sports Med. 2017;51(9):704–31.

13.	Higgins JP, Green S, editors. Cochrane Handbook for Systematic Reviews of Interventions [Internet]. 5.1.0. Cochrane Collaboration; 2011. Available from: www.handbook.cochrane.org

14.	Cochran Collaboration. Review Manager (RevMan). Copenhagen: The Nordic Cochrane Centre, The Cochrane Collaboration; 2014.

15.   Steinvil A, Chundadze T, Zeltser D, Rogowski O, Halkin A, Galily Y, et al. Mandatory electrocardiographic screening of athletes to reduce their risk for sudden death proven fact or wishful thinking? J Am Coll Cardiol. 2011;57(11):1291–6.

16.   Maron BJ, Haas TS, Doerer JJ, Thompson PD, Hodges JS. Comparison of U.S. and Italian experiences with sudden cardiac deaths in young competitive athletes and implications for preparticipation screening strategies. Am J Cardiol. 2009;104(2):276–80.

17.   Corrado D, Basso C, Pavei A, Michieli P, Schiavon M, Thiene G. Trends in sudden cardiovascular death in young competitive athletes after implementation of a preparticipation screening program. JAMA. 2006;296(13):1593–601.

18.   Abacherli R, Schmid R, Kobza R, Franz F, Schmid J, Erne P. Pre-paricitation ECG screening preventing SCD-insight from the Swiss Army. Hear Rhythm. 2014;11(5):S224.

19.   Rembold C. Number needed to screen: development of a statistic for disease screening. BMJ. 1998;317:307–12.

20.   Van Brabandt H, Desomer A, Gerkens S, Neyt M. Harms and benefits of screening young people to prevent sudden cardiac death. BMJ. 2016;353(April):1–5.

21.   Malhotra A, Dhutia H, Finocchiaro G, Gati S, Beasley I, Clift P, et al. Outcomes of Cardiac Screening in Adolescent Soccer Players. N Engl J Med. 2018;379(6):524–34.

22.   Berge HM, Andersen TE, Bahr R. Cardiovascular incidents in male professional football players with negative preparticipation cardiac screening results: an 8-year follow-up. Br J Sports Med [Internet]. 2018 Oct 1;bjsports-2018-099845. Available from: http://bjsm.bmj.com/content/early/2018/09/30/bjsports-2018-099845.abstract

23.   Harmon KG, Asif IM, Maleszewski JJ, Owens DS, Prutkin JM, Salerno JC, et al. Incidence

and Etiology of Sudden Cardiac Arrest and Death in High School Athletes in the United

States. Mayo Clin Proc [Internet]. 2016;91(11 PG-1493–1502):1493–502. Available from:

http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:27692971&id=doi:1

0.1016%2Fj.mayocp.2016.07.021&issn=0025-

6196&isbn=&volume=91&issue=11&spage=1493&date=2016&title=Mayo+Clinic+Procee

dings&atitle=Incidence+and+Etiology+of+Sudden+Cardiac+Arrest

24.   Maron BJ, Haas TS, Murphy CJ, Ahluwalia A, Rutten-Ramos S. Incidence and causes of

sudden death in U.S. college athletes. J Am Coll Cardiol. 2014;63(16):1636–43.

25.   Drezner JA, Harmon KG, Asif IM, Marek JC. Why cardiovascular screening in young

athletes can save lives: a critical review. Br J Sports Med [Internet]. 2016;bjsports-2016-

096606. Available from: http://bjsm.bmj.com/lookup/doi/10.1136/bjsports-2016-096606

26.   Drezner JA, Owens DS, Prutkin JM, Salerno JC, Harmon KG, Prosise S, et al.

Electrocardiographic Screening in National Collegiate Athletic Association Athletes. Am J

Cardiol [Internet]. 2016;118(5):754–9. Available from:

http://dx.doi.org/10.1016/j.amjcard.2016.06.004

27.   Maron BJ, Friedman RA, Kligfield P, Levine BD, Viskin S, Chaitman BR, et al. Assessment of

the 12-lead electrocardiogram as a screening test for detection of cardiovascular disease

in healthy general populations of young people (12-25 years of age): A scientific

statement from the american heart association and the american colleg. J Am Coll

Cardiol. 2014;64(14):1479–514.

28.   Williams EA, Pelto HF, Toresdahl BG, Prutkin JM, Owens DS, Salerno JC, et al.

Performance of the American Heart Association (AHA) 14-Point Evaluation Versus

Electrocardiography for the Cardiovascular Screening of High School Athletes: A Prospective Study. J Am Heart Assoc. 2019;8(14):1–9.

29. Maron BJ, Thompson PD, Ackerman MJ, Balady G, Berger S, Cohen D, et al. Recommendations and considerations related to preparticipation screening for cardiovascular abnormalities in competitive athletes: 2007 update: a scientific statement from the American Heart Association Council on Nutrition, Physical Activity, and Metabol. Circulation. 2007;115(12):1455–643.

30. Fudge J, Harmon KG, Owens DS, Prutkin JM, Salerno JC, Asif IM, et al. Cardiovascular screening in adolescents and young adults: A prospective study comparing the Pre-participation Physical Evaluation Monograph 4th Edition and ECG. Br J Sports Med. 2014;48(15):1172–8.

31. Maron BJ, Thompson PD, Maron MS. There is No Reason to Adopt ECGs and Abandon American Heart Association/American College of Cardiology History and Physical Screening for Detection of Cardiovascular Disease in the Young. J Am Heart Assoc. 2019;8(14):1–4.

32. Drezner JA, Sharma S, Baggish A, Papadakis M, Wilson MG, Prutkin JM, et al. International criteria for electrocardiographic interpretation in athletes: Consensus statement. Br J Sports Med. 2017;51(9):704–31.

Figures

PRISMA Flow Diagram

Figure 1: Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) flow sheet

Study or Subgroup	ECG Screen Events	ECG Screen Total	No ECG Screen Events	No ECG Screen Total	Weight	Risk Ratio M–H, Random, 95% CI
Corrado 2006	41	2613353	14	334129	54.0%	0.37 [0.20, 0.69]
Steinvil 2011	11	433071	13	488721	46.0%	0.95 [0.43, 2.13]
Total (95% CI)		3046424		822850	100.0%	**0.58 [0.23, 1.45]**
Total events	52		27			

Heterogeneity: Tau² = 0.32; Chi² = 3.41, df = 1 (P = 0.06); I² = 71%
Test for overall effect: Z = 1.17 (P = 0.24)

Figure 2: Meta-analysis of studies reporting data on outcomes of sudden cardiac death in athletes screened with ECG (experimental), compared to athletes not screened with ECG (control)