

Global incidence of sudden cardiac arrest in young athletes and military members: a systematic review and meta-analysis

Aaron Lear MD, MSc; Cleveland Clinic Akron General, Akron, OH USA
Niraj Patel, DO HeyDoctor by GoodRx, San Francisco, CA USA
Chanda Mullen, PhD; Cleveland Clinic Akron General, Akron, OH USA
Marian Simonson, MS; Cleveland Clinic Foundation, Cleveland, OH USA
Vince Leone, MD; Northeast Ohio Medical University (NEOMED), Akron, OH USA
Constantinos Koshiaris, PhD; University of Oxford, Oxford, UK
David Nunan, PhD; University of Oxford, Oxford, UK

Word Count: 3124

Contact author:

Aaron Lear

1 Akron General Way

Ctr for Fam Medicine

Akron, OH 44307 USA

ORCID iD: 0000-0002-1353-7785

leara@ccf.org

office: 330.344.6047 (Angela Watkins assistant)

Abstract

Objectives: The goals of this review are to evaluate the quality of the evidence on epidemiology of SCA/D in athletes and military members; and to synthesize existing data to estimate annual incidence of SCA/D.

Design: Systematic review and meta-analysis.

Data Sources: MEDLINE, Embase, Cochrane CENTRAL, Web of Science, BIOSIS, Scopus, SPORT discus, PEDro, were searched from inception to date searched between 2/21/19 and 3/1/19, and Clinicaltrials.gov on 7/29/19. All databases were searched without limits on language.

Eligibility for selecting studies: Studies which report incidence, or data from which incidence can be calculated, of SCA/D or both in athletes, or military members age 40.

Results: 40 total studies were included. Overall quality of the evidence was judged to be poor. 35 of 40 included studies reported 1,902 events of SCD over 404,268,083 person years and a pooled incidence rate of 1.39 events per 100,000 person years (95% CI: 0.80, 2.43) with high heterogeneity ($I^2=99\%$). 11 studies reported 219 events of SCA over 13,450,396 person years and a pooled incidence rate of 3.05 events per 100,000 person years (95% CI: 1.42, 6.58) with high heterogeneity ($I^2=96\%$).

Conclusion: The worldwide incidence of sudden cardiac death and arrest in young otherwise healthy athletes and military personnel is low. Available subgroup analysis showed incidence rates were highest in males; military members, elite athletes; and soccer athletes. Findings are tempered by unexplained heterogeneity and risk of bias in included studies.

PROSPERO Registration: CRD42019125560

Keywords: athletes, military, sudden cardiac arrest, sudden cardiac death, incidence

Background

Sudden cardiac arrest (SCA) leading to death has been shown to be the leading medical cause of death in young competitive athletes(1). It is believed that exercise and physical activity amongst athletes can lead to episodes of arrhythmia leading to sudden death(2). The topic of screening athletes with electrocardiograms in order to prevent sudden cardiac arrest and death (SCA/D) has become a controversial topic in sports medicine(3,4). Authors have pointed out that the overall burden of sudden cardiac death (SCD) remains low, around 1 event per 100,000 athletes/year or less (3), while others believe the incidence of SCA/D to be chronically undercounted (5), and that screening may decrease the incidence of arrest and death. A lack of consensus on methods to attribute sudden deaths to cardiovascular causes compounds the issue further. Surveys of high schools and universities (6,7); search of newspaper and web based reports (8,9); use of national health and autopsy records (10–12); catastrophic insurance records (13,14); and reporting on non-traumatic deaths in the military (15–17) have all been performed to determine incidence rates of SCA/D. Resulting estimates have varied widely, and whilst the validity of these different attempts has been questioned(3,4,18), existing data and their quality have not been adequately assessed systematically.

One systematic review has previously been published on the incidence of ‘sports related’ sudden cardiac death (19). The review was not pre-registered, did not report its methods according to existing reporting guidelines and assessment of the quality of evidence was not performed, resulting in more questions than answers. We are not aware of a comprehensive, pre-registered, peer-reviewed systematic review of the evidence base for the incidence of SCA

and SCD in young people regularly participating in regular exercise. Understanding the existing evidence and it's quality on the epidemiology of SCA and SCD is important when considering professional body recommendations(18,20–24) for screening athletes with electrocardiograms (ECG) for conditions which may cause such events.

Our objective is to identify publications reporting on the incidence of SCA/D in athletes and military members 40 years of age or younger, assess the quality of the evidence, and synthesize a population level incidence of both SCA and SCD.

Methods

This review is one portion of a systematic review with two objectives: Identifying the incidence of SCA and SCD in young athletes and military members; and the effect on SCA/D of ECG screening this population. The project was performed according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)(25) guidelines, and was registered on PROSPERO March 18th 2019 (CRD42019125560) (26).

Data Sources and Searches

The search strategy was designed in conjunction with a medical librarian experienced in systematic reviews (MS). The search strategy combined the dual objectives of the project into a single search. Searches were performed in MEDLINE, Embase, Cochrane CENTRAL, Web of Science, BIOSIS, Scopus, SPORT discus, PEDro, between 2/21/19 and 3/1/19, and Clinicaltrials.gov on 7/29/19 for available articles. Recent review articles and position statements were hand searched for eligible articles (5,18,19), and articles not identified in the

online search which reported incidence rates were selected for screening. There was no limitation on language of publication or date. Translation of foreign language articles was accomplished with Google Translate (translate.google.com, Google, LLC; Mountain View, CA) and physicians speaking Russian, German, French, Spanish and Chinese. Conference abstracts, published abstracts, and full text publications were included in the review.

Study Selection

Cohort studies, historically controlled trials, and survey studies which reported incidence of SCA, SCD or both in athletes, or military members age 40 and younger, as either the primary or secondary outcome were eligible for inclusion. Studies which did not report the incidence rate as the primary outcome were included if they reported a rate, or data from which a rate could be calculated, as part of their outcomes. In cases where the study reported events confirmed as SCA or SCD, and suspected cases of SCA or SCD, only the confirmed cases were included in data syntheses. In cases where the total number of person years included in the rate calculation was not reported, we used the following formula to calculate the total number of person years: $(\text{total number of events} / \text{total number person years}) \times 100,000 = \text{incidence rate per } 100,000 \text{ person years}$. In cases where the number of person years in an eligible athlete group is given as a percentage of the total number of person years included in the study, the percentage was used to calculate the number of person years in the group eligible for our study.

Independent dual screening and selection of studies was performed by a team of four reviewers (AL, NP, CM, VL). The primary author (AL) screened all abstracts/titles, and full text articles, and was involved in all resolutions of disagreements at each stage of study selection. 100% of abstracts and titles were dual screened. 90% of full text articles were reviewed by two reviewers due to resources and timing. In cases where multiple reports were published on one data set, only the most recent or complete report was included. In cases where different authors used the same databases, the article with the most complete overall database was included. Authors were contacted electronically with questions about screened texts; no author contact resulted in addition of further data sources.

Data Extraction and Quality Assessment

Data extraction, and evaluation of risk of bias (ROB) was performed in all eligible articles by the lead author (AL); 21 of the 40 eligible articles were independently assessed and extracted by a second person (NP, VL, CM), the remaining 19 individually assessed and extracted articles were checked by a senior author (DN). In cases of disagreement, these were resolved by discussion of the two involved parties. ROB was assessed with a validated tool developed to assess prevalence studies (27), and customized for the purposes of this review. Both the ROB tool, and the data extraction guide, are included in the supplemental appendix.

Data Synthesis and Analysis

The primary outcome was the reported incidence rate of SCA and SCD in competitive athletes, and active duty military members age 40 and under. Competitive athletes were considered

those participating in athletics of any form or level, including scholastic, recreational, club, collegiate/university level, and professional in or out of season. Active duty military members were included due to the high level of activity that is required of them, and the similar age profile, and interest in preventing SCA/D. Pre-specified subgroup analysis was performed by age, sex, race, and by type of sport or military member, as well as level of sport (definitions provided in subgroup analysis in supplementary appendix). Analysis was performed with the Metafor package (28) in the R statistical software(29). A random effects generalized linear mixed model(30) for both primary, and sub-group analysis was used for meta-analysis due to the high volume of zero outcome studies included in the review. Summary findings are presented as events of SCA and SCD per 100,000 person years (py) with 95% confidence intervals.

Heterogeneity was assessed with I^2 and Chi^2 ; pre-specified levels of heterogeneity for I^2 were: <30% considered low; 30-70% considered moderate; and >70% was considered high. A p-value of 0.10 or lower for Chi^2 statistic was indicated statistical heterogeneity. Assessment of publication bias of the studies included was planned using funnel plots when 10 or more studies were pooled. Sensitivity analyses were planned, if necessary, to evaluate heterogeneity within the results.

Role of the Funding Source

No funding was received for this project.

Results

After removal of 10,780 duplicates, 20,048 records were identified by the search, and 11 further articles were added by hand search. 20,059 abstracts were screened and full text of 322 were assessed for inclusion of which 40 studies met the criteria (Figure 1). Detailed characteristics of included studies is provided in the data supplement with selected details given in table one.

[FIGURE ONE HERE]

34 studies were included on athletes, ranging from youth to professional levels. Articles included were on athletes from the U.S.(n=15)(6,9,13,14,31–40), Europe (n=13)(10,12,41–51), International (n=2)(52,53) Canada (n=1)(11), Argentina (n=1)(54), Israel (n=1)(8), and Australia (n=1)(55). Five included papers were performed on military members, four in the United States (U.S.), and one in Finland. (15–17,56,57). A study focusing on firefighters in the U.S. (58) was also included due to their similarity to military members, and this was grouped with military studies.

25 (63%) included studies employed retrospective cohort designs; 12 were prospective designs (30%) and 3 were surveys, 2 cross-sectional, and one a recurring annual survey. 11 of the above studies reported extractable data only in men (32,33,42,44,50,52,55,57,58). 5 studies were included with a large proportion of the included cohort were men (15–17,51,56). Two studies did not have details about gender included in their methodology but are presumed to

be either all men, or a high proportion of men (53,54). Further details including ages, and levels of sport are included in table one.

1st Author and year of publication	Methodology	Open vs closed cohort	Exertional only or all cardiac deaths	Method of case identification	SCA or SCD Reported	Category of sport or activity	Included sport or activity	ECG screened	Ages (yrs)	Sex	Study Length (yrs)	ROB
Koskenvuo 1976	Retrospective Cohort	Known	Not described	Combined autopsy/other official record such as medical, military, athletic body	SCD only	Military	Military	No	18-24	M	6	Moderate
Phillips 1986	Retrospective Cohort	Known	Exertional	Autopsy reports	SCD only	Military	Military	No	17-28	M & F	6	Low
Assanelli 1995	Prospective Cohort	Not Described	Not described	Medical records	SCA and SCD	Multiple	Multiple	Yes	unclear	M & F	5	High
Jordaens 1996	Retrospective Cohort	Open	Not described	Media reports	SCD only	Cycling	Elite	Not described	unknown	M & F	2	High
Fuller 1997	Prospective Cohort	Known	Not described	Not reported	SCA and SCD	Multiple	Scholastic	Yes	<20	M & F	8	High
Maron 1998	Retrospective Cohort	Open	Exertional	Autopsy reports	SCD only	Multiple	Scholastic	No	13-19	M & F	1	Moderate
Young 1999	Retrospective Cohort	Open	Exertional	Autopsy reports	SCD only	Australian football	Competitive	No	15-37	M	3	Moderate
Eckart 2004	Retrospective Cohort	Known	All	Combined autopsy/other official record such as medical, military, athletic body	SCD only	Military	Military	No	17-35	M & F	3.3	High
Drezner 2005	Survey	Open	Not described	Facility event report (ex: school records)	SCA and SCD	Multiple	Elite	Not described	17-24	M & F	0.5	Moderate
Corrado 2006	Prospective Cohort	Open	All	Combined media reports/autopsy and medical records	SCD only	Multiple	Competitive	Portion	12-35	M & F	24	Low
Maron 2009	Retrospective Cohort	Open	All	Combined media reports/autopsy and medical records	SCA and SCD	Lacrosse	Competitive	No	14-23	M & F	32	High
Drezner 2009	Survey	Open	Not described	Facility event report (ex: school records)	SCA and SCD	Multiple	Scholastic	No	14-19	M & F	10	High
Chevalier 2009	Prospective Cohort	Open	Exertional	Medical records	SCD only	Multiple	Multiple	No	< 35	M & F	21	High
Solberg 2010	Retrospective Cohort	Open	Exertional	Combined autopsy/other official record such as medical, military, athletic body	SCD only	Multiple	Multiple	No	15-34	M	10	High

Holst 2010	Retrospective Cohort	Open	Exertional	Death certificate	SCD only	Soccer	Competitive	Not described	12-35	M & F	12	Low
Steinvil 2011	Retrospective Cohort	Open	Not described	Media reports	SCA and SCD	Multiple	Competitive	Portion	unclear	M & F	15	High
Marijon 2011	Prospective Cohort	Open	Exertional	Combined media reports/autopsy and medical records	SCD only	Multiple	Competitive	No	10 -35	M & F	26	Low
Eckart 2011	Prospective Cohort	Known	All	Combined autopsy/other official record such as medical, military, athletic body	SCD only	Military	Military	No	18-34	M & F	1	Low
Durakovic 2012	Retrospective Cohort	Open	Not described	Combined media reports/autopsy and medical records	SCD only	Multiple	Competitive	Portion	17-29	M	8	High
Biasco 2013	Retrospective Cohort	Known	Not described	Not reported	SCA and SCD	Soccer	Elite	Yes	Mean 23.6	M	5	High
Roberts 2013	Retrospective Cohort	Open	Exertional	Insurance records	SCD only	Multiple	Scholastic	No	12 -19	M & F	19	High
Maron 2013	Retrospective Cohort	Open	All	Combined media reports/autopsy and medical records	SCD only	Multiple	Scholastic	No	12 -18	M & F	3	Moderate
Boden 2013	Prospective Cohort	Open	Not described	Combined eyewitness interview and medical/autopsy record	SCD only	American Football	Multiple	Not described	14-25	M	6	High
Toresdahl 2014	Prospective Cohort	Known	All	Interview family/eyewitness	SCA and SCD	Multiple	Scholastic	No	14-19	M & F	11	Moderate
Risgaard 2014	Retrospective Cohort	Open	Exertional	Combined media reports/autopsy and medical records	SCD only	Multiple	Competitive	No	12-35	M & F	9	Low
Farioli 2015	Retrospective Cohort	Open	All	Combined autopsy/other official record such as medical, military, athletic body	SCD only	Military	Military	No	18-34	M	25	Low
Goutteborge 2015	Prospective Cohort	Open	All	Athletic organization records	SCD only	Soccer	Elite	Not described	18-45	Not reported	11	High
Harmon 2015	Retrospective Cohort	Open	All	Combined media reports/autopsy and medical records	SCD only	Multiple	University	Not described	17-24	M & F	26	Low
Harmon 2016	Prospective Cohort	Open	All	Media reports	SCA and SCD	Multiple	Scholastic	No	14-19	M & F	10	Moderate
Berge 2018	Retrospective Cohort	Known	All	Media reports	SCA and SCD	Soccer	Elite	Yes	18-40	M	15	Moderate
Smallman 2016	Retrospective Cohort	Known	Exertional	Combined autopsy/other official	SCD only	Military	Military	No	<35	M & F	10	Low

				record such as medical, military, athletic body								
Grani 2016	Retrospective Cohort	Open	Exertional	Autopsy reports	SCD only	Multiple	Multiple	Portion	10-39	M	7	Moderate
Maron 2016	Retrospective Cohort	Open	All	Combined media reports/autopsy and medical records	SCD only	Multiple	Competitive	Not described	15-24	M & F	10	High
Maron 2016a	Retrospective Cohort	Open	All	Autopsy reports	SCD only	Multiple	Multiple	No	14-23	M & F	20	Moderate
Landry 2017	Retrospective Cohort	Open	Exertional	Combined autopsy/other official record such as medical, military, athletic body	SCA and SCD	Multiple	Multiple	Not described	12-35	M & F	32	Low
Santos-Lozano 2017	Retrospective Cohort	Open	Exertional	Media reports	SCD only	Soccer	Elite	Portion	16-34	Not reported	21	High
Maurice 2018	Survey	Open	Exertional	Facility event report (ex: school records)	SCA and SCD	Rugby	Competitive	No	unknown	M & F	1	High
Malhotra 2018	Prospective Cohort	Known	All	Combined autopsy/other official record such as medical, military, athletic body	SCD only	Soccer	Elite	Yes	15-17 at time of screen	M & F	15	Low
Kucera 2018	Prospective Cohort	Open	All	Combined eyewitness interview and medical/autopsy record	SCD only	American Football	Competitive	No	<40	M	25	High
Endres 2019	Retrospective Cohort	Open	Exertional	Media reports	SCD only	Multiple	Multiple	No	6-17	M & F	27	High

Table 1: Characteristics of included studies

SCD=Sudden cardiac death, SCA=Sudden cardiac arrest, M=Male, F=Female

Athlete categories: Elite=professional or Division 1 NCAA athletes in U.S.; Competitive= athletes participating in organized sport with practices and competitions; University=athletes competing in all levels at a university setting; Scholastic=competing for schools between middle and high school equivalents in the U.S. (approximately ages 12-19); Multiple=combination of different levels; Military=active duty military, and one study including firefighters

12 (30%) studies were overall determined to present a low overall ROB (10,11,56,58,12,15–17,31,43,47,51); 10 (25%) were judged to have moderate risk of bias (6,9,13,37,39,40,49,50,55,57); and 18 (45%) were judged to have a high risk of bias (7,14,44–46,48,52–54,32–35,37,38,41,42). Details on ROB judgements, and table with results of ROB judgements are included in the supplemental appendix.

35 of 40 included studies provided data which was extracted to report a rate of SCD in athletes or military members. Four publications (32–34,39) are subsets of, or use data from a dataset included in the Maron et. al. (2016) article, and were included only in subgroup analysis; Smallman et. al (2016) overlapped 60% of their observation time with Eckart, et. al. (2011), and for this reason was included only with a narrative description. Included studies had a median follow up of 1,148,120 py (25th: 333,771 py; 75th: 3,084,817 py). The pooled incidence rate of SCD was 1.39 per 100,000 py (95%CI: 0.80; 2.43). There was substantial heterogeneity (I^2 99%; Chi^2 2911.11 with $p < 0.01$). 11 of these studies included data on SCA. The pooled incidence rate of SCA was 3.05 per 100,000 py (95%CI: 1.42, 6.58). This rate also exhibited substantial heterogeneity (I^2 96%; Chi^2 137.5, $p < 0.01$).

[INSERT FIGURE 2 HERE]

[INSERT FIGURE 3 HERE]

Pre-specified subgroup data synthesis was performed in cases where greater than three of the included articles had extractable data on each subgroup, and are included in table 3. For SCD

these include subjects less than 25 years old; males and females; athletes grouped by level of competition including elite, scholastic, competitive, and military; only soccer and American football included enough data extractable data amongst specific sports. For SCA these include subjects less than 25 years of age, scholastic and competitive athletes only. Forest plots for SCA/D subgroups are included in supplementary appendix. Narrative description is provided in supplementary appendix for subgroups in which no more than three articles included extractable data including university athletes, race, and further sport specific information.

Subgroup	Number of Studies	*Incidence (95% CI)	Heterogeneity I ²
Age < 25 SCD	16	0.75 (0.34, 1.65)	99%,
Age < 25 SCA	7	2.71 (1.14, 6.45)	94%,
Male SCD	16	1.69 (0.93, 3.08)	98%
Female SCD	7	0.18 (0.06, 0.52)	85%
Elite athletes SCD	8	4.38 (1.41,13.64)	97%
Scholastic athletes SCD	9	0.51 (0.33, 0.79)	80%
Scholastic athletes SCA	4	1.99 (1.08, 3.64)	81%
Competitive athletes SCD	11	0.99 (0.55, 1.76)	97%
Competitive athletes SCA	4	1.56 (1.06, 2.29)	58%
Military SCD	5	4.56 (2.80, 7.41)	93%
Soccer SCD	6	3.78, (1.60, 8.92)	89%
American Football SCD	4	0.94 (0.48, 1.85)	92%

Table 2: Results of subgroup meta-analyses

*Incidence listed per 100,000 person-years

SCD=sudden cardiac death; SCA=sudden cardiac arrest

Athlete categories: Elite=professional or Division 1 NCAA athletes in U.S.; Competitive= athletes participating in organized sport with practices and competitions; University=athletes competing in all levels at a university setting; Scholastic=competing for schools between middle and high school equivalents in the U.S. (approximately ages 12-19); Military=active duty military including recruits and activated reservists, and one study including firefighters

Categories with small number of studies are in supplementary appendix: Age ≥ 25 SCD and SCA; male and female SCA; elite athlete SCA, university athlete SCD and SCA, recreational athletes SCD and SCA, Military SCA; Race SCD and SCA; Soccer SCA; American football SCA; SCD and SCA in Australian football, baseball, basketball, cycling, lacrosse, rugby, swimming, track/cross country, volleyball.

With high heterogeneity found in both the primary and subgroup analyses, three separate sensitivity analyses were performed to investigate our findings. These were: inclusion of the studies with a cohort known to the authors, considered a ‘closed’ cohort, allowing a more accurate count of cases of SCD (10 studies) and SCA (four studies) in the numerator, and the at risk population in the denominator; an analysis of the 11 studies found to be at low risk of bias was performed on SCD only, due to a single low ROB study reporting SCA data; and an analysis including large cohorts not limited to single sports, nor single genders was performed on both SCD (21 studies), and SCA (six studies). None of the analyses performed were able to explain the heterogeneity with all I^2 values at 80% or greater, but did somewhat alter the point estimates of SCD and SCA (table three). Sensitivity analysis of the statistical model was also performed by removing outliers and zero count studies, and the point estimates were found to be stable.

Sensitivity Analysis	Number of Studies	Rate per 100,000 (95% CI)	Heterogeneity
Closed Cohort SCD	10	1.91 (0.70, 5.22)	$I^2 = 98\%$
Closed Cohort SCA	4	3.54 (0.68, 18.43)	$I^2 = 94\%$
Low ROB SCD	11	2.12 (1.24, 3.93)	$I^2 = 98\%$
Large cohorts SCD	21	0.99 (0.53, 1.82)	$I^2 = 99\%$
Large Cohorts SCA	6	1.71 (1.15, 2.56)	$I^2 = 80\%$

Table 3: Results of sensitivity analyses performed. Only 1 study with low ROB had SCA data extracted, and meta-analysis was not performed.

ROB=risk of bias; SCD=sudden cardiac death; SCA=sudden cardiac arrest

Visual assessment of funnel plot (included in supplementary appendix) for publication bias showed asymmetry in the direction of larger studies with larger incidence rates was noted in the funnel plot of all included studies. This finding is likely related to the heterogeneity present in the included studies.

Discussion

We believe ours to be the first full systematic review on the subject of sudden cardiac arrest and death in athletes, and in military members. The meta-analysis of all included studies shows a rate of SCD of 1.39 per 100,000 py and a rate of SCA of 3.05 events per 100,000 py.

Subgroup analyses for SCD showed males (n=19 studies) had higher rates of events (1.69 events per 100,000 py) than females (0.18 events per 100,000 py, n=7), but comparisons are limited by the greater number of studies reporting only on males. Scholastic athletes in the U.S., generally 12-19 years old, appear to have a lower rate of SCD than the overall estimate (n=9); this also appears true when considering those under 25 (n=16) compared to the overall estimate. This data appears fairly reliable given the number of studies analyzed, and indicates that as participants age, the risk of SCD increases.

Elite athletes had a higher rate of SCD events (4.38 per 100,000 py, n=8) compared with scholastic (0.52 per 100,000 py, n=9), and competitive athletes (0.89 per 100,000 py, n=11). The elite athlete estimate is influenced by two zero event studies in professional soccer(42,50) which give elevated point estimates due to the statistical model used, as well one professional

cycling study in men(41) which has an estimate of 89.05 cases of SCD per 100,000 py. The remaining five studies with extractable elite athlete samples have estimates ranging from 1.04 in the largest study, to 10.99 per 100,000 py. It is also true that most of the studies including elite athletes primarily include men, likely over-estimating the occurrence of SCD at a population level. With a broader sample, including other sports, and a greater number of studies including women, the rate would likely be lower. It is not clear based on the data presented here that elite athletes are necessarily at higher risk than others.

Caution should also be used interpreting the subgroup results in soccer athletes, who appear to have a higher event rate of SCD (3.78; 95%CI: 1.60, 8.92); this analysis also included the zero event studies discussed above, and is primarily male. This subgroup does include four other studies, three of whom have event rates above the point estimate of 3.78 per 100,000 py (4.22-10.99 per 100,000), with one study under the point estimate (1.04). The rate of SCD in American football relies on two studies(32,33) using database built on voluntary reporting of events. We believe this likely undercounts the number of events in this sport, and that the number reported here is an underestimate of the true incidence.

Estimates in military personnel show a high rate of SCD when compared with other athletes (4.56 per 100,000 py; 95%CI: 2.80, 7.41). This estimate includes studies dating into the 1970s, and with differences in health behavior, there is likely confounding present in this estimate. Smallman, et. al. (2016), which was not included in the data synthesis, reports that of cases of SCD in the U.S. military less than 35 years of age, 19% were smokers, 42% were overweight, and

26% obese, factors one would not expect to find at such a prevalence in athletes. While it is uncertain how traditional risk factors for coronary artery disease affects the risk of SCA/D in young active populations, it has been demonstrated that with age comes greater risk for SCA/D, and for sudden death from atherosclerotic heart disease. This estimate is higher than those seen in most athlete subgroups, and with population characteristics one would not expect to see in athletes, we believe may suggest caution when including athletes and military in the same analyses, and reviews in the future.

We were unable to perform a subgroup analysis of race but inspection of data in three studies (1,36,56) reporting some data on this suggest that black athletes and military members seem to have generally higher rates of SCD than non-black participants. Harmon, et. al. (2015) reported a rate of 1 SCD in black male basketball players per 4380 athlete years at the elite university level. The finding that the risk of SCD is higher in populations of black athletes is consistent with other data published which was not extractable (15,37,59).

We believe the primary strength of our review lies in the breadth of the search, leading to the ability of this review to draw data from multiple, large population level studies, as well as studies focusing specifically on scholastic and college athletes, and studies focusing on military members. We believe this review provides the most up-to-date and precise estimates of population level incidence, as well as estimates for a multitude of subgroups. The summary estimates of SCD and SCA appear to be stable under sensitivity analyses using different statistical methodologies despite the heterogeneity present in the results.

Limitations of our review include substantial clinical and statistical heterogeneity in the published data that was not explained by subgroup or sensitivity analyses. Many of the studies included only males, and single sports and were judged to be at moderate or high risk of bias, tempering the robustness of their findings. Most of the data included in this review is drawn from studies in Europe and the United States, potentially limiting the applicability of our findings outside of these regions. Due to available resources and timing, not all included studies were double extracted or assessed for risk of bias.

Conclusion

This systematic review provides the most comprehensive estimate of the incidence of SCA and SCD to date and shows both events to be a rare occurrence in the active young, with an incidence of SCA of 3.05 events per 100,000 py, and SCD of 1.39 events per 100,000 py. Our findings suggest higher incidence of both SCD and SCA with advancing age, male sex, and with higher level of sport or military involvement. Based on the observational nature of the studies included, the varied methodology for identifying and including cases, differences in populations, our confidence is low that the incidence rates presented may not be altered by high quality research. We believe that prospective research including a broad population of athletes at all levels is the best way to answer the question of population level incidence. As has been suggested previously(60), a uniform reporting system of sudden deaths in the young active population would benefit the understanding of this condition, and advance a more reliable understanding of the incidence of both sudden cardiac arrest and death.

Acknowledgements: This work was previously presented in abstract form at the AMSSM annual meeting 2020, and as a master's dissertation at Oxford University.

Competing Interests: No authors report competing interests with this work

Funding: No funding was received for this work.

What is already known:

- Estimated annual incidence of sudden cardiac death in athletes varies, and has been reported at between 1.0 and 4.0 per 100,000 athlete years by various narrative reviews and position statements from professional organizations, though some authors believe that the incidence is chronically undercounted due to poor methods.
- The incidence of sudden cardiac death is important when considering recommendations for or against screening electrocardiograms in young healthy individuals.
- Incidence of sudden cardiac death has been shown to be higher with advancing age, male sex, and black ethnicity.

What are the new findings:

- While there are high quality studies focusing on specific populations, the overall quality of the evidence regarding sudden cardiac arrest and death in young athletes and military members is poor, with both high clinical and statistical heterogeneity present in the published
- Meta-analysis shows the incidence both sudden cardiac arrest and death remain rare events. The incidence of sudden cardiac arrest was found to be more common at 3.05 per 100,000 athlete years (95%CI: 1.42, 6.58), and sudden cardiac death was found to be 1.39 events per 100,000 athlete years (95%CI: 0.80, 2.43).

- Subgroup analysis showed males having much higher rates than females; elite athletes, and military demonstrating higher rates of sudden death than non-elite athletes; and older subjects demonstrating higher rates of sudden death than younger athletes.

References

1. Harmon KG, Asif IM, Maleszewski J, Owens DS, Prutkin JM, Salerno JC, et al. Incidence, Etiology, and Comparative Frequency of Sudden Cardiac Death in NCAA Athletes: A Decade in Review. *Circulation*. 2015;132(1):10–9.
2. Drezner JA, O'Connor FG, Harmon KG, Fields KB, Asplund CA, Asif IM, et al. AMSSM Position Statement on Cardiovascular Preparticipation Screening in Athletes: Current Evidence, Knowledge Gaps, Recommendations and Future Directions. *Curr Sports Med Rep*. 2016;15(5):359–75.
3. Van Brabandt H, Desomer A, Gerkens S, Neyt M. Harms and benefits of screening young people to prevent sudden cardiac death. *BMJ*. 2016;353(April):1–5.
4. Drezner JA, Harmon KG, Asif IM, Marek JC. Why cardiovascular screening in young athletes can save lives: a critical review. *Br J Sports Med*. 2016;50(22):1376–8.
5. Harmon KG, Drezner JA, Wilson MG, Sharma S. Incidence of sudden cardiac death in athletes: A state-of-the-art review. *Br J Sports Med*. 2014;48(15):1185–92.
6. Drezner JA, Rogers KJ, Zimmer RR, Sennett BJ. Use of automated external defibrillators at NCAA Division I universities. *Med Sci Sport Exerc* [Internet]. 2005;37(9 PG-1487–92):1487–92. Available from: <http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:16177599&id=doi:&issn=0195-9131&isbn=&volume=37&issue=9&spage=1487&date=2005&title=Medicine+%26+Science+in+Sports+%26+Exercise&atitle=Use+of+automated+external+defibrillators+at+NCAA+Division+>

7. Drezner JA, Rao AL, Heistand J, Bloomingdale MK, Harmon KG. Effectiveness of emergency response planning for sudden cardiac arrest in United States high schools with automated external defibrillators. *Circulation* [Internet]. 2009;120(6 PG-518–25):518–25. Available from: <http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:19635968&id=doi:10.1161%2FCIRCULATIONAHA.109.855890&issn=0009-7322&isbn=&volume=120&issue=6&spage=518&date=2009&title=Circulation&atitle=Effectiveness+of+emergency+response+planning+for+sudden>
8. Steinvil A, Chundadze T, Zeltser D, Rogowski O, Halkin A, Galily Y, et al. Mandatory electrocardiographic screening of athletes to reduce their risk for sudden death proven fact or wishful thinking? *J Am Coll Cardiol*. 2011;57(11):1291–6.
9. Harmon KG, Asif IM, Maleszewski JJ, Owens DS, Prutkin JM, Salerno JC, et al. Incidence and Etiology of Sudden Cardiac Arrest and Death in High School Athletes in the United States. *Mayo Clin Proc* [Internet]. 2016;91(11 PG-1493–1502):1493–502. Available from: <http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:27692971&id=doi:10.1016%2Fj.mayocp.2016.07.021&issn=0025-6196&isbn=&volume=91&issue=11&spage=1493&date=2016&title=Mayo+Clinic+Proceedings&atitle=Incidence+and+Etiology+of+Sudden+Cardiac+Arrest>
10. Marijon E, Tafflet M, Celermajer DS, Dumas F, Perier MC, Mustafic H, et al. Sports-related sudden death in the general population. *Circulation* [Internet]. 2011;124(6 PG-672–81):672–81. Available from: <http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:21788587&id=doi:1>

- 0.1161%2FCIRCULATIONAHA.110.008979&issn=0009-7322&isbn=&volume=124&issue=6&spage=672&date=2011&title=Circulation&atitle=Sports-related+sudden+death+in+the+general+population.&
11. Landry CH, Allan KS, Connelly KA, Cunningham K, Morrison LJ, Dorian P. Sudden Cardiac Arrest during Participation in Competitive Sports. *N Engl J Med* [Internet]. 2017;377(20):1943–53. Available from: <http://www.nejm.org/doi/10.1056/NEJMoa1615710>
 12. Holst AG, Winkel BG, Theilade J, Kristensen IB, Thomsen JL, Ottesen GL, et al. Incidence and etiology of sports-related sudden cardiac death in Denmark--implications for preparticipation screening. *Heart Rhythm* [Internet]. 2010;7(10 PG-1365–71):1365–71. Available from: <http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:20580680&id=doi:10.1016%2Fj.hrthm.2010.05.021&issn=1547-5271&isbn=&volume=7&issue=10&spage=1365&date=2010&title=Heart+Rhythm&atitle=Incidence+and+etiology+of+sports-related+sudden+cardiac+death>
 13. Maron BJ, Gohman TE, Aeppli D. Prevalence of sudden cardiac death during competitive sports activities in Minnesota high school athletes. *J Am Coll Cardiol* [Internet]. 1998;32(7 PG-1881–4):1881–4. Available from: <http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:9857867&id=doi:&issn=0735-1097&isbn=&volume=32&issue=7&spage=1881&date=1998&title=Journal+of+the+American+College+of+Cardiology&atitle=Prevalence+of+sudden+cardiac+death+during+comp>

etitive+spo

14. Roberts WO, Stovitz SD. Incidence of sudden cardiac death in Minnesota high school athletes 1993-2012 screened with a standardized pre-participation evaluation. *J Am Coll Cardiol*. 2013;62(14):1298–301.
15. Eckart RE, Shry EA, Burke AP, McNear JA, Appel DA, Castillo-Rojas LM, et al. Sudden death in young adults: an autopsy-based series of a population undergoing active surveillance. *J Am Coll Cardiol* [Internet]. 2011;58(12 PG-1254–61):1254–61. Available from:
<http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:21903060&id=doi:10.1016%2Fj.jacc.2011.01.049&issn=0735-1097&isbn=&volume=58&issue=12&spage=1254&date=2011&title=Journal+of+the+American+College+of+Cardiology&atitle=Sudden+death+in+young+adult>
16. Smallman DP, Webber BJ, Mazuchowski EL, Scher AI, Jones SO, Cantrell JA. Sudden cardiac death associated with physical exertion in the US military, 2005-2010. *Br J Sports Med* [Internet]. 2016;50(2 PG-118–23):118–23. Available from:
<http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:26612844&id=doi:10.1136%2Fbjjsports-2015-094900&issn=0306-3674&isbn=&volume=50&issue=2&spage=118&date=2016&title=British+Journal+of+Sports+Medicine&atitle=Sudden+cardiac+death+associated+with+p>
17. Phillips M, Robinowitz M, Higgins JR. Sudden cardiac death in Air Force recruits. A 20-year review. *J Am Med Assoc* [Internet]. 1986;256(19 PG-2696–2699):2696–9. Available from:

<http://resolver.ebscohost.com/openurl?sid=OVID:embase&id=pmid:3773175&id=doi:10.1001%2Fjama.256.19.2696&issn=0098->

[7484&isbn=&volume=256&issue=19&spage=2696&date=1986&title=Journal+of+the+Am
erican+Medical+Association&atitle=Sudden+cardiac+death+in+Air+Forc](http://resolver.ebscohost.com/openurl?sid=OVID:embase&id=pmid:3773175&id=doi:10.1001%2Fjama.256.19.2696&issn=0098-7484&isbn=&volume=256&issue=19&spage=2696&date=1986&title=Journal+of+the+American+Medical+Association&atitle=Sudden+cardiac+death+in+Air+Forc)

18. Drezner JA, O'Connor FG, Harmon KG, Fields KB, Asplund CA, Asif IM, et al. AMSSM Position Statement on Cardiovascular Preparticipation Screening in Athletes: Current evidence, knowledge gaps, recommendations and future directions.[Erratum appears in Br J Sports Med. 2018 Mar 6;; PMID: 29510974]. Br J Sports Med. 2017;51(3):153–67.
19. Mohananey D, Masri A, Desai RM, Dalal S, Phelan D, Kanj M, et al. Global Incidence of Sports-Related Sudden Cardiac Death. J Am Coll Cardiol. 2017;69(21):2672–3.
20. Corrado D, Pelliccia A, Bjornstad HH, Vanhees L, Biffi A, Borjesson M, et al. Cardiovascular pre-participation screening of young competitive athletes for prevention of sudden death: proposal for a common European protocol. Consensus Statement of the Study Group of Sport Cardiology of the Working Group of Cardiac Rehabilitation and. Eur Heart J. 2005;26(5):516–24.
21. Ingersoll CD. The periodic health evaluation of elite athletes: A consensus statement from the international olympic committee. J Athl Train. 2009;44(5):453.
22. Exeter D, Kuah D, Carbon R, Shawdon A, Bolzonello D. Australasian College of Sport and Exercise Physicians (ACSEP) Position Statement on Pre- Participation Cardiac Evaluation in Young Athletes [Internet]. 2018. p. 1–4. Available from: <https://www.acsep.org.au>
23. Hainline B, Drezner J, Baggish A, Harmon KG, Emery MS, Myerburg RJ, et al. Interassociation consensus statement on cardiovascular care of college student-athletes.

- J Athl Train. 2016;51(4):344–57.
24. Maron BJ, Levine BD, Washington RL, Baggish AL, Kovacs RJ, Maron MS, et al. Eligibility and Disqualification Recommendations for Competitive Athletes With Cardiovascular Abnormalities: Task Force 2: Preparticipation Screening for Cardiovascular Disease in Competitive Athletes: A Scientific Statement From the American Heart Associ. Circulation. 2015;132(22):e267-72.
 25. PRISMA [Internet]. 2009 [cited 2019 Jul 16]. p. Flow Diagram. Available from: <http://prisma-statement.org>
 26. Lear A. A systematic review to determine the rate of sudden cardiac arrest and death in athletes, and to determine the benefits and harms of the electrocardiogram screening of athletes [Internet]. PROSPERO. 2019. Available from: http://www.crd.york.ac.uk/PROSPERO/display_record.php?ID=CRD42019125560
 27. Hoy D, Brooks P, Woolf A, Blyth F, March L, Bain C, et al. Assessing risk of bias in prevalence studies: Modification of an existing tool and evidence of interrater agreement. J Clin Epidemiol [Internet]. 2012;65(9):934–9. Available from: <http://dx.doi.org/10.1016/j.jclinepi.2011.11.014>
 28. Viechtbauer W. Conducting meta-analyses in R with the metafor package. J Stat Softw. 2010;36(3):1–48.
 29. R Core Team. R: A language and environment for statistical computing. [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2013. Available from: <http://www.r-project.org/>
 30. Stijnen T, Hamza TH, Özdemir P. Random effects meta-analysis of event outcome in the

- framework of the generalized linear mixed model with applications in sparse data. *Stat Med*. 2010;29(29):3046–67.
31. Harmon KG, Asif IM, Maleszewski JJ, Owens DS, Prutkin JM, Salerno JC, et al. Incidence, Cause, and Comparative Frequency of Sudden Cardiac Death in National Collegiate Athletic Association Athletes: A Decade in Review. *Circulation* [Internet]. 2015;132(1 PG-10–9):10–9. Available from:
<http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:25977310&id=doi:10.1161%2FCIRCULATIONAHA.115.015431&issn=0009-7322&isbn=&volume=132&issue=1&spage=10&date=2015&title=Circulation&atitle=Incidence%2C+Cause%2C+and+Comparative+Frequency+of+Sudde>
 32. Kucera KL, Klossner D, Colgate B, Cantu RC. Annual survey of football injury research [Internet]. Chapel Hill, NC: University of North Carolina, Chapel Hill; 2018. Available from:
NS -
 33. Boden BP, Breit I, Beachler JA, Williams A, Mueller FO. Fatalities in high school and college football players. *Am J Sports Med* [Internet]. 2013;41(5 PG-1108–16):1108–16. Available from:
<http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:23477766&id=doi:10.1177%2F0363546513478572&issn=0363-5465&isbn=&volume=41&issue=5&spage=1108&date=2013&title=American+Journal+of+Sports+Medicine&atitle=Fatalities+in+high+school+and+college+fo>
 34. Maron BJ, Doerer JJ, Haas TS, Estes NA, Hodges JS, Link MS. Commotio cordis and the epidemiology of sudden death in competitive lacrosse. *Pediatrics* [Internet]. 2009;124(3

PG-966–71):966–71. Available from:

[http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:19706581&id=doi:10.1542%2Fpeds.2009-0167&issn=0031-4005&isbn=&volume=124&issue=3&spage=966&date=2009&title=Pediatrics&atitle=Co](http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:19706581&id=doi:10.1542%2Fpeds.2009-0167&issn=0031-4005&isbn=&volume=124&issue=3&spage=966&date=2009&title=Pediatrics&atitle=Commotio+cordis+and+the+epidemiology+of+sudden+death+in+competitive)

[mmotio+cordis+and+the+epidemiology+of+sudden+death+in+competitive](http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:19706581&id=doi:10.1542%2Fpeds.2009-0167&issn=0031-4005&isbn=&volume=124&issue=3&spage=966&date=2009&title=Pediatrics&atitle=Commotio+cordis+and+the+epidemiology+of+sudden+death+in+competitive)

[mmotio+cordis+and+the+epidemiology+of+sudden+death+in+competitive](http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:19706581&id=doi:10.1542%2Fpeds.2009-0167&issn=0031-4005&isbn=&volume=124&issue=3&spage=966&date=2009&title=Pediatrics&atitle=Commotio+cordis+and+the+epidemiology+of+sudden+death+in+competitive)

35. Endres BD, Kerr ZY, Stearns RL, Adams WM, Hosokawa Y, Huggins RA, et al. Epidemiology of Sudden Death in Organized Youth Sports in the United States, 2007-2015. *J Athl Train* [Internet]. 2019;54(4 PG-349–355):349–55. Available from: NS -

36. Maron BJ, Haas TS, Ahluwalia A, Murphy CJ, Garberich RF. Demographics and Epidemiology of Sudden Deaths in Young Competitive Athletes: From the United States National Registry. *Am J Med* [Internet]. 2016;129(11 PG-1170–1177):1170–7. Available from:

[http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:27039955&id=doi:10.1016%2Fj.amjmed.2016.02.031&issn=0002-](http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:27039955&id=doi:10.1016%2Fj.amjmed.2016.02.031&issn=0002-9343&isbn=&volume=129&issue=11&spage=1170&date=2016&title=American+Journal+of+Medicine&atitle=Demographics+and+Epidemiology+of+Sudden+D)

[9343&isbn=&volume=129&issue=11&spage=1170&date=2016&title=American+Journal+](http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:27039955&id=doi:10.1016%2Fj.amjmed.2016.02.031&issn=0002-9343&isbn=&volume=129&issue=11&spage=1170&date=2016&title=American+Journal+of+Medicine&atitle=Demographics+and+Epidemiology+of+Sudden+D)

[of+Medicine&atitle=Demographics+and+Epidemiology+of+Sudden+D](http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:27039955&id=doi:10.1016%2Fj.amjmed.2016.02.031&issn=0002-9343&isbn=&volume=129&issue=11&spage=1170&date=2016&title=American+Journal+of+Medicine&atitle=Demographics+and+Epidemiology+of+Sudden+D)

37. Maron BJ, Haas TS, Duncanson ER, Garberich RF, Baker AM, Mackey-Bojack S. Comparison of the Frequency of Sudden Cardiovascular Deaths in Young Competitive Athletes Versus Nonathletes: Should We Really Screen Only Athletes? *Am J Cardiol*. 2016;117(8):1339–41.
38. Fuller CM, McNulty CM, Spring DA, Arger KM, Bruce SS, Chryssos BE, et al. Prospective screening of 5,615 high school athletes for risk of sudden cardiac death. *Med Sci Sport*

- Exerc. 1997;29(9):1131–8.
39. Maron BJ, Haas TS, Ahluwalia A, Rutten-Ramos SC. Incidence of cardiovascular sudden deaths in Minnesota high school athletes. *Hear Rhythm*. 2013;10(3):374–7.
 40. Toresdahl BG, Rao AL, Harmon KG, Drezner JA. Incidence of sudden cardiac arrest in high school student athletes on school campus. *Hear Rhythm* [Internet]. 2014;11(7 PG-1190–4):1190–4. Available from:
<http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:24732370&id=doi:10.1016%2Fj.hrthm.2014.04.017&issn=1547-5271&isbn=&volume=11&issue=7&spage=1190&date=2014&title=Heart+Rhythm&atitle=Incidence+of+sudden+cardiac+arrest+in+high+school+student+at>
 41. Jordaens L, Van Lierde J, Van De Velde D, De Backer G. Competitive cycling in Belgium is associated with an increased risk of sudden death. *Pacing Clin Electrophysiol* [Internet]. 1996;19(4 PG-602–603):602–3. Available from:
<http://search.ebscohost.com/login.aspx?direct=true&AuthType=ip,cookie,url,uid&db=s3h&AN=17417795&site=ehost-live> NS -
 42. Biasco L, Cristoforetti Y, Castagno D, Giustetto C, Astegiano P, Ganzit G, et al. Clinical, electrocardiographic, echocardiographic characteristics and long-term follow-up of elite soccer players with J-point elevation. *Circ Arrhythmia Electrophysiol*. 2013;6(6):1178–84.
 43. Risgaard B, Winkel BG, Jabbari R, Glinge C, Ingemann-Hansen O, Thomsen JL, et al. Sports-related sudden cardiac death in a competitive and a noncompetitive athlete population aged 12 to 49 years: data from an unselected nationwide study in Denmark. *Hear Rhythm* [Internet]. 2014;11(10 PG-1673–81):1673–81. Available from:

- <http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:24861446&id=doi:10.1016%2Fj.hrthm.2014.05.026&issn=1547-5271&isbn=&volume=11&issue=10&spage=1673&date=2014&title=Heart+Rhythm&atitle=Sports-related+sudden+cardiac+death+in+a+competitive+and+a+>
44. Solberg EE, Gjertsen F, Haugstad E, Kolsrud L. Sudden death in sports among young adults in Norway. *Eur J Cardiovasc Prev Rehabil* [Internet]. 2010;17(3 PG-337–41):337–41. Available from:
<http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:20038839&id=doi:10.1097%2FHJR.0b013e328332f8f7&issn=1741-8267&isbn=&volume=17&issue=3&spage=337&date=2010&title=European+Journal+of+Cardiovascular+Prevention+%26+Rehabilitation&atitle=Sudden+d>
45. Assanelli D, Marconi M, Cazzamalli L, Poeta ML, Bortolotti M, Restori M, et al. Sudden death in seven young athletes during sports activity. [Italian]. *Int J Sport Cardiol* [Internet]. 1995;4(1 PG-25–29):25–9. Available from:
<http://resolver.ebscohost.com/openurl?sid=OVID:embase&id=pmid:&id=doi:&issn=0393-6066&isbn=&volume=4&issue=1&spage=25&date=1995&title=International+Journal+of+Sports+Cardiology&atitle=ARRESTO+CARDIACO+IN+SETTE+GIOVANI+ATLETI+DURANTE+ATTIVITA+SPORTIVA&aula>
46. Chevalier L, Hajjar M, Douard H, Cherief A, Dindard JM, Sedze F, et al. Sports-related acute cardiovascular events in a general population: a French prospective study. *Eur J Cardiovasc Prev Rehabil* [Internet]. 2009;16(3 PG-365–70):365–70. Available from:

- <http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:19318955&id=doi:10.1097%2FHJR.0b013e3283291417&issn=1741-8267&isbn=&volume=16&issue=3&spage=365&date=2009&title=European+Journal+of+Cardiovascular+Prevention+%26+Rehabilitation&atitle=Sports-r>
47. Corrado D, Basso C, Pavei A, Michieli P, Schiavon M, Thiene G. Trends in sudden cardiovascular death in young competitive athletes after implementation of a preparticipation screening program. *JAMA*. 2006;296(13):1593–601.
 48. Durakovic Z, Durakovic MM, Skavic J. Sudden cardiac death due to physical exercise in Croatia in a 27-year period. *Med Sci [Internet]*. 2012;37(PG-19-51):19–51. Available from: NS - Durakovic 2012
 49. Grani C, Chappex N, Fracasso T, Vital C, Kellerhals C, Schmied C, et al. Sports-related sudden cardiac death in Switzerland classified by static and dynamic components of exercise. *Eur J Prev Cardiol [Internet]*. 2016;23(11 PG-1228–36):1228–36. Available from: <http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:26915579&id=doi:10.1177%2F2047487316632967&issn=2047-4873&isbn=&volume=23&issue=11&spage=1228&date=2016&title=European+Journal+of+Preventive+Cardiology&atitle=Sports-related+sudden+cardiac+dea>
 50. Berge HM, Andersen TE, Bahr R. Cardiovascular incidents in male professional football players with negative preparticipation cardiac screening results: an 8-year follow-up. *Br J Sports Med [Internet]*. 2018 Oct 1;bjsports-2018-099845. Available from: <http://bjsm.bmj.com/content/early/2018/09/30/bjsports-2018-099845.abstract>
 51. Malhotra A, Dhutia H, Finocchiaro G, Gati S, Beasley I, Clift P, et al. Outcomes of Cardiac

- Screening in Adolescent Soccer Players. *N Engl J Med*. 2018;379(6):524–34.
52. Gouttebarga V, Ooms W, Tummers T, Inklaar H. Mortality in international professional football (soccer): a descriptive study. *J Sports Med Phys Fitness* [Internet]. 2015;55(11 PG-1376–1382):1376–82. Available from:
<http://resolver.ebscohost.com/openurl?sid=OVID:embase&id=pmid:25289716&id=doi:&issn=0022-4707&isbn=&volume=55&issue=11&spage=1376&date=2015&title=The+Journal+of+sports+medicine+and+physical+fitness&atitle=Mortality+in+international+professional+football+%25>
53. Santos-Lozano A, Martin-Hernandez J, Baladron C, Angeles Turrado-Sevilla M, Arraras-Flores A, Villacastin JP, et al. Sudden Cardiac Death in Professional Soccer Players. *J Am Coll Cardiol*. 2017;70(11 PG-1420–1421):1420–1.
54. Maurice MF, Di Tommaso F, Barros Pertuz MC, Mendoza WA, Spagnuolo D, Lucas V. Sudden cardiac death in rugby clubs. *Rev Argent Cardiol* [Internet]. 2018;86(1 PG-40–44):40–4. Available from:
<http://resolver.ebscohost.com/openurl?sid=OVID:embase&id=pmid:&id=doi:10.7775%2Frac.v86.i1.12263&issn=0034-7000&isbn=&volume=86&issue=1&spage=40&date=2018&title=Revista+Argentina+de+Cardiologia&atitle=Muerte+subita+en+clubes+deportivos+de+rugby&aulast=Mau>
55. Young MC, Fricker PA, Thomson NJ, Lee KA. Sudden death due to ischaemic heart disease in young aboriginal sportsmen in the Northern Territory, 1982-1996. *Med J Aust* [Internet]. 1999;170(9 PG-425–8):425–8. Available from:

[\[729X&isbn=&volume=170&issue=9&spage=425&date=1999&title=Medical+Journal+of+Australia&atitle=Sudden+death+due+to+ischaemic+heart+disease+in+young+aboriginal+sportsmen\]\(http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:10341774&id=doi:&issn=0025-729X&isbn=&volume=170&issue=9&spage=425&date=1999&title=Medical+Journal+of+Australia&atitle=Sudden+death+due+to+ischaemic+heart+disease+in+young+aboriginal+sportsmen\)](http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:10341774&id=doi:&issn=0025-</p></div><div data-bbox=)

56. Eckart RE, Scoville S, Campbell C, Shry EA, Stajduhar KC, Potter RN, et al. Sudden death in young adults : A 25-year review of autopsies in Annals of Internal Medicine Article. Ann Intern Med. 2004;141(11).

57. Koskenvuo K. Sudden deaths among Finnish conscripts. Br Med J [Internet]. 1976;2(6049 PG-1413–5):1413–5. Available from:

[http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:1009356&id=doi:&issn=0007-](http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:1009356&id=doi:&issn=0007-1447&isbn=&volume=2&issue=6049&spage=1413&date=1976&title=British+Medical+Journal&atitle=Sudden+deaths+among+Finnish+conscripts.&aulast=Koskenvuo)

[1447&isbn=&volume=2&issue=6049&spage=1413&date=1976&title=British+Medical+Journal&atitle=Sudden+deaths+among+Finnish+conscripts.&aulast=Koskenvuo](http://resolver.ebscohost.com/openurl?sid=OVID:medline&id=pmid:1009356&id=doi:&issn=0007-1447&isbn=&volume=2&issue=6049&spage=1413&date=1976&title=British+Medical+Journal&atitle=Sudden+deaths+among+Finnish+conscripts.&aulast=Koskenvuo) NS -
Koskenvuo 1976

58. Farioli A, Christophi CA, Quarta CC, Kales SN. Incidence of Sudden Cardiac Death in a Young Active Population. J Am Heart Assoc [Internet]. 2015;4 (6) (no(1818 PG-). Available from:

[http://resolver.ebscohost.com/openurl?sid=OVID:embase&id=pmid:26066031&id=doi:10.1161%2FJAHA.115.001818&issn=2047-](http://resolver.ebscohost.com/openurl?sid=OVID:embase&id=pmid:26066031&id=doi:10.1161%2FJAHA.115.001818&issn=2047-9980&isbn=&volume=4&issue=6&spage=&date=2015&title=Journal+of+the+American+Heart+Association&atitle=Incidence+of+Sudden+Cardiac+Death+in+a+Yo)

[9980&isbn=&volume=4&issue=6&spage=&date=2015&title=Journal+of+the+American+Heart+Association&atitle=Incidence+of+Sudden+Cardiac+Death+in+a+Yo](http://resolver.ebscohost.com/openurl?sid=OVID:embase&id=pmid:26066031&id=doi:10.1161%2FJAHA.115.001818&issn=2047-9980&isbn=&volume=4&issue=6&spage=&date=2015&title=Journal+of+the+American+Heart+Association&atitle=Incidence+of+Sudden+Cardiac+Death+in+a+Yo)

59. Maron BJ, Haas TS, Murphy CJ, Ahluwalia A, Rutten-Ramos S. Incidence and causes of sudden death in U.S. college athletes. *J Am Coll Cardiol.* 2014;63(16):1636–43.
60. Solberg EE, Borjesson M, Sharma S, Papadakis M, Wilhelm M, Drezner JA, et al. Sudden cardiac arrest in sports - Need for uniform registration: A Position Paper from the Sport Cardiology Section of the European Association for Cardiovascular Prevention and Rehabilitation. *Eur J Prev Cardiol.* 2016;23(6):657–67.

Figures/Tables

PRISMA Flow Diagram (25)




Figure 1: PRISMA flow diagram




Figure 2: Meta-analysis of incidence rate of sudden cardiac death in athletes/military on 35 of 40 included studies.




Figure 3: Meta-analysis of incidence rate of sudden cardiac arrest in athletes/military on 11 included studies reporting sudden cardiac arrest.