Improved estimation of time-varying reproduction numbers at low case incidence and between epidemic waves

Kris V Parag[†, 1]

[1]MRC Centre for Global Infectious Disease Analysis, Imperial College London, London, W2 1PG, UK

[†]Email: k.parag@imperial.ac.uk

Abstract

We construct a recursive Bayesian smoother, termed EpiFilter, for estimating the effective reproduction number, R, from the incidence of an infectious disease in real time and retrospectively. Our approach borrows from Kalman filtering theory, is quick and easy to compute, generalisable, deterministic and unlike many current methods, requires no change-point or window size assumptions. We model R as a flexible, hidden Markov state process and exactly solve forward-backward algorithms, to derive R estimates that incorporate all available incidence information. This unifies and extends two popular methods, EpiEstim, which considers past incidence, and the Wallinga-Teunis method, which looks forward in time. We find that this combination of maximising information and minimising assumptions significantly reduces the bias and variance of R estimates. Moreover, these properties make EpiFilter more statistically robust in periods of low incidence, where existing methods can become destabilised. As a result, EpiFilter offers improved inference of time-varying transmission patterns that are especially advantageous for assessing the risk of second waves of infection in real time and at various spatial scales.

Key-words: Bayesian filters, reproduction numbers, epidemic models, COVID-19, infectious diseases.

INTRODUCTION

During an unfolding epidemic, one of the most commonly available and useful types of surveillance data is the daily (or weekly) number of newly reported cases. This time-series of case counts, also known as the incidence curve, not only measures the epidemic size and burden, but also provides information about trends or changes in its transmissibility [1], [2]. These trends are captured by the time-varying effective or instantaneous reproduction number, denoted R_s at time s, which defines how the number of secondary cases generated per primary case varies across the outbreak [3]. Broadly, when $R_s > 1$ we can expect and prepare for growing incidence, whereas sustained $R_s < 1$ signifies that the epidemic is waning and likely to enter a more controlled phase [4].

Inferring changes in R_s given an observed incidence curve is therefore crucial, both to understanding transmissibility and to forecasting upcoming case loads, especially for an ongoing epidemic, where it can help inform

policymaking and intervention choices or predict healthcare demands [1], [5]. Real-time and retrospective R_s estimates have been used to characterise rates and patterns of spread in various diseases such as malaria [6] and Ebola virus disease [7]. Such estimates have proven valuable throughout the COVID-19 pandemic, providing updating synopses of global transmission [8] and evidencing the impact of past control actions (e.g. lockdowns and social distancing) [9] or the likelihood of a resurgence in infections when those controls are relaxed [10].

Most studies that infer R_s or related quantities either apply the Wallinga-Teunis (WT) method [2] or the Cori *et al* method, known as EpiEstim [11]. Both methods take complementary viewpoints on how incidence data inform on transmissibility and hence have diverse use-cases. The WT method reconstructs the average number of new cases caused by infectious individuals at s and so requires incidence data beyond time s for its estimate. It computes the case or cohort reproduction number, R_s^c, which is a function of R_{s+j} for future times $j \geq 0$, and is suited for retrospective analyses [12]. Alternatively, EpiEstim infers how past infections propagate to form the incidence observed at s, only requiring data prior to time s. EpiEstim directly computes instantaneous reproduction numbers, R_s, and is preferred for real-time investigations [3].

While both methods provide useful and important estimators of transmission, they are not perfect. Two main limitations exist. First, each suffers from data censoring or edge-effects [3]. Because the WT method is forward-looking i.e., depends on data later than s, its estimates are right censored when s is close to the last observed time point [12]. In contrast, EpiEstim looks backward in time and suffers edge-effects when s is near the first observed time point [11]. Estimates in the vicinity of the start and end of the incidence time-series are therefore unreliable under EpiEstim and the WT method, respectively. Techniques have been proposed to limit this unreliability [5], [13], but the problem is intrinsic, and inevitable near the actual start and end times of an epidemic, where there is necessarily no or sparse data.

This leads into the second key limitation: inference in periods of small incidence. This presents a fundamental challenge for any R_s estimation approach and effectively creates additional edge-effects. When few or no case counts are available to constrain inference, methods are largely driven by their inherent prior distributions and assumptions. This can result in misleading or unreliable estimates and mask subcritical (i.e., $R_s < 1$) transmission patterns [11], [15]. Small incidence may occur during sustained periods of interventions, when pathogens invade new susceptible regions and also naturally arise by division when analyses are performed at smaller spatial scales (e.g., regional or community levels). Understanding how to best mediate the trade-off between prior assumptions and data when incidence is small is of both statistical and epidemiological significance.

Following a period of low incidence, two important outcomes are possible: either the epidemic continues to exhibit small or zero case counts until it goes extinct, or a resurgence in infections, also termed a second wave, occurs. Inferring, in real time, which of these conditions is likely presents a key challenge for infectious disease

Fig. 1: **EpiFilter algorithm and relationship to other methods.** In the left panels we consider three ways of inferring the instantaneous or effective reproduction number at time s, R_s, from the incidence curve, I_1^t (blue dots). The filtering solution produces the posterior distribution $\mathbf{p_s}$ from all data prior to time s. EpiEstim approximates this solution by using the subset of data in a window of size k into the past. Reverse-filtering considers the complementary part of the incidence curve, leading to $\mathbf{r_s}$, which utilises data beyond s. The WT method, with future window k, approximates this type of solution. Smoothing uses all information from I_1^t to generate $\mathbf{q_s}$, which is precisely computed by EpiFilter. Blue windows show the portions of I_1^t that inform on R_s for each of $\mathbf{p_s}$, $\mathbf{r_s}$ and $\mathbf{q_s}$ while red windows highlight the subsets used by EpiEstim and the WT method. Double arrows indicate data used for constructing various posterior distributions, while square arrows pinpoint instances of those distributions at the edges of I_1^t. In the right panels we summarise the construction of EpiFilter. We outline the main assumptions (the model box) and computations (the algorithm box) necessary for realising EpiFilter, which allow us to obtain the most informative (and minimum MSE) smoothing posterior distribution $\mathbf{q_s}$. See the main text for the specific equations employed in our implementation [14].

epidemiology given the information bottleneck at low incidence [16]. Better inference of transmission under these

conditions is currently considered central to designing data-informed COVID-19 intervention exit or relaxation

strategies [17]. With many countries facing multiple resurgent waves in this pandemic, estimating fluctuations in

transmission during suspected epidemic troughs could be essential to achieving sustained control [10].

Here we present and develop a novel method, termed EpiFilter, for reliably estimating R_s in real time, which

ameliorates the above limitations. We take an engineering inspired approach and construct an exact, recursive

and deterministic (i.e., EpiFilter produces the same output for fixed input data and requires no Monte Carlo

steps) inference algorithm that is quick and easy to compute both across an unfolding outbreak and in retrospect.

Our method solves what is called the smoothing problem in control engineering [14]. This means we compute instantaneous reproduction number, R_s, estimates that formally integrate both forward and backward looking information. This unifies the WT method and EpiEstim, and largely nullifies their edge-effect issues.

Further, EpiFilter only makes a minimal Markov assumption for R_s, which allows it to avoid the strong prior window size and change-point assumptions that existing methods may apply to infer shifts in transmission [11], [9]. Using simulated and empirical data, we show that EpiFilter accurately tracks changes in R_s and provides reliable one-step-ahead incidence predictions. Moreover, we find that EpiFilter is appreciably more robust and statistically efficient than even optimised versions of EpiEstim [13]. Specifically, it does not easily destabilise when performing real-time inference in periods of low incidence, such as in lulls between epidemic waves, and generally it minimises the mean squared error of R_s estimates, while maintaining good coverage and predictive performance.

We illustrate the practical utility of EpiFilter on the COVID-19 incidence curve of New Zealand, which exhibits a second wave that was seeded during a prolonged low incidence period, finding stark improvements in the transmission patterns it uncovers. Our method, which is outlined in Fig. 1, provides a straightforward yet formally optimal (in mean squared error sense [18]) solution to real-time and retrospective instantaneous reproduction number estimation. Because it couples minimal prior assumptions with maximum information extraction, it more gracefully handles periods with scarce data. Hopefully our approach will serve as a useful inference tool for investigating the risk of resurgence in COVID-19 and other epidemics. Matlab and R implementations of EpiFilter are available at https://github.com/kpzoo/EpiFilter and its mechanics are explored and validated in the Appendix.

METHODS

Renewal models and inference problems

We consider an infectious disease epidemic observed over some time period $1 \leq s \leq t$ in a homogeneous and well-mixed population. While epidemics actually spread on dynamic networks involving stratified contact structures, homogeneous models can provide useful real-time insight into key transmission patterns and are more easily fit and verified with routine surveillance data such as incidence curves [19]. If the incidence or number of newly infected cases at time s is I_s then the set $I_1^t = \{I_1, I_2, \ldots I_t\}$ is the incidence curve of the epidemic. We assume that incidence is available on a daily scale so that I_1^t is a vector of t daily counts but weeks or months could be used instead. A common problem in infectious disease is the inference of the transmissibility of the epidemic given this curve. The renewal model [1], [20] presents a general and popular framework for investigating this problem and its estimates, in some instances, can even approximate those from detailed network models [21].

The renewal model posits that epidemic transmissibility, summarised by the effective or instantaneous reproduction number, R_s, generates the observed incidence as in Eq. (1). We assume incidence counts are Poisson distributed

with Pois indicating Poisson noise. Here $\overset{d}{=}$ signifies equality in distributions and $\mid$ means 'conditioned on'.

$$\mathbb{P}(I_s \,|\, R_s, \, I_1^{s-1}) \overset{d}{=} \text{Pois}(\Lambda_s R_s) \tag{1}$$

While Eq. (1) does not directly model how susceptible individuals become infected, these effects are encoded in the reproduction number R_s, which measures the secondary cases generated per effective primary case at s [3]. The quantity $\Lambda_s := \sum_{u=1}^{s-1} I_{s-u} w_u$, known as the total infectiousness, counts how many effective past cases are still infectious at s i.e., it describes the number of circulating cases that can actively transmit.

The generation time distribution of the epidemic, $w_1^{s-1} = \{w_1, \, w_2, \, \ldots w_{s-1}\}$, controls how past incidence influences Λ_s, with w_u as the probability that a primary case takes between $u-1$ and u days to generate a secondary case [1]. We make the standard assumption that w_1^{∞} is well approximated by the serial interval distribution of the epidemic of interest, which is known [11]. The serial interval is the time between symptom onset of a primary and its secondary case. Onset times are more practically measurable than actual infection times. While this approximation can limit inference (for example, serial intervals often have larger variances than generation times), methods are being developed to improve its quality [22].

We focus on inferring the complete set of R_s values, denoted $R_1^t = \{R_1, \, R_2, \, \ldots R_t\}$, given the incidence curve I_1^t with t as the last recorded time. Estimating R_1^t is important because changes in the values of instantaneous reproduction numbers often signify key transitions in epidemic transmissibility, which might be due to the imposition or relaxation of interventions. Instantaneous reproduction numbers are also the basis of other transmissibility metrics, such as cohort reproduction numbers or growth rates [3]. We define three main inference problems, based on how information in I_1^t is recruited to construct every R_s estimate. We represent these problems in terms of the posterior distribution their solutions induce over possible R_s values. Estimates are functions of these posteriors.

The first is called filtering, where we sequentially compute the filtering posterior $\mathbf{p_s} = \mathbb{P}(R_s \,|\, I_1^s)$ for every $s \leq t$ [23]. Filtering only uses incidence data up to time s, I_1^s, for inferring R_s. Solving this problem is fundamental to real-time inference [24]. The second problem, which we call reverse-filtering, is the complement of the first. The reverse-filtering posterior $\mathbf{r_s} = \mathbb{P}(R_s \,|\, I_s^t)$, is important for retrospective or backward-looking estimates and infers R_s from incidence beyond s i.e., I_s^t [25]. Filtering solutions are commonly employed for inferring instantaneous reproduction numbers. Practical R_s calculations do not involve reverse-filtering. Instead, the information used by $\mathbf{r_s}$ is implicit to deriving cohort reproduction numbers, $R_s^c = \sum_{j=0}^{t-s} R_{s+j} w_{j+1}$ [12]. Later we show that estimates from EpiEstim and the WT method are related to the $\mathbf{p_s}$ and $\mathbf{r_s}$ distributions, respectively.

The last problem, which is termed smoothing, is our main interest. It asks the question: how can we construct an R_s, at every $s \leq t$, that integrates all available incidence information from I_1^t. To solve this problem we must formulate the smoothing posterior distribution $\mathbf{q_s} = \mathbb{P}(R_s \,|\, I_1^t)$ [14]. Functions of this posterior would then yield

maximally informed instantaneous reproduction number estimates (and cohort reproduction number estimates by extension). Note that $\mathbf{p_s}$, $\mathbf{r_s}$ and $\mathbf{q_s}$ depend on the choices of state space model, which describes the dynamics of R_s across time, and observation model, which explains how changes in R_s lead to trends in observed incidence data [18]. These models encode our assumptions about the epidemic of interest and determine how estimates trade off assumptions against data. We next explore how posterior distribution selection determines performance, especially when data are scarce, and examine how EpiEstim and the WT methods fit within this framework.

Inference methods and low incidence

We mostly detail EpiEstim as instantaneous reproduction number, R_s, estimates are the main focus of this work. EpiEstim assumes that the estimate of R_s at time s depends on a rolling past window of data, $\tau(s) = \{s, s-1, \ldots, s-k+1\}$, of size k [11]. Consequently, at time s, only size k subsets of the incidence, I_{s-k+1}^s, and total infectiousness, $\Lambda_{s-k+1}^s = \{\Lambda_s, \Lambda_{s-1}, \ldots \Lambda_{s-k+1}\}$, are considered informative about R_s. Their sums over this window are $i_{\tau(s)} = \sum_{u \in \tau(s)} I_u$ and $\lambda_{\tau(s)} = \sum_{u \in \tau(s)} \Lambda_u$. When $k = 1$, EpiEstim only uses the most recent case count, I_s, (and Λ_s) to estimate R_s. While this maximises flexibility, it usually results in over-fitting and so larger windows are often employed to trade off estimate variance with bias [11], [26].

EpiEstim therefore effectively solves a filtering problem, as discussed in the previous section. The filtering posterior distribution produced by EpiEstim is restricted to the informative window $\tau(s)$ and denoted $\mathbf{p_{\tau(s)}} = \mathbb{P}(R_s \,|\, I_{s-k+1}^s)$. This is parametrised by the shape-scale gamma posterior distribution in Eq. (2).

$$\mathbf{p_{\tau(s)}} \overset{d}{=} \mathrm{Gam}(a + i_{\tau(s)}, \, (c + \lambda_{\tau(s)})^{-1}) \tag{2}$$

The posterior $\mathbf{p_{\tau(s)}}$ results from combining a gamma prior distribution on R_s, $\mathbb{P}(R_s) \overset{d}{=} \mathrm{Gam}(a, c^{-1})$, with a Poisson observation likelihood for the incidence (see Eq. (1)), as detailed in [11], [13]. We never explicitly include Λ_s terms in our notation (e.g., we could write $\mathbf{p_{\tau(s)}}$ as $\mathbb{P}(R_s \,|\, I_{s-k+1}^s, \Lambda_{s-k+1}^s)$) since they appear naturally when using Eq. (1) and the key difference among our inference problems relate to I_s terms.

The posterior mean estimate from Eq. (2) is constructed as $\tilde{R}_{\tau(s)} = \int \mathbf{p_{\tau(s)}} R_s \, dR_s = \mathbb{E}[R_s \,|\, I_{s-k+1}^s] = (a + i_{\tau(s)})(c + \lambda_{\tau(s)})^{-1}$. The variance around this estimate is $(a + i_{\tau(s)})(c + \lambda_{\tau(s)})^{-2}$. The observation model is given by Eq. (1) but the state space model of EpiEstim is not explicit. However, if $R_{\tau(s)}$ is the assumed average reproduction number in $\tau(s)$ (which is used to estimate R_s) then $\lambda_{\tau(s)} R_{\tau(s)} = \sum_{u \in \tau(s)} \Lambda_u R_u$ [13]. Thus, EpiEstim somewhat incorporates a linear moving average state space model, and assumes that the filtering distribution $\mathbf{p_s} \approx \mathbf{p_{\tau(s)}}$ by deeming data outside $\tau(s)$, I_1^{s-k}, as effectively uninformative [12]. Since $\mathbf{p_{\tau(s)}}$ can be computed sequentially across an ongoing epidemic, EpiEstim provides real-time inference.

The WT method takes a complementary approach to EpiEstim, computing transmissibility over a forward-looking

window $\gamma(s) = \{s, s+1, \ldots, s+k-1\}$ [2]. Often $k = t - s + 1$ i.e., the window extends to the last observed incidence. The WT method uses the observation model of Eq. (1) and has an implicit moving average state model $R_{\gamma(s)} = \sum_{u \in \gamma(s)} R_u w_{u-s+1}$, which leads to its cohort reproduction number estimates [3]. As this method effectively uses future information [12], it implicitly involves approximating the reverse-filtering distribution (see previous section) as $\mathbf{r_s} \approx \mathbf{r_{\gamma(s)}} = \mathbb{P}(R_s \,|\, I_s^{s+k-1})$. We illustrate the information windows employed by the WT method and EpiEstim, as well as the complete filtering and reverse-filtering windows in Fig. 1. The goodness of the windowed distributions $\mathbf{p}_{\tau(\mathbf{s})}$ and $\mathbf{r}_{\gamma(\mathbf{s})}$ as approximations to the general posterior distributions $\mathbf{p_s}$ and $\mathbf{r_s}$ will depend on k and the appropriateness of the state model underlying each method [13].

While EpiEstim and the WT methods are powerful tools for inferring transmissibility in real-time and in retrospect, they have two main and related limitations, which necessarily reduce the reliability of their outputs [12]. First, their performance degrades as s gets close to 1 for EpiEstim and t for the WT method [11]. These edge or censoring effects correspond, at the extreme, to $\mathbf{p_1} = \mathbf{p}_{\tau(1)} = \mathbb{P}(R_1 \,|\, I_1)$ and $\mathbf{r_t} = \mathbf{r}_{\gamma(\mathbf{t})} = \mathbb{P}(R_t \,|\, I_t)$, which are weakly informed posterior distributions. As a result, at the beginning of the incidence curve EpiEstim can be unreliable (and even unidentifiable). The WT method suffers similarly at the end of that curve [1] (see Fig. 1).

The second limitation occurs in phases of the epidemic where incidence is low for a prolonged period of time [26], [17]. In these periods data are sparse and the quality of estimates depend on how well the method of choice mediates between the little available information and its inherent assumptions [15]. We illustrate this with EpiEstim, using the mean and variance of the posterior estimate from Eq. (2), $\tilde{R}_{\tau(s)}$, which are defined above. If incidence is small over the window $\tau(s)$ then the sum of incidence, $i_{\tau(s)}$, and the total infectiousness $\lambda_{\tau(s)}$ shrink, meaning that the prior hyperparameters, a and c, strongly influence the resulting estimate mean and variance. This contrasts the data-rich scenario when an epidemic is large, where $i_{\tau(s)}$ and $\lambda_{\tau(s)}$ overpower a and c.

Further, should a sequence of $n \geq k$ zero-incidence days occur, then $i_{\tau(s)} = 0$ and $\lambda_{\tau(s)} \to 0$ as n increases, with a rate controlled by the serial interval of the epidemic. Here the shape parameter of $\mathbf{p}_{\tau(\mathbf{s})}$ is exactly that of the prior distribution $\mathbb{P}(R_s)$. As $\lambda_{\tau(s)}$ decays, $\mathbf{p}_{\tau(\mathbf{s})} \to \mathbb{P}(R_s)$ (see Eq. (2)), R_s becomes statistically unidentifiable from the window of data and inference is completely prior driven [27], [26]. While lack of data is a fundamental limitation, the point at which we lose inferential power is not fixed, and depends on the window size, k. Studies that formally optimised k for estimate reliability, found that small k is needed to infer sharp changes in transmissibility (e.g. due to lockdowns) [13], indicating that these issues can be acute. Analogous effects occur in the WT method if there are few incident cases across its forward-looking window $\gamma(s)$ [12].

These prior-driven scenarios are realistic for epidemics in waning or tail phases, and can precede either elimination (i.e., epidemic extinction) or resurgence [28]. While some estimate degradation is guaranteed for any R_s inference method when faced with either edge-effects or low incidence, robustness can still be improved. Edge-effects can be

largely overcome by constructing the smoothed posterior distribution for estimating the instantaneous reproduction number R_s, denoted $\mathbf{q_s}$. Solving the smoothing problem melds the advantages of the opposite looking windows of EpiEstim and the WT method, removing the vulnerability near the ends of the incidence curve I_1^t. This follows as $\mathbf{q_1} = \mathbf{r_1}$ and $\mathbf{q_t} = \mathbf{p_t}$ (see Fig. 1). Further, by maximising the information used for inferring every R_s and by minimising our state model assumptions, we can ameliorate the impact of low incidence. We next develop a method, termed EpiFilter, to realise these improvements.

Bayesian (forward) recursive filtering

We reformulate the inference problem of estimating instantaneous reproduction numbers R_s from past incidence I_1^s as an optimal Markov state filtering problem. Filtering describes a general class of engineering problems aimed at optimally, usually in a mean squared error (MSE) sense, inferring some hidden state in real time from noisy observations [18], [14]. Given some functions f_s and g_s, which describe the state (R_s in our case) space dynamics and the process of generating noisy observations (the I_s here), the filtering problem tries to construct the posterior distribution $\mathbf{p_s}$ (see previous section) [23], which EpiEstim approximates. The conditional mean estimate $\tilde{R}_s = \mathbb{E}[R_s \,|\, I_1^s]$ leads to the minimum MSE of $\mathbb{E}[(R_s - \tilde{R}_s)^2]$ [23], which depends on all the past information.

The famed Kalman filter [24] was the genesis of these methods. Here we focus on Bayesian recursive filters for models with noisy count observations. These generalise the Kalman filter [23] and have been successfully applied to similar problems in phylodynamics and computational biology [29], [30], [31]. We reconsider our renewal model inference problem within this engineering state-observation framework, as described in Eq. (3) [14].

$$R_s = f_s(R_{s-1}, \epsilon_{s-1}), \quad I_s = g_s(R_s, \nu_s) \tag{3}$$

Here R_s is the hidden Markov state that we wish to infer. It dynamically depends on the previous state R_{s-1} and a noise term ϵ_{s-1} via f_s. Observation I_s is then elicited due to R_s and a noise term ν_s, according to g_s [32].

We develop our filter under two very mild assumptions. First, we define some closed space, $\mathcal{R}$, over which R_s is valid. For a given resolution m, extrema $R_{\min}$ and $R_{\max}$, and grid size $\delta = m^{-1}(R_{\max} - R_{\min})$ then $\mathcal{R} := \{R_{\min}, R_{\min} + \delta, \ldots, R_{\max}\}$. This means the instantaneous reproduction number R_s must take a discrete value in $\mathcal{R}$, the i^{th} element of which is denoted $\mathcal{R}[i]$. We formalise this notion in Eq. (4a).

$$\sum_{i=1}^m \mathbb{P}(R_s = \mathcal{R}[i]) = 1, \, 1 \leq s \leq t \tag{4a}$$

$$R_s = R_{s-1} + (\eta\sqrt{R_{s-1}})\,\epsilon_{s-1} \tag{4b}$$

This is not restrictive since we can compute our filter for large m if needed and usually we are only interested in R_s on a coarse scale (e.g., policymakers may only want to know if $R_s \leq 1$ or not). Other approaches, which

depend on MCMC or related sampling methods (e.g., [9] and some implementations of EpiEstim), all implicitly assume some sort of discretisation [29]. In Fig. A1 we show that often convergence occurs at small m.

Second, we propose a linear model for f_s, as defined in Eq. (4b). There ϵ_{s-1} is a standard white noise term i.e., $\mathbb{P}(\epsilon_{s-1}) \stackrel{d}{=} \mathrm{Norm}(0,1)$ with Norm signifying a normal distribution and η as a free parameter. We assume that a noisy linear projection of states over consecutive time-points provides a good approximation of the state trajectory. Not only is this assumption standard in engineering [23] and epidemiology [33] but it is also more flexible than the state model inherent to EpiEstim and the WT method. We scale the noise of this projection by a fraction, $\eta < 1$ of the magnitude of R_{s-1}. This parameter controls the correlation among successive instantaneous reproduction numbers (and hence the state noise) but ensures R_s is a-priori non-negative.

Our observation model, g_s, is implicit and leads to the probability law in Eq. (1). As a result, both our observations and state models are discrete (see Fig. 1 for summaries). Because the state model governing R_s in Eq. (4b) is stochastic, Eq. (1) actually describes an over-dispersed (doubly stochastic) Poisson incidence curve. Consequently, η, allows us to better model some of the heterogeneity in transmissibility, and may increase robustness to violations of the well-mixed assumption inherent to renewal models [21]. We can optimise our choice of η value by minimising the incidence one-step-ahead predictions that result from our observation model [13].

We now define the Bayesian recursive filtering procedure, which is a main contribution of this work, and can be solved exactly, in real-time and with minimal computational effort. We adapt general recursive filtering equations from [32], [14], [25], [18], which are valid for various types of observation and state models, to our renewal model inference problem. The proof of the equations we employ can be found in these works. While we solve discrete, univariate problems (our state model is one dimensional), extensions to continuous-time, multivariate problems also exist [30], [23]. These recursive equations can also be approximately solved using particle filters [32], [14].

Recursive filtering involves two steps: prediction and correction. The first, given in Eq. (5a), constructs a sequential prior predictive distribution, $p_s = \mathbb{P}(R_s \mid I_1^{s-1})$. This is informed by past incidence data I_1^{s-1} and the last state R_{s-1}. The second step then corrects or updates this prior prediction into a posterior filtering distribution, $\mathbf{p_s}$, which constrains p_s using the latest observation, I_s, according to Eq. (5b).

$$p_s = \int \mathbb{P}(R_s \mid R_{s-1}, I_1^{s-1})\, \mathbf{p_{s-1}}\, \mathrm{d}R_{s-1} \tag{5a}$$

$$\mathbf{p_s} \propto \mathbb{P}(I_s \mid R_s, I_1^{s-1})\, p_s \tag{5b}$$

Here $\mathbb{P}(R_s \mid R_{s-1}, I_1^{s-1}) \stackrel{d}{=} \mathrm{Norm}(R_{s-1}, \eta^2 R_{s-1})$ is the state model from Eq. (4b), $\mathbb{P}(I_s \mid R_s, I_1^{s-1})$ is the observation model from Eq. (1) and the constant of proportionality for Eq. (5b) is simply a normalising factor.

Solving Eq. (5) iteratively and simultaneously over the grid of $\mathcal{R}$ leads to our novel real-time estimate of the time-varying effective reproduction number. We initialise this process with a uniform prior distribution over $\mathcal{R}$ for $\mathbf{p_1}$ and

note that p_s and $\mathbf{p_s}$ are m element vectors that sum to 1, with i^{th} term corresponding to when $R_s = \mathcal{R}[i]$. Eq. (5) forms the first half of EpiFilter, is flexible and can be adapted to many related problems [14]. A key difference between EpiFilter and the EpiEstim-type methods [11], [13] is that the latter approximate the distributions $\mathbb{P}(R_s \mid I_1^s)$ and $\mathbb{P}(R_s \mid I_1^{s-1})$ with $\mathbb{P}(R_s \mid I_{s-k+1}^s)$ and $\mathbb{P}(R_s)$, respectively. Estimators based on these approximations can be suboptimal, especially when data (i.e., cases) are scarce.

Bayesian (backward) recursive smoothing

While Eq. (5) provides a complete real-time solution to the filtering problem, it is necessarily limited at the starting edge of the incidence curve, where past data are limited or unavailable. Further, because it does not update past estimates as new data accumulate, it cannot provide optimal retrospective estimates. Here we develop the second half of EpiFilter, which involves solving the optimal smoothing problem i.e., computing the smoothing posterior distribution $\mathbf{q_s} = \mathbb{P}(R_s \mid I_1^t)$, which provides maximally informed estimates of the instantaneous reproduction number R_s given the complete incidence curve I_1^t. To our knowledge, smoothing has not yet been explicitly considered in infectious disease epidemiology (either exactly or approximately).

We specialise the general methodology from [14], [25] to obtain the recursive smoother of Eq. (6a). This equation uses the filtering distribution, $\mathbf{p_s} = \mathbb{P}(R_s \mid I_1^s)$ and the predictive distributions $p_{s+1} = \mathbb{P}(R_{s+1} \mid I_1^s)$, which we obtain from Eq. (5). Our state model means that $\mathbb{P}(R_{s+1} \mid R_s, I_1^s) \overset{d}{=} \text{Norm}(R_s, \eta^2 R_s)$.

$$\mathbf{q_s} = \mathbf{p_s} \int \mathbb{P}(R_{s+1} \mid R_s, I_1^s) \, \mathbf{q_{s+1}} \, p_{s+1}^{-1} \, \mathrm{d}R_{s+1} \tag{6a}$$

$$\mathbf{q_s} \propto \mathbf{r_s} \, \mathbf{p_s} \, \mathbb{P}(R_s)^{-1}, \quad \text{if } \mathbf{r_s} \approx \mathbb{P}(R_s \mid I_{s+1}^t) \tag{6b}$$

We realise Eq. (6a) exactly by taking a forward-backward algorithmic approach (this is the backward pass whereas Eq. (5) is the forward one). We solve this equation by noting that $\mathbf{q_t} = \mathbf{p_t}$ and iterating backwards in time to obtain the first smoothing distribution $\mathbf{q_1}$. The integrals become sums over our grid $\mathcal{R}$ and distributions are m element vectors. Eq. (6a) sequentially updates our earlier filtering solutions to include future data, forms the second half of EpiFilter and can also be approximately solved using particle smoothers [14].

This approach neatly links the filtering and smoothing distributions $\mathbf{p_s}$ and $\mathbf{q_s}$. If we assume that the reverse-filtering distribution $\mathbf{r_s}$ is reasonably approximated by $\mathbb{P}(R_s \mid I_{s+1}^t)$ then we can also apply the two-filter smoothing solution of [25] to get Eq. (6b). If either future (I_{s+1}^t) or past (I_1^{s-1}) incidence is uninformative then either $\mathbf{r_s}$ or $\mathbf{p_s}$ will reduce to the prior $\mathbb{P}(R_s)$, leading to $\mathbf{q_s} \propto \mathbf{p_s}$ or $\mathbf{q_s} \propto \mathbf{r_s}$, respectively. The end and beginning of the epidemic provide important examples of each of these scenarios. Consequently, Eq. (6b) shows how smoothing connects EpiEstim and the WT methods, and explains why EpiFilter, which can be used for both real-time and retrospective inference, better overcomes edge-effects and periods of low data.

Further, the smoothed posterior $\mathbf{q_s}$ yields the conditional mean estimate $\hat{R}_s = \mathbb{E}[R_s \,|\, I_1^t]$, which is known to significantly improve on the MSE of the filtered equivalent $\tilde{R}_s$ (see previous section) [23]. While filtering provides the minimum MSE estimator of every instantaneous reproduction number R_s given past knowledge, smoothing provides the minimum given all knowledge. This relationship is formal, with filtered and smoothed MSE values mapping to the amount of mutual information that I_1^t provides about R_1^t [34]. Extracting the maximum information from the incidence curve I_1^t should engender estimates that are more robust and statistically efficient in periods of low incidence. We summarise the EpiFilter algorithm in Fig. 1.

While our main interest is on optimised and rigorous real-time and retrospective estimates of transmissibility, which are completely defined by the smoothing distribution $\mathbf{q_s}$, we may also want to predict future incidence, for informing epidemic preparedness plans and for validating past R_s estimates [35], [13]. We compute the filtered one-step-ahead posterior predictive distribution as in Eq. (7) (integrals are over $\mathcal{R}$ [14].

$$\mathbb{P}(I_{s+1} \,|\, I_1^s) = \int \mathbb{P}(I_{s+1} \,|\, R_s, \, I_1^s) \, \mathbf{p_s} \, \mathrm{d}R_s \tag{7}$$

We assume, as in [36], that $\mathbb{P}(I_{s+1} \,|\, R_s, \, I_1^s) \overset{d}{=} \mathrm{Pois}(\Lambda_{s+1} R_s)$. Replacing $\mathbf{p_s}$ with $\mathbf{q_s}$ yields the smoothed equivalent of Eq. (7). We will use Eq. (7) to compare EpiFilter against APEestim, which is the prediction-optimised version of EpiEstim, developed in [13]. Since predictions depend strongly on $\mathbf{p_s}$ or $\mathbf{q_s}$, optimising these distributions can be crucial, for example, to forecasting second waves of infection.

RESULTS

Improved estimation at low incidence

The reliable estimation of time-varying effective reproduction numbers, R_s, at low incidence, I_s, is a key challenge limiting our understanding of transmission [17]. Periods with small counts of new cases contain little information and so present necessary statistical difficulties [26]. Here we compare EpiFilter, which allows exact inference over a state grid $\mathcal{R}$, with EpiEstim and APEestim at these data-poor settings. We use EpiEstim with weekly and monthly windows. Weekly windows are the default recommendation in [11]. We include monthly ones as long windows can improve robustness at low incidence [28]. APEestim, developed in [13], optimises the window choice of EpiEstim to minimise one-step-ahead prediction errors. The assumptions and choices inherent in estimation methods become important and visible when data are scarce, and can bias inference or support spurious predictions [15].

We generate epidemics using Eq. (1) under the serial interval distribution of Ebola virus disease described in [37]. We examine three diverse scenarios in Fig. 2. These describe (A) rapidly controlled epidemics (R_s step changes from 2 to 0.5 at $s = 100$), (B) small outbreaks with exponentially rising and falling R_s (change-point at $s = 30$ and rates of 0.02 and -0.008 per time unit) and (C) medium outbreaks that are initially controlled (R_s changes

from 4 to 0.6 at $s = 40$) then resurge (R_s rebounds to 2 at $s = 80$) into large epidemics, before finally being

suppressed ($R_s = 0.2$ from $s = 150$). These scenarios are similar to ones investigated in [28], [12] and describe

various epidemic dynamics that culminate in elimination. We simulate 200 incidence curves for each scenario and

apply APEestim (optimal window k^*), EpiEstim ($k = 7$ and 31) and EpiFilter (state noise $\eta = 0.1$) to estimate R_s

and sequentially predict I_s (given data up to $s - 1$) for each curve.

For the first two methods we compute mean one-step-ahead incidence predictions ($\tilde{I}_{\tau(s)}$) as in [13] and instanta-

neous reproduction number estimates ($\tilde{R}_{\tau(s)}$) from Eq. (2) with $\tau(s)$ delimiting the window times used. We obtain

smoothed EpiFilter estimates ($\hat{R}_s$) and filtered predictions ($\tilde{I}_s$) from Eq. (6) and Eq. (7). We could use smoothed

predictions ($\hat{I}_s$), which implicitly include data beyond $s - 1$, to assess fit but as we want to test model adequacy $\tilde{I}_s$

is more appropriate [38]. Our main focus is on real-time performance so we do not investigate the WT method. See

[11], [12] for comparisons of the WT method and EpiEstim. Panels A-C of Fig. 2 provide representative single runs

(we only show $k = 31$ for EpiEstim as $k = 7$ is often similar to APEestim) while panel D presents reproduction

number MSE and one-step-ahead predicted incidence MSE (PMSE) distributions from the 200 runs. The left panels

of Fig. A3 in the Appendix provide distributions of the coverage statistics from these runs (i.e., the probability that

the true R_s or I_s is contained within estimated or predicted 95% confidence intervals).

In the Methods we showed that since EpiEstim-type methods group data over some window into the past, they

revert to their prior distribution as the total cases in this window becomes small. During low incidence periods, e.g.,

when the epidemic is waning, using long windows makes sense [28]. However, this reduces predictive accuracy as

fluctuations in R_s are underfit. This is the use-case for APEestim, which optimises for prediction. The consequences

of these trade-offs are made clear in A-C of Fig. 2 where APEestim provides the best one-step-ahead I_s predictions

but quickly reverts to its prior (seen as wide confidence intervals) when cases are few. Longer window EpiEstim

improves on this destabilisation but cannot track salient fluctuations in R_s. Panel D of Fig. 2 confirms these trends.

APEstim has the smallest PMSE but often worse MSE than the larger k EpiEstim choices. However, as seen in

Fig. A3, both EpiEstim approaches have smaller coverage than APEestim.

Interestingly, EpiFilter is able to jointly optimise both instantaneous reproduction number tracking and incidence

predictions (computed via Eq. (7)). In A-C of Fig. 2 we see that EpFilter maintains stable and accurate estimates of

R_s and only slowly reverts to its prior (which has the same support as that of EpiEstim and APEesim). However,

unlike long window methods it does not sacrifice prediction fidelity. The improvement in MSE shown in panel

D of Fig. 2 is stark (the numerical reduction in MSE when compared to the next best method is on average at

least 2-fold and often 10-fold). The PMSE, while larger than that of APEestim (which is optimised for predictions)

is still good. Moreover, the coverage of both the true R_s and I_s of EpiFilter is superior to all other approaches

as shown in Fig. A3. EpiFilter therefore combines the advantages of APEestim and long-window EpiEstim and,

further, is able to detect transmission change-points automatically.

Fig. 2: **Small or waning epidemics.** We compare reproduction number estimates ($\tilde{R}_{\tau(s)}$ or $\hat{R}_s$) and one-step-ahead incidence predictions ($\tilde{I}_{\tau(s)}$ or $\tilde{I}_s$) from APEestim with optimal window k^*, EpiEstim with window k and EpiFilter with state noise η. We simulate 200 epidemics with low daily case numbers or long tails (long sequences of zero cases) using the standard renewal model (Eq. (1)) for three scenarios, representative examples of which are given in A-C. The true R_s and I_s are in black. All mean estimates or predictions are in red and blue with 95% confidence intervals. APEestim and EpiEstim use a $\mathrm{Gam}(1, 2)$ prior distribution and EpiFilter a grid with $m = 2000$, $R_{\min} = 0.01$ and $R_{\max} = 10$. In D we provide statistics of the MSE of these estimates (relative to R_s) and the PMSE of these predictions (relative to I_s) for all 200 runs. We find that EpiFilter is more robust to small incidence (better uncertainty), whereas the other approaches can quickly decay to their prior distribution. It achieves significantly smaller MSE (2-10 fold reductions), comparable PMSE to APEestim (which is optimised for prediction) and superior coverage of the true R_s and I_s (see Fig. A3).

Fig. 3: **Epidemics with multiple waves.** We compare reproduction number estimates ($\tilde{R}_{\tau(s)}$ or $\hat{R}_s$) and one-step-ahead incidence predictions ($\tilde{I}_{\tau(s)}$ or $\tilde{I}_s$) from APEestim with optimal window k^*, EpiEstim with window k and EpiFilter with state noise η. We simulate 200 epidemics with multiple waves of infection using the standard renewal model (Eq. (1)) for three scenarios, representative examples of which are given in A-C. The true R_s and I_s are in black. All mean estimates or predictions are in red and blue with 95% confidence intervals. APEestim and EpiEstim use a $\mathrm{Gam}(1, 2)$ prior distribution and EpiFilter a grid with $m = 2000$, $R_{\min} = 0.01$ and $R_{\max} = 10$. In D we provide statistics of the MSE of these estimates (relative to R_s) and the PMSE of these predictions (relative to I_s) for all 200 runs. We find EpiFilter is best able to negotiate troughs between epidemic peaks and hence infer resurging infectious dynamics, achieving significantly smaller MSE (2-10 fold reductions), comparable PMSE to APEestim (which is optimised for prediction) and superior coverage of the true R_s and I_s (see Fig. A3).

Improved estimation between epidemic waves

Maintaining robust instantaneous reproduction number, R_s, estimation when incidence, I_s, becomes small is not just statistically important. Two possible outcomes may follow periods of small I_s: either the epidemic goes extinct (elimination occurs, as in the previous section), or an additional wave of infection surfaces (resurgence) e.g., due to imports or unmonitored local transmission. Predicting which outcome is likely, rapidly and in real-time, is of major global concern as countries aim to relax interventions during the ongoing COVID-19 pandemic, while also minimising the risks of further resurgence [17], [39]. As changes in instantaneous reproduction numbers predate and signal corresponding variations in incidence, reliably identifying and inferring R_s trends in the trough preceding potential new peaks is crucial for preparedness and both timely and effective epidemic control [10].

Reliable estimation of R_s between epidemic waves depends on the prior assumptions of the inference method used and on how that method relies on those assumptions when data are scarce [40], [41]. Here we examine this dependence and investigate cases where resurgence follows a low-incidence period. As in the above section, we compare EpiFilter ($\eta = 0.1$) with APEestim (optimal window k^*) and EpiEstim (weekly, $k = 7$, and monthly, $k = 31$, windows) over 200 simulated epidemics under the serial interval of Ebola virus. We explore scenarios depicting (A) epidemics that are initially controlled (R_s falls from 2.5 to 0.5 at $s = 70$) but which resurge just as quickly (R_s returns to 2.5 from $s = 230$), (B) periodic or seasonal transmission (R_s is sinusoidal with magnitude 1.3 $\pm$1.2 and period of 120 time units) and (C) outbreaks with exponentially rising and then falling transmissibility (change-points at $s = 40$ and 190 and exponent rates 0.03, -0.015 and 0.02).

These examples are similar to some in [13] and describe diverse epidemics with multiple peaks and troughs. We provide representative runs of each scenario in A-C and collect MSE (for R_s) and PMSE (relative to I_s) distributions over the 200 runs of every scenario in D of Fig. 3. Coverage distributions are shown in right panels of Fig. A3 of the Appendix. We observe a similar pattern in performance among the methods as in our previous analysis of Fig. 2. APEestim is best able to predict upcoming incidence and achieves the best PMSE as expected. While the monthly window EpiEstim is less useful in these cases (since prior reversion does not occur as often, though could be an issue for more extended troughs), the weekly window version loses predictive fidelity for minor MSE improvements. Its coverage is, however, generally worse than that of APEestim.

EpiFilter once again combines the advantages of the other approaches. For every scenario in Fig. 3 it provides accurate tracking of changes in R_s with stable confidence intervals and a MSE that is at least $\frac{1}{2}$ and sometimes even $\frac{1}{10}$ that of the next best method. Concurrently, its incidence PMSE rivals that of APEestim. Importantly, it also achieves the best coverage of the true R_s and I_s values. EpiFilter is therefore a powerful tool for detecting resurgence. We also find that the $\eta = 0.1$ parameter value seems to be an all-purpose heuristic, meaning that usage of EpiFilter can be somewhat simpler than EpiEstim and other window or change-point based methods. The

superior capability of EpiFilter in these scenarios likely results from its minimalistic assumptions (see Eq. (4)) and its increased information extraction. We next test our method on empirical incidence data.

COVID-19 in New Zealand and H1N1 influenza in the USA

The previous sections confirmed EpiFilter as a powerful inference and prediction tool, especially in data-poor conditions, using simulated epidemics. We now confront our method with empirical data from the 1918 H1N1 influenza pandemic in Baltimore (USA) [42] and the ongoing COVID-19 pandemic in New Zealand (up to 17 August 2020) [43]. The H1N1 dataset has been well-studied and so we first use this to benchmark EpiFilter. We clean this dataset by applying a 5-day moving average filter as recommended in [42]. Previous work [11] analysed this dataset with EpiEstim and found that sensible instantaneous reproduction number, R_s, estimates are obtained when a weekly window ($k = 7$) is applied. However, more recent work, using APEestim [13], showed that while $k = 7$ provides stable estimates for this epidemic, it is a poor predictor of the incidence data. Instead, an optimised window of 2 days ($k^* = 2$) yields good predictions but the resulting R_s estimates are noisy.

We reproduce the instantaneous reproduction number estimates ($\tilde{R}_{\tau(s)}$) and incidence predictions ($\tilde{I}_{\tau(s)}$) from both studies in Fig. 4 and compare them against EpiFilter with $\eta = 0.1$ ($\hat{R}_s$ and $\tilde{I}_s$). Top and middle rows of Fig. 4 illustrate the aforementioned trade-off between estimate stability and prediction accuracy. The bottom row confirms the power of EpiFilter. Our R_s estimates are of comparable stability to those of EpiEstim at $k = 7$, yet our prediction fidelity matches that of APEestim. Our improved inference again benefits from using more information (i.e., the backward pass in Fig. 1) and making less restrictive prior assumptions. We see the latter from the R_s confidence intervals over $40 \leq s \leq 60$. There EpiEstim seems overconfident, and this results in a rigid overestimation of incidence. However, EpiFilter mediates its estimate uncertainty to a level similar to APEestim.

We explore COVID-19 transmission patterns in New Zealand using incidence data up to 17 August 2020 from [43]. New Zealand presents an insightful case study because officials combined swift lockdowns with intensive testing to achieve and sustain very low incidence levels that eventually led to local elimination of COVID-19 [44]. However, an upsurge in cases in early August inspired concerns about a second wave (which led to new interventions and is why we do not consider data beyond 17 August). Here we investigate time-varying transmission in New Zealand to see if this uptick suggests that the epidemic was resurfacing in mid-August. We believe smoothing can confer important inferential advantages in exactly these types of low incidence scenarios.

We make the common assumptions that case under-reporting is constant [11], which seems reasonable given the intensive surveillance employed by New Zealand [45]. We ignore reporting delays, which are known to be small [46] and use the COVID-19 serial interval distribution from [47]. We do not explicitly distinguish imported from local cases in our analysis. The latter could bias our study [28], [39] but our focus is on demonstrating differences

Fig. 4: **H1N1 influenza transmission in Baltimore (1918).** We compare APEestim (top), EpiEstim with recommended weekly window (middle) (both with $\mathrm{Gam}(1,2)$ prior distribution) and EpiFilter (with $m = 2000$, $\eta = 0.1$, $R_{\min} = 0.01$ and $R_{\max} = 10$) on the H1N1 influenza dataset from [42]. We use a 5-day moving average filter, as in [42], to remove known sampling biases. Estimates of reproduction numbers, R_s, and corresponding 95% confidence intervals are in red. One-step-ahead predictions of incidence, I_s, (with 95% confidence intervals) are in blue with the actual incidence in black. We find that EpiFilter combines the benefits of APEestim and EpiEstim, achieving both good estimates and predictions.

between filtering and smoothing on instantaneous reproduction number, R_s, trends and not on providing detailed R_s estimates during this period. We plot the results of our exploration in Fig. 5. More involved analyses can be performed by pre-processing the data if introductions and case ascertainment fractions are known [12], [6].

We apply EpiFilter and obtain filtered ($\tilde{R}_s$) and smoothed ($\hat{R}_s$) conditional mean estimates together with their 95% confidence intervals. These are in the left panels of Fig. 5 and computed from Eq. (5) and Eq. (6) respectively. The times of lockdown and release are included for reference. Interestingly, we see a notable difference in the quality of inference between $\tilde{R}_s$ and $\hat{R}_s$. The former, as expected, is unreliable at the beginning of the incidence curve and features wider uncertainty and noisier trends. The smoothed $\hat{R}_s$, by using both forward and backward-looking data largely overcomes these issues and clarifies transmission dynamics. In the right panels we compare EpiFilter with EpiEstim (weekly windows) and APEestim. The difference is striking. Neither APEestim nor EpiEstim recovers a clear trend and both are appreciably worse than even the filtered estimate $\tilde{R}_s$.

Our smoothed analysis suggests that R_s has resurged and supports the re-implementation of measures around 14 August. We recover suppression of the initial wave in April, likely associated with the implementation of key interventions, including lockdown [46]. Following this, we infer a prolonged period of subcritical transmission, where $\mathbb{P}(R_s \leq 1) \approx 1$. Low and then zero incidence over this period strikingly destabilises both APEestim and

Fig. 5: **COVID-19 transmission in New Zealand.** We compute smoothed and filtered reproduction number estimates, $\hat{R}_s$ and $\tilde{R}_s$ respectively, from the COVID-19 incidence curve for New Zealand (available at [43]) in the left panels. We use EpiFilter with $m = 2000$, $\eta = 0.1$, $R_{min} = 0.01$ and $R_{max} = 10$ with a uniform prior distribution over the grid $\mathcal{R}$. The top sub-figure shows conditional mean estimates and 95% confidence intervals for $\hat{R}_s$ (red) and $\tilde{R}_s$ (blue). Vertical lines indicate the start and end of lockdown, a major intervention that was employed to halt transmission. The additional 'future' information used in smoothing has a notable effect. The bottom sub-figure provides smoothed one-step-ahead predictions $\hat{I}_s$ (blue, with 95% confidence) of the actual reported cases I_s (black). The inset gives the estimated probability of $R_s \leq 1$. We observe a clear trend of subcritical transmission that eventually seeds a second wave by August. In the right panels we compare EpiFilter with EpiEstim (using weekly windows) and APEestim (both with $\mathrm{Gam}(1, 2)$ priors) with all left subfigures presenting R_s estimates and right ones providing filtered I_s predictions. We observe that both APEestim and EpiEstim lead to largely unusable estimates that mask transmission trends, in stark contrast to EpiFilter.

389 EpiEstim and conceals the rise in R_s that occurs after July. EpiFilter provides early-warning of this upsurge, which

390 continually grows until August, where a second wave becomes likely. We also provide one-step-ahead predictions

391 (which are from the smoothed $\hat{R}_s$ hence the notation $\hat{I}_s$) and their 95% confidence intervals against the reported

392 incidence from [43]. These verify that $\hat{R}_s$ reasonably describes the data (we also present $\tilde{I}_s$ when comparing

393 methods). We find that $\mathbb{P}(R_s \leq 1) \approx 0$, further supporting this resurgence hypothesis.

394   DISCUSSION

Estimating time-varying trends in the instantaneous or effective reproduction number, R_s, reliably and in real-

time is an important and popular problem in infectious disease epidemiology [5]. As the COVID-19 pandemic

has unfolded, the interest in solving this problem has only elevated with R_s playing a central role both in aiding

situational awareness [8] and informing policymaking [48]. Initially, interest was on understanding how changes

in R_s may correlate with interventions such as lockdowns and social distancing [10], [9]. However, as countries

have entered waning phases of the pandemic and vaccine deployment has begun, focus has shifted to charactering

how existing interventions can be relaxed with minimum risk [49]. The literature on intervention exit strategies is, however, still in development, and several challenges remain to modelling transmission.

One key challenge lies in understanding and inferring transmissibility during periods when the incidence of new cases is small [17]. Such periods may occur under sustained control measures and necessarily contain limited data, which make inferences difficult. Moreover, it is in these lulls that information on transmission may be crucial, helping to determine if the removal of interventions will lead to resurgence or if elimination is realistic by maintaining controls [28], [39]. While reproduction numbers are not the only analytic for assessing these outcomes, they do provide an important real-time diagnostic since upticks in R_s generally precede elevations in case loads. Unfortunately, current approaches to estimating R_s become underpowered, unstable or prior-constrained in these data-limited conditions [11], [26], [49]. These problems are only magnified when finer-scale analyses (where cases are fewer by division) are of interest (e.g., regional versus national level estimation).

In this paper we re-examined existing methodology for inferring instantaneous reproduction numbers, R_s, from an engineering perspective. We observed that two of the most useful and popular inference approaches, EpiEstim [11] and the WT method (this computes cohort reproduction numbers, which are functions of R_s) [2], only capitalise on a portion of the data available, deeming either upcoming or past incidence to be informative (see Fig. 1) [12]. This informative portion is directly controlled by prior assumptions on the speed of possible R_s changes, which are often characterised by a window of size k. Other methods also apply similarly strong change-point or state assumptions on R_s, explicitly linking its variations with specific dates or events, for example [26], [9]. When data are scarce these assumptions can unduly control or bias inference.

In control engineering a common problem, known as filtering, involves optimally (in a MSE sense) estimating hidden Markov states, in real-time, from noisy and uncertain observations [18]. A related problem termed smoothing provides accompanying and optimal retrospective inferences [14]. By reinterpreting R_s as a Markov state (Eq. (4)) observed through a noisy renewal process (Eq. (1)) and defining R_s on a predetermined grid $\mathcal{R}$, we were able to construct exact filtering (Eq. (5)) and smoothing (Eq. (6)) solutions. This led to EpiFilter, which is our central contribution. Generally, filtering and smoothing can be involved and require sophisticated sequential Monte Carlo techniques [32]. However, because we make only minimal assumptions about R_s, modelling it as a simple diffusion, we were able to solve these problems exactly and without complex sampling algorithms [29].

Our solutions are computationally simple, often executing in a few minutes (see Fig. A1), and deterministic i.e., precisely reproducible given the same data and settings. Our method replaces strong change-point or window size assumptions with one free parameter, η, which allows us to model some heterogeneity in transmission and sets the correlation among successive R_s values. We find that $\eta = 0.1$ serves as a general heuristic, providing good estimates and automatically detecting change-points and salient R_s dynamics over diverse scenarios (see Fig. 2,

Fig. 3 and Fig. A2). This heuristic is also statistically justified by its good one-step-ahead predictive performance and its coverage of true simulated values (see Fig. A3) [13].

Importantly, EpiFilter is able to look both forward and backward through the incidence data, and so maximise the information extracted at every time point [34]. This property means it combines advantages from both EpiEstim and the WT method (see Fig. 1) and largely ameliorates their edge-effect issues [12]. These benefits, which also hold at large incidence, make EpiFilter a useful and robust tool for both real-time and retrospective R_s inference. We confirmed the advantages of EpiFilter by comparing it to EpiEstim and APEestim (a prediction optimised analogue to EpiEstim) on many simulated examples with periods of low incidence and epidemic resurgences (Fig. 2 and Fig. 3). Interestingly, we found EpiFilter was able to achieve significant 2-10 fold reductions in the MSE of R_s estimates without compromising predictive power. Further, its estimate and prediction confidence intervals contained the true R_s and I_s values with the highest probabilities (see Fig. A3).

EpiFilter was especially better at negotiating periods of low incidence, offering a graceful degradation to its prior distribution or assumptions without sacrificing predictive accuracy. When incidence is low, it can be beneficial to use longer windows with EpiEstim [28]. This keeps R_s estimates reasonably stable but often leads to poor predictions [13]. APEestim, which optimises window size for prediction fidelity, showed that in many of the simulated scenarios short windows are necessary for describing transmission patterns. Consequently, we have a trade-off between estimate robustness and prediction accuracy. We found that EpiFilter overcomes this trade-off, concurrently achieving good estimates and predictions. In doing so, it revealed subcritical transmission trends and unmasked important signals of resurgence from noisy data in those periods.

We verified the practical utility and performance of EpiFilter on empirical data from the H1N1 pandemic of 1918 (see Fig. 4) and COVID-19 in New Zealand (see Fig. 5). In the first, which is a standard dataset that has been used to test previous R_s methods, we found that EpiFilter integrated the benefits of EpiEstim and APEestim to achieve simultaneously good estimates and predictions. A key use-case for EpiFilter is in signalling resurgence during low incidence. The COVID-19 epidemic in New Zealand featured precisely those dynamics [46]. While EpiEstim and APEestim were destabilised and unable to extract clear transmission trends, EpiFilter inferred subcritical R_s values and forewarned of resurgence by signalling an uptick in R_s just before a second wave become apparent. Recent, more involved COVID-19 analyses [39], have confirmed EpiFilter as a useful early-warning method.

Balancing the assumptions inherent to a model against the data it is applied on, to produce reliable inference is a non-trivial problem that is still under active investigation in several fields [15], [40], [41]. EpiFilter, by maximising the information extracted from available incidence data and minimising its state space model assumptions, appears to strike this balance as an estimator of instantaneous or effective reproduction numbers. Consequently, it performs strongly on a wide range of problems, including those involving sparse data, where other methods might struggle.

Given its demonstrated advantages, straightforward formulation and theoretical underpinning, we hope that EpiFilter will be useful as a diagnostic tool for reliably warning about second waves of infection over multiple scales and more generally for assessing dynamical patterns in transmission both in real time and retrospectively. EpiFilter is freely available at https://github.com/kpzoo/EpiFilter.

ACKNOWLEDGMENTS

Thanks to Christl A Donnelly for thoughtful comments.

FUNDING

This work is jointly funded under grant reference MR/R015600/1 by the UK Medical Research Council (MRC) and the UK Department for International Development (DFID) under the MRC/DFID Concordat agreement and is also part of the EDCTP2 programme supported by the European Union.

REFERENCES

[1] C. Fraser, "Estimating individual and household reproduction numbers in an emerging epidemic," *PLOS One*, vol. 8, p. e758, 2007.

[2] J. Wallinga and P. Teunis, "Different epidemic curves for severe acute respiratory syndrome reveal similar impacts of control measures," *Am. J. Epidemiol*, vol. 160, no. 6, pp. 509–16, 2004.

[3] H. Nishiura and G. Chowell, "The effective reproduction number as a prelude to statistical estimation of time-dependent epidemic trends," in *Mathematical and statistical estimation approaches in epidemiology*, pp. 103–21, Springer, 2009.

[4] R. Anderson and R. May, *Infectious diseases of humans: dynamics and control*. Oxford University Press, 1991.

[5] S. Cauchemez, P. Boelle, G. Thomas, *et al.*, "Estimating in real time the efficacy of measures to control emerging communicable diseases," *Am. J. Epidemiol*, vol. 164, no. 6, pp. 591–7, 2006.

[6] T. Churcher, J. Cohen, N. Ntshalintshali, *et al.*, "Measuring the path toward malaria elimination," *Science*, vol. 344, no. 6189, pp. 1230–32, 2014.

[7] WHO Ebola Response Team, "Ebola virus disease in West Africa – the first 9 months of the epidemic and forward projections," *N. Engl. J. Med*, vol. 371, no. 16, pp. 1481–95, 2014.

[8] S. Bhatia, A. Cori, K. Parag, *et al.*, "Short-term forecasts of COVID-19 deaths in multiple countries.."

[9] S. Flaxman, S. Mishra, A. Gandy, *et al.*, "Estimating the effects of non-pharmaceutical interventions on COVID-19 in Europe," *Nature*, vol. 584, pp. 257–261, 2020.

[10] X. Hao, S. Cheng, D. Wu, *et al.*, "Reconstruction of the full transmission dynamics of COVID-19 in Wuhan," *Nature*, vol. 584, pp. 420–4, 2020.

[11] A. Cori, N. Ferguson, C. Fraser, *et al.*, "A new framework and software to estimate time-varying reproduction numbers during epidemics," *Am. J. Epidemiol*, vol. 178, no. 9, pp. 1505–12, 2013.

[12] K. Gostic, L. McGough, E. Baskerville, *et al.*, "Practical considerations for measuring the effective reproductive number, rt," *PLOS Comput. Biol*, vol. 16, no. 12, p. e1008409, 2020.

[13] K. Parag and C. Donnelly, "Using information theory to optimise epidemic models for real-time prediction and estimation," *PLOS Comput. Biol*, vol. 16, no. 7, p. e1007990, 2020.

[14] S. Sarrka, *Bayesian Filtering and Smoothing*. Cambridge, UK: Cambridge University Press, 2013.

[15] K. Parag, O. Pybus, and C. Wu, "Are skyline plot-based demographic estimates overly dependent on smoothing prior assumptions?," *BioRxiv*, vol. 920215, 2020.

[16] T. Britton, T. House, A. Lloyd, *et al.*, "Five challenges for stochastic epidemic models involving global transmission," *Epidemics*, vol. 10, pp. 54–7, 2015.

[17] R. Thompson, D. Hollingsworth, V. Isham, *et al.*, "Key questions for modelling COVID-19 exit strategies," *Proc. R. Soc. B*, vol. 287, no. 1932, p. 20201405, 2020.

[18] K. Astrom and R. Murray, *Feedback Systems: An Introduction for Scientists and Engineers*. New Jersey: Princeton University Press, 2008.

[19] S. Riley, K. Eames, V. Isham, *et al.*, "Five challenges for spatial epidemic models," *Epidemics*, vol. 10, no. 68-71, 2015.

[20] J. Wallinga and M. Lipsitch, "How generation intervals shape the relationship between growth rates and reproductive numbers," *Proc. R. Soc. B*, vol. 274, pp. 599–604, 2007.

[21] Q. Liu, M. Ajelli, A. Aleta, *et al.*, "Measurability of the epidemic reproduction number in data-driven contact networks," *PNAS*, vol. 115, no. 50, pp. 12680–85, 2018.

[22] T. Britton and G. Tomba, "Estimation in emerging epidemics: biases and remedies," *J. R. Soc. Interface*, vol. 16, no. 150, p. 20180670, 2019.

[23] D. Snyder and M. Miller, *Random Point Processes in Time and Space*. Springer-Verlag, 2 ed., 1991.

[24] R. Kalman, "A new approach to linear filtering and prediction problems," *Trans. ASME J. Basic Eng*, vol. 82, pp. 35–45, 1960.

[25] B. Anderson and I. Rhodes, "Smoothing algorithms for nonlinear finite-dimensional systems," *Stochastics*, vol. 9, pp. 139–65, 1983.

[26] K. Parag and C. Donnelly, "Adaptive estimation for epidemic renewal and phylogenetic skyline models," *Syst. Biol*, vol. 69, no. 6, pp. 1163–79, 2020.

[27] T. Rothenberg, "Identification in parametric models," *Econometrica*, vol. 39, no. 3, pp. 577–91, 1971.

[28] K. Parag, C. Donnelly, R. Jha, *et al.*, "An exact method for quantifying the reliability of end-of-epidemic declarations in real time," *PLOS Comput. Biol*, vol. 16, no. 11, p. e1008478, 2020.

[29] K. Parag and O. Pybus, "Exact bayesian inference for phylogenetic birth-death models," *Bioinformatics*, vol. 34, no. 21, pp. 3638–45, 2018.

[30] K. Parag and G. Vinnicombe, "Point Process Analysis of Noise in Early Invertebrate Vision," *PLOS Comput. Biol*, vol. 13, no. 10, p. e1005687, 2017.

[31] C. Zechner and H. Koeppl, "Uncoupled analysis of stochastic reaction networks in fluctuating environments," *PLOS Comput. Biol*, vol. 10, no. 12, p. e1003942, 2014.

[32] Z. Chen, "Bayesian filtering: From Kalman filters to particle filters, and beyond," *Statistics*, vol. 182, no. 1, pp. 1–69, 2003.

[33] L. Allen, "A primer on stochastic epidemic models: Formulation, numerical simulation, and analysis," *Infect. Dis. Model*, vol. 2, pp. 128–142, 2017.

[34] D. Guo, D. Shamai, and S. Verdu, "Mutual information and minimum mean-square error in Gaussian channels," *IEEE Trans. Info. Theo*, vol. 51, no. 4, pp. 1261–82, 2005.

[35] S. Funk, A. Camacho, A. Kucharski, *et al.*, "Assessing the performance of real-time epidemic forecasts: A case study of Ebola in the western area region of Sierra Leone, 2014-15," *PLOS Comput. Biol*, vol. 15, no. 2, p. e1006785, 2019.

[36] P. Nouvellet, A. Cori, T. Garske, *et al.*, "A simple approach to measure transmissibility and forecast incidence," *Epidemics*, vol. 22, pp. 29–35, 2018.

[37] M. Van Kerkhove, A. Bento, H. Mills, *et al.*, "A review of epidemiological parameters from Ebola outbreaks to inform early public health decision-making," *Sci. Data*, vol. 2, p. 150019, 2015.

[38] E. Wagenmakers, P. Grunwald, and M. Steyvers, "Accumulative prediction error and the selection of time series models," *J. Math. Psychol*, vol. 50, pp. 149–166, 2006.

[39] K. Parag, B. Cowling, and C. Donnelly, "Deciphering early-warning signals of the elimination and resurgence potential of SARS-CoV-2 from limited data at multiple scales," *medRxiv*, vol. 2020.11.23.20236968, 2021.

[40] E. Volz and X. Didelot, "Modeling the growth and decline of pathogen effective population size provides insight into epidemic dynamics and drivers of antimicrobial resistance," *Syst. Biol*, vol. 67, no. 4, pp. 719–28, 2018.

[41] J. Faulkner, A. Magee, B. Shapiro, *et al.*, "Horseshoe-based Bayesian nonparametric estimation of effective population size trajectories," *Biometrics*, pp. 1–14, 2020.

[42] W. Frost and E. Sydenstricker, "Influenza in Maryland: preliminary statistics of certain localities," *Public Health Rep*, vol. 34, pp. 491–504, 1919.

[43] WHO, "WHO coronavirus disease (COVID-19) dashboard," 2020.

[44] S. Cousins, " New Zealand eliminates COVID-19," *Lancet*, vol. 395, p. 1474, 2020.

[45] M. Roser, H. Ritchie, E. Ortiz-Ospina, *et al.*, "Coronavirus pandemic (COVID-19)," 2020.

[46] S. Jefferies, N. French, C. Gilkison, *et al.*, "COVID-19 in New Zealand and the impact of the national response: a descriptive epidemiological study," *Lancet Public Health*, vol. 5, no. 11, pp. e612–23, 2020.

[47] N. Ferguson, D. Laydon, G. Nedjati-Gilani, *et al.*, "Impact of non-pharmaceutical interventions (NPIs) to reduce COVID- 19 mortality and healthcare demand," tech. rep., Imperial College London, 2020.

[48] P. Bruce, R. Anderson, C. Bangham, *et al.*, "Reproduction number (R) and growth rate (r) of the COVID-19 epidemic in the UK: methods of estimation, data sources, causes of heterogeneity, and use as a guide in policy formulation," tech. rep., The Royal Society, 2020.

[49] Y. Li, H. Campbell, D. Kulkarni, *et al.*, "The temporal association of introducing and lifting non-pharmaceutical interventions with the time-varying reproduction number (R) of SARS-CoV-2: a modelling study across 131 countries," *Lancet Infect. Dis*, vol. 21, no. 2, pp. 193–202, 2020.

APPENDIX

Validating the mechanics of EpiFilter

Having constructed EpiFilter (see Methods) we now investigate its mechanics and performance. First we show how the choice of grid resolution m affects the accuracy and computational complexity of our smoothed instantaneous reproduction number estimates, $\hat{R}_s$. Our recursive implementation is also known as a grid smoother, since its estimates are optimal (in a MSE sense) but conditioned on the grid $\mathcal{R}$ [14], [29]. In the left panels of Fig. A1 we reconsider the empirical H1N1 and COVID-19 examples, from the main text, at various m. We find that EpiFilter has linear complexity in m but that $\hat{R}_s$ (averaged over 50 replicate runs) converges quickly, both in mean and variance (latter not shown). Execution times, which include both R_s estimation and incidence (I_s) prediction, are only a few minutes, generally. While the R_s smoother is always fast, the I_s prediction block may be slower for very large epidemics (e.g., 1000s of cases per day), resulting in ~ 10 minute run times.

As a rule-of-thumb setting $m \geq 10^3$ safely guarantees convergence, and still executes in a few minutes. Note that estimates at small m are still valid. They represent coarser assumptions on reproduction numbers and provide

Fig. A1: **EpiFilter mechanics.** We examine the computational complexity and accuracy of EpiFilter. Left panels indicate how increasing grid resolution, m, leads to a linear increase in execution time (blue with 95% confidence intervals) for the empirical case studies of the main text. These times are averaged over 50 replicate runs (using the R implementation) and include the latency in computing both reproduction number, R_s, estimates and incidence, I_s, predictions. The mean estimate of R_s, averaged over time and the replicates, is in grey, and converges quickly. The variance of these estimates, while not shown, converges equally fast. Right panels provide an illustrative example of how estimates $\hat{R}_s$ (red with 95% confidence intervals) change with m and choices of process noise parameter, η. Even at coarse grid sizes (small m) the estimates are sensible. Reducing η sharpens confidence intervals but can potentially underfit. Increasing η adds state noise. We find that $\eta = 0.1$ serves as a good general heuristic. All simulations are at $R_{\min} = 0.01$ and $R_{\max} = 10$ and the true R_s (a seasonal epidemic) is in black.

minimised MSE estimates under those assumptions. We illustrate a coarse estimate and how it converges for a

simulated example in the right panels of Fig. A1. There, we also test the behaviour under different choices of the

process or state noise parameter η. This parameter is vaguely similar to the window size parameter, k, in EpiEstim

approaches, but is easier to select. It controls the autocorrelation among successive R_s and allows some modelling

of heterogeneity (hence the term state noise). Fig. A1, shows how too small η might lead to underfitting, while

overly large η (where we would not consider $\eta \geq 1$ given Eq. (4b)) could promote overfitting.

However, while the optimal window size in EpiEstim varies with the form of the true R_s (e.g., large k is needed

for stable R_s time-series and small k for strongly fluctuating ones [13]), we find that a fixed $\eta = 0.1$ works

well across diverse epidemic scenarios and enables automatic detection of transmission change-points. This is an

advantage over approaches requiring explicit change-point modelling [9]. We examine this further in Fig. A2. There

we compare R_s estimates from APEestim (with optimal window k) [13] to EpiFilter ($\eta = 0.1$, $m = 2000$) for

simulated epidemic scenarios. We generate epidemics via the standard renewal model (Eq. (1)) using the Ebola

virus serial interval distribution from [37] and ensure the prior distributions of both methods have the same support.

The epidemics simulated span the range of possibilities (e.g., they may have large or small incidence).

We find EpiFilter outperforms APEestim, despite our using a single η value, with improvements especially stark for the simulations with subcritical transmission (as detailed in main text). This single η contrasts the discordant optimal k values and allows EpiFilter to recover the salient dynamics for epidemics with stable trajectories, seasonal variations and rapidly or gradually fluctuating transmission. In none of these scenarios does the true R_s have either the state model of Eq. (4b) or the sliding-window relationship in EpiEstim. Hence, these simulations also indicate robustness of the different inherent state assumptions in both approaches to moderate model mismatch. We use single runs to showcase how a practical epidemic would be processed in real-time. Neither method depends on random algorithms (e.g., MCMC) so their estimates will remain constant under the same parameters.

Fig. A2: **Validation against APEestim.** We compare estimates from EpiFilter (right panels) with optimised ones from APEestim (left panels [13]) over various general epidemic scenarios (which may have large or small incidence). These are simulated using Eq. (1) and not restricted to epidemics that may face elimination or resurgence as in Fig. 2 and Fig. 3. We use $m = 2000$, $R_{\min} = 0.01$ and $R_{\max} = 10$ for EpiFilter. To be fair we use a $\mathrm{Gam}(1, 2)$ prior distribution with APEestim, which sums to 1 by $R_{\max}$. Here k is the optimal window choice in APEestim (under predictive error) and $\eta = 0.1$ is fixed for EpiFilter inferences. True R_s is in black and conditional mean estimates with 95% confidence intervals are in red. EpiFilter is more stable, smoother and minimises estimate uncertainty in all scenarios. While the optimal k changes considerably among epidemics, $\eta = 0.1$ is able to automatically detect salient changes in transmission for all examples considered.

Last, we complete our batch analysis of EpiFilter performance by examining coverage statistics. In Fig. 2 and Fig. 3 a range of waning and resurging epidemic examples were explored. There we found that EpiFilter considerably improves the MSE when inferring instantaneous reproduction numbers, R_s, relative to EpiEstim and APEestim. APEestim, expectedly, had the best PMSE when predicting incidence, I_s, but EpiFilter was only marginally worse. In Fig. A3 we plot the distribution (over the 200 runs of each scenario from Fig. 2 and Fig. 3) of true coverage i.e., the probability that the R_s estimate or I_s prediction confidence intervals of a given method contain the true R_s and I_s values from the simulations. Interestingly, we find that EpiFilter generally has the best coverage (closest

to 1) across all scenarios for both R_s and I_s, confirming its superior performance.

Fig. A3: **Coverage statistics.** We provide the probability that the true values of instantaneous reproduction numbers, R_s, and incidence, I_s, from each scenario in Fig. 2 and Fig. 3 (over 200 runs) are contained within the 95% estimate and prediction confidence intervals generated by EpiFilter ($\eta = 0.1$), APEestim ($k = k^*$) and EpiEstim (weekly, $k = 7$ and monthly, $k = 31$ windows). All simulation and method parameters follow from the main text and Fig. 2 and Fig. 3 (y axes indicate corresponding scenarios). We find that EpiFilter usually achieves the highest coverage of all methods (1 indicates perfect coverage). APEestim consistently outperforms fixed-window EpiEstim methods, showing the importance of adaptive window selection.