First detection of SARS-CoV-2 genetic material in the vicinity of COVID-19 isolation centre through wastewater surveillance in Bangladesh

Firoz Ahmed[1], Md. Aminul Islam[1], Manish Kumar[2], Maqsud Hossain[3], Prosun Bhattachariya[4], Md. Tahmidul Islam[4], **Foysal Hossen[1],** Md. Shahadat Hossain[5], **Md. Main Uddin[5], Md. Nur Islam[5], Newaz Mohammed Bahadur[6], Md. Didar-ul-Alam[7], Hasan Mahmud Reza[8], Md. Jakariya[9]**

[1]Department of Microbiology, Noakhali Science and Technology University, Noakhali- 3814, Bangladesh

[2]Discipline of Earth Science, Indian Institute of Technology, Gandhinagar, Gujarat 382 355, India

[3]Department of Biochemistry and Microbiology, North South University, & NSU Genome Research Institute (NGRI), North South University, Bashundhara, Dhaka-1229, Bangladesh

[4]COVID-19 Research@KTH, Department of Sustainable Development, Environmental Science and Engineering, KTH Royal Institute of Technology, Teknikringen 10B, SE-10044 Stockholm, Sweden

[5]Department of Biotechnology and Genetic Engineering, Noakhali Science and Technology University, Noakhali- 3814

[6]Department of Applied Chemistry and Chemical Engineering, Noakhali Science and Technology University, Noakhali- 3814

[7]Professor and Honorable Vice-Chancellor, Noakhali Science and Technology University, Noakhali- 3814

[8]Department of Pharmaceutical Sciences, North South University, Bashundhara, Dhaka-1229, Bangladesh

[9]Department of Environmental Science and Management, North South University, Bashundhara, Dhaka-1229, Bangladesh

Corresponding Author:
Dr. Firoz Ahmed
Professor and Chairman
&
Focal Point
COVID-19 Diagnostic Lab
Department of Microbiology
Noakhali Science and Technology University
Noakhali- 3814, Bangladesh
Email: firoz@nstu.edu.bd; firoz19701016@gmail.com
Mobile: +8801713195658

HIGHLIGHTS

☐ First detection report of SARS-CoV-2 RNA in the wastewaters of Bangladesh.

☐ We traced the genetic load in the vicinity of the isolation center with 200 COVID-19 patients.

☐ About 75% of positive samples were found during the monitoring period.

☐ Main sewer and canal exhibit temporal accumulation of genetic load of SARS-CoV-2

☐ In general, shredding shows a variable trend during the sampling tenure.

46
47

ABSTRACT:

In the course of a COVID-19 pandemic, 0.33 million people got infected in Bangladesh, we made the first and successful attempt to detect SARS-CoV-2 viruses' genetic material in the vicinity wastewaters of an isolation centre i.e. Shaheed Bhulu Stadium, situated at Noakhali. The idea was to understand the genetic loading variation, both temporal and distance-wise in the nearby wastewater drains when the number of infected COVID-19 patients is not varying much. Owing to the fact that isolation center, in general, always contained a constant number of 200 COVID-19 patients, the prime objective of the study was to check if several drains carrying RNA of coronavirus are actually getting diluted or accumulated along with the sewage network. Our finding suggested that while the temporal variation of the genetic load decreased in small drains over the span of 50 days, the main sewer exhibited accumulation of SARS-CoV-2 RNA. Other interesting finding displays that probably distance of sampling location in meters is not likely to have a significant impact on gene detection concentration, although the quantity of the RNA extracted in the downstream of the drain was higher. These findings are of immense value from the perspective of wastewater surveillance of COVID-19, as they largely imply that we do not need to monitor every wastewater system, and probably major drains monitoring may illustrate the city health. Perhaps, we are reporting the accumulation of SARS-CoV-2 genetic material along the sewer network i.e. from primary to tertiary drains. The study sought further data collection in this line to simulate conditions prevailed in the most of south Asian country and to shed further light

80   on the temporal variation and decay/accumulation processes of the genetic load of the SARS-COV-

81   2.

83   **Keywords:** SARS-COV-2, Environmental surveillance, Sewage waste, Isolation centre, COVID-19.

1. Introduction:

Severe acute respiratory syndrome coronavirus 2 (**SARS-CoV-2**) is the strain of coronavirus that

causes coronavirus disease 2019 (COVID-19), that are now being frequently reported in specimens

collected from the wastewater treatment plants around the world owing to the shredding of the

symptomatic/asymptomatic COVID-19 infection (Ahmed et al., 2020; Haramoto et al., 2020;

Kumar et al., 2020; La Rosa et al., 2020; Medema et al., 2020; Nemudryi et al., 2020; Or et al.,

2020; Randazzo et al., 2020; Rimoldi et al., 2020; Wu et al., 2020b; Wurtzer et al., 2020). In Milano,

Italy, ORF1ab assay and N protein assay was used to detect SARS-CoV-2 RNA from untreated

wastewater and river water samples (Rimoldi et al., 2020). However, the wastewater surveillance

of COVID-19 (WWSoC-19) has been mostly focused on wastewater treatment plants around the

globe, and there is slow progress on ambient waters and especially in the sewer system. This is

critical because several developing countries like India, Bangladesh, Pakistan, and others do not

have proper wastewater systems even in their class-I cities. As a result, the policymakers in these

countries are in confusion pertaining to the national scale implementation of WWSoC-19.

Further, while the infectivity issues of SARS-CoV-2 RNA present in the wastewater as well as

ambient environmental waters are not yet neglected or proved in the scholarly world, the public

around the world are very skeptical about the wastewaters coming from the isolation centers

mainly that are not equipped with wastewater treatment systems. There have been some decay

reports of genetic loading of SARS-CoV-2 (Ahmed et al., 2020b, Kumar et al., 2020b) in the

wastewater system, but sewer systems are not yet investigated. In addition, there is a lack of

explicit understanding of either decay or accumulation of Covid-19 genetic material along with the

sewer systems with distance and the networking from small drain to larger drains followed by the

canal and main sewer system.

Furthermore, it is a general observation that most of the WWSoC-19 studies reported worldwide

have either correlated their C_t-value or detected gene copies with the total infected person in the

corresponding monitoring city or country. Uncertainty is very much evident on the average

amount of gene copies shredded by an infected person, and its relationship with the number of

genes detected during WWSoC-19. While we already know about the variations that exist in the

length of viral shedding in wastewater from the various studies (Wu et al., 2020b; Xu et al., 2020),

the magnitude of the shedding ranges between 10^2 and 10^8 copies of RNA per gram of human

waste (Lescure et al., 2020; Pan et al., 2020; Wölfel et al., 2020). The general trend has been to

see the fluctuation in the Ct value and then estimate the corresponding increase or decrease of

the COVID-19 patient in a given vicinity of treatment plants. However, there has been a complete

lack of WWSoC-19 study where infected individuals information are known and then a variation in

genetic loading has been studied.

Accordingly, we conducted a preliminary detection of SARS-CoV-2 RNA in wastewater samples

from the sewage network in the isolation centre at Noakhali, Bangladesh. The aim of the study

was to: i) understand the genetic load in the vicinity of the isolation centre with an almost constant

number of COVID-19 patients. ii) analyse the distance impact on genetic loading and tracing the

decay/accumulation of the same along with the sewage network, iii) and temporal variation in

genetic material shedding from the infected patient.

2. Material and Methods:

2.1. Sampling:

Wastewater samples were collected from sewage waste tank, passage drain, and toilets near

Shaheed Bhulu Stadium at Noakhali, Bangladesh (**Figure 1**). The sampling location for this study

was selected based on the fact that Shaheed Bhulu Stadium is the largest detention Centre for

COVID-19 patients at Noakhali district, Bangladesh. This facility has been established to

accommodate more than two hundred COVID-19 positive patients for isolation purposes but kept

around 200 patients all the time during the monitoring period. This preliminary study has been

carried out with samples collected from the three different drains that are coming out of the

stadium and can be considered as primary drains, which connects to a canal (secondary drainage

system) and eventually meets the main sewer system (tertiary drains).

In order to understand the distance impact on genetic loading along the drains, we collected

samples at various distance i.e. 100m, 200m, 300m, and 400m as presented in **Table 1**. Specimens

were aseptically collected in a 50 ml sterile falcon tube, transported in the laboratory keeping

inside the ice-box, refrigerated at 4°C during preparatory activities, and were analyzed on the

same day. Sterile falcon tubes for sampling with identical blanks were analyzed to determine any

possible contamination during the transport. All analyses were done at the Microbiology

Laboratory of the Department of Microbiology, Noakhali Science and Technology University

(NSTU), Bangladesh.

154
155
Figure 1. Study area location depicting the isolation camp established at the Shaheed Bhulu

Stadium at Noakhali along with the schematic diagram of sampling drains.

2.2. Sample preparation, and procedure for the RNA extraction and concentration:

We almost followed the same procedure of extraction described in Kumar et al (Kumar et al.,

2020). Briefly, sewage samples (50 mL) were centrifuged (Thermo Scientific) at 4500×g for 30 min

followed by filtration of supernatant using 0.22-micron filters (Himedia). Further, each sewage

filtrate was concentrated using polyethylene glycol (PEG) method. In this method, PEG 6000 (80

g/L) and NaCl (17.5 g/L) were mixed in 25 ml filtrate, which was then incubated at 17°C in 100 rpm

shaking for overnight. The next day, the mixture was centrifuged at 13000×g for 90 min. After

centrifugation, the supernatant was discarded and the pellet was resuspended in 300 µL RNase

free water. This was further used as a sample for RNA isolation. RNA isolation was carried out using

a commercially available kit according to the manufacturer's instructions. RNA concentrations

were measured by NanoDrop (Thermo Scientific™ NanoDrop 2000 and 2000c, BioRad) and were

stored at -70 °C until further use.

2.3. RT-PCR Analysis:

RNAs were analyzed for the detection of SARS-CoV-2 by RT-PCR (CFX96, BioRad) using the Sansure

RT-PCR kit (Sansure Biotech Inc., China). Technical procedures carried out as described in the

product manual and interpretations of results were analyzed as instructed in the manual. In brief,

we set the samples layout with RT-PCR protocol covering 45 cycles containing FAM fluorescence

select for ORF1ab, ROX for N gene as well as CY5 for Internal control. As quality control measures,

one positive control and one negative control were also run for validation of the test procedure.

3. Results:

We collected 16 specimens between 10th July and 29th August 2020 from the drain, sewage, and

toilets near Shaheed Bhulu Stadium Detention Centre (DC) for COVID-19 patient at Noakhali,

Bangladesh. A summary of the results of amplification cycles (Ct) of various sampled water along

with the distance of sampling in the vicinity of isolation center with 50 days temporal resolution

i.e. in between 10th July and 29th August 2020 has been presented in **Table 1** and amplification

plots obtained through RT-PCR illustrating temporal variation through Ct value in various sampling

drains, (a) Drain 1 (b) Drain 2 (c) Drain 3, and (d) Main sewer is presented in **Figure 2**.

Table 1: Summary of the results of amplification cycles (Ct) of various sampled wastewater along
with the distance of sampling in the vicinity of isolation center with 50 days temporal resolution
i.e. in between 10th July and 29th August 2020.
193

Source	Distance from Isolation Center (m)	Date of Sample collection	RNA Quantity		Ct Value (Sansure Kit)		
			Concentration (ng/µl)	Ratio (260/280)	FAM (ORF1ab)	ROX (Ngene)	Cy5 (RnaseP)
Drain 1	100	18/08/2020	12.42	1.97	34.69	32.25	37.91
		29/08/2020	25.5	1.95	ND	ND	36.42
	200	17/08/2020	12.37	1.99	40.72	ND	36.95
	300	07/08/2020	24.68	1.92	ND	ND	ND
	400	26/08/2020	12.23	2.00	39.43	ND	ND
Drain 2	**200**	18/08/2020	13.18	1.92	36.94	35.78	37.57
	400	26/08/2020	14.37	2.01	ND	ND	36.14
Drain 3	**200**	17/08/2020	17.47	1.97	36.78	35.82	39.08
		18/08/2020	22.61	1.98	22.35	20.42	32.30
	300	17/08/2020	15.97	1.92	31.03	29.03	37.08
		26/08/2020	22.16	1.99	32.85	30.47	40.13
	400	26/07/2020	15.31	1.95	32.84	30.35	34.52
Canal	**NTS**	10/07/2020	27.31	1.97	40.77	39.38	38.79
		26/08/2020	18.14	2.00	32.18	30.18	35.32
Main Sewer	**NTS**	07/08/2020	24.63	1.96	ND	ND	ND
		18/08/2020	16	1.94	34.85	33.24	37.94

194   *NTS= Distance not to scale; ND: Not Detected*
195

Table 1 summarizes the Ct values obtained during the monitoring which ranged between 20.42 to

39.38 for N genes, and 22.35 to 40.72 for ORF1ab genes, implying a huge variation in gene copies

of SARS-CoV-2. Interestingly as their lowest and highest values belong to corresponding samples

of the same date i.e. 17th and 26th August, it seems sewer systems played a critical factor WWSoC-

19. Other than this anomaly 17th August samples exhibited higher loading of genetic material than

29th August 2020, while the number of patients in the containment remained the same during the

monitoring period. This emphasizes on the fact that just becoming COVID-19 positive is not a

measure of the viruses shedding by the infected person, but perhaps the state of infection

matters. It is easy to speculate that with each day passing owing to aggressive testing and capacity

building of carrying out the tests, early detections of COVID-19 positive people were happening,

and thus probably it is reflecting on the genetic load. Moreover, temporal environmental

variations due to huge rainfall with temperature and humidy fluctuations along with inadequate

sweage treatment might have significant impact on the quantitative variations of SARS-CoV-2 viral

genetic material.

Figure 2: Amplification plots obtained through RT-PCR illustrating temporal variation through C$_t$ value in various sampling drains; (a) Drain 1 (b) Drain 2 (c) Drain 3, and (d) Main sewer.

As far as different characteristics of sampled drains are concerned, as depicted in **Figure 2**, drain 3 seems carrying the heavier RNA load of SARS-CoV-2 followed by drain 2 and drain 1. Althoguh dissimilarities observed between the primary drainage line i.e. drain samples and secondary drainage system i.e. canal and tertiary drainage system i.e. the main sewer, however, trend of higher genetic material loading in the secondary and tertiary system was also found on twilight tenure. This is a unique finding where gene accumulation has been observed instead of the general expectation of dilution in the larger sewer system. The probable reasoning, other than the accumulation of loading from various drains of the isolation centre, in support of this observation can be the additional contribution of RNA excreted from the asymptomatic patient as well as yet to be diagnosed people into the sewer system.

Figure 3: (a) Drain outside DC (b) Drain water (c) Sewage outside DC (d) Sewage inside DC (e) Main

Drain

We employed the distance factor in our sampling strategies and the results are presented in **Figure**

3. While drain 3 in general showed the increase in the ORF1ab and N genes, drain 1 was not

showing a consistent trend in genetic material loading of SARS-CoV-2 along with the distance. We

232 tend to propose that probably distance in meters is not likely of a critical factor capable of

233 producing a trend.

235 Table 2. Comparative positive sample and the range of Ct value reported of ORF-1ab genes of
236 SARS-CoV-2 in the wastewater of various countries.

Country	Period of Examination	Positive sample from total samples	Ct	Reference
Italy	14/04/2020 - 22/04/2020	4(12)	-	(Rimoldi et al., 2020)
Spain	06/04/2020 - 21/04/2020	7(7)	33.61 – 39.60	(Balboa et al., 2020)
Turkey	21/04/2020-25/04/2020	7(9)	34.67 – 39.54	(Kocamemi et al., 2020)
Netherlands	05/02/2020 - 16/03/2020	10(24)	-	(Medema et al., 2020)
France	05/03/2020 - 23/04/2020	100 % of samples	-	(Wurtzer et al., 2020)
Australia	20/03/2020-01/04/2020	2(9)	37.5 - 39	(Ahmed et al., 2020)
U.S.A.	08/01/2020 - 25/03/2020	10(12)	33.87 - 38.39	(Wu et al., 2020a)
India	0/05/2020 - 28/05/2020	100 % of samples	32.65 – 34.18	(Kumar et al., 2020)
Israel	10/03/2020-21/04/2020	10(26)	32.76 – 38.5	(Or et al., 2020)
Bangladesh	10/07/2020 – 29/08/2020	12 (16)	20.42 – 40.77	Present study (2020)

Table 2 shows a comparative analysis of monitoring period, percentage of positive samples

detected, and range of Ct-value along with their reference (Ahmed et al., 2020; Balboa et al., 2020;

Kocamemi et al., 2020; Kumar et al., 2020; Medema et al., 2020; Or et al., 2020; Rimoldi et al.,

2020; Wu et al., 2020b; Wurtzer et al., 2020). One finding stands out on this comparison is the Ct

value of 20.42 which corresponds to much higher genetic material loading of SARS-CoV-2 than any

other study reported. This may be because we sampled in the vicinity of the isolation centre, while

other studies compared in **Table 2** have reported the values from the wastewater treatment plant.

Overall, we successfully detected ORF1ab, and N protein genes, from the wastewater samples of

Bangladesh which is for the first time, reported the data from the containment centre. Our study

provides an opportunity to produce a realistic coefficient in the future for the conversion of

genetic material loading with the number of infected people in the community.

Further, referring to the limitations of the present study, surrogate samples were sent to other

laboratories to check how precisely the Sansure kit data matches with others. We also tested the

filtration methods to check with the efficiency of PEG method used in this study. Although, Hata

et al. reported a high efficiency of the PEG method in Japanese wastewater, and the same has

been found true by Kumar et al. while testing the sample concentration efficiency between PEG

and filtration (Hata and Honda, 2020; Kumar et al., 2020). While lack of supply always poses critical

challenges in any research, during lockdown we could not find a supply of MS2, as used by Kumar

et al. or could perform the whole process control (WPC) together with MPC as recommended by

Haramoto et al. (Haramoto et al., 2020; Kumar et al., 2020). We had to use a swab sample of a

symptomatic person as our control as indicated by CY-5 as quality control of our analyses, which

makes sense and provides a low-cost control for such analyses, yet it may be controversial, less

precise and at times comes negative as the case in a few samples in our case (drain 1 samples at

300 and 400m and main sewer sample on 07[th] and 26[th] August). By applying fluorescence

quantitative RT-PCR technology, this test utilized the novel Covid-19 ORF1ab and the specific

conserved sequence of coding nucleocapsid protein N gene as the target regions, which were

designed for conserved sequences of the double target genes to achieve detection of wastewater

samples RNA through fluorescent signal changes. The PCR detection system used the positive

internal control, which monitors the presence of PCR inhibitors in test specimens by detecting

whether the internal control signal is normal, to avoid a false negative result when used for human

RT-PCR experiments. In our experiments with wastewater, few specimens were negative for CY5

indicated that the human gene RNase P gene was missing in the samples. Hence it has been noted

that the human RNase P gene is more vulnerable to degradation than Covid-19 viral genes.

We also refrained to put here any semi-quantitative calculation of gene copies owing to lack of

enough replicates, kit designed for the human sample as well as uncertainties involving RT-PCR

(Stuart et al 2014) and will induct such calculation in our future study when there will be a method

at hand for precise copy calculations using suitable methods. Nevertheless, the bottom line is that

the patterns of obtained Ct values suggest successful detection of SARS-CoV-2 RNA from the

wastewater samples in Bangladesh, and their increasing abundance in tertiary drain demonstrated

that it is not difficult to employ the COVID-19 surveillance through wastewater in the sewer

systems to know the community health as we probably do not need dense sampling but the major

drains may be enough to use the capability of wastewater-based epidemiology in south-Asian

settings.

4. Conclusion:

While the wastewater surveillance of COVID-19 has been focused on wastewater treatment plants

around the world, we have opted for drain waters monitoring in the vicinity of the isolation center,

which is first of its kind. Apart from this being the first detection report of SARS-CoV-2 RNA in the

wastewaters of Bangladesh, the uniqueness of the study has been the tracing the genetic load in

the vicinity of the isolation center that contains almost the constant number with 200 COVID-19

patients, which takes the variable of the number of infected persons out of the equation. This has

been the key feature of this study as most of the study reported worldwide has either reported

total infected person in the city or country. There has been a complete lack of infected person

information that is contributing to the total genetic load to the sampled wastewater. We have

found about 75% of our sampled water positive based on ORF1ab gene absence or presence.

However, the critical observation has been the temporal variation where small drains showed an

easing of the loading of genetic load, the bigger canal, and main sewer city exhibited temporal

accumulation of SARS-CoV-2 RNA. On the other hand, the distance of sampling location in meters

appears to be insignificant from the perspective of wastewater surveillance of COVID-19. We

intend to analyze further samples taken in June, July, and August and preserved to shed further

light on the temporal variation and decay/accumulation processes of the genetic load of the SARS-

COV-2.

Ethics Statement

The work did not involve any human subject and animal experiments.

Acknowledgement:

309 We acknowledge the sincere help extended by several COVID-19 Diagnostic Lab Volunteers to

310 name a few are: Md. Sydul Islam, Md. Shariful Islam, Md. Sahedul Islam, MD. Amzad Hossain,

311 Mahmudul Islam Rakib, Golam Shamdani, Amor Chandra Nath, and Md. Julker Nyne, of Noakhali

312 Science and Technology University, Bangladesh.

References:

Ahmed W, Angel N, Edson J, Bibby K, Bivins A, O'Brien JW, et al. First confirmed detection of SARS-CoV-2 in untreated wastewater in Australia: A proof of concept for the wastewater surveillance of COVID-19 in the community. Science of The Total Environment 2020: 138764.

Balboa S, Mauricio-Iglesias M, Rodríguez S, Martínez-Lamas L, Vasallo FJ, Regueiro B, et al. The fate of SARS-CoV-2 in wastewater treatment plants points out the sludge line as a suitable spot for incidence monitoring. medRxiv 2020.

Haramoto E, Malla B, Thakali O, Kitajima M. First environmental surveillance for the presence of SARS-CoV-2 RNA in wastewater and river water in Japan. medRxiv 2020.

Hata A, Honda R. Potential Sensitivity of Wastewater Monitoring for SARS-CoV-2: Comparison with Norovirus Cases. Environmental Science & Technology 2020; 54: 6451-6452.

Kocamemi BA, Kurt H, Hacioglu S, Yarali C, Saatci AM, Pakdemirli B. First Data-Set on SARS-CoV-2 Detection for Istanbul Wastewaters in Turkey. medRxiv 2020.

Kumar M, Patel AK, Shah AV, Raval J, Rajpara N, Joshi M, et al. First proof of the capability of wastewater surveillance for COVID-19 in India through detection of genetic material of SARS-CoV-2. Science of The Total Environment 2020: 141326.

La Rosa G, Iaconelli M, Mancini P, Ferraro GB, Veneri C, Bonadonna L, et al. First detection of SARS-CoV-2 in untreated wastewaters in Italy. Science of The Total Environment 2020: 139652.

Lescure F-X, Bouadma L, Nguyen D, Parisey M, Wicky P-H, Behillil S, et al. Clinical and virological data of the first cases of COVID-19 in Europe: a case series. The Lancet Infectious Diseases 2020.

Medema G, Heijnen L, Elsinga G, Italiaander R, Brouwer A. Presence of SARS-Coronavirus-2 RNA in Sewage and Correlation with Reported COVID-19 Prevalence in the Early Stage of the Epidemic in The Netherlands. Environmental Science & Technology Letters 2020; 7: 511-516.

341   Nemudryi A, Nemudraia A, Wiegand T, Surya K, Buyukyoruk M, Cicha C, et al. Temporal
342   detection and phylogenetic assessment of SARS-CoV-2 in municipal wastewater. Cell
343   Reports Medicine 2020: 100098.
344   Or IB, Yaniv K, Shagan M, Ozer E, Erster O, Mendelson E, et al. Regressing SARS-CoV-2 sewage
345   measurements onto COVID-19 burden in the population: a proof-of-concept for
346   quantitative environmental surveillance. medRxiv 2020.
347   Pan Y, Zhang D, Yang P, Poon LLM, Wang Q. Viral load of SARS-CoV-2 in clinical samples.
348   The Lancet Infectious Diseases 2020; 20: 411-412.
349   Randazzo W, Truchado P, Cuevas-Ferrando E, Simón P, Allende A, Sánchez G. SARS-CoV-2
350   RNA in wastewater anticipated COVID-19 occurrence in a low prevalence area. Water
351   Research 2020: 115942.
352   Rimoldi SG, Stefani F, Gigantiello A, Polesello S, Comandatore F, Mileto D, et al. Presence and
353   vitality of SARS-CoV-2 virus in wastewaters and rivers. medRxiv 2020.
354   Wölfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, Müller MA, et al. Virological
355   assessment of hospitalized patients with COVID-2019. Nature 2020; 581: 465-469.
356   Wu F, Zhao S, Yu B, Chen Y-M, Wang W, Song Z-G, et al. A new coronavirus associated with
357   human respiratory disease in China. Nature 2020a; 579: 265-269.
358   Wu Y, Guo C, Tang L, Hong Z, Zhou J, Dong X, et al. Prolonged presence of SARS-CoV-2 viral
359   RNA in faecal samples. The lancet Gastroenterology & hepatology 2020b; 5: 434-435.
360   Wurtzer S, Marechal V, Mouchel J-M, Moulin L. Time course quantitative detection of SARS-
361   CoV-2 in Parisian wastewaters correlates with COVID-19 confirmed cases. MedRxiv
362   2020.
363   Xu Y, Li X, Zhu B, Liang H, Fang C, Gong Y, et al. Characteristics of pediatric SARS-CoV-2
364   infection and potential evidence for persistent fecal viral shedding. Nature medicine 2020;
365   26: 502-505.
366