The Effects of India's COVID-19 Lockdown on Critical Non-COVID Health Care and Outcomes:
Evidence from a Retrospective Cohort Analysis of Dialysis Patients

Authors: Radhika Jain, Pascaline Dupas[1]

Working Paper Under Review

Abstract: India's COVID-19 lockdown, one of the most severe in the world, is widely believed to have disrupted critical non-COVID health services. However, linking these disruptions to effects on health outcomes has been difficult due to the lack of reliable, up-to-date health outcomes data. We identified all dialysis patients under a statewide health insurance program in Rajasthan, India, and conducted surveys to examine the effects of the lockdown on care access, morbidity, and mortality. 63% of patients experienced a disruption to their care. Transport barriers, hospital service disruptions, and difficulty obtaining medicines were the most common causes. We compared monthly mortality in the four months after the lockdown with pre-lockdown mortality trends, as well as with mortality trends for a similar cohort in the previous year. Mortality in May 2020, after a month of exposure to the lockdown, was 1.70 percentage points or 64% (p=0.01) higher than in March 2020 and total excess mortality between April and July was estimated to be 22%. Post-lockdown morbidity, hospitalization, and mortality were strongly positively associated with lockdown-related disruptions to care, providing further evidence that the uptick in mortality was driven by the lockdown. Females, socioeconomically disadvantaged groups, and patients living far from the health system faced worse outcomes. The results highlight the unintended consequences of the lockdown on critical, life-saving non-COVID health services that must be taken into account in the implementation of future policy efforts to control the spread of pandemics.

Main Text:
On 24 March 2020, the Government of India ordered one of the most stringent nationwide COVID-19 lockdowns in the world to control virus spread (Hale et al 2020). The lockdown was announced with four hours' notice, barred people from leaving their homes, required non-essential commercial establishments and transport services to close, and was enforced strictly with penalty of arrest (Ministry of Home Affairs 2020). National restrictions were eased 10 weeks later, at the beginning of June, but localized restrictions continued in areas with high case counts.

Although critical health services were officially exempt from the lockdown, the media reported widespread disruptions to routine and emergency non-COVID care due to transport and curfew barriers for patients and health workers, hospitals turning patients away, and supply chain disruptions that affected medicine access and costs (IndiaSpend 2020, Indian Express 2020, New York Times 2020). Medical services under government health insurance programs across the country decreased by 51% and the national Health Management Information System (HMIS) ,

---

which collates monthly reports from the public health system, reported dramatic decreases in preventive and curative care (Smith et al 2020, IndiaSpend 2020).

Careful measurement of the indirect morbidity and mortality effects of the lockdown is critical to understanding the full consequences of the pandemic and how to prepare health systems better for future disease outbreaks (Kiang et al 2020, Beaney et al 2020, Setel et al 2020).

However, quantifying these impacts has been difficult in the Indian context due to the unavailability of reliable and high frequency data measuring health outcomes and cause-specific mortality. Because the COVID-19 lockdown may have reduced deaths from some causes, such as road accidents, evaluating its effects requires disaggregated cause-specific mortality. The vast majority of deaths in India occur at home, rather than at health facilities, are not included in the national Civil Registration System, and have no certified cause of death (Jha et al 2005). While government health insurance programs collect real-time data on hospital services provided, they record no details on patient morbidity or mortality. The HMIS data are typically of low quality, underrepresent care in the private sector, and do not provide complete morbidity or mortality outcomes (Sharma et al 2016). The Sample Registration System, which estimates age-specific death rates through population surveys, does not provide details on cause of death and the most recent report only provides total mortality estimates through 2018 (SRS Bulletin).

Given the dearth of reliable, updated, and publicly available data on health outcomes in India, evaluating the effect of the COVID-19 lockdown on non-elective health services requires the identification of patients in need of such care and the collection of primary data. We used insurance claims filed under a largescale government health insurance program to identify patients requiring dialysis, a form of non-elective chronic care, during the lockdown. We conducted phone surveys with their households to estimate the effects of the lockdown on health care access, morbidity, and mortality in the four months following its imposition. To our knowledge, we provide some of the first quantitative measures of the effects of India's lockdown on excess mortality from non-COVID health conditions.

DIALYSIS IN INDIA
We focused on dialysis, a form of life-sustaining long-term hospital care for patients with end stage chronic kidney disease (CKD). CKD is the 17th leading cause of deaths globally and 8th in India (GBD 2015). Although access to dialysis treatment is relatively low, the expansion of government health insurance has increased its reach and a 2019 study estimated that there are approximately 175,000 CKD patients on dialysis in India (Jha et al 2019). Dialysis removes waste, salts, and excess water to prevent their build up in the body; regulates levels of potassium, sodium and bicarbonate in the blood; and controls blood pressure. The typical patient requires three- to four-hour sessions, two to four times each week for the duration of their life or until they get a kidney transplant. Disruptions to dialysis treatment result in the accumulation of fluids and toxins in the body, and can cause extreme swelling, nausea and vomiting, difficulty in breathing and urinating, and other symptoms. Missing dialysis visits or shortening their duration is associated with large increases in hospitalization and mortality (Saran et al 2003, Anderson et al 2009, Zoraster et al 2007, Abdel-Kadar et al 2009). Recent work documents substantial

disruptions to dialysis care in India during the lockdown, but does not link them to health outcomes (Smith et al 2020, Ramachandran and Jha 2020, Prasad et al 2020).

METHODS
Study Population and Data

We conducted a retrospective cohort analysis to study care-seeking and health outcomes before and after the COVID-19 lockdown. The study population was all dialysis patients enrolled in a government health insurance program that covers the poorest two-thirds of Rajasthan's population. The Ayushman Bharat Mahatma Gandhi Rajasthan Swasthya Bima Yojana (AB-MGRSBY), is a statewide government health insurance program in Rajasthan, India. It entitles approximately 50 million low-income individuals to free secondary and tertiary care at over 1200 empaneled hospitals. Household eligibility is based on state poverty lists. All members are automatically enrolled and face no premium, deductible, or copay. Hospitals file claims in real-time for every patient visit through the government's electronic filing system. These claims data, which include patient name, phone number, address, hospital visited, and services received, provide one of the only ways of directly identifying patients utilizing hospital care in Rajasthan.

We obtained access to all administrative claims data filed under the program from its launch through October 2019. Using the last 3 months of these data, we identified all patients on dialysis under insurance between August and October 2019. Because dialysis is long-term and non-elective care, patients on dialysis in late 2019 would continue to require it through the COVID-19 lockdown if still alive. Therefore, these patients provide an ideal group to study the effects of the lockdown on critical chronic care.

To collect data on health care access and outcomes through the lockdown period, we conducted phone surveys using patient contact numbers in the administrative records. The survey was conducted with the patient or the person in the household most knowledgeable of their care if the patient was unwell or dead, and collected data on dialysis visits in the month prior to the lockdown, disruptions to care due to the lockdown, morbidity, hospitalization, the date and cause of death, and basic demographic and socioeconomic characteristics. We also embedded open-ended questions into the surveys to collect qualitative descriptions from patients about care disruptions faced. We completed the first round of surveys between late May 2020 and mid-June 2020, and follow-up surveys in July and August with all patients alive at the time of the first survey to track complete mortality through July. To help put survey results into context, we also conducted structured interviews with 15 hospitals from 10 of Rajasthan's 33 districts.

Outcomes and Measures

We measured disruptions to dialysis care by asking whether patients faced each of the following problems during the lockdown: their dialysis hospital was closed; it was open but refused to provide them services; they could not travel to their hospital due to lack of transport or curfews; they had to switch to a different hospital from their primary dialysis hospital; their hospital increased charges over the typical payment; they had difficulties obtaining their dialysis medicines. Reported dialysis visits were used to create an indicator for any decrease in visits between the month before and after the lockdown. We created an individual care disruptions

index using the first component of a principal-components analysis (PCA) of all of these indicators, and standardized it over the study sample for ease of interpretation.

To track changes in monthly mortality over the months before and after the lockdown, we created a mortality time series between December 2019 and July 2020. We determined the month of death for all dead patients by cross-checking two measures - the exact month of death and the number of months that had lapsed since the death at the time of survey - as well as the cause and location of death to ensure it was not unrelated to dialysis. Monthly mortality, or the likelihood of death, was calculated as the number of deaths each month as a share of people alive at the beginning of that month.

Morbidity and hospitalization were reported for the four weeks prior to the survey, in each round of the survey. A morbidity index was constructed from a PCA of indicators for whether the patient experienced the following symptoms that are known to follow disruptions to dialysis care and can be reported by patients: swelling of the face, hands, legs, or body; vomiting or nausea; extreme tiredness or weakness; difficulty breathing; difficulty urinating; and muscle cramps. The index was standardized over the sample. Hospitalization was an indicator for any in-patient hospital visit.

To examine heterogeneity in outcomes by socioeconomic and demographic characteristics, we created binary classifications for female, age under 45 years, lower caste group (scheduled caste or tribe), and low assets. Patients were classified as low asset if they had a below median score on an asset index created from a PCA of a list of assets they own, such as a motorcycle, television, or air-conditioner. Additionally, we geocoded all dialysis hospitals in the AB-MGRSBY program as well as the residence locations for 88% of surveyed patients using addresses in the administrative data that were verified by survey. We calculated the distance from patient residence to the district administrative headquarters, the largest city in the district, as a proxy for remoteness, and to the closest dialysis hospital as a proxy for distance from the health system. For heterogeneity analysis, we created binary classifications for above and below the median distance in the study sample, which was 45km for distance to the city and 35km for distance to a dialysis hospital.

Comparison Cohort
Because the COVID-19 lockdown was implemented across all of India at the same time, there are no comparable "untreated" populations in 2020. Therefore, our analysis primarily focused on changes in mortality trends within our study cohort over the four months before and after the lockdown was imposed. In addition, we constructed a historical comparison cohort to allow us to account for potential seasonal trends or monthly fluctuations in mortality and to examine whether 2020 mortality follows historical trends. Using the same administrative insurance claims data, we identified all patients on dialysis under insurance between August and October 2018. Because phone surveys with these patients would likely suffer from attrition and recall bias, we instead used the share of patients that permanently dropped out from dialysis care each month to proxy for the share of people that died in that month. We have previously validated this measure of mortality through surveys and confirmed that 80% of patients that dropped out of claims had died and the remaining 20% had dropped out for reasons besides death (exiting the insurance program, out of state migration, kidney transplant, or discontinuation of treatment for financial

reasons). Monthly mortality in the historical cohort was, therefore, calculated as the 80% of the share of all patients on dialysis each month that dropped out of treatment in that month.

Statistical Analysis

To examine changes in monthly mortality before and after the lockdown, we estimated a non-parametric discrete-time logistic model with the probability of death as the outcome and binary indicators for each month from December 2019 to July 2020. We excluded October and November 2019 from the analysis because survey dropout was likely to be highest in these months and could bias mortality estimates, but complete death counts for the full period are provided in the supplement. Adjusted models included indicators for age above 45 years, female sex, low caste group, and high assets, and continuous measures of months on dialysis and total dialysis visits at baseline. To examine historical mortality trends, we ran the same unadjusted model on the comparison cohort with indicators for each month from December 2018 to July 2019, as well as a model adjusted for age above 45 years and total dialysis visits prior to enrollment. Standard errors were clustered at the patient level in all models to account for autocorrelation.

To analyze the association between care disruptions and post-lockdown health outcomes, we restricted the sample to patients alive at the end of April and exposed to at least one month of the lockdown, and ran linear OLS regressions, with heteroskedasticity-robust standard errors, of post-lockdown morbidity, hospitalization, and death on the care disruptions index. Adjusted models included indicators for age above 45 years, female sex, low caste group, and low assets, as well as continuous measures of months on dialysis and total dialysis visits at baseline. Sensitivity to using logistic regression models was also assessed.

We also examined whether the lockdown had differential effects on care-seeking by sex, age (above 45 years), low caste, and low assets, we estimated bivariate and multivariate logistic regressions of an indicator for having experienced any disruption on binary indicators for each subgroup characteristic. To study heterogeneity in effects on mortality, we calculated the percentage point change in monthly mortality between March and May for each subgroup from logistic regressions controlling for all other subgroup characteristics and dialysis history. Lastly, we examined the association between care disruptions and post-lockdown health outcomes by running separate OLS regressions with heteroskedasticity-robust standard errors for each subgroup of post-lockdown morbidity, hospitalization, and death on the care disruptions index.

RESULTS

We identified 3,183 patients on dialysis under insurance between August and October 2019 across Rajasthan. Of these, 2,234 (70.2%) had a reachable phone number and confirmed having an eligible dialysis patient, and 94% of them consented to participating, resulting in a study sample of 2,110 patients (supplementary table S1). Attrition may be due to numbers being entered incorrectly in the insurance data, households changing numbers, or unused numbers being deactivated and reassigned to new households. Of the 1,392 patients alive at the time of the first survey, 1,177 (85%) completed a follow-up survey. Successfully surveyed patients were disproportionately male (69%), had a mean age of 46 years, and had been on dialysis for a year, with 5 visits per month on average, when they were enrolled into the study (Table 1). Surveyed

patients were almost identical in these characteristics to patients not reached, but had been on dialysis for slightly longer, increasing our confidence that attrition did not meaningfully bias our sample (supplement table S2). On average, patients lived 25km from the nearest hospital providing dialysis care. The vast majority of patients (83%) were visiting a private hospital for their dialysis treatments.

Disruptions to Dialysis Care

Over 62% of patients reported a disruption in access to dialysis care during the lockdown (Figure 1). 42% of patients reported being unable to reach their hospital due to travel barriers. Open-ended interviews indicated that travel barriers were largely due to difficulties finding transport and obtaining official exemptions from district administrative and health authorities to travel to the hospital visits. 15% of patients found the hospital was closed or refused to provide care, 11% faced increased hospital charges, and 23% had to switch to a different hospital from the one they typically visit. Interviews suggest supply chains disruptions reduced medicine availability and increased prices. As a result, 17% of patients could not obtain their necessary medicines. Patients faced a 172% increase in payments per visit, driven largely by increased charges at private hospitals, and a 6% average decline in monthly visits between March and April. Hospital interviews confirmed patient reports, but also suggested that hospitals faced their own constraints.

Seven of the 15 hospitals reported having to during the lockdown due to staffing and supply shortages. Those that continued operating reported a substantial drop in patient visits, which they credited to transportation barriers, and larger hospitals reported receiving patients displaced from nearby hospitals that had closed.

Changes in Mortality Trends

Monthly mortality declined steadily from December to March in the 2019-2020 cohort (Figure 2), due to higher early mortality among the most vulnerable patients, such as the elderly and lower caste (supplement Figure S1). Mortality in May 2020, after a full month of exposure to the lockdown, increased sharply to 4.37%, a 1.70pp or 63.60% (p=0.01) increase relative to 2.67% mortality in March, prior to the lockdown. Controlling for sociodemographic characteristics and dialysis history increased this difference (supplement FigS2). Mortality declined in June and July 2020, but never dropped below March levels, indicating that deaths in May were not simply a displacement of mortality from the subsequent two months. Because we did not reach all households in the follow-up survey, true mortality in this period may be higher than measured. Mortality in the comparison cohort from the previous year followed a statistically similar trend between December and April, but exhibited no increase in May or the subsequent months. Comparison of the two trends suggests that the sharp increase in May 2020 is not explained by seasonal fluctuations and that, absent the lockdown, monthly mortality would have remained similar to or slightly below March levels. Only four patients who died in June and July were reported testing positive for COVID-19. Excluding them reduces mortality to 3.16% in June and 2.69% in July.

To estimate total excess mortality in the four months following the COVID-19 lockdown, we calculated the difference between observed mortality between April and July 2020 and what mortality would have been if the March 2020 mortality probability of 2.67% had applied each

month. Out of 1,532 patients alive at the end of March, 192 had died by the end of July (12.54% mortality), whereas 157 would have died had the March rate (10.25% mortality), resulting in 22.3% total estimated excess mortality in the four months following the lockdown (supplement table S4). This is likely to be a conservative estimate, as both pre-lockdown trends and the historical comparison suggest that mortality would have continued to decline in the absence of the lockdown rather than remain at March levels.

Association Between Care Disruptions and Post-lockdown Outcomes
Among patients alive at the end of April and exposed to at least one month of the lockdown, a 1SD increase in the care disruptions index was associated with a 0.17SD increase in the morbidity index (p=0.000), 3.1pp increase in the probability of hospitalization (p=0.002), and 2.1pp increase in the probability of death (p=0.013) in the period from May to July (Table 2). These effects are sizeable relative to an overall 14.2% hospitalization and 10.8% mortality hazard over those months. Controlling for sociodemographic characteristics and dialysis history did not change these relationships. They also remain robust to using logistic instead OLS regression models for the binary hospitalization and death outcomes (supplemental table S5).

Subgroup Heterogeneity in Lockdown Effects
Although the lockdown was universal, its effects on care-seeking were worse for vulnerable and remote households. Being lower caste, poorer, and living further away from a city or a dialysis hospital had large and significant positive associations with the likelihood of facing any disruption (Table 3). Poverty had the strongest relationship and remained significant in multivariate regressions controlling for other socioeconomic characteristics and dialysis history.

Figure 3 presents the change in mortality in May relative to March by subgroup. Underlying data are presented in supplemental table S6. The increase in mortality was largest and significant for females (3.56pp, p=0.022), patients 45 years or younger (2.94pp, p=0.006), poorer patients (2.81pp, p=0.020), those living further from the city (3.51pp, p=0.013) and those living further from a dialysis hospital (5.64pp, p=0.031). The association between the care disruptions index and post-lockdown morbidity, hospitalization, and death were also stronger for these subgroups (supplement figure S3). The greater disruptions to care faced by poorer and remote patients noted above could partly explain their worse outcomes. However, we find no evidence that females experienced greater disruptions. One explanation for high female mortality may be lower care-seeking by households, consistent with prior literature on gender bias in health care in India (Anonymous 2021, Kapoor et al 2019, Shaikh et al 2018). Females had fewer monthly dialysis visits at baseline (4.8 visits) relative to males (5.1 visits, p=0.062), which could contribute to worse health prior to the lockdown. Additionally, care disruptions were associated with increased morbidity for both males and females, but with increased hospitalization only for males, and larger increases in mortality for females, suggesting that households may have been less likely to seek hospital care for females that faced complications (supplement figure S3).

DISCUSSION
We studied the indirect effects of India's COVID-19 lockdown on critical, chronic, non-COVID care and outcomes among a large cohort of low-income dialysis patients. Dialysis patients faced substantial disruptions to their care during the lockdown. Travel barriers, due to the closure of

public transport and administrative hurdles in obtaining exemptions for medical travel, were the most common cause. Hospital closures, service refusals, and increased charges forced patients to switch hospitals and miss or delay treatment.

Mortality in May 2020, after a month of exposure to the lockdown, increased sharply. The timing and size of the increase in mortality relative to pre-lockdown trends in 2020, as well as to trends in estimated mortality for a similar cohort in the previous year, strongly suggests that the observed increase was due to the nationwide COVID-19 lockdown. Lockdown-related disruptions to dialysis care were strongly positively associated with post-lockdown morbidity and mortality, providing additional support for a causal interpretation. Estimated overall excess mortality in the four months after imposition of the lockdown was 22%.

This excess mortality is unlikely to be due to COVID-19 infections. Although patients on dialysis are at higher risk of COVID-19 infection and complications (Gansevoort and Hilbrands 2020), if this were driving mortality, we would expect deaths to increase over time as the virus spread and to affect older patients more. However, the increase in mortality was largest in May, soon after the lockdown and before the virus had spread widely, and was greater among patients under age 45 than among older patients who have higher COVID-19 mortality risk. Therefore, we believe our estimates are capturing the indirect effects of the pandemic through lockdown related disruptions to care. Nevertheless, given that COVID-19 testing rates in India were relatively low at the time of the study, we cannot rule out the possibility that it contributed to the deaths we measured.

Importantly, we find that the effects of the lockdown on care-seeking and mortality were more severe among females, patients of low socioeconomic status, and those in remote locations underserved by the health system. These results indicate that a universal lockdown policy is likely to have larger adverse effects on already vulnerable subpopulations, consistent with the literature on the unequal distribution of both direct and indirect effects of the pandemic (Bambra et al 2020, Marmot and Allen 2020, Cash and Patel 2020). It is critical that future pandemic control efforts take these distributional consequences into account and put in place extra protections for populations that have limited financial and geographic access to health services.

While our analysis is restricted to one type of critical chronic care in one state, our findings are likely to be indicative of the serious but largely undocumented health effects of severe disruptions to a range of similar critical chronic care services in other states in India. A 2019 study estimated that there are approximately 175,000 patients on chronic dialysis across India (Jha et al 2019). Based on recent research and media reports, it is likely that a large share of these patients suffered life-threatening disruptions to their care (Ramachandran and Jha 2020, Prasad et al 2020). Furthermore, national government health insurance programs targeting low-income households have seen a 6% reduction in dialysis care (similar to our results), a 64% decline in oncology care, and an 80% decrease in critical cardiovascular surgeries (Smith et al 2020).

A strength of this study is the use of existing administrative government health insurance data to rapidly identify and remotely survey a large sample of poor dialysis patients in need of critical care during the lockdown. However, a limitation of this approach is the substantial dropout

between identified patients and those reached for survey by phone due to patient phone numbers entered incorrectly in the claims data, households switching phone numbers, and unused numbers being deactivated. Surveyed patients were almost identical to those not reached, increasing confidence that attrition did not bias our study sample. Deactivation, which happens after 6 months, are most likely to affect patients that died soon after October 2019 and is unlikely to explain the increased mortality in May 2020. Another limitation is that our outcomes are based on self-reported data and not clinical measures, which were infeasible due to the pandemic. However, our primary outcome is mortality, which is likely to be reliably reported, and the recall period for all outcomes was relatively short. Finally, as we only measure mortality through July, our estimates may be a lower bound on the full health costs of the lockdown. We found that disruptions to dialysis were lower but persistent in July and August, after the lockdown was eased, suggesting that adverse health effects may continue to unfold over the coming months (supplement figure S4).

Lockdowns can reduce COVID-19 transmission (Flaxman et al 2020, Figueiredo et al 2020, Islam et al 2020). However, they have substantial indirect health and economic costs that may outweigh their benefits relative to other pandemic control strategies in some contexts (Douglas et al 2020, Modesti et al 2020, Haug et al, 2020, Cash and Patel 2020). In India, these costs were exacerbated by the hasty imposition of the lockdown, which gave the health system and households little time to prepare (The Lancet 2020). Our findings highlight the unintended consequences of India's lockdown on critical non-COVID health services that must be taken into account in the implementation of future policy efforts to control the spread of pandemics.

Figure 1: Disruptions to dialysis care during the COVID-19 lockdown
The figure presents the share of patients that reported experiencing disruptions to their dialysis care due to the lockdown between imposition of the lockdown in March 2020 and the survey conducted in May-June 2020.

Figure 2: Monthly dialysis mortality in the surveyed and historical cohorts
The solid line presents monthly likelihood of death, or the share of people still alive that die in each month, for the surveyed cohort, from an unadjusted discrete time logistic regression model with indicators for each month from December 2019 through July 2020. Vertical bars represent 95% confidence intervals. The dashed line presents the monthly hazard for the comparison cohort of patients on dialysis in August-October 2018 from a similar model with indicators for each month from December 2018 through July 2019. Mortality estimates have been converted into percentage terms for ease of interpretation. Covariate-adjusted models in the online supplement look almost identical.

Figure 3: Changes in mortality between March and May by subgroup. Bars represent the percentage point change in mortality between March and May 2020 for the full sample and each subgroup. Each model is adjusted for all other characteristics (indicators for age, sex, caste, and asset group), as well as total dialysis visits prior to enrollment. Unadjusted models are in supplemental figure S4. Lines represent 90% and 95% confidence intervals. The grey figures on the right present the increase between March and May as a percentage of March mortality.

	Mean	SD
Age (yrs)	46.0	14.7
Female (%)	31.1	
Scheduled caste/tribe (%)	18.7	
Education (yrs)	11.3	2.6
Distance to district HQ (km)	48.8	38.8
Distance to dialysis hospital (km)	19.7	17.7
Dialysis history at baseline		
Private dialysis hospital	83.2	
Monthly dialysis visits	5.0	3.5
Months on dialysis	12.6	11.5
Observations	2110	

Table 1: Study Sample Characteristics

Dialysis history is drawn from administrative claims data on all dialysis treatment received prior to enrollment in the study in October 2019. Distance to subdistrict HQ is the distance in km to administrative center of the subdistrict, a proxy for distance to the closest city. Distance to dialysis hospital is the distance to the closest hospital providing dialysis under insurance, a proxy for distance from the health system.

	Morbidity Index			Hospitalization			Death		
	Estimate	95% CI	P	Estimate	95% CI	P	Estimate	95% CI	P
A. Unadjusted									
Care Disruptions Index	0.172	[0.125 - 0.219]	<0.001	0.031	[0.012 - 0.051]	0.002	0.021	[0.004 - 0.038]	0.013
B. Adjusted									
Care Disruptions Index	0.171	[0.125 - 0.219]	<0.001	0.030	[0.010 - 0.050]	0.003	0.025	[0.008 - 0.042]	0.004

Table 2: Association between care disruptions and post-lockdown health outcomes

The table presents linear regressions of each health outcome on a standardized index of lockdown-related disruptions to dialysis care for the 1,489 patients alive thorough April and exposed to at least one full month of the lockdown. Morbidity is a standardized index of symptoms known to follow disruptions to dialysis. Hospital and death are binary outcomes. All three outcomes are measured over the period May through July 2020. Covariates in adjusted regressions include age, sex, caste group, month of dialysis initiation, and lifetime dialysis visits at baseline.

	Outcome: Faced Any Care Disruption					
	Bivariate			Multivariate		
	OR	95% CI	P	OR	95% CI	P
Female	0.974	0.780 - 1.216	0.815	0.979	0.768 - 1.247	0.862
Lower caste	1.143	0.932 - 1.402	0.200	1.063	0.850 - 1.330	0.593
45yrs or younger	1.333	1.003 - 1.773	0.048	1.152	0.849 - 1.563	0.363
Low assets	1.618	1.315 - 1.990	<0.001	1.495	1.191 - 1.876	0.001
Far from city	1.302	1.038 - 1.635	0.023	1.212	0.962 - 1.528	0.103
Far from hospital	1.360	0.995 - 1.861	0.054	1.269	0.923 - 1.744	0.142

Table 3. Association of Patient Characteristics and Likelihood of Facing Care Disruptions
The table presents estimates from bivariate and multivariate logistic regressions of an indicator for facing any lockdown-related care disruption on indicators for patient characteristics. The sample is 1,574 patients that were alive through March, when the lockdown was imposed.

REFERENCES

1. Hale T, Petherick A, Phillips T, Webster S. Variation in government responses to COVID-19. Blavatnik school of government working paper. 2020 May 27;31.
2. Ministry of Home Affairs Order No 40-3/2020-D dated 24.03.2020. https://www.mha.gov.in/sites/default/files/Guidelines.pdf.
3. "How Healthcare Became Unaffordable For Non-COVID Patients During The Pandemic," IndiaSpend, 19 June 2020; https://www.indiaspend.com/how-healthcare-became-unaffordable-for-non-covid-patients-during-the-pandemic/.
4. "Private hospitals can't refuse critical patients or insist on testing all for COVID-19: Centre," Indian Express, 28 April 2020; https://www.newindianexpress.com/nation/2020/apr/28/private-hospitals-cant-refuse-critical-patients-or-insist-on-testing-all-for-covid-19-centre-2136540.html.
5. "8 hospitals in 15 hours: A pregnant woman's crisis in the pandemic," New York Times, 12 July 2020; https://www.nytimes.com/2020/06/21/world/asia/coronavirus-india-hospitals-pregnant.html.
6. Smith, O., Naib, P., Sehgal, P., Chhabra, S. PM-JAY Under Lockdown: Evidence on Utilization Trends. PM-JAY Policy Brief 8. National Health Authority 2020.
7. "COVID-19 disrupted India's routine health services," IndiaSpend, 27 August 2020; https://www.indiaspend.com/covid-19-disrupted-indias-routine-health-services.
8. Kiang MV, Irizarry RA, Buckee CO, Balsari S. Every body counts: measuring mortality from the COVID-19 pandemic. Annals of Internal Medicine. 2020 Sep 11.
9. Beaney T, Clarke JM, Jain V, Golestaneh AK, Lyons G, Salman D, Majeed A. Excess mortality: the gold standard in measuring the impact of COVID-19 worldwide?. Journal of the Royal Society of Medicine. 2020 Sep;113(9):329-34.
10. Setel P, AbouZahr C, Atuheire EB, Bratschi M, Cercone E, Chinganya O, Clapham B, Clark SJ, Congdon C, de Savigny D, Karpati A. Mortality surveillance during the COVID-19 pandemic. Bulletin of the World Health Organization. 2020 Jun 1;98(6):374.)
11. Jha, P., Gajalakshmi, V., Gupta, P.C., et al. Prospective study of one million deaths in India: rationale, design, and validation results. *PLoS Med* 2005;*3*(2),p.e18.
12. Sharma, A., Rana, S. K., Prinja, S., & Kumar, R. Quality of health management information system for maternal & child health care in Haryana state, India. *PLoS One* 2016;*11*(2), e0148449.
13. Sample Registration System (SRS) Bulletins, Office of the Registrar General and Census Commissioner; https://censusindia.gov.in/vital_statistics/SRS_Bulletins/Bulletins.html.
14. GBD 2015 Mortality and Causes of Death Collaborators. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. Lancet. 2016; 388: 1459-1544.
15. Jha V, Ur-Rashid H, Agarwal SK, Akhtar SF, Kafle RK, Sheriff R. The state of nephrology in South Asia. Kidney international. 2019 Jan 1;95(1):31-7.
16. Saran R, Bragg-Gresham JL, Rayner HC, et al. Nonadherence in hemodialysis: associations with mortality, hospitalization, and practice patterns in the DOPPS. *Kidney International* 2003 Jul 1;64(1):254-62.
17. Anderson AH, Cohen AJ, Kutner NG, et al. Missed dialysis sessions and hospitalization in hemodialysis patients after Hurricane Katrina. *Kidney International* 2009 Jun1;75(11):1202-8.

18. Zoraster R, Vanholder R, Sever MS. Disaster management of chronic dialysis patients. *American Journal of Disaster Medicine* 2007 Mar 1;2(2):96-106.

19. Abdel-Kader K, Unruh ML. Disaster and end-stage renal disease: targeting vulnerable patients for improved outcomes. *Kidney International* 2009 Jun 1;75(11):1131-3.

20. Prasad N, Bhatt M, Agarwal SK, et al. The adverse effect of COVID pandemic on the care of patients with kidney diseases in India. *Kidney International Reports* 2020 Jul 6.

21. Ramachandran R, Jha V. Adding insult to injury: kidney replacement therapy during COVID-19 in India. Kidney International. 2020 Apr 25.

22. Anonymous. Gender disparities in government health insurance in India. *Working paper* 2020.

23. Shaikh M, Peters SA, Woodward M, et al. Sex differences in utilisation of hospital care in a state-sponsored health insurance programme providing access to free services in South India. *BMJ Global Health* 2018 Jun 1;3(3):e000859.

24. Kapoor M, Agrawal D, Ravi S, et al. Missing female patients: an observational analysis of sex ratio among outpatients in a referral tertiary care public hospital in India. *BMJ open* 2019 Aug 1;9(8):e026850.

25. Gansevoort RT, Hilbrands LB. CKD is a key risk factor for COVID-19 mortality. Nature Reviews Nephrology. 2020 Dec;16(12):705-6.

26. Bambra C, Riordan R, Ford J, Matthews F. The COVID-19 pandemic and health inequalities. J Epidemiol Community Health. 2020 Jun 12.

27. Marmot M, Allen J. COVID-19: exposing and amplifying inequalities. J Epidemiol Community Health. 2020 Sep 1;74(9):681-2.

28. Flaxman S, Mishra S, Gandy A, Unwin HJ, Mellan TA, Coupland H, Whittaker C, Zhu H, Berah T, Eaton JW, Monod M. Estimating the effects of non-pharmaceutical interventions on COVID-19 in Europe. Nature. 2020 Aug;584(7820):257-61.

29. Figueiredo AM, Daponte Codina A, Figueiredo M, Saez M, Cabrera León A. Impact of lockdown on COVID-19 incidence and mortality in China: an interrupted time series study. Bull World Health Organ. 2020.

30. Islam N, Sharp SJ, Chowell G, Shabnam S, Kawachi I, Lacey B, Massaro JM, D'Agostino RB, White M. Physical distancing interventions and incidence of coronavirus disease 2019: natural experiment in 149 countries. bmj. 2020 Jul 15;370.

31. Douglas M, Katikireddi SV, Taulbut M, McKee M, McCartney G. Mitigating the wider health effects of covid-19 pandemic response. Bmj. 2020 Apr 27;369.

32. Modesti PA, Wang J, Damasceno A, Agyemang C, Van Bortel L, Persu A, Zhao D, Jarraya F, Marzotti I, Bamoshmoosh M, Parati G. Indirect implications of COVID-19 prevention strategies on non-communicable diseases. BMC medicine. 2020 Dec;18(1):1-6.

33. Haug N, Geyrhofer L, Londei A, Dervic E, Desvars-Larrive A, Loreto V, Pinior B, Thurner S, Klimek P. Ranking the effectiveness of worldwide COVID-19 government interventions. Nature human behaviour. 2020 Nov 16:1-0.

34. Cash R, Patel V. Has COVID-19 subverted global health?. The Lancet. 2020 May 30;395(10238):1687-8.

35. The Lancet. India under COVID-19 lockdown. Lancet (London, England). 2020 Apr 25;395(10233):1315.

Acknowledgments: We are grateful to Ambika Chopra, Mustufa Patel, and Mantasha Hussain for their excellent and dedicated research assistance and to Putul Gupta for outstanding research management. The preliminary results in this paper were presented at the Development Economics meetings of the NBER Summer Institute in July 2020, as well as in a Webinar organized by JPAL South Asia in August 2020. **Funding:** This research was funded by a grant from the Bill and Melinda Gates Foundation, awarded through the J-PAL CaTCH Initiative's special COVID-19 funding window. The funder had no role in the design or execution of the research; **Author contributions:** R.J. and P.D. jointly conceived of and designed the research. R.J. performed the data analyses and produced the figures and tables. R.J. and P.D. wrote the paper. All authors contributed to the scientific discussions and preparation of the manuscript; **Competing interests:** Authors declare no competing interests; **Data and materials availability:** The insurance claims data is subject to a non-disclosure agreement. Replication code and deidentified survey data will be made available at the time of publication on the authors' website. The institutional review boards at IFMR and at Stanford University (protocol no. 41683) approved this research.

The Effects of India's COVID-19 Lockdown on Critical Non-COVID Health Care and Outcomes
Supplementary Materials

Radhika Jain, Pascaline Dupas

Correspondence to: rjain19@stanford.edu, pdupas@stanford.edu

Table S1. Survey Success Rate

First survey	Percent	Count	Observations
Sampled		3183	3183
Reached by phone	0.835	2659	3183
Reached and eligible dialysis patient confirmed	0.702	2234	3183
Consented to survey	0.944	2110	2234
Surveyed in first round		2110	2110
Follow-up survey			
Alive at first survey		1392	1392
Reached and consented to follow-up survey	0.846	1177	1392

Table S2. Baseline Characteristics of Patients by Survey Status

	Surveyed		Not Surveyed		Total		Difference
	Mean	SD	Mean	SD	Mean	SD	p
Age	45.95	14.65	45.21	16.02	45.70	15.13	0.20
Female	0.31	0.46	0.31	0.46	0.31	0.46	0.70
Visits per month	4.98	3.53	5.04	3.56	4.99	3.54	0.67
Months on dialysis	12.65	11.50	9.26	11.04	11.51	11.45	0.00
Observations	2110		1073		3183		3183

Table S3. Complete Death Counts in Survey and Comparison Cohorts

	2019-2020 Survey Cohort			2018-2019 Comparison Cohort		
	(1)	(2)	(3)	(4)	(5)	(6)
Month	Individuals Alive	Deaths	Mortality	Individuals Alive	Deaths	Mortality
Enrolled Aug-Oct	2110	210	0.0995	2118	213	0.1006
November	1900	89	0.0468	1852	111	0.0599
December	1811	107	0.0591	1713	94	0.0549
January	1704	70	0.0411	1596	63	0.0395
February	1634	60	0.0367	1517	59	0.0389
March	1574	42	0.0267	1443	38	0.0263
April	1532	43	0.0281	1395	38	0.0272
May	1489	65	0.0437	1347	38	0.0282
June	1424	46	0.0323	1300	31	0.0238
July	1378	38	0.0276	1261	33	0.0262

Table S4. Excess Mortality Estimates

	2020 Observed Data			Counterfactual Data Using March Mortality		
	(1)	(2)	(3)	(4)	(5)	(6)
	Individuals Alive	Deaths	Mortality	Individuals Alive	Deaths	Mortality
April	1532	43	0.0280	1532	41	0.0267
May	1489	65	0.0440	1491	40	0.0267
June	1424	46	0.0320	1451	39	0.0267
July	1378	38	0.0280	1413	38	0.0267
April-July	**1532**	**192**	**0.1253**	**1532**	**157**	**0.1025**
April-July Excess Mortality						**0.2222**

Table S5. Sensitivity of Association Between Care Disruptions and Health Outcomes to Logistic Models

	Hospitalization			Death		
	OR	95% CI	P	OR	95% CI	P
A. Unadjusted						
Care Disruptions Index	1.262	1.103 - 1.444	0.001	1.222	1.053,1.418	0.008
B. Adjusted						
Care Disruptions Index	1.264	1.096 - 1.458	0.001	1.293	1.104 - 1.515	0.001

The table presents similar regressions as in Table 2 of the paper of health outcomes on the care disruptions index for the 1,489 patients alive thorough April, but estimates logistic instead of OLS models to account for the fact that hospitalization and death are binary outcomes. Covariates in adjusted regressions include age, sex, caste group, month of dialysis initiation, and lifetime dialysis visits at baseline.

Table S6. Changes in mortality between March and May by subgroup

	March Mortality (%)	May Mortality (%)	PP Difference	% Difference	Difference p-value	95% CIL	95% CIU
Male	3.04	4.91	1.87	61.7%	0.044	0.053	3.693
Female	3.27	6.83	3.56	108.7%	0.022	0.512	6.601
Above 45yrs	3.78	5.57	1.79	47.5%	0.133	-0.548	4.135
45yrs or younger	2.46	5.41	2.94	119.9%	0.006	0.847	5.050
Upper caste	2.87	5.18	2.30	80.2%	0.007	0.644	3.964
Lower caste	4.36	7.16	2.80	64.3%	0.226	-1.741	7.369
High assets	3.06	5.06	2.01	65.7%	0.06	-0.085	4.103
Low assets	3.17	5.98	2.81	88.8%	0.02	0.445	5.184
Close to city	3.15	5.14	1.99	63.3%	0.078	-0.221	4.208
Far from city	3.64	7.15	3.51	96.6%	0.013	0.745	6.277
Close to hospital	3.28	5.38	2.09	63.7%	0.025	0.266	3.919
Far from hospital	3.80	9.44	5.64	148.6%	0.031	0.516	10.766

Figure S1. Unadjusted mortality trends by subgroup

Mortality by Age Group

Mortality by Sex

Mortality by Social Group

Mortality by Wealth Status

The figures present the full mortality trends over time from unadjusted models of the likelihood of death on the interaction of indicators for month and subgroup.

Figure S2. Covariate-adjusted monthly dialysis mortality

The figure presents mortality estimates adjusted for age, sex, caste group, month on dialysis, and lifetime dialysis visits at baseline for the 2019-2020 cohort and age, sex, months on dialysis, and lifetime dialysis visits at baseline (variables available in the claims data) for the 2018-2019 cohort.

Figure S3. Associations between disruptions and health outcomes by subgroup

The figure presents associations between the care disruptions index and post-lockdown health outcomes by subgroup for the 1,489 patients alive through the end of April and exposed to at least one month of the lockdown. Each bar is the coefficient on a indicator for the subgroup characteristic from an OLS regression with robust standard errors and controls for all other characteristics (age, sex, caste group, and asset class) and dialysis history at baseline. The outcomes are the standardized morbidity index, and indicators for hospitalization, death in May, and death between May and July.

Figure S4. Disruptions to Dialysis Care in July and August

The figure presents the share of people that reported experiencing disruptions in the four weeks prior to the survey, for all patients re-interviewed in July and August. These results suggest that disruptions to dialysis care continued well after the nationwide lockdown was released and may continue to have effects on morbidity and mortality in the coming months.

Supplementary Text
Qualitative interviews with patients and hospitals
To gain a more nuanced understanding of patients' experiences, we built open-ended questions into the survey. Surveyors were instructed to provide prompts, such as "Have you experienced any other health problems or difficulties in getting dialysis care due to the coronavirus lockdown? Can you describe them to me?" and to encourage respondents to answer freely. The research team listened to voice recordings of these answers and compiled them into structured summaries. The qualitative results confirmed that the list of barriers to care-seeking we included in the survey were comprehensive and covered the key proximate causes of care disruptions: hospital closure or service refusal, increased costs of care, travel/transport barriers, having to switch from their primary hospital to a different one, and difficulty obtaining medicines. They also reveal that many patients experienced a series of different barriers that had compounded effects on their care. Excerpts from some open-ended answers:

"The hospital was closed on visits and refused treatment as well. […] We did not have a mode of transport to reach the hospital, I had to try very hard to arrange one. […] As a result, I could only get 3 dialysis visits in total instead of 8."

"There was no doctor in the hospital. No one gave us any information where to find one, there was just one ward boy who did not know anything. […] For getting a treatment that my life depends on, I had to get slips from doctors, permissions from District Collector, stand in long queues..."

"There is this particular medicine that is required for dialysis procedure and that is not currently available in the market. We have not been able to find it for the last few days. […] Private dialysis centers have increased their rates - it has become unaffordable for us. We don't have money to seek treatment in a private hospital, we have lost our jobs so don't have enough money to go get treated in a private hospital, hence I had to take my wife very far to find a public hospital for dialysis. […] Every time we change a hospital, they issue a new set of tests and we have to get them because they will not give us dialysis otherwise. […] Due to shutdown of public transport, I have to now take my wife to dialysis. Because of that neither can I work nor I can be with my wife during her visits."

"I went to two private hospitals, both of them were closed, hence I had to go to a public hospital and got my dialysis there. […] The cost of acquiring my dialysis medicines almost doubled, as they were not available at my regular pharmacist, my hospital was closed, I had to look everywhere and they were available in limited quantities only."

"I have been advised to get three dialysis visits per week. Now because of the lockdown, I have only been able to get two per week. Hence, water fills in my lungs and I have difficulty breathing if I do any physical activity."

Family member of dead patient: *"We live in [District A] and the hospital he regularly visited was in [District B]. When the lockdown was imposed and there was a curfew, he had to find a smaller, alternative facility for dialysis. We were asked to get permissions and multiple slips from govt officials. He didn't get the care he usually received. That smaller hospital didn't do his*

dialysis properly and that's why he died. [...] The medicines that were prescribed to him were only available in [District B] and not his local area due to the lockdown. He missed his medicines for BP and diabetes for 20 days."

Family member of dead patient: *"Due to the lockdown, the patient couldn't travel to [District B] on time and wasn't able to find a hospital in the vicinity, so he died on the way to the hospital itself."*

Ethical considerations

The research protocol and survey instruments were approved by the Institutional Review Boards of the Institute for Financial Management and Research (IFMR) in India and Stanford University in the United States. All data collection was managed by a team employed by JPAL South Asia at IFMR, which has extensive research conducting field and phone-based research. Standard research protocols were followed. Participants were informed of the nature and possible consequences of the study and were given the opportunity to discontinue participation at any point of their choosing.

Given that data were being collected during a pandemic that may have severely adversely affected households, the households in our sample may have been particularly vulnerable due to their socioeconomic status and illness, and the health and security of our data team was also a concern, we took several additional precautions to reduce any adverse consequences. To reduce the burden on households, we designed the surveys to be short: among surveyed households, the first round of the survey took 25 minutes and the second follow-up round took 13 minutes, on average. We trained surveyors to very clearly offer households the option not to participate, to be sensitive to households during the survey, particularly those that had faced substantial difficulties or deaths, and to provide all households with information on local COVID-19 and hospital health service helplines to consult if they experience medical problems. To protect the surveyors, all surveys were conducted over the phone from the security of their homes. Additional software was installed on the surveyor tablets to mask phone numbers from them and to automatically upload all data to secure cloud services without storing it locally