

LOW PSYCHOLOGICAL WELL-BEING IN MEN WHO HAVE SEX WITH MEN (MSM) DURING THE SHELTER-IN-PLACE ORDERS TO PREVENT THE COVID-19 SPREAD: RESULTS FROM A NATIONWIDE STUDY

Emerson Lucas Silva Camargo¹; Bruna Isabela Adolpho de Oliveira²; Igor Fessina Siffoni¹; Anderson Reis de Sousa³; Jules Ramon Brito Teixeira³; Isabel Amélia Costa Mendes¹; Álvaro Francisco Lopes de Sousa⁴

1. Human Exposome and Infectious Diseases Network (HEID), Escola de Enfermagem de Ribeirão Preto, Universidade de São Paulo, Brazil.
2. Universidade de Ribeirão Preto, São Paulo, Brazil.
3. Universidade Federal da Bahia, Salvador, Brazil.
4. Global Health and Tropical Medicine, Instituto de Higiene e Medicina Tropical, Universidade Nova de Lisboa, Lisboa, Portugal.

Corresponding author:

Álvaro Francisco Lopes de Sousa

Avenida dos Bandeirantes, 3900

Campus Universitário - Bairro Monte Alegre

Ribeirão Preto - SP - Brasil

CEP: 14040-902

Email: sousa.alvaromd@gmail.com

ABSTRACT

Introduction: Little is known about how sheltering in place to contain the spread of COVID-19 over extended periods affects individuals' psychological well-being. This study's objective was to analyze the factors associated with MSM's low psychological well-being in the COVID-19 pandemic context.

Method: This cross-sectional study was conducted online in the entire Brazilian territory (26 states and federal district) in April and May 2020. The participants were recruited using an adapted version of Respondent-Driven Sampling (RDS). Data were collected using social media and MSM dating apps. We estimated the prevalence, crude prevalence ratio (PR), and respective confidence intervals (CI95%).

Results: Only 44.4% of the sample presented high levels of well-being. Low psychological well-being appeared associated with the youngest age group (PR: 2.76; CI95%: 1.90-4.01), polyamorous relationships (PR: 2.78; CI95%: 1.51-5.11), not complying with social isolation measures (PR: 6.27; CI95%: 4.42-8.87), not using the social media to find partners (PR: 1.63; CI95%: 1.06-2.53), having multiple sexual partners (PR: 1.80; CI95%: 1.04-3.11), having reduced the number of partners (PR: 2.67; CI95%: 1.44-4.95), and group sex (PR: 1.82; CI95%: 1.23-2.69)

Conclusion: The well-being of MSM living in Brazil was negatively affected during the social distancing measures intended to control the spread of COVID-19.

Policy Implications: Planning and implementing public policies and actions to promote psychological well-being are needed to improve MSM's resilience through the adoption of safe strategies and behavior.

Keywords: Psychological well-being; well-being; Men who have sex with men; Gay man; COVID-19; SARS-CoV-2.

Introduction

The COVID-19 pandemic is configured as the century's health challenge as more than 15 million people were infected worldwide up to July 2020 (WHO, 2020). Its impact is apparent in all

countries, and the populations' psychosocial health has been affected (Serafini, Parmigiani, Americo, Aguglia, Sher & Amore, 2020), especially among the more vulnerable ones. Brazil is currently facing a disheartening situation. It ranks third in the number of COVID-19 cases and has recorded more than 120,000 deaths (Brazil, 2020).

The COVID-19 pandemic context is permeated by high transmissibility rates, clinical complications, sudden changes in life, changes in the flow of cities, human behavior, and socio-affective relationships. These changes affect the individuals' psychological well-being, significantly decreasing the quality of life (Serafini et al., 2020). These problems are heightened by mandatory social isolation measures, intended to control the virus spread. The damage caused to human well-being is undeniable and has drawn the attention of researchers, lawmakers, health authorities, and public managers, intending to minimize the impact.

The pandemic's harmful impact, heightened by social isolation, in the individuals' physical, mental, social, and sexual dimensions, is quite noticeable (Forte, Favieri, Tambelli & Casagrande, 2020). In this context, sexual and gender minorities are subject to greater vulnerability given the historically conditioning inequality they face in terms of health and violence. That is, these groups are relegated to a marginal position within the society, and, as a consequence, a safety net system and psychosocial support are not always provided or available to them.

The results reported by a global survey performed by Hornet (News, 2020), a Gay social network, commissioned by the Thomson Reuters Foundation to be conducted among men who have sex with men (MSM), reveal considerable levels of loneliness and depression associated with shelter-in-place measures, emphasizing their broader impact on the individuals' mental health, which may imply greater impairment in terms of psychological and social well-being throughout the pandemic. The literature has addressed the pandemic's impact on the maintenance of affective and sexual practices, including the establishment of barriers impeding access to health services, necessary to maintain therapeutic processes and care technologies. From this perspective, there is a lack of concrete

governmental actions directed to MSM during the COVID-19 pandemic (Sousa, Oliveira, Schneider, Queiroz, Carvalho, Fronteira, 2020; Carvalho, et al, 2020; Torres et al., 2020).

Given this population's context, coupled with the repercussions of the novel socio-historical phenomenon caused by the COVID-19 worldwide on the individuals' psychological well-being, this study's objective was to analyze the factors associated with low psychological wellbeing among MSM living in Brazil during the COVID-19 pandemic.

Methods

Study design

This cross-sectional study addresses data from the “40TENA” project, conducted in all the 26 Brazilian states and the Federal District from May to April 2020, when restrictive public health measures such as the social isolation and shelter-in-place measures were in force.

Population, sample, and eligibility criteria

A total of 2,646 Brazilian MSM participated in this study. The participants were recruited using Respondent-Driven Sampling (RDS) adapted to the virtual environment. In this method, the participants themselves are responsible for recruiting other individuals within the same condition via social media. To meet the method's criteria, 15 MSM with different characteristics were selected, namely: place of residence (one of the five Brazilian regions); Race (Caucasian/non-Caucasian); age (young, adult, or elderly); and educational level. These were the first participants and were called 'seeds'. Each participant received a link to the survey and was instructed to invite other MSM from their social network until a significant sample was obtained. The seeds were identified through two geo-location-based dating applications (Grindr and Hornet) via direct chat with online users, following established methods (Queiroz, Sousa et al., 2019; Queiroz et al., 2019 (2); Sousa et al., 2019).

The researchers also boosted the survey on Facebook, directing it to the MSM population aged from 18 to 60 years old (Facebook imposed the age restriction), using a fixed post on the official page of the survey (<https://www.facebook.com/taafimdeque/>) accompanied by an electronic link that granted access to a free and informed consent form and the survey's questionnaire.

Only individuals who identified themselves as men (cisgender or transgender) and aged 18+ years old were included. Tourists and non-Portuguese speaking individuals were excluded.

Data collection instrument

The instrument used to collect data was developed by this study's authors, considering the research variables and the participants' characterization. It was content validated by a panel of judges specialized in the topic and method. The instrument was divided into four sections with 46 questions. Most were multiple-choice questions, and some were mandatory; otherwise, the participant could not proceed with the questionnaire. The questions addressed:

1. Sociodemographic data (age, gender identity, education, sexual orientation, type of relationship, country, state, place of residence);
2. Psychological well-being and coping strategies used during the shelter-in-place orders.

The 5-item World Health Organization Well-Being Index (WHO-5), validated for the Brazilian context (Souza & Hidalgo, 2012), was used to assess psychological well-being. It comprises five items rated on a 5-point Likert scale with a total score ranging from 0 to 25. WHO-5 was designed to measure psychological well-being in the previous two weeks. Scores below 20 indicate the presence of depressive disorder (Souza & Hidalgo, 2012), and scores equal to or lower than 13 indicate impaired well-being (Bech, 2004). Thus, psychological wellbeing was categorized as low (≤ 13 points), moderate (14 to 19 points) and high (≥ 20 points).

3. Sexual practice and activities during the pandemic (i.e., casual sex, sex with the consumption of drugs, use of condoms, protection strategies, strategy used to search for partners, frequency of sexual encounters, and protective measures against COVID-19);
4. Sexual practice and activities before the pandemic (Use of HIV Pre-Exposure Prophylaxis and Post-Exposure Prophylaxis; strategies to search for partners; knowledge regarding STIs and testing);

For security reasons, the form used to collect data was hosted by a specific site that enabled only one response per IP (internet protocol).

Data analysis

Data Analysis and Statistical Software (STATA), version 12.0 was used for data analysis. Descriptive and inferential statistics were used. Bivariate analysis was performed using the Chi-square test, estimating prevalence and crude prevalence ratio (PR) with a 95% confidence interval.

Multivariate analysis was performed using Poisson's Regression analysis with robust variation. The outcome variable (psychological well-being) was included in the analysis, along with each of the associated independent variables with $p\text{-value} \leq 0.20$. The stepwise procedure was used. The sequence in which each term was inserted in the model was determined by a mutual analysis of theoretical relevance criteria and statistical significance obtained in the bivariate analysis. Each term was added or removed from the model after identifying statistical significance ($p\text{-value} < 0.05$), stability of power of association, and Akaike Information Criterion (AIC), defining the useful subset of terms. All the variables with a 5% statistical significance remained in the final model. To determine the best final model, the one with the lowest AIC value was selected.

Ethical and legal aspects

The research project was approved by the Institutional Review Board at the Universidade Nova de Lisboa and the University of São Paulo. All the users signed free and informed consent forms before proceeding with the questionnaire.

Results

A total of 2,646 men who had sex with other men participated. Most men presented moderate psychological well-being (47.7%), followed by those with high well-being (44.4%), while a prevalence ratio of 7.9% experiencing low well-being was found. Most were aged from 18 to 29 years old (69.9%), 31 years old on average (Min. 18 and Max. 60; $SD\pm 8.4$), single (67.2%), fully complying with shelter-in-place orders (72.1%), with a duration of social isolation between 30 and 45 days (58.1%), and decreased consumption of alcohol (43.1%) (Table 1).

Table 1 – Distribution of men according to sociodemographic characteristics, social isolation compliance, sexual behavior, and COVID-19-related characteristics. Brazil, 2020. (N=2,646).

Variables	N	%
Sociodemographic characteristics		
Age group		
18 to 29 years old	1,850	69.9
30 years old or older	796	30.1
Relationship status		
Stable relationship	802	30.3
Polyamorous relationship	66	2.5
Single	1,778	67.2
Social isolation characteristics		

Social isolation

Total	1,908	72.1
Partial	664	26.1
No	74	2.8

Duration of social isolation (N=2,586)

<30 days	483	16.7
30 to 45 days	1,504	58.1
>45 days	599	23.2

Alcohol consumption

Decreased	1,140	43.1
Did not change	974	36.8
Increased	532	20.1

Sexual Behavior

Live with a partner (N=2,011)

No	1,723	85.7
Yes	288	14.3

Use social media to find partners (N=2,478)

Yes	2,164	87.3
No	314	12.7

Frequency of sexual practice

Did not change	493	18.6
Increased	427	16.2
Decreased	1,726	65.2

Number of partners

None	719	27.2
1	921	34.8
2 or more	1,006	38.0
Variation in the number of partners (N=2,645)		
Did not change	1,395	52.7
Decreased	921	34.8
Increased	329	12.5
Casual sex		
No	1,239	46.8
Yes	1,407	53.2
Group sex		
No	2,195	83.0
Yes	451	17.0
Chemsex Practice		
No	1,612	60.9
Yes	1,034	39.1
COVID-19-related characteristics		
Presence of symptoms		
No	1,282	48.4
Yes	1,364	51.6
COVID-19 testing		
Yes	218	8.2
No	2,428	91.8
Tested positive for COVID-19		

No	2,520	95.2
Yes	126	4.8

COVID-19: Coronavirus Disease 2019.

The bivariate analysis of factors associated with psychological well-being showed statistical association with age ($p < 0.001$), relationship status ($p < 0.001$), and social isolation ($p < 0.001$).

Being 18 to 29 years old, in a polyamorous relationship, and not complying with shelter in place orders increased three, twice, and seven times, respectively, the prevalence of low psychological well-being. Social isolation increased the frequency of low psychological well-being by 47%. Other findings are presented in Table 2.

Table 2 – Prevalence (%), prevalence ratio (PR), 95% confidence interval, low psychological well-being according to sociodemographic characteristics, social isolation compliance, sexual behavior, and COVID-19-related characteristics. Brazil, 2020.

Variables	n	P (%)	p-value*	PR	CI 95%
Sociodemographic characteristics					
Age group					
18 to 29 years old (N=1,849)	183	9.9	<0.001	3.03	2.03-4.5
30 years old or older (N=796)	26	3.3		1.00	-
Relationship status					
Stable relationship (N=801)	79	9.9	<0.001	1.00	-
Polyamorous relationship (N=66)	13	19.7		2.00	1.18-3.39
Single (N=1,778)	117	6.6		0.67	0.51-0.88
Social isolation characteristics					

Social isolation

Total (N=1,907)	117	6.1	<0.001	1.00	-
Partial (N=664)	60	9.0		1.47	1.10-2.0
Not complying (N=74)	32	43.2		7.05	5.15-9.65

Duration of social isolation

<30 days (N=483)	38	7.9	0.158	1.00	-
30 to 45 days (N=1,503)	97	6.5		0.82	0.57-1.18
>45 days (N=599)	30	5.0		0.64	0.40-1.01

Alcohol consumption

Decreased (N=1,139)	103	9.0	0.148	1.00	-
Did not change (N=974)	71	7.3		0.81	0.60-1.08
Increased (N=532)	35	6.6		0.73	0.50-1.05

Social behavior

Live with a partner

Yes (N=288)	31	10.8	0.032	1.00	-
No (N=1,722)	123	7.1		1.51	1.04-2.19

Use social media to find partners

Yes (N=2,164)	152	7.0	0.009	1.00	-
No (N=313)	35	11.2		1.59	1.12-2.25

Frequency of sexual practice

Increased (N=427)	39	9.1	0.091	1.00	-
Did not change (N=493)	48	9.7		0.77	0.55-1.09
Decreased (N=1,725)	122	7.1			

Number of partners

None (N=719)	38	5.3	0.009	1.00	-
1 (N=920)	80	8.7		1.64	1.13-2.39
2 or more (N=1,006)	91	9.1		1.71	1.19-2.47
Variation in the number of partners					
Did not change (N=1,394)	134	9.6	0.002	1.00	-
Decreased (N=921)	59	6.4		1.32	0.77-2.56
Increased (N=329)	16	4.9		1.98	1.19-3.27
Casual sex					
No (N=1,239)	84	6.8	0.045	1.00	-
Yes (N=1,406)	125	8.9		1.31	1.00-1.71
Group sex					
No (N=2,194)	154	4.0	<0.001	1.00	-
Yes (N=451)	55	12.2		1.74	1.30-2.32
Chemsex Practice					
No (N=1,612)	114	7.1	0.048	1.00	-
Yes (N=1,033)	95	9.2		1.30	1.00-1.69
COVID-19- characteristics					
Presence of symptoms					
No (N=1,281)	115	9.0	0.047	1.00	-
Yes (N=1,364)	94	6.9		0.77	0.59-0.99
COVID-19 testing					
Yes (N=218)	8	3.7	0.016	1.00	-
No (N=2,427)	201	8.3		2.26	1.13-4.51
Tested positive for COVID-19					

No (N=2,519)	203	8.1	0.181	1.00	-
Yes (N=126)	6	4.8		0.59	0.27-1.30

*p-value obtained with the Chi-square test.

COVID-19: Coronavirus Disease 2019; P: Prevalence; PR: prevalence ratio; CI95%: Confidence interval of 95%.

The variable ‘type of partner’ was excluded from the multivariate analysis because it presented collinearity with relationship status (VIF=7.37). The association between the presence of COVID-19 symptoms and testing for COVID-19 was not statistically significant but remained in the model as confounding adjustment because the model presented a lower AIC.

The following variables were associated with low psychological well-being: being in the youngest age group (PR: 2.76; CI95%: 1.90-4.01); having a polyamorous relationship (PR: 2.78; CI95%: 1.51-5.11); not complying with social isolation measures (RP: 6.27; CI95%: 4.42-8.87); not using the social media to find partners (PR: 1.63; CI95%: 1.06-2.53); having a higher number of sexual partners (PR: 1.80; CI95%: 1.04-3.11); having reduced the number of sexual partners (PR: 2.67; CI95%: 1.44-4.95); and group sex (PR: 1.82; CI95%: 1.23-2.69) (Table 3).

Table 3 – Prevalence ratios and confidence intervals (95%) of the factors associated with low psychological well-being in the context of the COVID-19 pandemic. Brazil, 2020.

Variables	PR	CI95%
Sociodemographic characteristics		
Age group		
18 to 29 years old	2.76	1.90-4.01
30 years old or older	1.00	-
Relationship status		

Stable relationship	1.00	-
Polyamorous relationship	2.78	1.51-5.11
Single	0.99	0.64-1.53
Social isolation characteristics		
Social isolation		
Yes	1.00	-
No	6.27	4.42-8.87
Use social media to find partners		
Yes	1.00	-
No	1.63	1.06-2.53
Sexual behavior		
Number of partners		
None	1.00	-
1	1.36	0.82-2.24
2 or more	1.80	1.04-3.11
Variation in the number of partners		
Did not change	1.00	-
Decreased	2.67	1.44-4.95
Increased	2.42	1.47-4.01
Group sex		
No	1.00	-
Yes	1.82	1.23-2.69
AIC		0.4935822

Model adjusted for the presence of COVID-19 symptoms and COVID-19 testing.

PR: prevalence ratio; CI 95%: Confidence interval of 95%.

Discussion

MSM living in Brazil experienced worse well-being during the shelter-in-place orders implemented to restrain the spread of COVID-19. The psychological well-being of more than half of the participants presented some level of impairment, while a high prevalence of low well-being was found. As for the variables that contributed to this outcome, those related to relationships (relationship status/sexual relationships), not complying with social isolation, strategies used to search for sexual partners, number of sexual partners, and group sex stood out. These results reinforce the need to promote and acquire in-depth knowledge regarding the role of affective and sexual dimensions within relationships and partnerships in the psychological well-being of LGBT populations when facing a crisis, such as pandemics.

Thus far, little has been published regarding the real impact of the need to deal with the COVID-19 pandemic and its duration on the sexual practices and loving relationships of populations (Sanchez, Zlotorzynska, Rai, & Baral, 2020), especially among LGBT populations. Few studies address the pandemic from the perspective of the quality or lack of social interactions and the role it plays in the psychological well-being of individuals by intensifying unfavorable triggers. Hence, there are various complex issues such as the historical denial and lack of commitment to LGBT people. As a consequence, this population is seldom addressed by scientific studies, which negatively influence the development of sensitive indicators able to properly support strategic and contingency actions, as well as health promotion and prevention of diseases, as well as risk factors (Jungmann & Witthoft, 2020; Esterwood & Saeed, 2020; Sahni & Sharma, 2020).

Other issues refer to the weak production of healthcare in terms of psychosocial care targeting MSM, especially in the pandemic context. Epidemiological indicators already show a context in which these individuals' mental health has been affected, and they are more frequently exposed to risks.

Nonetheless, this context has not been more accurately investigated in most countries, especially in Brazil, as the government leaders have not even discussed these problems in the pandemic context (Sousa, Carvalho et al., 2020).

The declining indicators of psychological well-being among MSM are worth noting. In a cataclysmic context, such as that of a pandemic, access to psychosocial support networks and the individuals' socio-affective network is considerably weakened and even absent in some contexts, which may intensify the vulnerability of these individuals from the perspective of mental health and social well-being.

The sudden rupture in socio-affective networks caused by the COVID-19 pandemic, for instance, not having access to nightlife, leisure, meeting friends and groups of belonging, intimate and sexual encounters, and affective relationships, restricts interactions to the virtual environment or small groups.

This context is likely to cause anxiety, decision conflicts, decreased self-esteem and interest in life, suicidal behavior, eating and sleep disorders, and abusive alcohol or drug consumption among MSM. These individuals may even become involved in practices in which they expose themselves to the Coronavirus for not bearing the prolonged social distancing measures, boredom, and the lack of social interaction with people, or restricted mobility within the cities. Coupled with these is the fact that difficulty in searching for affective and sexual partners during the period of social isolation may also decrease psychological well-being, impacting the health of these men (Sousa, Oliveira et al, 2020; Carvalho et al., 2020).

In this study, a greater prevalence of low psychological well-being was found among MSM who did not report the use of social media to find partners, revealing that even though the social media is considered to be associated with sex, in times of social distancing, it may play a protective role by enabling social interactions, even if remotely. The results from a multi-center study show that 95% of its MSM sample was (totally or partially) complying with social distancing measures, which resulted in

a drop in the number of partners and relationships, but, at the same time, increased online interactions (Sousa, Oliveira et al., 2020). The study also reports that MSM devised strategies to keep interactions during the social distancing measures, reinforcing this study's findings that interactions, whether to seek or exchange content, are essential for MSM's psychological well-being. The study mentioned above addressed only sexual interactions though, so inferences concerning other types of relationships are impossible.

Interestingly, having a polyamorous relationship (three people or more) was associated with low psychological well-being. One hypothesis is that polyamorous relationships do not always occur under the same roof, implying that those involved may have to deal with distancing measures. Additionally, polyamorous relationships allow for new people to be included, which in this case implies exposure among all those involved, resulting in fear, and decreased psychological well-being. Consensual non-monogamy or polyamorous relationships have become an increasingly accepted affective and sexual experience, which does not comply with monogamous relationship patterns, usually considered ideal in stable and committed relationships (Costa & Ribeiro, 2020; Jowett, 2020). This context has been intensified among sexual minorities, including MSM. We believe that the fact these individuals are involved in polyamorous relationships and are currently experiencing the impossibility to maintain the frequency of affective and sexual meetings during the pandemic, that is, impaired affective and sexual dynamics are coupled with diminished leisure opportunities these relationships usually promote, decrease these men's psychological well-being.

One cannot rule out the fact that MSM in polyamorous relationships have to live with stigma. In the context of this pandemic, this experience may be heightened considering the disruptive phenomena caused by the stigmatization process such as labels, judgment, withdrawing, and discredit, negatively impacting their physical and mental health, and consequently their psychological well-being (Conley, Piemonte, Guskova & Rubin, 2018; Conley, 2017).

In line with this hypothesis, this study shows that having a higher number of sexual partners and practicing group sex was associated with low psychological well-being, which reinforces the hypothesis that having sex with casual partners implies exposure that includes the risk of acquiring sexually transmissible infections and COVID-19. On the other hand, MSM are motivated by desire, libido, and sexual lust, satisfying their needs through group sex and an increased number of partners. Exposure to often unknown partners, or familiar partners who do not effectively adopt preventive measures, including those against COVID-19, increases the fear of being contaminated or anxiety with the possibility of presenting signs and symptoms of the disease, leading to low psychological well-being (Newman & Guta, 2020; Suen, Chan & Wong, 2020; Brennan, Card, Collicot, Jolimore & Lachowsky, 2020).

Considering the disciplinary role of affective and sexual practices, which coupled with the advent of a novel infectious disease, for which an effective treatment or cure has not been devised thus far, keeping multiple sexual partners and/or having group sex may increase the levels of anxiety, stress, uncertainty, worry, and fear among MSM. Additionally, the COVID-19 epidemiological curve in Brazil at the time of data collection was ascending, with an expressive number of new cases and deaths. Concomitantly, there was an increase in the amount of information provided by the media, exposing the complexity of the disease and virulence of SARS-CoV-2, which also tends to influence the individuals' psychological well-being negatively.

The prevalence of low psychological well-being was approximately six times greater among MSM who did not comply or could not comply with the social isolation measures, reinforcing the hypothesis that even though the individuals may not comply with social distancing measures to seek interactions, exposure still may cause concerns (Sousa, Oliveira et al., 2020; Newman & Guta, 2020; Sanchez et al, 2020).

Nonetheless, having a small number of partners also affected these individuals' psychological well-being negatively. If, on the one hand, the individuals want to decrease exposure to protect

themselves, on the other hand, not having the social support provided by relationships and partners, which would decrease the pandemic's harmful effects, could negatively affect these individuals' mental health (Sanchez et al., 2020; Carrico, et al., 2020; Newman & Guta, 2020).

This variation in the number of affective and sexual partners has compromised MSM's psychological well-being. They have not received the psychosocial support provided by the measures institutionalized in Brazil to protect them from or minimize the pandemic context's impact on mental health. In Brazil, stress has been intensified among minority social groups as their 'invisibility' has been exposed along with denial of rights, cuts on budgets previously designated to the LGBTQIA+ population, and naturalized institutional homophobia (Folha de São Paulo, 2020).

A lack of directive public policies mainly affects the youngest population, which in this study presented the highest prevalence of low psychological well-being, almost three times higher than among older MSM. The World Health Organization has shown concern with the youngest population within the context of the COVID-19 pandemic, mainly because this is the most vulnerable in not adhering to social isolation measures. An increase in the number of new COVID-19 cases, agglomerations, and non-adherence to public health measures to control the spread of Sars-CoV-2, has been found in countries already in the summer season (Young mind, 2020; Cohen & Bosk, 2020).

The Inter-Agency Standing Committee (IASC) has proposed specific guidelines to promote mental health and strengthen psychosocial support in emergencies, as is the case of the COVID-19 pandemic. These guidelines focus on promoting people's psychological well-being and preventing and/or treating mental diseases (IASC, 2007). These goals are achieved by employing multiple approaches that take into account biological, sociocultural, educational, and community characteristics, which should focus on reducing harm, promoting human rights and equality, social participation, and valorization of local resources and competencies, the work via integrated support systems, and rehabilitation (Dong, Du, & Gardner, 2020; Fukuti et al., 2020).

Socio-affective networks are a significant source of access to protection, sociability, exchange, and maintenance of MSM's psychological well-being and health. In the pandemic context, and even after the pandemic, supporting, strengthening, and disseminating these networks may be an effective strategy to improve the levels of psychological well-being, decreasing the impact caused by social isolation, such as feelings of loneliness, heightened anxiety, and stress.

Limitations include the study design. Cross-sectional studies do not allow for cause and effect conclusions; that is, longitudinal studies would allow for better evidence of associations. Additionally, due to the need to comply with social distancing measures, population-based face-to-face interviews were not possible, except for the MSM who did not have access to the Internet and could not complete the electronic form. Finally, a probabilistic sampling procedure was not adopted here, which impedes the generalization of results to other contexts or countries, though the results provide a situational diagnosis of the MSM's psychological well-being in the pandemic context.

Conclusion

The factors associated with the MSM's low psychological well-being were: belonging to the youngest age group; being in a polyamorous relationship; not complying with the social isolation measures; not using social media to find partners; having multiple sexual partners; having reduced the number of partners, and group sex.

The planning and implementation of public policies and actions to promote psychological well-being and mental health in general, intended to increase resilience among MSM are needed. Safe sexual strategies and behavior need to be promoted, along with the use of digital technologies to facilitate social relationships by enabling remote interactions. These measures can alleviate the harmful effects of the intensified psychological needs accruing from the pandemic context, social isolation measures, and maladaptive coping strategies.

Compliance with Ethical Standards

Disclosure of potential conflicts of interest

The authors report not having any potential conflicts of interest (financial or non-financial)

Research involving Human Participants and/or Animals

All ethical processes involving human beings were followed and the research project was approved by the Institutional Review Board at the Universidade Nova de Lisboa and the University of São Paulo, before data collection begins.

Informed consent

All the users signed free and informed consent forms online before proceeding with the questionnaire.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Acknowledgements

Not applicable.

Authors' contributions

Sousa AFL and IAMCM conceptualised the study. Sousa AFL, IAMCM and ELSC implemented the study and supported data collection. Sousa AFL, IAMCM, ELSC, BIAO, ARS and JRBT analysed the data and drafted the manuscript. All authors write, read and approved the final manuscript.

Funding

Conselho Nacional de Ensino e Pesquisa – CNPq

Process: 159908/2019-1

References

- Badahdah, A.M., Khamis, F. & Mahyjarib, N.A. (2020). The psychological well-being of physicians during COVID-19 outbreak in Oman. *Psychiatry Res*,289, 113053. doi: 10.1016/j.psychres.2020.113053
- Brennan, D.J., Card, K.G., Collict, D., Jollimore, J. & Lachowsky, N.J.(2020). How Might Social Distancing Impact Gay, Bisexual, Queer, Trans and Two-Spirit Men in Canada? *AIDS Behav*, 30,1–3. doi: 10.1007/s10461-020-02891-5 [Epub ahead of print]
- Brennan, D. J., Card, K. G., Collict, D., Jollimore, J., & Lachowsky, N. J. (2020). How Might Social Distancing Impact Gay, Bisexual, Queer, Trans and Two-Spirit Men in Canada?. *AIDS Behav*, 1.
- Cardoso, J. G. M., Paz, B. M., Rocha, K. B., & Pizzinato, A. (2019). Imagem, corpo e linguagem em usos do aplicativo Grindr. *Psicologia USP*, 30. <http://dx.doi.org/10.1590/0103-6564e180160>.
- Carrico, A. W., Horvath, K. J., Grov, C., Moskowitz, J. T., Pahwa, S., Pallikkuth, S., & Hirshfield, S. (2020). Double jeopardy: methamphetamine use and HIV as risk factors for COVID-19. *AIDS Behav*, 1.
- Carvalho, H. E. F., Schneider, G., de Sousa, A. R., Camargo, E. L. S., Nunes, R. V., Possani, M. A., ... & Sousa, Á. F. L.(2020) Síndrome gripal suspeita de covid-19 em homens que se envolveram em sexo casual. [*scielo pre print*]
- Carville, O., & Lanxon, N. (2020) How to date online in the age of COVID-19. Bloomberg.
- Chau, E. Dating app usage up thanks to COVID-19, study suggests. Toronto Sun. 2020. <https://torontosun.com/life/relationships/dating-app-usage-up-thanks-to-covid-19study-suggests>.
- Cohen, R. I. S., & Bosk, E. A. (2020). Vulnerable youth and the COVID-19 pandemic. *Pediatrics*, 146(1).
- Comitê Permanente entre Agências (IASC) (2007). Diretrizes da IASC sobre Saúde Mental e Apoio Psicossocial em Ambientes de Emergência. Genebra: IASC.

- Conley, T.D., Matsick, J.L., Moors, A.C., & Ziegler, A. (2017). Investigation of consensually nonmonogamous relationships. *Perspectives on Psychological Science*, 12, 205-32. DOI: 10.1177/1745691616667925.
- Conley, T.D., Piemonte, J.L., Gusakova, S., & Rubin, J.D. (2018). Sexual satisfaction among individuals in monogamous and consensually non-monogamous relationships. *Journal of Social and Personal Relationships*. 35, 509-531. DOI: 10.1177/0265407517743078.
- Correio do Povo. LGBTIs vivem acirramento de violência familiar em isolamento social Dificuldade de denunciar agressores próximos aumenta durante a pandemia. São Paulo, 2020. Available from: <https://www.correiodopovo.com.br/not%C3%ADcias/geral/lgbtis-vivem-acirramento-de-viol%C3%A2ncia-familiar-em-isolamento-social-1.423485>
- Costa, P.A, & Ribeiro-Gonçalves, J.A. (2020). Não monogamia consensual: atitudes e experiências de pessoas heterossexuais, homossexuais e plurissexuais. *Psic Saúde & Doenças*, 21(1), 104-110. <http://dx.doi.org/10.15309/20psd210116>.
- Dong, E., Du, H., & Gardner, L. (2020). Um painel interativo baseado na Web para rastrear o COVID-19 em tempo real. *Lancet Infect Dis*. 20 (5), 533-4. [https://doi.org/10.1016/S1473-3099\(20\)30120-1](https://doi.org/10.1016/S1473-3099(20)30120-1)
- Donna Redação. Aplicativos para tirar a vida sexul da rotina. Porto Alegre, 2019. Available from: <https://gauchazh.clicrbs.com.br/donna/sexo-e-relacionamento/noticia/2016/06/sexytech-8-aplicativos-para-tirar-a-vida-sexual-da-rotina-cjpl7floa00iowscnxrc1lf11.html>
- Esterwood, E., & Saeed, S. A. (2020). Past Epidemics, Natural Disasters, COVID19, and Mental Health: Learning from History as we Deal with the Present and Prepare for the Future. *Psychiatric Quarterly*, 1-13.
- Folha de São Paulo. Relatos de pessoas que estiveram com presidente na epidemia descrevem momentos de tensão. São Paulo, 2020. Available from:

<https://www1.folha.uol.com.br/colunas/monicabergamo/2020/07/mascara-e-coisa-de-v-dizia-bolsonaro-na-frente-de-visitas.shtml>

Forte, G., Favieri, F., Tambelli, R., & Casagrande, M. (2020) COVID-19 Pandemic in the Italian Population: Validation of a Post-Traumatic Stress Disorder Questionnaire and Prevalence of PTSD Symptomatology. *Int J Environ Res Public Health*, 10;17(11),4151. doi: 10.3390/ijerph17114151.

Fukuti, P., Uchôa, C. L. M., Mazzoco, M. F., Corchs, F., Kamitsuji, C. S., Rossi, L. D., ... & Miguel, E. C. (2020). How Institutions Can Protect the Mental Health and Psychosocial Well-Being of Their Healthcare Workers in the Current COVID-19 Pandemic. *Clinics*, 75.

G1. Mapeamento mostra aumento do consumo de mídia online no Brasil durante a quarentena. São Paulo, 2020. Available from: <https://g1.globo.com/economia/midia-e-marketing/noticia/2020/04/16/mapeamento-mostra-aumenta-consumo-de-midia-online-no-brasil-durante-a-quarentena.ghtml>

Greenhalgh, H. One in three gay men feel unsafe at home during coronavirus. Available from: <https://news.trust.org/item/20200512170907-1h2mg/>

GRINDR. 2020. Disponível em: <https://www.grindr.com/>

Hart, T. A., Noor, S. W., Adam, B. D., Vernon, J. R., Brennan, D. J., Gardner, S., ... & Myers, T. (2017). Number of psychosocial strengths predicts reduced HIV sexual risk behaviors above and beyond syndemic problems among gay and bisexual men. *AIDS Behav*, 21(10), 3035-3046. <https://www.bloomberg.com/news/articles/2020-03-20/online-dating-in-a-pandemic-coronavirus-keeps-singles-apart>

Jowett, A. (2020). The Psychological Impact of Social Distancing on Gender, Sexuality and Relationship Diverse populations. *Psychology of Sexualities Section Review*, 11(1), 6-8.

- Jungmann, S. M., & Witthöft, M. (2020). Health anxiety, cyberchondria, and coping in the current COVID-19 pandemic: Which factors are related to coronavirus anxiety?. *Journal of Anxiety Disorders*, 102239.
- Lai, J., Ma, S., & Wang, Y. (2020) Fatores associados a resultados de saúde mental entre profissionais de saúde expostos à doença de Coronavírus 2019. *JAMA Netw*, 3 (3) doi: 10.1001 / jamanetworkopen.2020.3976.
- Ko, N. Y., Lu, W. H., Chen, Y. L., Li, D. J., Wang, P. W., Hsu, S. T., ... & Yen, C. F. (2020). COVID-19-related information sources and psychological well-being: An online survey study in Taiwan. *Brain Behavior, and Immunity*. Doi: 10.1016 / j.bbi.2020.05.019
- Newman, P. A., & Guta, A. (2020). How to Have Sex in an Epidemic Redux: Reinforcing HIV Prevention in the COVID-19 Pandemic. *AIDS Behav*, 1.
- O Estado de São Paulo. Usuários de apps de paquera se reinventam na quarentena para driblar carência. São Paulo, 2020. Available from: <https://emails.estadao.com.br/noticias/comportamento,usuarios-de-apps-de-paquera-se-reinventam-na-quarentena-para-driblar-carencia,70003267446>
- Portal UOL. Sexo a três: app te ajuda a realizar a fantasia em tempos de coronavírus. São Paulo, 2020. Available from: <https://sexting.blogosfera.uol.com.br/2020/04/03/sexo-a-tres-app-te-ajuda-a-realizar-a-fantasia-em-tempos-de-coronavirus/?cmpid=copiaecola>
- Queiroz, A. A. F. L., de Sousa, A. F. L., Brignol, S., Araújo, T. M. E., & Reis, R. K. (2019). Vulnerability to HIV among older men who have sex with men users of dating apps in Brazil. *Braz J Infect Dis*, 23(5), 298-306. <https://doi.org/10.1016/j.bjid.2019.07.005>
- Queiroz, A. A. F. L. N., Sousa, Á. F. L., Matos, M. C. B., de Araújo, T. M. E., Brignol, S., Reis, R. K., ... & Moura, M. E. B. (2019). Factors associated with self-reported non-completion of the hepatitis B vaccine series in men who have sex with men in Brazil. *BMC infectious diseases*, 19(1), 335.

- Queiroz, A. A. F. L., Sousa, A. F. L., Brignol, S., Araújo, T. M. E., & Reis, R. K. (2019). Vulnerability to HIV among older men who have sex with men users of dating apps in Brazil. *Braz J Infect Dis*, 23(5), 298-306.
- Sahni, H., & Sharma, H. (2020). Role of social media during the COVID-19 pandemic: Beneficial, destructive, or reconstructive?. *International Journal of Academic Medicine*, 6(2), 70.
- Sanchez, T. H., Zlotorzynska, M., Rai, M., & Baral, S. D. (2020). Characterizing the Impact of COVID-19 on men who have sex with men across the United States in April, 2020. *AIDS and Behavior*, 1-9. doi: 10.1007/s10461-020-02894-2 [Epub ahead of print
- Saraiva, L. A. S., Santos, L. T. D., & Pereira, J. R. (2020). Heteronormatividade, Masculinidade e Preconceito em Aplicativos de Celular: O Caso do Grindr em uma Cidade Brasileira. *BBR. Brazilian Business Review*, 17(1), 114-131. <https://doi.org/10.15728/bbr.2020.17.1.6>.
- Serafini, G., Parmigiani, B., Amerio, A., Aguglia, A., Sher, L., & Amore, M. (2020). The psychological impact of COVID-19 on the mental health in the general population.
- Sousa, A. F. L., Oliveira, L. B., Schneider, G., Queiroz, A. A. F. L., de Carvalho, H. E. F., de Araujo, T. M. E., ... & Fronteira, I. (2020). Casual sex among MSM during the period of social isolation in the COVID-19 pandemic: Nationwide study in Brazil and Portugal. medRxiv.doi: <https://doi.org/10.1101/2020.06.07.20113142>
- Sousa, A. F. L., Queiroz, A. A. F. L. N., Fronteira, I., Lapão, L., Mendes, I. A. C., & Brignol, S. (2019). HIV Testing Among Middle-Aged and Older Men Who Have Sex With Men (MSM): A Blind Spot?. *American journal of men's health*, 13(4), 1557988319863542. DOI: 10.1177/1557988319863542 journals.sagepub.com/home/jm
- Sousa, A. R., Carvalho, E. S. S., Santana, T. S., Sousa, A. F. L., Figueiredo, T. F. G., Escobar, O. J. V., Mota, T. N., Pereira, A. (2020) Sentimentos e emoções de homens no enquadramento da doença Covid-19. *Ciência & Saúde Coletiva*.

- Suen, Y. T., Chan, R. C. H., & Wong, E. M. Y. (2020). Effects of general and sexual minority-specific COVID-19-related stressors on the mental health of lesbian, gay and bisexual people in Hong Kong. *Psychiatry Research*, 113365.
- Sullivan A. Love in the time of coronavirus: COVID-19 changes the game for online dating. Deutsche Welle. 2020. Available from: <https://www.dw.com/en/love-in-the-time-of-coronaviruscovid-19-changes-the-game-for-online-dating/a-52933001>.
- Torres, T. S., Hoagland, B., Bezerra, D. R., Garner, A., Jalil, E. M., Coelho, L. E., ... & Veloso, V. G. (2020). Impact of COVID-19 Pandemic on Sexual Minority Populations in Brazil: An Analysis of Social/Racial Disparities in Maintaining Social Distancing and a Description of Sexual Behavior. *AIDS Behav*, 1-12.
- World Health Organization. (2020). WHO Coronavirus Disease (COVID-19) Dashboard. Available from: <https://covid19.who.int>
- Coronavirus Brasil. (2020). COVID19 Painel Coronavírus. Available from <https://covid.saude.gov.br>
- Young Minds. (2020). Coronavirus: Impact on young people with mental health needs. Available from: <https://youngminds.org.uk/media/3904/coronavirus-report-summer-2020-final.pdf>