Molecular Architecture of Early Dissemination and Massive Second Wave of the SARS-CoV-2 Virus in a Major Metropolitan Area

Running Title: Two waves of COVID-19 disease in Houston, Texas

S. Wesley Long,[a,b,1] Randall J. Olsen,[a,b,1] Paul A. Christensen,[a,1] David W. Bernard,[a,b] James J. Davis,[c,d] Maulik Shukla,[c,d] Marcus Nguyen,[c,d] Matthew Ojeda Saavedra,[a] Prasanti Yerramilli,[a] Layne Pruitt,[a] Sishir Subedi,[a] Hung-Che Kuo,[e] Heather Hendrickson,[a] Ghazaleh Eskandari,[a] Hoang A. T. Nguyen,[a] J. Hunter Long,[a] Muthiah Kumaraswami,[a] Jule Goike,[e] Daniel Boutz,[f] Jimmy Gollihar,[a,f] Jason S. McLellan,[e] Chia-Wei Chou,[e] Kamyab Javanmardi,[e] Ilya J. Finkelstein,[e,g], and James M. Musser[a,b#]

[a]Center for Molecular and Translational Human Infectious Diseases Research, Department of Pathology and Genomic Medicine, Houston Methodist Research Institute and Houston Methodist Hospital, 6565 Fannin Street, Houston, Texas 77030

[b]Departments of Pathology and Laboratory Medicine, and Microbiology and Immunology, Weill Cornell Medical College, 1300 York Avenue, New York, New York 10065

[c]Consortium for Advanced Science and Engineering, University of Chicago, 5801

South Ellis Avenue, Chicago, Illinois, 60637

[d]Computing, Environment and Life Sciences, Argonne National Laboratory, 9700

South Cass Avenue, Lemont, Illinois 60439

[e]Department of Molecular Biosciences and Institute of Molecular Biosciences,

The University of Texas at Austin, Austin, Texas 78712

[f]CCDC Army Research Laboratory-South, University of Texas, Austin, Texas 78712

[g]Center for Systems and Synthetic Biology, University of Texas at Austin, Austin,

Texas 78712

[1]S.W.L., R.J.O., and P.A.C. contributed equally to this article. The order of co-first

authors was determined by discussion and mutual agreement between the three

co-first authors.

[#]Address correspondence to: James M. Musser, M.D., Ph.D., Department of

Pathology and Genomic Medicine, Houston Methodist Research Institute, 6565

Fannin Street, Suite B490, Houston, Texas 77030. Tel: 713.441.5890, E-mail:

jmmusser@houstonmethodist.org.

44   Health, New Jersey; and David M. Morens, National Institute of Allergy and

45   Infectious Diseases, National Institutes of Health, Maryland.

ABSTRACT   We sequenced the genomes of 5,085 SARS-CoV-2 strains

causing two COVID-19 disease waves in metropolitan Houston, Texas, an

ethnically diverse region with seven million residents. The genomes were

from viruses recovered in the earliest recognized phase of the pandemic in

Houston, and an ongoing massive second wave of infections. The virus

was originally introduced into Houston many times independently. Virtually

all strains in the second wave have a Gly614 amino acid replacement in the

spike protein, a polymorphism that has been linked to increased

transmission and infectivity. Patients infected with the Gly614 variant

strains had significantly higher virus loads in the nasopharynx on initial

diagnosis. We found little evidence of a significant relationship between

virus genotypes and altered virulence, stressing the linkage between

disease severity, underlying medical conditions, and host genetics. Some

regions of the spike protein - the primary target of global vaccine efforts -

are replete with amino acid replacements, perhaps indicating the action of

selection. We exploited the genomic data to generate defined single amino

acid replacements in the receptor binding domain of spike protein that,

importantly, produced decreased recognition by the neutralizing

monoclonal antibody CR30022. Our study is the first analysis of the

molecular architecture of SARS-CoV-2 in two infection waves in a major

metropolitan region. The findings will help us to understand the origin,

composition, and trajectory of future infection waves, and the potential

effect of the host immune response and therapeutic maneuvers on SARS-

CoV-2 evolution.

IMPORTANCE   There is concern about second and subsequent waves of

COVID-19 caused by the SARS-CoV-2 coronavirus occurring in

communities globally that had an initial disease wave. Metropolitan

Houston, Texas, with a population of 7 million, is experiencing a massive

second disease wave that began in late May 2020. To understand SARS-

CoV-2 molecular population genomic architecture, evolution, and

relationship between virus genotypes and patient features, we sequenced

the genomes of 5,085 SARS-CoV-2 strains from these two waves. Our study

provides the first molecular characterization of SARS-CoV-2 strains

causing two distinct COVID-19 disease waves.

KEYWORDS: SARS-CoV-2, COVID-19 disease, genome sequencing, molecular

population genomics, evolution

[Introduction]

Pandemic disease caused by the severe acute respiratory syndrome

coronavirus 2 (SARS-CoV-2) virus is now responsible for massive human

morbidity and mortality worldwide (1-5). The virus was first documented to cause

severe respiratory infections in Wuhan, China, beginning in late December 2019

(6-9). Global dissemination occurred extremely rapidly and has affected major

population centers on most continents (10, 11). In the United States, the Seattle

and the New York City (NYC) regions have been especially important centers of

COVID-19 disease caused by SARS-CoV-2. For example, as of August 19,

2020, there were 227,419 confirmed SARS-CoV-2 cases in NYC, causing 56,831

hospitalizations and 19,005 confirmed fatalities and 4,638 probable fatalities (12).

Similarly, in Seattle and King County, 17,989 positive patients and 696 deaths

have been reported as of August 18, 2020 (13).

The Houston metropolitan area is the fourth largest and most ethnically

diverse city in the United States, with a population of approximately 7 million

(14, 15). The 2,400-bed Houston Methodist health system has seven hospitals

and serves a large, multiethnic, and socioeconomically diverse patient population

throughout greater Houston (13, 14). The first COVID-19 case in metropolitan

Houston was reported on March 5, 2020 with community spread occurring one

week later (16). Many of the first cases in our region were associated with

national or international travel in areas known to have SARS-CoV-2 virus

outbreaks (16). A central molecular diagnostic laboratory serving all Houston

Methodist hospitals and our very early adoption of a molecular test for the SARS-

CoV-2 virus permitted us to rapidly identify positive patients and interrogate

genomic variation among strains causing early infections in the greater Houston

area. Our analysis of SARS-CoV-2 genomes causing disease in Houston has

continued unabated since early March and is ongoing. Genome sequencing and

related efforts were expanded extensively in late May as we recognized that a

prominent second wave was underway (**Figure 1**).

Here, we report that SARS-CoV-2 was introduced to the Houston area

many times, independently, from diverse geographic regions, with virus

genotypes representing genetic clades causing disease in Europe, Asia, South

America and elsewhere in the United States. There was widespread community

dissemination soon after COVID-19 cases were reported in Houston. Strains with

a Gly614 amino acid replacement in the spike protein, a polymorphism that has

been linked to increased transmission and *in vitro* cell infectivity, increased

significantly over time and caused virtually all COVID-19 cases in the massive

second disease wave. Patients infected with strains with the Gly614 variant had

significantly higher virus loads in the nasopharynx on initial diagnosis. Some

naturally occurring single amino acid replacements in the receptor binding

domain of spike protein resulted in decreased reactivity with a neutralizing

monoclonal antibody, consistent with the idea that some virus variants arise due

to host immune pressure.

RESULTS

Description of metropolitan Houston. Houston, Texas, is located in the

southwestern United States, 50 miles inland from the Gulf of Mexico. It is the

most ethnically diverse city in the United States (14). Metropolitan Houston is

comprised predominantly of Harris County plus parts of eight contiguous

surrounding counties. In the aggregate, the metropolitan area includes 9,444

square miles**.** The estimated population size of metropolitan Houston is 7 million

(https://www.houston.org/houston-data).

Epidemic curve characteristics over two disease waves. The first

confirmed case of COVID-19 in the Houston metropolitan region was reported on

March 5, 2020 (16), and the first confirmed case diagnosed in Houston Methodist

hospitals was reported on March 6, 2020. The epidemic curve indicated a first

wave of COVID-19 cases that peaked around April 11-15, followed by a decline

in cases until May 11. Soon thereafter, the slope of the case curve increased with

a very sharp uptick in confirmed cases beginning on June 12 (**Figure 1B**). We

consider May 11 as the transition between waves, as this date is the inflection

point of the cumulative new cases curve and had the absolute lowest number of

new cases in the mid-May time period. Thus, for the data presented herein, wave

1 is defined as March 5 through May 11, 2020, and wave 2 is defined as May 12

through July 7, 2020. Epidemiologic trends within the Houston Methodist Hospital

population were mirrored by data from Harris County and the greater

metropolitan Houston region (**Figure 1A**). Through the 7[th] of July, 25,366

COVID-19 cases were reported in Houston, 37,776 cases in Harris County, and

53,330 in metropolitan Houston, including 9,823 cases in Houston Methodist

facilities (inpatients and outpatients) (https://www.tmc.edu/coronavirus-

updates/infection-rate-in-the-greater-houston-area/ and

https://harriscounty.maps.arcgis.com/apps/opsdashboard/index.html#/c0de71f8e

a484b85bb5efcb7c07c6914).

During the first wave (early March through May 11), 11,476 COVID-19

cases were reported in Houston, including 1,729 cases in the Houston Methodist

Hospital system. Early in the first wave (from March 5 through March 30, 2020),

we tested 3,080 patient specimens. Of these, 406 (13.2%) samples were positive

for SARS-CoV-2, representing 40% (358/898) of all confirmed cases in

metropolitan Houston during that time period. As our laboratory was the first

hospital-based facility to have molecular testing capacity for SARS-CoV-2

available on site, our strain samples are likely representative of COVID-19

infections during the first wave.

For the entire study period (March 5 through July 7, 2020), we tested

68,418 specimens from 55,800 patients. Of these, 9,121 patients (16.4%) had a

positive test result, representing 17.1% (9,121/53,300) of all confirmed cases in

metropolitan Houston. Thus, our strain samples are also representative of those

responsible for COVID-19 infections in the massive second wave.

To test the hypothesis that, on average, the two waves affected different

groups of patients, we analyzed individual patient characteristics (hospitalized

and non-hospitalized) in each wave. Consistent with this hypothesis, we found

significant differences in the COVID-19 patients in each wave (**Table S1**). For

example, patients in the second wave were significantly younger, had fewer

comorbidities, were more likely to be Hispanic/Latino (by self-report), and lived in

zip codes with lower median incomes (**Table S1**). A detailed analysis of the

characteristics of patients hospitalized in Houston Methodist facilities in the two

waves has recently been published (17).

SARS-CoV-2 genome sequencing and phylogenetic analysis. To

investigate the genomic architecture of the virus across the two waves, we

sequenced the genomes of 5,085 SARS-CoV-2 strains dating to the earliest time

of confirmed COVID-19 cases in Houston. Analysis of SARS-CoV-2 strains

causing disease in the first wave (March 5 through May 11) identified the

presence of many diverse virus genomes that, in the aggregate, represent the

major clades identified globally to date (**Figure 1B**). Clades G, GH, GR, and S

were the four most abundantly represented phylogenetic groups (**Figure 1B**).

Strains with the Gly614 amino acid variant in spike protein represented 82% of

the SARS-CoV-2 strains in wave 1, and 99.9% in wave 2 ($p<0.0001$; Fisher's

exact test) (**Figure 1B**). This spike protein variant is characteristic of clades G,

GH, and GR. Importantly, strains with the Gly614 variant represented only 71%

of the specimens sequenced in March, the early part of wave 1 (**Figure 1B**). We

attribute the decrease in strains with this variant observed in the first two weeks

of March (**Figure 1B**) to fluctuation caused by the relatively fewer COVID-19

cases occurring during this period.

Relating spatiotemporal genome analysis with virus genotypes over

two disease waves. We examined the spatial and temporal mapping of genomic

data to investigate community spread during wave 1 (**Figure 2**). Rapid and

widespread community dissemination occurred soon after the initial COVID-19

cases were reported in Houston. The heterogenous virus genotypes present very

early in wave 1 indicate that multiple strains independently entered metropolitan

Houston, rather than introduction and spread of a single strain. An important

observation was that strains of most of the individual subclades were distributed

over broad geographic areas (**Figure S1**). These findings are consistent with the

known ability of SARS-CoV-2 to spread very rapidly from person to person.

Relationship between virus clades, clinical characteristics of infected

patients, and additional metadata. It is possible that SARS-CoV-2 genome

subtypes have different clinical characteristics, analogous to what is believed to

have occurred with Ebola virus (18-20) and known to occur for other pathogenic

microbes (21). As an initial examination of this issue in SARS-CoV-2, we tested

the hypothesis that patients with disease severe enough to warrant

hospitalization were infected with a non-random subset of virus genotypes. We

also examined the association between virus clades and disease severity based

on overall mortality, highest level of required care (intensive care unit,

intermediate care unit, inpatient or outpatient), need for mechanical ventilation,

and length of stay. There was no simple relationship between virus clades and

disease severity using these four indicators. Similarly, there was no simple

relationship between virus clades and other metadata, such as sex, age, or

ethnicity (**Figure S2**).

Machine learning analysis. Machine learning models can be used to

identify complex relationships not revealed by statistical analyses. We built

machine learning models to test the hypothesis that virus genome sequence can

predict patient outcomes including mortality, length of stay, level of care, ICU

admission, supplemental oxygen use, and mechanical ventilation. Models to

predict outcomes based on virus genome sequence alone resulted in low F1

scores less than 50% (0.41 – 0.49) and regression models showed similarly low

R^2 values (-0.01 – -0.20) (**Table S2**). F1 scores near 50% are indicative of

classifiers that are performing similarly to random chance. The use of patient

metadata alone to predict patient outcome improved the model's F1 scores by 5-

10% (0.51 – 0.56) overall. The inclusion of patient metadata with virus genome

sequence data improved most predictions of outcomes, compared to genome

sequence alone, to 50% to 55% F1 overall (0.42 – 0.55) in the models (**Table

S2**). The findings are indicative of two possibilities that are not mutually

exclusive. First, patient metadata, such as age and sex, may provide more signal

for the model to use and thus result in better accuracies. Second, the model's

use of single nucleotide polymorphisms (SNPs) may have resulted in overfitting.

Most importantly, no SNP predicted a significant difference in outcome. A table of

classifier accuracy scores and performance information is provided in **Table S2**.

Patient outcome and metadata correlations. Overall, very few metadata

categories correlated with patient outcomes (**Table S3**). Mortality was

independently correlated with both Rh factor-positive blood type and increasing

age, with Pearson correlation coefficients (PCC) equal to 0.22 and 0.27,

respectively. This means that 22-27% of the variation in mortality can be

predicted from Rh factor-positive blood type and patient age, respectively. Length

of stay correlated independently with increasing age (PCC=0.20). Positive Rh

factor also correlated with ICU length of stay (PCC=0.20). All other patient

metadata correlations to outcomes had PCC less than 0.20 (**Table S3**).

We further analyzed outcomes correlated to isolates from wave 1 and 2,

and the presence of the Gly614 variant in spike protein. Being in wave 1 was

independently correlated with mechanical ventilation days, overall length of stay,

and ICU length of stay, with PCC equal to 0.20, 0.18, and 0.14, respectively.

Importantly, the presence of the Gly614 variant did not correlate with patient

outcomes (**Table S3**).

Analysis of the *nsp12* polymerase gene. The SARS-CoV-2 genome

encodes an RNA-dependent RNA polymerase (RdRp, also referred to as Nsp12)

used in virus replication (22-25). Two amino acid substitutions (Phe479Leu and

Val556Leu) in RdRp each confer significant resistance *in vitro* to remdesivir, an

adenosine analog (26). Remdesivir is inserted into RNA chains by RdRp during

replication, resulting in premature termination of RNA synthesis and inhibition of

virus replication. This compound has shown prophylactic and therapeutic benefit

against MERS-CoV and SARS-CoV-2 experimental infection in rhesus

macaques (27, 28). Recent reports indicate that remdesivir has therapeutic

benefit in some COVID-19 hospitalized patients (29-33), leading it to be now

widely used in patients worldwide. Thus, it may be important to understand

variation in RdRp in large strain samples.

To acquire data about allelic variation in the *nsp12* gene, we analyzed our

5,085 virus genomes. The analysis identified 265 SNPs, including 140

nonsynonymous (amino acid-altering) SNPs, resulting in amino acid

replacements throughout the protein (**Table 1, Figure 3, Figure 4, Figure S3,**

and **Figure S4**). The most common amino acid change was Pro322Leu,

identified in 4,893 of the 5,085 (96%) patient isolates. This amino acid

replacement is common in genomes from clades G, GH, and GR, which are

distinguished from other SARS-CoV-2 clades by the presence of the Gly614

amino acid change in the spike protein. Most of the other amino acid changes in

RdRp were present in relatively small numbers of strains, and some have been

identified in other isolates in a publicly available database (34). Five prominent

exceptions included amino acid replacements: Ala15Val in 138 strains, Met462Ile

in 59 strains, Met600Ile in 75 strains, Thr907Ile in 45 strains, and Pro917Ser in

80 strains. All 75 Met600Ile strains were phylogenetically closely related

members of clade G, and also had the Pro322Leu amino acid replacement

characteristic of this clade (**Figure S3**). These data indicate that the Met600Ile

change is likely the evolved state, derived from a precursor strain with the

Pro322Leu replacement. Similarly, we investigated phylogenetic relationships

among strains with the other four amino acid changes noted above. In all cases,

the vast majority of strains with each amino acid replacement were found among

individual subclades of strains (**Figure S3**).

Importantly, none of the observed amino acid polymorphisms in RdRp

were located precisely at two sites known to cause *in vitro* resistance to

remdesivir (26). Most of the amino acid changes are located distantly from the

RNA-binding and catalytic sites (**Figure S4** and **Table 1**). However,

replacements at six amino acid residues (Ala442Val, Ala448Val, Ala553Pro/Val,

Gly682Arg, Ser758Pro, and Cys812Phe) may potentially interfere with either

remdesivir binding or RNA synthesis. Four (Ala442Val, Ala448Val,

Ala553Pro/Val, and Gly682Arg) of the six substitution sites are located

immediately above the nucleotide-binding site, that is comprised of Lys544,

Arg552, and Arg554 residues as shown by structural studies (**Figure 4**). The

positions of these four variant amino acid sites are comparable to Val556 (**Figure**

4), for which a Val556Leu mutation in SARS-CoV was identified to confer

resistance to remdesivir *in vitro* (26). The other two substitutions (Ser758Pro and

Cys812Phe) are inferred to be located either at, or in the immediate proximity of,

the catalytic active site, that is comprised of three contiguous residues (Ser758,

Asp759, and Asp760). A proline substitution we identified at Ser758 (Ser758Pro)

is likely to negatively impact RNA synthesis. Although Cys812 is not directly

involved in the catalysis of RNA synthesis, it is only 3.5 Å away from Asp760.

The introduction of the bulkier phenylalanine substitution at Cys812 (Cys812Phe)

may impair RNA synthesis. Consequently, these two substitutions are expected

to detrimentally affect virus replication or fitness.

Analysis of the gene encoding the spike protein. The densely glycosylated

spike protein of SARS-CoV-2 and its close coronavirus relatives binds directly to

host-cell angiotensin-converting enzyme 2 (ACE2) receptors to enter host cells

(35-37). Thus, the spike protein is a major translational research target, including

intensive vaccine and therapeutic antibody (35-64). Analysis of the gene

encoding the spike protein identified 470 SNPs, including 285 that produce

amino acid changes (**Table 2, Figure 5**). Forty-nine of these replacements

(V11A, T51A, W64C, I119T, E156Q, S205A, D228G, L229W, P230T, N234D,

I235T, T274A, A288V, E324Q, E324V, S325P, S349F, S371P, S373P, T385I,

A419V, C480F, Y495S, L517F, K528R, Q628E, T632I, S708P, T719I, P728L,

S746P, E748K, G757V, V772A, K814R, D843N, S884A, M902I, I909V, E918Q,

S982L, M1029I, Q1142K, K1157M, Q1180R, D1199A, C1241F, C1247G, and

V1268A) are not represented in a publicly available database (34) as of August

19, 2020. Interestingly, 25 amino acid sites have three distinct variants (that is,

the reference amino acid plus two additional variant amino acids), and five amino

acid sites (amino acid positions 21, 27, 228, 936, and 1050) have four distinct

variants represented in our sample of 5,085 genomes (**Table 2, Figure 5**).

We mapped the location of amino acid replacements onto a model of the

full-length spike protein (35, 65) and observed that the substitutions are found in

each subunit and domain of the spike (**Figure 6**). However, the distribution of

amino acid changes is not uniform throughout the protein regions. For example,

compared to some other regions of the spike protein, the receptor binding

domain (RBD) has relatively few amino acid changes, and the frequency of

strains with these substitutions is low, each occurring in fewer than 10 isolates.

This finding is consistent with the functional constraints on RBD to mediate

interaction with ACE2. In contrast, the periphery of the S1 subunit NTD contains

a dense cluster of substituted residues, with some single amino acid

replacements found in 10–20 isolates (**Table 2, Figure 5, Figure 6**). Clustering

of amino acid changes in a distinct region of the spike protein may be a signal of

positive selection. Inasmuch as infected patients make antibodies against the

NTD, we favor the idea that host immune selection is one force contributing to

some of the amino acid variation in this region. One NTD substitution, H49Y, was

found in 142 isolates. This position is not well exposed on the surface of the NTD

and is likely not a result of immune pressure. The same is true for another highly

represented substitution, F1052L. This substitution was observed in 167 isolates,

and F1052 is buried within the core of the S2 subunit. The substitution observed

most frequently in the spike protein in our sample is D614G, a change observed

in 4,895 of the isolates. As noted above, strains with the Gly614 variant

significantly increased in wave 2 compared to wave 1.

As observed with RdRp, the majority of strains with each single amino

acid change in the spike protein were found on a distinct phylogenetic lineage

(**Figure S5**), indicating identity by descent. A prominent exception is the

Leu5Phe replacement that is present in all major clades, suggesting that this

amino acid change arose multiple times independently or very early in the course

of SARS-CoV-2 evolution. Finally, we note that examination of the phylogenetic

distribution of strains with multiple distinct amino acid replacements at the same

site (e.g., Arg21Ile/Lys/Thr, Ala27Ser/Thr/Val, etc.) revealed that they were

commonly found in different genetic branches, consistent with independent origin

(**Figure S5**).

Cycle threshold (Ct) comparison of SARS-CoV-2 strains with either

the Asp614 or Gly614 amino acid replacements in spike protein. It has been

reported that patients infected with strains having spike protein Gly614 variant

have, on average, higher virus loads on initial diagnosis (66-70). To determine if

this is the case in Houston strains, we examined the cycle threshold (Ct) for

every sequenced strain that was detected from a patient specimen using the

SARS-CoV-2 Assay done by the Hologic Panther instrument. We identified a

significant difference ($p<0.0001$) between the mean Ct value for strains with an

Asp614 ($n=102$) or Gly614 ($n=812$) variant of the spike protein (**Figure 7**).

Strains with Gly614 had a Ct value significantly lower than strains with the

Asp614 variant, indicating that patients infected with the Gly614 strains had, on

average, higher virus loads on initial diagnosis than patients infected by strains

with the Asp614 variant (**Figure 7**). This observation is consistent with the

conjecture that, on average, strains with the Gly614 variant are better able to

disseminate.

Characterization of recombinant proteins with single amino acid

replacements in the receptor binding domain region of spike protein. The

RBD of spike protein binds the ACE2 surface receptor and is also targeted by

neutralizing (36, 37, 41, 43-46, 48-62, 71). Thus, single amino acid replacements

in this domain may have functional consequences that enhance virus fitness. To

begin to test this idea, we expressed spike variants with the Asp614Gly

replacement and 13 clinical RBD variants identified in our genome sequencing

studies (**Figure 8, Table S4, S5**). All RBD variants were cloned into an

engineered spike protein construct that stabilizes the perfusion state and

increases overall expression yield (spike-6P, here referred to as spike) (64).

We first assessed the biophysical properties of spike-Asp614Gly, an

amino acid polymorphism that is common globally and increased significantly in

our wave 2 strain isolates. Pseudotyped viruses expressing spike-Gly614 have

higher infectivity for host cells *in vitro* than spike-Asp614 (66, 67, 69, 72, 73). The

higher infectivity of spike-Gly614 is correlated with increased stability and

incorporation of the spike protein into the pseudovirion (73). We observed a

higher expression level (**Figure 8A, B**) and increased thermostability for the

spike protein construct containing this variant (**Figure 8C, D**). The size exclusion

chromatography (SEC) elution profile of spike-Asp614 was indistinguishable from

spike-Gly614, consistent with a trimeric conformation (**Figure 8A**). These results

are broadly consistent with higher-resolution structural analyses of both spike

variants.

Next, we purified and biophysically characterized 13 RBD mutants that

each contain Gly614 and one additional single amino acid replacement we

identified by genome sequencing our clinical samples (**Table S6**). All variants

eluted as trimers, indicating the global structure, remained intact (**Figure 8** and

Figure S6). However, several variants had reduced expression levels and

virtually all had decreased thermostability relative to the variant that had only a

D614G single amino acid replacement (**Figure 8D**). The A419V and A522V

mutations were especially deleterious, reducing yield and precluding further

downstream analysis (**Figure 8B**). We next assayed the affinity of the 11 highest-

expressing spike variants for ACE2 and the neutralizing monoclonal antibody

CR3022 via enzyme-linked immunosorbent assays (ELISAs) (**Figure 8E-G** and

Table S6). Most variants retained high affinity for the ACE2 surface receptor.

However, importantly, three RBD variants (F338L, S373P, and R408T) had

substantially reduced affinity for CR3022, a monoclonal antibody that disrupts the

spike protein homotrimerization interface (63, 74). Notably, the S373P mutation

is one amino acid away from the epitope recognized by CR3022. These results

are consistent with the interpretation that some RBD mutants arising in COVID-

19 patients may have increased ability to escape humoral immune pressure, but

otherwise retain strong ACE2 binding affinity.

DISCUSSION

In this work we analyzed the molecular population genomics, sociodemographic,

and medical features of two waves of COVID-19 disease occurring in

metropolitan Houston, Texas, between early March and early July 2020. We also

studied the biophysical and immunologic properties of some naturally occurring

single amino acid changes in the spike protein RBD identified by sequencing the

5,085 genomes. We discovered that the first COVID-19 wave was caused by a

heterogenous array of virus genotypes assigned to several different clades. The

majority of cases in the first wave are related to strains that caused widespread

disease in European and Asian countries, as well as other localities. We

conclude that the SARS-CoV-2 virus was introduced into Houston many times

independently, likely by individuals who had traveled to or from different parts of

the world, including other communities in the United States. In support of this

conclusion, the first cases in metropolitan Houston were associated with a travel

history to a known COVID-19 region (16). The data are consistent with the fact

that Houston is a large international city characterized by a multi-ethnic

population and is a prominent transport hub with direct flights to major cities

globally.

The second wave of COVID-19 cases also is characterized by SARS-

CoV-2 strains with diverse genotypes. Virtually all cases in the second and

ongoing disease wave were caused by strains with the Gly614 variant of spike

protein (**Figure 1B**). Our data unambiguously demonstrate that strains with the

Gly614 variant increased significantly in frequency in wave 2 relative to wave 1 in

the Houston metropolitan region. This shift occurred very rapidly in a matter of

just a few months. Amino acid residue Asp614 is located in subdomain 2 (SD-2)

of the spike protein and forms a hydrogen bond and electrostatic interaction with

two residues in the S2 subunit of a neighboring protomer. Replacement of

aspartate with glycine would eliminate both interactions, thereby substantively

weakening the contact between the S1 and S2 subunits. We previously

speculated (75) that this weakening produces a more fusogenic spike protein, as

S1 must first dissociate from S2 before S2 can refold and mediate fusion of virus

and cell membranes. Stated another way, virus strains with the Gly614 variant

may be better able to enter host cells, potentially resulting in enhanced spread.

Consistent with this idea, Korber et al. (66) showed that the Gly614 variant grows

to higher titer as pseudotyped virions. On initial diagnosis infected individuals had

lower RT-PCR cycle thresholds suggesting higher upper respiratory tract viral

loads. Our data (**Figure 7**) are fully consistent with that finding Zhang et al. (73)

reported that pseudovirus with the 614Gly variant infected ACE2-expressing cells

more efficiently than the 614Asp. Similar results have been described by Hu et

al. (67) and Lorenzo-Redondo et al. (68). Plante et al. (76) recently studied

isogenic mutant SARS-CoV-2 strains with either the 614Asp or 614Gly variant

and found that the 614Gly variant virus had significantly increased replication in

human lung epithelial cells *in vitro* and increased infectious titers in nasal and

trachea washes obtained from experimentally infected hamsters. These results

are consistent with the idea that the 614Gly variant bestows increased virus

fitness in the upper respiratory tract (76).

Additional work is needed to investigate the potential biomedical relevance

and public health importance of the Asp614Gly polymorphism, including but not

limited to virus dissemination, overall fitness, impact on clinical course and

virulence, and development of vaccines and therapeutics. Although it is possible

that stochastic processes alone may account for the rapid increase in COVID-19

disease frequency caused by viruses containing the Gly614 variant, we do not

favor that interpretation in part because of the cumulative weight of the

epidemiologic, human RT-PCR diagnostics data, *in vitro* experimental findings,

and animal infection studies using isogenic mutant virus strains. In addition, if

stochastic processes solely are responsible, we believe it is difficult to explain

essentially simultaneous increase in frequency of the Gly614 variant in

genetically diverse viruses in three distinct clades (G, GH, and GR) in a

geographically large metropolitan area with 7 million ethnically diverse people.

Regardless, more research on this important topic is warranted.

Several groups have analyzed the relationship between human genetics,

blood types, and patient outcomes (33, 34, 77-80). In this regard, we identified a

correlation between Rh factor-positive blood type and increased mortality and

length of ICU stay. (Supplemental Table 3). Thus, our data are consistent with

studies suggesting that there are host genetic factors that contribute to disease

severity and outcome. More research is required to address the important

possibility.

The diversity present in our 1,026 virus genomes from the first disease

wave contrasts somewhat with data reported by Gonzalez-Reiche et al., who

studied 84 SARS-CoV-2 isolates causing disease in patients in the New York

City region (11). Those investigators concluded that the vast majority of disease

was caused by progeny of strains imported from Europe. Similarly, Bedford et al.

(10) reported that much of the COVID-19 disease in the Seattle, Washington

area was caused by strains that are progeny of a virus strain recently introduced

from China. Some aspects of our findings are similar to those reported recently

by Lemieux et al. based on analysis of strains causing disease in the Boston

area (81). Our findings, like theirs, highlight the importance of multiple

importation events of genetically diverse strains in the epidemiology of COVID-19

disease in this pandemic. Similarly, Icelandic and Brazilian investigators

documented that SARS-CoV-2 was imported by individuals traveling to or from

many European and other countries (82, 83).

The virus genome diversity and large sample size in our study permitted

us to test the hypothesis that distinct virus clades were nonrandomly associated

with hospitalized COVID-19 patients or disease severity. We did not find

evidence to support this hypothesis, but our continuing study of COVID-19 cases

accruing in the second wave will further improve statistical stratification.

We used machine learning classifiers to identify if any SNPs contribute to

increased infection severity or otherwise affect virus-host outcome. The models

could not be trained to accurately predict these outcomes from the available virus

genome sequence data. This may be due to sample size or class imbalance.

However, we do not favor this interpretation. Rather, we think that the inability to

identify particular virus SNPs predictive of disease severity or infection outcome

likely reflects the substantial heterogeneity in underlying medical conditions and

treatment regimens among COVID-19 patients studied herein. An alternative but

not mutually exclusive hypothesis is that patient genotypes play an important role

in determining virus-human interactions and resulting pathology. Although some

evidence has been presented in support of this idea (84, 85), available data

suggest that in the aggregate, host genetics does not play an overwhelming role

in determining outcome in the great majority of adult patients, once virus infection

is established.

Remdesivir is a nucleoside analog reported to have activity against

MERS-CoV, a coronavirus related to SARS-CoV-2. Recently, several studies

have reported that remdesivir shows promise in treating COVID-19 patients (29-

33), leading the FDA to issue an emergency use authorization. Because *in vitro*

resistance of SARS-CoV to remdesivir has been reported to be caused by either

of two amino acid replacements in RdRp (Phe479Leu or Val556Leu), we

interrogated our data for polymorphisms in the *nsp12* gene. Although we

identified 140 different inferred amino acid replacements in RdRp in the 5,085

genomes analyzed, none of these were located precisely at the two positions

associated with *in vitro* resistance to remdesivir. Inasmuch as remdesivir is now

being deployed widely to treat COVID-19 patients in Houston and elsewhere, our

findings suggest that the majority of SARS-CoV-2 strains currently circulating in

our region should be susceptible to this drug.

The amino acid replacements Ala442Val, Ala448Val, Ala553Pro/Val, and

Gly682Arg that we identified occur at sites that, intriguingly, are located directly

above the nucleotide substrate entry channel and nucleotide binding residues

Lys544, Arg552, and Arg554 (22, 23) (**Figure 4**). One possibility is that

substitution of the smaller alanine or glycine residues with the bulkier side chains

of Val/Pro/Arg may impose structural constraints for the modified nucleotide

analog to bind, and thereby disfavor remdesivir binding. This, in turn, may lead to

reduced incorporation of remdesivir into the nascent RNA, increased fidelity of

RNA synthesis, and ultimately drug resistance. A similar mechanism has been

proposed for a Val556Leu change (23).

We also identified one strain with a Lys477Asn replacement in RdRp. This

substitution is located close to a Phe479Leu replacement reported to produce

partial resistance to remdesivir *in vitro* in SARS-CoV patients from 2004,

although the amino acid positions are numbered differently in SARS-CoV and

SARS-CoV-2. Structural studies have suggested that this amino acid is surface-

exposed, and distant from known key functional elements. Our observed

Lys477Asn change is also located in a conserved motif described as a finger

domain of RdRp (**Figure 3** and **4**). One speculative possibility is that Lys477 is

involved in binding a yet unidentified cofactor such as Nsp7 or Nsp8, an

interaction that could modify nucleotide binding and/or fidelity at a distance.

These data warrant additional study in larger patient cohorts, especially in

individuals treated with remdesivir.

Analysis of the gene encoding the spike protein identified 285 polymorphic

amino acid sites relative to the reference genome, including 49 inferred amino

acid replacements not present in available databases as of August 19, 2020.

Importantly, 30 amino acid sites in the spike protein had two or three distinct

replacements relative to the reference strain. The occurrence of multiple variants

at the same amino acid site is one characteristic that may suggest functional

consequences. These data, coupled with structural information available for

spike protein, raise the possibility that some of the amino acid variants have

functional consequences, for example including altered serologic reactivity and

shown here. These data permit generation of many biomedically relevant

hypotheses now under study.

A recent study reported that RBD amino acid changes could be selected

in vitro using a pseudovirus neutralization assay and sera obtained from

convalescent plasma or monoclonal antibodies (86). The amino acid sites

included positions V445 and E484 in the RBD. Important to note, variants G446V

and E484Q were present in our patient samples. However, these mutations

retain high affinity to CR3022 **(Figure 8F, G)**. The high-resolution structure of the

RBD/CR3022 complex shows that CR3022 makes contacts to residues 369-386,

380-392, and 427-430 of RBD (74). Although there is no overlap between

CR3022 and ACE2 epitopes, CR3022 is able to neutralize the virus through an

allosteric effect. We found that the Ser373Pro change, which is located within the

CR3022 epitope, has reduced affinity to CR3022 (**Figure 8F, G**). The F338L and

R408T mutations, although not found directly within the interacting epitope, also

display reduced binding to CR3022. Other investigators (86) using *in vitro*

antibody selection identified a change at amino acid site S151 in the N-terminal

domain, and we found mutations S151N and S151I in our patient samples. We

also note that two variant amino acids (Gly446Val and Phe456Leu) we identified

are located in a linear epitope found to be critical for a neutralizing monoclonal

antibody described recently by Li et al. (87).

In the aggregate, these findings suggest that mutations emerging within

the spike protein at positions within and proximal to known neutralization

epitopes may result in escape from antibodies and other therapeutics currently

under development. Importantly, our study did not reveal that these mutant

strains had disproportionately increased over time. The findings may also bear

on the occurrence of multiple amino acid substitutions at the same amino acid

site that we identified in this study, commonly a signal of selection. In the

aggregate, the data support a multifaceted approach to serological monitoring

and biologics development, including the use of monoclonal antibody cocktails

(46, 47, 88).

CONCLUDING STATEMENT

Our work represents analysis of the largest sample to date of SARS-CoV-2

genome sequences from patients in one metropolitan region in the United States.

The investigation was facilitated by the fact that we had rapidly assessed a

SARS-CoV-2 molecular diagnostic test in January 2020, more than a month

before the first COVID-19 patient was diagnosed in Houston. In addition, our

large healthcare system has seven hospitals and many facilities (e.g., outpatient

care centers, emergency departments) located in geographically diverse areas of

the city. We also provide reference laboratory services for other healthcare

entities in the Houston area. Together, our facilities serve patients of diverse

ethnicities and socioeconomic status. Thus, the data presented here likely reflect

a broad overview of virus diversity causing COVID-19 infections throughout

metropolitan Houston. We previously exploited these features to study influenza

and *Klebsiella pneumoniae* dissemination in metropolitan Houston (89, 90). We

acknowledge that every "twig" of the SARS-CoV-2 evolutionary tree in Houston is

not represented in these data. The samples studied are not comprehensive for

the entire metropolitan region. For example, it is possible that our strain samples

are not fully representative of individuals who are indigent, homeless, or of very

low socioeconomic groups. In addition, although the strain sample size is

relatively large compared to other studies, the sample represents only about 10%

of all COVID-19 cases in metropolitan Houston documented in the study period.

In addition, some patient samples contain relatively small amounts of virus

nucleic acid and do not yield adequate sequence data for high-quality genome

analysis. Thus, our data likely underestimate the extent of genome diversity

present among SARS-CoV-2 causing COVID-19 and will not identify all amino

acid replacements in the virus in this geographic region. It will be important to

sequence and analyze the genomes of additional SARS-CoV-2 strains causing

COVID-19 cases in the ongoing second massive disease wave in metropolitan

Houston, and these studies are underway. Data of this type will be especially

important to have if a third and subsequent waves were to occur in metropolitan

Houston, as it could provide insight into molecular and epidemiologic events

contributing to them.

The genomes reported here are an important data resource that will

underpin our ongoing study of SARS-CoV-2 molecular evolution, dissemination,

and medical features of COVID-19 in Houston. As of August 19, 2020, there

were 135,866 reported cases of COVID-19 in metropolitan Houston, and the

number of cases is increasing daily. Although the full array of factors contributing

to the massive second wave in Houston is not known, it is possible that the

potential for increased transmissibility of SARS-CoV-2 with the Gly614 may have

played a role, as well as changes in behavior associated with the Memorial Day

and July 4[th] holidays, and relaxation of some of the social constraints imposed

during the first wave. The availability of extensive virus genome data dating from

the earliest reported cases of COVID-19 in metropolitan Houston, coupled with

the database we have now constructed, may provide critical insights into the

origin of new infection spikes and waves occurring as public health constraints

are further relaxed, schools and colleges re-open, holidays occur, commercial air

travel increases, and individuals change their behavior because of COVID-19

"fatigue." The genome data will also be useful in assessing ongoing molecular

evolution in spike and other proteins as baseline herd immunity is generated,

either by natural exposure to SARS-CoV-2 or by vaccination. The signal of

potential selection contributing to some spike protein diversity and identification

of naturally occurring mutant RBD variants with altered serologic recognition

warrant close attention and expanded study.

MATERIALS AND METHODS

Patient specimens. All specimens were obtained from individuals who

were registered patients at Houston Methodist hospitals, associated facilities

(e.g., urgent care centers), or institutions in the greater Houston metropolitan

region that use our laboratory services. Virtually all individuals met the criteria

specified by the Centers for Disease Control and Prevention to be classified as a

person under investigation.

SARS-CoV-2 molecular diagnostic testing. Specimens obtained from

symptomatic patients with a high degree of suspicion for COVID-19 disease were

tested in the Molecular Diagnostics Laboratory at Houston Methodist Hospital

using an assay granted Emergency Use Authorization (EUA) from the FDA

(https://www.fda.gov/medical-devices/emergency-situations-medical-

devices/faqs-diagnostic-testing-sars-cov-2#offeringtests). Multiple testing

platforms were used, including an assay that follows the protocol published by

the WHO (https://www.who.int/docs/default-source/coronaviruse/protocol-v2-

1.pdf) using the EZ1 virus extraction kit and EZ1 Advanced XL instrument or

QIASymphony DSP Virus kit and QIASymphony instrument for nucleic acid

extraction and ABI 7500 Fast Dx instrument with 7500 SDS software for reverse

transcription RT-PCR, the COVID-19 test using BioFire Film Array 2.0

instruments, the Xpert Xpress SARS-CoV-2 test using Cepheid GeneXpert

Infinity or Cepheid GeneXpert Xpress IV instruments, the SARS-CoV-2 Assay

using the Hologic Panther instrument, and the Aptima SARS-CoV-2 Assay using

the Hologic Panther Fusion system. All assays were performed according to the

manufacturer's instructions. Testing was performed on material obtained from

nasopharyngeal or oropharyngeal swabs immersed in universal transport media

(UTM), bronchoalveolar lavage fluid, or sputum treated with dithiothreitol (DTT).

To standardize specimen collection, an instructional video was created for

Houston Methodist healthcare workers

(https://vimeo.com/396996468/2228335d56).

Epidemiologic curve. The number of confirmed COVID-19 positive cases

was obtained from USAFacts.org (https://usafacts.org/visualizations/coronavirus-

covid-19-spread-map/) for Austin, Brazoria, Chambers, Fort Bend, Galveston,

Harris, Liberty, Montgomery, and Waller counties. Positive cases for Houston

Methodist Hospital patients were obtained from our Laboratory Information

System and plotted using the documented collection time.

SARS-CoV-2 genome sequencing. Libraries for whole virus genome

sequencing were prepared according to version 1 or version 3 of the ARTIC

nCoV-2019 sequencing protocol (https://artic.network/ncov-2019). Long reads

were generated with the LSK-109 sequencing kit, 24 native barcodes (NBD104

and NBD114 kits), and a GridION instrument (Oxford Nanopore). Short reads

were generated with the NexteraXT kit and NextSeq 550 instrument (Illumina).

SARS-CoV-2 genome sequence analysis. Consensus virus genome

sequences from the Houston area isolates were generated using the ARTIC

nCoV-2019 bioinformatics pipeline. Publicly available genomes and metadata

were acquired through GISAID on August 19, 2020. GISAID sequences

containing greater than 1% N characters, and Houston sequences with greater

than 5% N characters were removed from consideration. Identical GISAID

sequences originating from the same geographic location with the same

collection date were also removed from consideration to reduce redundancy.

Nucleotide sequence alignments for the combined Houston and GISAID strains

were generated using MAFFT version 7.130b with default parameters (91).

Sequences were manually curated in JalView (92) to trim the ends and to

remove sequences containing spurious inserts. Phylogenetic trees were

generated using FastTree with the generalized time-reversible model for

nucleotide sequences (93). CLC Genomics Workbench (QIAGEN) was used to

generate the phylogenetic tree figures.

Geospatial mapping. The home address zip code for all SARS-CoV-2

positive patients was used to generate the geospatial maps. To examine

geographic relatedness among genetically similar isolates, geospatial maps were

filtered to isolates containing specific amino acid changes.

Time series. Geospatial data were filtered into wave 1 (3/5/2020-

5/11/2020) and wave 2 (5/12/2020-7/7/2020) time intervals to illustrate the

spread of confirmed SARS-CoV-2 positive patients identified over time.

Machine learning. Virus genome alignments and patient metadata were

used to build models to predict patient metadata and outcomes using both

classification models and regression. Metadata considered for prediction in the

classification models included age, ABO and Rh blood type, ethnic group,

ethnicity, sex, ICU admission, IMU admission, supplemental oxygen use, and

ventilator use. Metadata considered for prediction in regression analysis included

ICU length of stay, IMU length of stay, total length of stay, supplemental oxygen

use, and ventilator use. Because sex, blood type, Rh factor, age, age decade,

ethnicity, and ethnic group are features in the patient features and combined

feature sets, models were not trained for these labels using patient and

combined feature sets. Additionally, age, length of stay, IMU length of stay, ICU

length of stay, mechanical ventilation days, and supplemental oxygen days were

treated as regression problems and XGBoost regressors were built while the rest

were treated as classification problems and XGBoost classifiers were built.

Three types of features were considered for training the XGBoost

classifiers: alignment features, patient features, and the combination of alignment

and patient features. Alignment features were generated from the consensus

genome alignment such that columns containing ambiguous nucleotide bases

were removed to ensure the models did not learn patterns from areas of low

coverage. These alignments were then one-hot encoded to form the alignment

features. Patient metadata values were one-hot encoded with the exception of

age, which remained as a raw integer value, to create the patient features.

These metadata values consisted of age, ABO, Rh blood type, ethnic group,

ethnicity, and sex. All three types of feature sets were used to train models that

predict ICU length of stay, IMU length of stay, overall length of stay, days of

supplemental oxygen therapy, and days of ventilator usage while only alignment

features were used to train models that predict age, ABO, Rh blood type, ethnic

group, ethnicity, and sex.

A ten-fold cross validation was used to train XGBoost models (94) as

described previously (95, 96). Depths of 4, 8, 16, 32, and 64 were used to tune

the models, but accuracies plateaued after a depth of 16. SciKit-Learn's (97)

classification report and r2 score were then used to access overall accuracy of

the classification and regression models, respectively.

Patient metadata correlations. We encoded values into multiple columns

for each metadata field. For example, the ABO column was divided into four

columns for A, B, AB, and O blood type. Those columns were encoded with a 1

for the patients' ABO type, with all other columns encoded with 0. This was

repeated for all non-outcome metadata fields. Age, however, was not re-

encoded, as the raw integer values were used. Each column was then correlated

to the various outcome values for each patient (deceased, ICU length, IMU

length, length of stay, supplemental oxygen length, and ventilator length) to

obtain a Pearson coefficient correlation value for each metadata label and

outcome.

Analysis of the *nsp12* polymerase and S protein genes. The *nsp12*

virus polymerase and S protein genes were analyzed by plotting SNP density in

the consensus alignment using Python (Python v3.4.3, Biopython Package

v1.72). The frequency of SNPs in the Houston isolates was assessed, along with

amino acid changes for nonsynonymous SNPs.

Cycle threshold (Ct) comparison of SARS-CoV-2 strains with either

Asp614 or Gly614 amino acid replacements in the spike protein. The cycle

threshold (Ct) for every sequenced strain that was detected from a patient

specimen using the SARS-CoV-2 Assay on the Hologic Panther instrument was

retrieved from the Houston Methodist Hospital Laboratory Information System.

Statistical significance between the mean Ct value for strains with an aspartate

(*n*=102) or glycine (*n*=812) amino acid at position 614 of the spike protein was

determined with the Mann-Whitney test (GraphPad PRISM 8).

Creation and characterization of spike protein RBD variants. Spike

RBD variants were cloned into the spike-6P (HexaPro; F817P, A892P, A899P,

A942P, K986P, V987P) base construct that also includes the D614G substitution

(pIF638). Briefly, a segment of the gene encoding the RBD was excised with

EcoRI and NheI, mutagenized by PCR, and assembled with a HiFi DNA

Assembly Cloning Kit (NEB).

FreeStyle 293-F cells (Thermo Fisher Scientific) were cultured and

maintained in a humidified atmosphere of 37°C and 8% CO_2 while shaking at

110-125rpm. Cells were transfected with plasmids encoding spike protein

variants using polyethylenimine. Three hours post-transfection, 5µM kifunensine

was added to each culture. Cells were harvested four days after transfection and

the protein containing supernatant was separated from the cells by two

centrifugation steps: 10 min at 500rcf and 20 min at 10,000rcf. Supernatants

were kept at 4°C throughout. Clarified supernatant was loaded on a Poly-Prep

chromatography column (Bio-Rad) containing Strep-Tactin Superflow resin (IBA),

washed with five column volumes (CV) of wash buffer (100mM Tris-HCl pH 8.0,

150mM NaCl; 1mM EDTA), and eluted with four CV of elution buffer (100mM

Tris-HCl pH 8.0, 150mM NaCl, 1mM EDTA, 2.5mM d-Desthiobiotin). The eluate

was spin-concentrated (Amicon Ultra-15) to 600µL and further purified via size-

exclusion chromatography (SEC) using a Superose 6 Increase 10/300 column

(G.E.) in SEC buffer (2mM Tris pH 8.0, 200mM NaCl and 0.02% NaN_3). Proteins

were concentrated to 300µL and stored in SEC buffer.

The RBD spike mutants chosen for analysis were all RBD amino acid

mutants identified by our genome sequencing study as of June 15, 2020. We

note that the exact boundaries of the RBD domain varies depending on the paper

used as reference. We used the boundaries demarcated in Figure 1A of Cai et al.

Science paper 21 July) (98) that have K528R located at the RBD-CTD1 interface.

Differential scanning fluorimetry. Recombinant spike proteins were

diluted to a final concentration of 0.05mg/mL with 5X SYPRO orange (Sigma) in

a 96-well qPCR plate. Continuous fluorescence measurements (λex=465nm,

λem=580nm) were collected with a Roche LightCycler 480 II. The temperature

was increased from 22°C to 95°C at a rate of 4.4°C/min. We report the first

melting transition.

Enzyme-linked immunosorbent assays. ELISAs were performed to

characterize binding of S6P, S6P D614G, and S6P D614G-RBD variants to

human ACE2 and the RBD-binding monoclonal antibody CR3022. The ACE2-

hFc chimera was obtained from GenScript (Z03484), and the CR3022 antibody

was purchased from Abcam (Ab273073). Corning 96-well high-binding plates

(CLS9018BC) were coated with spike variants at 2µg/mL overnight at 4°C. After

washing four times with phosphate buffered saline + 0.1% Tween20 (PBST;

300µL/well), plates were blocked with PBS+2% milk (PBSM) for 2 h at room

temperature and again washed four times with PBST. These were serially diluted

in PBSM 1:3 seven times in triplicate. After 1 h incubation at room temperature,

plates were washed four times in PBST, labeled with 50μL mouse anti-human

IgG1 Fc-HRP (SouthernBlots, 9054-05) for 45 min in PBSM, and washed again

in PBST before addition of 50μL 1-step Ultra TMB-ELISA substrate (Thermo

Scientific, 34028). Reactions were developed for 15 min and stopped by addition

of 50μL 4M H_2SO_4. Absorbance intensity (450nm) was normalized within a plate

and EC_{50} values were calculated through 4-parameter logistic curve (4PL)

analysis using GraphPad PRISM 8.4.3.

ACKNOWLEDGMENTS

We thank Dr. Steven Hinrichs and colleagues at the Nebraska Public Health

Laboratory, and Dr. David Persse and colleagues at the Houston Health

Department for providing samples used to validate our initial SARS-CoV-2

molecular assay. We thank Drs. Jessica Thomas and Zejuan Li, Erika Walker,

Concepcion C. Cantu, the very talented and dedicated molecular technologists,

and the many labor pool volunteers in the Molecular Diagnostics Laboratory for

their dedication to patient care. We also thank Brandi Robinson, Harrold Cano,

Cory Romero, Brooke Burns, and Hayder Mahmood for technical assistance. We

are indebted to Drs. Marc Boom and Dirk Sostman for their support, and to many

very generous Houston philanthropists for their tremendous support of this

ongoing project, including but not limited to anonymous, Ann and John Bookout

III, Carolyn and John Bookout, Ting Tsung and Wei Fong Chao Foundation, Ann

and Leslie Doggett, Freeport LNG, the Hearst Foundations, Jerold B. Katz

Foundation, C. James and Carole Walter Looke, Diane and David Modesett, the

Sherman Foundation, and Paula and Joseph C. "Rusty" Walter III. We gratefully

acknowledge the originating and submitting laboratories of the SARS-CoV-2

genome sequences from GISAID's EpiFluTM Database used in some of the work

presented here. We also thank many colleagues for critical reading of the

manuscript and suggesting improvements, and Sasha Pejerrey, Adrienne

Winston, Heather McConnell, and Kathryn Stockbauer for editorial contributions.

We are especially indebted to Drs. Nancy Jenkins and Neal Copeland for their

scholarly suggestions to improve an early version of the manuscript.

Author contributions: J.M.M. conceptualized and designed the project; S.W.L,

R.J.O., P.A.C., D.W.B., J.J.D., M.S., M.N., M.O.S., C.C.C., P.Y., L.P., S.S., H.-C.

K., H.H., G.E., H.A.T.N., J.H.L., M.K., J.G., D.B., J.G., J.S.M., C.-W.C., K.J., and

I.F. performed research. All authors contributed to writing the manuscript.

Data and material availability: The spike-6P ("HexaPro") plasmid is available from

Addgene (ID: 154754) or from I.J.F. under a material transfer agreement with

The University of Texas at Austin. Additional plasmids are available upon request

from I.J.F.

This study was supported by the Fondren Foundation, Houston Methodist

Hospital and Research Institute (to J.M.M.), NIH grant AI127521 (to J.S.M.), NIH

grants GM120554 and GM124141 to I.J.F., the Welch Foundation (F-1808 to

I.J.F.), and the National Science Foundation (1453358 to I.J.F.). I.J.F. is a CPRIT

Scholar in Cancer Research. J.J.D., M.S., and M.N. are supported by the NIAID

Bacterial and Viral Bioinformatics resource center award (contract number

75N93019C00076). J.J.D. and M.N. are also funded by the United States

Defense Advanced Research Projects Agency Award iSENTRY Friend or Foe

program award (contract number HR0011937807).

REFERENCES

1.   2020. World Health Organization Coronavirus Disease 2019 (COVID-19) Situation Report.
890   https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200420-sitrep-91-
891   covid-19.pdf?sfvrsn=fcf0670b_4. Accessed April 21.
892   2.   Gorbalenya AE, Baker SC, Baric RS, de Groot RJ, Drosten C, Gulyaeva AA, Haagmans BL,
893   Lauber C, Leontovich AM, Neuman BW, Penzar D, Perlman S, Poon LLM, Samborskiy DV,
894   Sidorov IA, Sola I, Ziebuhr J, Coronaviridae Study Group of the International Committee on
895   Taxonomy of V. 2020. The species Severe acute respiratory syndrome-related coronavirus:
896   classifying 2019-nCoV and naming it SARS-CoV-2. Nature Microbiology 5:536-544.
897   3.   Wang C, Horby PW, Hayden FG, Gao GF. 2020. A novel coronavirus outbreak of global health
898   concern. Lancet 395:470-473.
899   4.   Perlman S. 2020. Another Decade, Another Coronavirus. New England Journal of Medicine
900   382:760-762.
901   5.   Allel K, Tapia-Muñoz T, Morris W. 2020. Country-level factors associated with the early spread
902   of COVID-19 cases at 5, 10 and 15 days since the onset. Glob Public Health
903   doi:10.1080/17441692.2020.1814835:1-14.
904   6.   Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X, Cheng Z, Yu T, Xia
905   J, Wei Y, Wu W, Xie X, Yin W, Li H, Liu M, Xiao Y, Gao H, Guo L, Xie J, Wang G, Jiang R,
906   Gao Z, Jin Q, Wang J, Cao B. 2020. Clinical features of patients infected with 2019 novel
907   coronavirus in Wuhan, China. Lancet 395:497-506.
908   7.   Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, Zhao X, Huang B, Shi W, Lu R, Niu P, Zhan F,
909   Ma X, Wang D, Xu W, Wu G, Gao GF, Tan W. 2020. A Novel Coronavirus from Patients with
910   Pneumonia in China, 2019. New England Journal of Medicine 382:727-733.
911   8.   Chan JF, Yuan S, Kok KH, To KK, Chu H, Yang J, Xing F, Liu J, Yip CC, Poon RW, Tsoi HW,
912   Lo SK, Chan KH, Poon VK, Chan WM, Ip JD, Cai JP, Cheng VC, Chen H, Hui CK, Yuen KY.
913   2020. A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating
914   person-to-person transmission: a study of a family cluster. Lancet 395:514-523.
915   9.   Wu F, Zhao S, Yu B, Chen YM, Wang W, Song ZG, Hu Y, Tao ZW, Tian JH, Pei YY, Yuan ML,
916   Zhang YL, Dai FH, Liu Y, Wang QM, Zheng JJ, Xu L, Holmes EC, Zhang YZ. 2020. A new
917   coronavirus associated with human respiratory disease in China. Nature 579:265-269.
918   10.   Bedford T, Greninger AL, Roychoudhury P, Starita LM, Famulare M, Huang M-L, Nalla A,
919   Pepper G, Reinhardt A, Xie H, Shrestha L, Nguyen TN, Adler A, Brandstetter E, Cho S, Giroux
920   D, Han PD, Fay K, Frazar CD, Ilcisin M, Lacombe K, Lee J, Kiavand A, Richardson M, Sibley
921   TR, Truong M, Wolf CR, Nickerson DA, Rieder MJ, Englund JA, Hadfield J, Hodcroft EB,
922   Huddleston J, Moncla LH, Müller NF, Neher RA, Deng X, Gu W, Federman S, Chiu C, Duchin J,
923   Gautom R, Melly G, Hiatt B, Dykema P, Lindquist S, Queen K, Tao Y, Uehara A, Tong S, et al.
924   2020. Cryptic transmission of SARS-CoV-2 in Washington State. medRxiv
925   doi:10.1101/2020.04.02.20051417:2020.04.02.20051417.
926   11.   Gonzalez-Reiche AS, Hernandez MM, Sullivan MJ, Ciferri B, Alshammary H, Obla A, Fabre S,
927   Kleiner G, Polanco J, Khan Z, Alburquerque B, van de Guchte A, Dutta J, Francoeur N, Melo BS,
928   Oussenko I, Deikus G, Soto J, Sridhar SH, Wang Y-C, Twyman K, Kasarskis A, Altman DR,
929   Smith M, Sebra R, Aberg J, Krammer F, García-Sastre A, Luksza M, Patel G, Paniz-Mondolfi A,
930   Gitman M, Sordillo EM, Simon V, van Bakel H. 2020. Introductions and early spread of SARS-
931   CoV-2 in the New York City area. Science 369:297-301.
932   12.   Health N. 2020. COVID-19 Data. https://www1.nyc.gov/site/doh/covid/covid-19-data.page.
933   Accessed August 19.
934   13.   County K. 2020. Daily COVID-19 outbreak summary.
935   https://www.kingcounty.gov/depts/health/covid-19/data/daily-summary.aspx. Accessed August
936   18.
937   14.   Cline M, Emerson M, bratter j, howell j, Jeanty P. 2012. Houston Region Grows More
938   Racially/Ethnically Diverse,With Small Declines in Segregation.A Joint Report Analyzing Census
939   Data from 1990, 2000, and 2010.
940   15.   Emerson M, Bratter J, Howell J, Jeanty P, Cline M. 2012. Houston Region Grows More
941   Racially/Ethnically Diverse, With Small Declines in Segregation. A Joint Report Analyzing

Census Data from 1990, 2000, and 2010. Kinder Institute for Urban Research & the Hobby
Center for the Study of Texas,

16.   Services THaH. 2020. Texas Health and Human Services. https://hhs.texas.gov/. Accessed
August 18.

17.   Vahidy FS, Drews AL, Masud FN, Schwartz RL, Askary BB, Boom ML, Phillips RA. 2020.
Characteristics and Outcomes of COVID-19 Patients During Initial Peak and Resurgence in the
Houston Metropolitan Area. Jama doi:10.1001/jama.2020.15301.

18.   Diehl WE, Lin AE, Grubaugh ND, Carvalho LM, Kim K, Kyawe PP, McCauley SM, Donnard E,
Kucukural A, McDonel P, Schaffner SF, Garber M, Rambaut A, Andersen KG, Sabeti PC, Luban
J. 2016. Ebola Virus Glycoprotein with Increased Infectivity Dominated the 2013-2016 Epidemic.
Cell 167:1088-1098.e6.

19.   Urbanowicz RA, McClure CP, Sakuntabhai A, Sall AA, Kobinger G, Müller MA, Holmes EC,
Rey FA, Simon-Loriere E, Ball JK. 2016. Human Adaptation of Ebola Virus during the West
African Outbreak. Cell 167:1079-1087.e5.

20.   Dietzel E, Schudt G, Krähling V, Matrosovich M, Becker S. 2017. Functional Characterization of
Adaptive Mutations during the West African Ebola Virus Outbreak. J Virol 91.

21.   Kachroo P, Eraso JM, Beres SB, Olsen RJ, Zhu L, Nasser W, Bernard PE, Cantu CC, Saavedra
MO, Arredondo MJ, Strope B, Do H, Kumaraswami M, Vuopio J, Grondahl-Yli-Hannuksela K,
Kristinsson KG, Gottfredsson M, Pesonen M, Pensar J, Davenport ER, Clark AG, Corander J,
Caugant DA, Gaini S, Magnussen MD, Kubiak SL, Nguyen HAT, Long SW, Porter AR, DeLeo
FR, Musser JM. 2019. Integrated analysis of population genomics, transcriptomics and virulence
provides novel insights into Streptococcus pyogenes pathogenesis. Nat Genet 51:548-559.

22.   Gao Y, Yan L, Huang Y, Liu F, Zhao Y, Cao L, Wang T, Sun Q, Ming Z, Zhang L, Ge J, Zheng
L, Zhang Y, Wang H, Zhu Y, Zhu C, Hu T, Hua T, Zhang B, Yang X, Li J, Yang H, Liu Z, Xu W,
Guddat LW, Wang Q, Lou Z, Rao Z. 2020. Structure of the RNA-dependent RNA polymerase
from COVID-19 virus. Science doi:10.1126/science.abb7498:eabb7498.

23.   Yin W, Mao C, Luan X, Shen D-D, Shen Q, Su H, Wang X, Zhou F, Zhao W, Gao M, Chang S,
Xie Y-C, Tian G, Jiang H-W, Tao S-C, Shen J, Jiang Y, Jiang H, Xu Y, Zhang S, Zhang Y, Xu
HE. 2020. Structural basis for inhibition of the RNA-dependent RNA polymerase from SARS-
CoV-2 by remdesivir. Science 368:1499-1504.

24.   Shannon A, Le NT, Selisko B, Eydoux C, Alvarez K, Guillemot JC, Decroly E, Peersen O, Ferron
F, Canard B. 2020. Remdesivir and SARS-CoV-2: Structural requirements at both nsp12 RdRp
and nsp14 Exonuclease active-sites. Antiviral Res 178:104793.

25.   Gordon CJ, Tchesnokov EP, Woolner E, Perry JK, Feng JY, Porter DP, Gotte M. 2020.
Remdesivir is a direct-acting antiviral that inhibits RNA-dependent RNA polymerase from severe
acute respiratory syndrome coronavirus 2 with high potency. J Biol Chem
doi:10.1074/jbc.RA120.013679.

26.   Agostini ML, Andres EL, Sims AC, Graham RL, Sheahan TP, Lu X, Smith EC, Case JB, Feng
JY, Jordan R, Ray AS, Cihlar T, Siegel D, Mackman RL, Clarke MO, Baric RS, Denison MR.
2018. Coronavirus Susceptibility to the Antiviral Remdesivir (GS-5734) Is Mediated by the Viral
Polymerase and the Proofreading Exoribonuclease. mBio 9.

27.   de Wit E, Feldmann F, Cronin J, Jordan R, Okumura A, Thomas T, Scott D, Cihlar T, Feldmann
H. 2020. Prophylactic and therapeutic remdesivir (GS-5734) treatment in the rhesus macaque
model of MERS-CoV infection. Proc Natl Acad Sci U S A 117:6771-6776.

28.   Williamson BN, Feldmann F, Schwarz B, Meade-White K, Porter DP, Schulz J, van Doremalen
N, Leighton I, Yinda CK, Pérez-Pérez L, Okumura A, Lovaglio J, Hanley PW, Saturday G, Bosio
CM, Anzick S, Barbian K, Cihlar T, Martens C, Scott DP, Munster VJ, de Wit E. 2020. Clinical
benefit of remdesivir in rhesus macaques infected with SARS-CoV-2. Nature 585:273-276.

29.   Grein J, Ohmagari N, Shin D, Diaz G, Asperges E, Castagna A, Feldt T, Green G, Green ML,
Lescure FX, Nicastri E, Oda R, Yo K, Quiros-Roldan E, Studemeister A, Redinski J, Ahmed S,
Bernett J, Chelliah D, Chen D, Chihara S, Cohen SH, Cunningham J, D'Arminio Monforte A,
Ismail S, Kato H, Lapadula G, L'Her E, Maeno T, Majumder S, Massari M, Mora-Rillo M, Mutoh
Y, Nguyen D, Verweij E, Zoufaly A, Osinusi AO, DeZure A, Zhao Y, Zhong L, Chokkalingam A,
Elboudwarej E, Telep L, Timbs L, Henne I, Sellers S, Cao H, Tan SK, Winterbourne L, Desai P,
et al. 2020. Compassionate Use of Remdesivir for Patients with Severe Covid-19. N Engl J Med
doi:10.1056/NEJMoa2007016.

30. Goldman JD, Lye DCB, Hui DS, Marks KM, Bruno R, Montejano R, Spinner CD, Galli M, Ahn
MY, Nahass RG, Chen YS, SenGupta D, Hyland RH, Osinusi AO, Cao H, Blair C, Wei X,
Gaggar A, Brainard DM, Towner WJ, Muñoz J, Mullane KM, Marty FM, Tashima KT, Diaz G,
Subramanian A. 2020. Remdesivir for 5 or 10 Days in Patients with Severe Covid-19. N Engl J
Med doi:10.1056/NEJMoa2015301.
31. Beigel JH, Tomashek KM, Dodd LE, Mehta AK, Zingman BS, Kalil AC, Hohmann E, Chu HY,
Luetkemeyer A, Kline S, Lopez de Castilla D, Finberg RW, Dierberg K, Tapson V, Hsieh L,
Patterson TF, Paredes R, Sweeney DA, Short WR, Touloumi G, Lye DC, Ohmagari N, Oh MD,
Ruiz-Palacios GM, Benfield T, Fätkenheuer G, Kortepeter MG, Atmar RL, Creech CB, Lundgren
J, Babiker AG, Pett S, Neaton JD, Burgess TH, Bonnett T, Green M, Makowski M, Osinusi A,
Nayak S, Lane HC. 2020. Remdesivir for the Treatment of Covid-19 - Preliminary Report. N Engl
J Med doi:10.1056/NEJMoa2007764.
32. Spinner CD, Gottlieb RL, Criner GJ, Arribas López JR, Cattelan AM, Soriano Viladomiu A,
Ogbuagu O, Malhotra P, Mullane KM, Castagna A, Chai LYA, Roestenberg M, Tsang OTY,
Bernasconi E, Le Turnier P, Chang SC, SenGupta D, Hyland RH, Osinusi AO, Cao H, Blair C,
Wang H, Gaggar A, Brainard DM, McPhail MJ, Bhagani S, Ahn MY, Sanyal AJ, Huhn G, Marty
FM. 2020. Effect of Remdesivir vs Standard Care on Clinical Status at 11 Days in Patients With
Moderate COVID-19: A Randomized Clinical Trial. Jama doi:10.1001/jama.2020.16349.
33. Olender SA, Perez KK, Go AS, Balani B, Price-Haywood EG, Shah NS, Wang S, Walunas TL,
Swaminathan S, Slim J, Chin B, De Wit S, Ali SM, Soriano Viladomiu A, Robinson P, Gottlieb
RL, Tsang TYO, Lee IH, Haubrich RH, Chokkalingam AP, Lin L, Zhong L, Bekele BN, Mera-
Giler R, Gallant J, Smith LE, Osinusi AO, Brainard DM, Hu H, Phulpin C, Edgar H, Diaz-Cuervo
H, Bernardino JI. 2020. Remdesivir for Severe COVID-19 versus a Cohort Receiving Standard of
Care. Clin Infect Dis doi:10.1093/cid/ciaa1041.
34. (CNCB) CNCfB. 2020. 2019 Novel Coronavirus Resource (2019nCoVR).
https://bigd.big.ac.cn/ncov/about?lang=en. Accessed August 19.
35. Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh C-L, Abiona O, Graham BS, McLellan JS.
2020. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. Science
367:1260-1263.
36. Walls AC, Park YJ, Tortorici MA, Wall A, McGuire AT, Veesler D. 2020. Structure, Function,
and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. Cell 181:281-292.e6.
37. Wang Q, Zhang Y, Wu L, Niu S, Song C, Zhang Z, Lu G, Qiao C, Hu Y, Yuen KY, Wang Q,
Zhou H, Yan J, Qi J. 2020. Structural and Functional Basis of SARS-CoV-2 Entry by Using
Human ACE2. Cell doi:10.1016/j.cell.2020.03.045.
38. Jackson LA, Anderson EJ, Rouphael NG, Roberts PC, Makhene M, Coler RN, McCullough MP,
Chappell JD, Denison MR, Stevens LJ, Pruijssers AJ, McDermott A, Flach B, Doria-Rose NA,
Corbett KS, Morabito KM, O'Dell S, Schmidt SD, Swanson PA, 2nd, Padilla M, Mascola JR,
Neuzil KM, Bennett H, Sun W, Peters E, Makowski M, Albert J, Cross K, Buchanan W, Pikaart-
Tautges R, Ledgerwood JE, Graham BS, Beigel JH. 2020. An mRNA Vaccine against SARS-
CoV-2 - Preliminary Report. N Engl J Med doi:10.1056/NEJMoa2022483.
39. Folegatti PM, Ewer KJ, Aley PK, Angus B, Becker S, Belij-Rammerstorfer S, Bellamy D, Bibi S,
Bittaye M, Clutterbuck EA, Dold C, Faust SN, Finn A, Flaxman AL, Hallis B, Heath P, Jenkin D,
Lazarus R, Makinson R, Minassian AM, Pollock KM, Ramasamy M, Robinson H, Snape M,
Tarrant R, Voysey M, Green C, Douglas AD, Hill AVS, Lambe T, Gilbert SC, Pollard AJ. 2020.
Safety and immunogenicity of the ChAdOx1 nCoV-19 vaccine against SARS-CoV-2: a
preliminary report of a phase 1/2, single-blind, randomised controlled trial. Lancet 396:467-478.
40. Zhu FC, Guan XH, Li YH, Huang JY, Jiang T, Hou LH, Li JX, Yang BF, Wang L, Wang WJ, Wu
SP, Wang Z, Wu XH, Xu JJ, Zhang Z, Jia SY, Wang BS, Hu Y, Liu JJ, Zhang J, Qian XA, Li Q,
Pan HX, Jiang HD, Deng P, Gou JB, Wang XW, Wang XH, Chen W. 2020. Immunogenicity and
safety of a recombinant adenovirus type-5-vectored COVID-19 vaccine in healthy adults aged 18
years or older: a randomised, double-blind, placebo-controlled, phase 2 trial. Lancet 396:479-488.
41. Brouwer PJM, Caniels TG, van der Straten K, Snitselaar JL, Aldon Y, Bangaru S, Torres JL, Okba
NMA, Claireaux M, Kerster G, Bentlage AEH, van Haaren MM, Guerra D, Burger JA, Schermer
EE, Verheul KD, van der Velde N, van der Kooi A, van Schooten J, van Breemen MJ, Bijl TPL,
Sliepen K, Aartse A, Derking R, Bontjer I, Kootstra NA, Wiersinga WJ, Vidarsson G, Haagmans

BL, Ward AB, de Bree GJ, Sanders RW, van Gils MJ. 2020. Potent neutralizing antibodies from
COVID-19 patients define multiple targets of vulnerability. Science 369:643-650.

42.   Chi X, Yan R, Zhang J, Zhang G, Zhang Y, Hao M, Zhang Z, Fan P, Dong Y, Yang Y, Chen Z,
Guo Y, Zhang J, Li Y, Song X, Chen Y, Xia L, Fu L, Hou L, Xu J, Yu C, Li J, Zhou Q, Chen W.
2020. A neutralizing human antibody binds to the N-terminal domain of the Spike protein of
SARS-CoV-2. Science 369:650-655.

43.   Wec AZ, Wrapp D, Herbert AS, Maurer DP, Haslwanter D, Sakharkar M, Jangra RK, Dieterle
ME, Lilov A, Huang D, Tse LV, Johnson NV, Hsieh C-L, Wang N, Nett JH, Champney E,
Burnina I, Brown M, Lin S, Sinclair M, Johnson C, Pudi S, Bortz R, Wirchnianski AS,
Laudermilch E, Florez C, Fels JM, O'Brien CM, Graham BS, Nemazee D, Burton DR, Baric RS,
Voss JE, Chandran K, Dye JM, McLellan JS, Walker LM. 2020. Broad neutralization of SARS-
related viruses by human monoclonal antibodies. Science 369:731-736.

44.   Zost SJ, Gilchuk P, Case JB, Binshtein E, Chen RE, Nkolola JP, Schäfer A, Reidy JX, Trivette A,
Nargi RS, Sutton RE, Suryadevara N, Martinez DR, Williamson LE, Chen EC, Jones T, Day S,
Myers L, Hassan AO, Kafai NM, Winkler ES, Fox JM, Shrihari S, Mueller BK, Meiler J,
Chandrashekar A, Mercado NB, Steinhardt JJ, Ren K, Loo YM, Kallewaard NL, McCune BT,
Keeler SP, Holtzman MJ, Barouch DH, Gralinski LE, Baric RS, Thackray LB, Diamond MS,
Carnahan RH, Crowe JE, Jr. 2020. Potently neutralizing and protective human antibodies against
SARS-CoV-2. Nature 584:443-449.

45.   Greaney AJ, Starr TN, Gilchuk P, Zost SJ, Binshtein E, Loes AN, Hilton SK, Huddleston J, Eguia
R, Crawford KH, Dingens AS, Nargi RS, Sutton RE, Suryadevara N, Rothlauf PW, Liu Z, Whelan
SP, Carnahan RH, Crowe JE, Bloom JD. 2020. Complete mapping of mutations to the SARS-
CoV-2 spike receptor-binding domain that escape antibody recognition. bioRxiv
doi:10.1101/2020.09.10.292078:2020.09.10.292078.

46.   Baum A, Copin R, Ajithdoss D, Zhou A, Lanza K, Negron N, Ni M, Wei Y, Atwal GS, Oyejide
A, Goez-Gazi Y, Dutton J, Clemmons E, Staples HM, Bartley C, Klaffke B, Alfson K, Gazi M,
Gonzales O, Dick E, Carrion R, Pessaint L, Porto M, Cook A, Brown R, Ali V, Greenhouse J,
Taylor T, Andersen H, Lewis MG, Stahl N, Murphy AJ, Yancopoulos GD, Kyratsous CA. 2020.
REGN-COV2 antibody cocktail prevents and treats SARS-CoV-2 infection in rhesus macaques
and hamsters. bioRxiv doi:10.1101/2020.08.02.233320:2020.08.02.233320.

47.   Baum A, Fulton BO, Wloga E, Copin R, Pascal KE, Russo V, Giordano S, Lanza K, Negron N, Ni
M, Wei Y, Atwal GS, Murphy AJ, Stahl N, Yancopoulos GD, Kyratsous CA. 2020. Antibody
cocktail to SARS-CoV-2 spike protein prevents rapid mutational escape seen with individual
antibodies. Science 369:1014-1018.

48.   Barnes CO, West AP, Jr., Huey-Tubman KE, Hoffmann MAG, Sharaf NG, Hoffman PR, Koranda
N, Gristick HB, Gaebler C, Muecksch F, Lorenzi JCC, Finkin S, Hägglöf T, Hurley A, Millard
KG, Weisblum Y, Schmidt F, Hatziioannou T, Bieniasz PD, Caskey M, Robbiani DF,
Nussenzweig MC, Bjorkman PJ. 2020. Structures of Human Antibodies Bound to SARS-CoV-2
Spike Reveal Common Epitopes and Recurrent Features of Antibodies. Cell 182:828-842.e16.

49.   Alsoussi WB, Turner JS, Case JB, Zhao H, Schmitz AJ, Zhou JQ, Chen RE, Lei T, Rizk AA,
McIntire KM, Winkler ES, Fox JM, Kafai NM, Thackray LB, Hassan AO, Amanat F, Krammer F,
Watson CT, Kleinstein SH, Fremont DH, Diamond MS, Ellebedy AH. 2020. A Potently
Neutralizing Antibody Protects Mice against SARS-CoV-2 Infection. J Immunol 205:915-922.

50.   Salazar E, Kuchipudi SV, Christensen PA, Eagar T, Yi X, Zhao P, Jin Z, Long SW, Olsen RJ,
Chen J, Castillo B, Leveque C, Towers D, Lavinder JJ, Gollihar J, Cardona JA, Ippolito GC,
Nissly RH, Bird I, Greenawalt D, Rossi RM, Gontu A, Srinivasan S, Poojary I, Cattadori IM,
Hudson P, Josleyn NM, Prugar L, Huie KE, Herbert AS, Bernard DW, Dye JM, Kapur V, Musser
JM. 2020. Convalescent plasma anti-SARS-CoV-2 spike protein ectodomain and receptor binding
domain IgG correlate with virus neutralization. The Journal of Clinical Investigation
doi:10.1172/JCI141206.

51.   Salazar E, Christensen PA, Graviss EA, Nguyen DT, Castillo B, Chen J, Lopez BV, Eagar TN, Yi
X, Zhao P, Rogers J, Shehabeldin A, Joseph D, Leveque C, Olsen RJ, Bernard DW, Gollihar J,
Musser JM. 2020. Treatment of COVID-19 Patients with Convalescent Plasma Reveals a Signal
of Significantly Decreased Mortality. Am J Pathol doi:10.1016/j.ajpath.2020.08.001.

52.   Starr TN, Greaney AJ, Hilton SK, Ellis D, Crawford KHD, Dingens AS, Navarro MJ, Bowen JE,
Tortorici MA, Walls AC, King NP, Veesler D, Bloom JD. 2020. Deep Mutational Scanning of

SARS-CoV-2 Receptor Binding Domain Reveals Constraints on Folding and ACE2 Binding. Cell
doi:10.1016/j.cell.2020.08.012.
53.   Steffen TL, Stone ET, Hassert M, Geerling E, Grimberg BT, Espino AM, Pantoja P, Climent C,
Hoft DF, George SL, Sariol CA, Pinto AK, Brien JD. 2020. The receptor binding domain of
SARS-CoV-2 spike is the key target of neutralizing antibody in human polyclonal sera. bioRxiv
doi:10.1101/2020.08.21.261727:2020.08.21.261727.
54.   Corbett KS, Flynn B, Foulds KE, Francica JR, Boyoglu-Barnum S, Werner AP, Flach B,
O'Connell S, Bock KW, Minai M, Nagata BM, Andersen H, Martinez DR, Noe AT, Douek N,
Donaldson MM, Nji NN, Alvarado GS, Edwards DK, Flebbe DR, Lamb E, Doria-Rose NA, Lin
BC, Louder MK, O'Dell S, Schmidt SD, Phung E, Chang LA, Yap C, Todd J-PM, Pessaint L, Van
Ry A, Browne S, Greenhouse J, Putman-Taylor T, Strasbaugh A, Campbell T-A, Cook A, Dodson
A, Steingrebe K, Shi W, Zhang Y, Abiona OM, Wang L, Pegu A, Yang ES, Leung K, Zhou T,
Teng I-T, Widge A, et al. 2020. Evaluation of the mRNA-1273 Vaccine against SARS-CoV-2 in
Nonhuman Primates. New England Journal of Medicine doi:10.1056/NEJMoa2024671.
55.   van Doremalen N, Lambe T, Spencer A, Belij-Rammerstorfer S, Purushotham JN, Port JR,
Avanzato VA, Bushmaker T, Flaxman A, Ulaszewska M, Feldmann F, Allen ER, Sharpe H,
Schulz J, Holbrook M, Okumura A, Meade-White K, Pérez-Pérez L, Edwards NJ, Wright D,
Bissett C, Gilbride C, Williamson BN, Rosenke R, Long D, Ishwarbhai A, Kailath R, Rose L,
Morris S, Powers C, Lovaglio J, Hanley PW, Scott D, Saturday G, de Wit E, Gilbert SC, Munster
VJ. 2020. ChAdOx1 nCoV-19 vaccine prevents SARS-CoV-2 pneumonia in rhesus macaques.
Nature doi:10.1038/s41586-020-2608-y.
56.   Wang C, Li W, Drabek D, Okba NMA, van Haperen R, Osterhaus A, van Kuppeveld FJM,
Haagmans BL, Grosveld F, Bosch BJ. 2020. A human monoclonal antibody blocking SARS-CoV-
2 infection. Nat Commun 11:2251.
57.   Ju B, Zhang Q, Ge J, Wang R, Sun J, Ge X, Yu J, Shan S, Zhou B, Song S, Tang X, Yu J, Lan J,
Yuan J, Wang H, Zhao J, Zhang S, Wang Y, Shi X, Liu L, Zhao J, Wang X, Zhang Z, Zhang L.
2020. Human neutralizing antibodies elicited by SARS-CoV-2 infection. Nature 584:115-119.
58.   Liu L, Wang P, Nair MS, Yu J, Rapp M, Wang Q, Luo Y, Chan JF, Sahi V, Figueroa A, Guo XV,
Cerutti G, Bimela J, Gorman J, Zhou T, Chen Z, Yuen KY, Kwong PD, Sodroski JG, Yin MT,
Sheng Z, Huang Y, Shapiro L, Ho DD. 2020. Potent neutralizing antibodies against multiple
epitopes on SARS-CoV-2 spike. Nature 584:450-456.
59.   Rogers TF, Zhao F, Huang D, Beutler N, Burns A, He W-t, Limbo O, Smith C, Song G, Woehl J,
Yang L, Abbott RK, Callaghan S, Garcia E, Hurtado J, Parren M, Peng L, Ramirez S, Ricketts J,
Ricciardi MJ, Rawlings SA, Wu NC, Yuan M, Smith DM, Nemazee D, Teijaro JR, Voss JE,
Wilson IA, Andrabi R, Briney B, Landais E, Sok D, Jardine JG, Burton DR. 2020. Isolation of
potent SARS-CoV-2 neutralizing antibodies and protection from disease in a small animal model.
Science 369:956-963.
60.   Hassan AO, Case JB, Winkler ES, Thackray LB, Kafai NM, Bailey AL, McCune BT, Fox JM,
Chen RE, Alsoussi WB, Turner JS, Schmitz AJ, Lei T, Shrihari S, Keeler SP, Fremont DH, Greco
S, McCray PB, Jr., Perlman S, Holtzman MJ, Ellebedy AH, Diamond MS. 2020. A SARS-CoV-2
Infection Model in Mice Demonstrates Protection by Neutralizing Antibodies. Cell 182:744-
753.e4.
61.   Chandrashekar A, Liu J, Martinot AJ, McMahan K, Mercado NB, Peter L, Tostanoski LH, Yu J,
Maliga Z, Nekorchuk M, Busman-Sahay K, Terry M, Wrijil LM, Ducat S, Martinez DR, Atyeo C,
Fischinger S, Burke JS, Slein MD, Pessaint L, Van Ry A, Greenhouse J, Taylor T, Blade K, Cook
A, Finneyfrock B, Brown R, Teow E, Velasco J, Zahn R, Wegmann F, Abbink P, Bondzie EA,
Dagotto G, Gebre MS, He X, Jacob-Dolan C, Kordana N, Li Z, Lifton MA, Mahrokhian SH,
Maxfield LF, Nityanandam R, Nkolola JP, Schmidt AG, Miller AD, Baric RS, Alter G, Sorger
PK, Estes JD, et al. 2020. SARS-CoV-2 infection protects against rechallenge in rhesus macaques.
Science 369:812-817.
62.   Mercado NB, Zahn R, Wegmann F, Loos C, Chandrashekar A, Yu J, Liu J, Peter L, McMahan K,
Tostanoski LH, He X, Martinez DR, Rutten L, Bos R, van Manen D, Vellinga J, Custers J,
Langedijk JP, Kwaks T, Bakkers MJG, Zuijdgeest D, Huber SKR, Atyeo C, Fischinger S, Burke
JS, Feldman J, Hauser BM, Caradonna TM, Bondzie EA, Dagotto G, Gebre MS, Hoffman E,
Jacob-Dolan C, Kirilova M, Li Z, Lin Z, Mahrokhian SH, Maxfield LF, Nampanya F,
Nityanandam R, Nkolola JP, Patel S, Ventura JD, Verrington K, Wan H, Pessaint L, Ry AV,

Blade K, Strasbaugh A, Cabus M, et al. 2020. Single-shot Ad26 vaccine protects against SARS-
CoV-2 in rhesus macaques. Nature doi:10.1038/s41586-020-2607-z.
63. Yuan M, Wu NC, Zhu X, Lee CD, So RTY, Lv H, Mok CKP, Wilson IA. 2020. A highly
conserved cryptic epitope in the receptor binding domains of SARS-CoV-2 and SARS-CoV.
Science 368:630-633.
64. Hsieh C-L, Goldsmith JA, Schaub JM, DiVenere AM, Kuo H-C, Javanmardi K, Le KC, Wrapp D,
Lee AG, Liu Y, Chou C-W, Byrne PO, Hjorth CK, Johnson NV, Ludes-Meyers J, Nguyen AW,
Park J, Wang N, Amengor D, Lavinder JJ, Ippolito GC, Maynard JA, Finkelstein IJ, McLellan JS.
2020. Structure-based design of prefusion-stabilized SARS-CoV-2 spikes. Science 369:1501-
1505.
65. Woo H, Park SJ, Choi YK, Park T, Tanveer M, Cao Y, Kern NR, Lee J, Yeom MS, Croll TI, Seok
C, Im W. 2020. Developing a Fully Glycosylated Full-Length SARS-CoV-2 Spike Protein Model
in a Viral Membrane. J Phys Chem B 124:7128-7137.
66. Korber B, Fischer WM, Gnanakaran S, Yoon H, Theiler J, Abfalterer W, Hengartner N, Giorgi
EE, Bhattacharya T, Foley B, Hastie KM, Parker MD, Partridge DG, Evans CM, Freeman TM, de
Silva TI, McDanal C, Perez LG, Tang H, Moon-Walker A, Whelan SP, LaBranche CC, Saphire
EO, Montefiori DC. 2020. Tracking Changes in SARS-CoV-2 Spike: Evidence that D614G
Increases Infectivity of the COVID-19 Virus. Cell 182:812-827.e19.
67. Hu J, He C-L, Gao Q-Z, Zhang G-J, Cao X-X, Long Q-X, Deng H-J, Huang L-Y, Chen J, Wang
K, Tang N, Huang A-L. 2020. D614G mutation of SARS-CoV-2 spike protein enhances viral
infectivity. bioRxiv doi:10.1101/2020.06.20.161323:2020.06.20.161323.
68. Lorenzo-Redondo R, Nam HH, Roberts SC, Simons LM, Jennings LJ, Qi C, Achenbach CJ,
Hauser AR, Ison MG, Hultquist JF, Ozer EA. 2020. A Unique Clade of SARS-CoV-2 Viruses is
Associated with Lower Viral Loads in Patient Upper Airways. medRxiv
doi:10.1101/2020.05.19.20107144:2020.05.19.20107144.
69. Cassia Wagner PR, Chris D. Frazar, Jover Lee, Nicola F. Müller, Louise H. Moncla, James
Hadfield, Emma B. Hodcroft, Benjamin Pelle, Matthew Richardson, Caitlin Behrens, Meei-Li
Huang, Patrick Mathias, Gregory Pepper, Lasata Shrestha, Hong Xie, Amin Addetia, Truong
Nguyen, Victoria M Rachleff, Romesh Gautom, Geoff Melly, Brian Hiatt, Philip Dykema,
Amanda Adler, Elisabeth Brandstetter, Peter D. Han, Kairsten Fay, Misja Ilcisin, Kirsten
Lacombe, Thomas R. Sibley, Melissa Truong, Caitlin R. Wolf, Karen Cowgill, Stephanie Schrag,
Jeff Duchin, Michael Boeckh, Janet A. Englund, Michael Famulare, Barry R. Lutz, Mark J.
Rieder, Matthew Thompson, Richard A. Neher, Geoffrey S. Baird, Lea M. Starita, Helen Y. Chu,
Jay Shendure, Scott Lindquist, Deborah A. Nickerson, Alexander L. Greninger, Keith R. Jerome,
Trevor Bedford. 2020. Comparing viral load and clinical outcomes in Washington State across
D614G substitution in spike protein of SARS-CoV-2. https://github.com/blab/ncov-wa-d614g.
Accessed September 8.
70. Volz EM, Hill V, McCrone JT, Price A, Jorgensen D, O'Toole A, Southgate JA, Johnson R,
Jackson B, Nascimento FF, Rey SM, Nicholls SM, Colquhoun RM, da Silva Filipe A, Shepherd
JG, Pascall DJ, Shah R, Jesudason N, Li K, Jarrett R, Pacchiarini N, Bull M, Geidelberg L,
Siveroni I, Goodfellow IG, Loman NJ, Pybus O, Robertson DL, Thomson EC, Rambaut A,
Connor TR. 2020. Evaluating the effects of SARS-CoV-2 Spike mutation D614G on
transmissibility and pathogenicity. medRxiv
doi:10.1101/2020.07.31.20166082:2020.07.31.20166082.
71. Lv Z, Deng Y-Q, Ye Q, Cao L, Sun C-Y, Fan C, Huang W, Sun S, Sun Y, Zhu L, Chen Q, Wang
N, Nie J, Cui Z, Zhu D, Shaw N, Li X-F, Li Q, Xie L, Wang Y, Rao Z, Qin C-F, Wang X. 2020.
Structural basis for neutralization of SARS-CoV-2 and SARS-CoV by a potent therapeutic
antibody. Science 369:1505-1509.
72. Yurkovetskiy L, Wang X, Pascal KE, Tomkins-Tinch C, Nyalile T, Wang Y, Baum A, Diehl WE,
Dauphin A, Carbone C, Veinotte K, Egri SB, Schaffner SF, Lemieux JE, Munro J, Rafique A,
Barve A, Sabeti PC, Kyratsous CA, Dudkina N, Shen K, Luban J. 2020. Structural and Functional
Analysis of the D614G SARS-CoV-2 Spike Protein Variant. bioRxiv
doi:10.1101/2020.07.04.187757:2020.07.04.187757.
73. Zhang L, Jackson CB, Mou H, Ojha A, Rangarajan ES, Izard T, Farzan M, Choe H. 2020. The
D614G mutation in the SARS-CoV-2 spike protein reduces S1 shedding and increases infectivity.
bioRxiv doi:10.1101/2020.06.12.148726:2020.06.12.148726.

74.   Huo J, Zhao Y, Ren J, Zhou D, Duyvesteyn HME, Ginn HM, Carrique L, Malinauskas T, Ruza
RR, Shah PNM, Tan TK, Rijal P, Coombes N, Bewley KR, Tree JA, Radecke J, Paterson NG,
Supasa P, Mongkolsapaya J, Screaton GR, Carroll M, Townsend A, Fry EE, Owens RJ, Stuart DI.
2020. Neutralization of SARS-CoV-2 by Destruction of the Prefusion Spike. Cell Host Microbe
doi:10.1016/j.chom.2020.06.010.

75.   Long SW, Olsen RJ, Christensen PA, Bernard DW, Davis JR, Shukla M, Nguyen M, Ojeda
Saavedra M, Cantu CC, Yerramilli P, Pruitt L, Subedi S, Hendrickson H, Eskandari G,
Kumaraswami M, McLellan JS, Musser JM. 2020. Molecular Architecture of Early Dissemination
and Evolution of the SARS-CoV-2 Virus in Metropolitan Houston, Texas. bioRxiv
doi:10.1101/2020.05.01.072652:2020.05.01.072652.

76.   Plante JA, Liu Y, Liu J, Xia H, Johnson BA, Lokugamage KG, Zhang X, Muruato AE, Zou J,
Fontes-Garfias CR, Mirchandani D, Scharton D, Bilello JP, Ku Z, An Z, Kalveram B, Freiberg
AN, Menachery VD, Xie X, Plante KS, Weaver SC, Shi P-Y. 2020. Spike mutation D614G alters
SARS-CoV-2 fitness and neutralization susceptibility. bioRxiv
doi:10.1101/2020.09.01.278689:2020.09.01.278689.

77.   Latz CA, DeCarlo C, Boitano L, Png CYM, Patell R, Conrad MF, Eagleton M, Dua A. 2020.
Blood type and outcomes in patients with COVID-19. Ann Hematol 99:2113-2118.

78.   Wu BB, Gu DZ, Yu JN, Yang J, Shen WQ. 2020. Association between ABO blood groups and
COVID-19 infection, severity and demise: A systematic review and meta-analysis. Infect Genet
Evol 84:104485.

79.   Zhao J, Yang Y, Huang H, Li D, Gu D, Lu X, Zhang Z, Liu L, Liu T, Liu Y, He Y, Sun B, Wei M,
Yang G, Wang X, Zhang L, Zhou X, Xing M, Wang PG. 2020. Relationship between the ABO
Blood Group and the COVID-19 Susceptibility. Clin Infect Dis doi:10.1093/cid/ciaa1150.

80.   Zietz M, Tatonetti NP. 2020. Testing the association between blood type and COVID-19 infection,
intubation, and death. medRxiv doi:10.1101/2020.04.08.20058073.

81.   Lemieux J, Siddle KJ, Shaw BM, Loreth C, Schaffner S, Gladden-Young A, Adams G, Fink T,
Tomkins-Tinch CH, Krasilnikova LA, Deruff KC, Rudy M, Bauer MR, Lagerborg KA,
Normandin E, Chapman SB, Reilly SK, Anahtar MN, Lin AE, Carter A, Myhrvold C, Kemball M,
Chaluvadi S, Cusick C, Flowers K, Neumann A, Cerrato F, Farhat M, Slater D, Harris JB, Branda
J, Hooper D, Gaeta JM, Baggett TP, O'Connell J, Gnirke A, Lieberman TD, Philippakis A, Burns
M, Brown C, Luban J, Ryan ET, Turbett SE, LaRocque RC, Hanage WP, Gallagher G, Madoff
LC, Smole S, Pierce VM, Rosenberg ES, et al. 2020. Phylogenetic analysis of SARS-CoV-2 in the
Boston area highlights the role of recurrent importation and superspreading events. medRxiv
doi:10.1101/2020.08.23.20178236:2020.08.23.20178236.

82.   Gudbjartsson DF, Helgason A, Jonsson H, Magnusson OT, Melsted P, Norddahl GL,
Saemundsdottir J, Sigurdsson A, Sulem P, Agustsdottir AB, Eiriksdottir B, Fridriksdottir R,
Gardarsdottir EE, Georgsson G, Gretarsdottir OS, Gudmundsson KR, Gunnarsdottir TR, Gylfason
A, Holm H, Jensson BO, Jonasdottir A, Jonsson F, Josefsdottir KS, Kristjansson T, Magnusdottir
DN, le Roux L, Sigmundsdottir G, Sveinbjornsson G, Sveinsdottir KE, Sveinsdottir M,
Thorarensen EA, Thorbjornsson B, Löve A, Masson G, Jonsdottir I, Möller AD, Gudnason T,
Kristinsson KG, Thorsteinsdottir U, Stefansson K. 2020. Spread of SARS-CoV-2 in the Icelandic
Population. N Engl J Med doi:10.1056/NEJMoa2006100.

83.   Candido DS, Claro IM, de Jesus JG, Souza WM, Moreira FRR, Dellicour S, Mellan TA, du
Plessis L, Pereira RHM, Sales FCS, Manuli ER, Thézé J, Almeida L, Menezes MT, Voloch CM,
Fumagalli MJ, Coletti TM, da Silva CAM, Ramundo MS, Amorim MR, Hoeltgebaum HH, Mishra
S, Gill MS, Carvalho LM, Buss LF, Prete CA, Ashworth J, Nakaya HI, Peixoto PS, Brady OJ,
Nicholls SM, Tanuri A, Rossi ÁD, Braga CKV, Gerber AL, de C. Guimarães AP, Gaburo N,
Alencar CS, Ferreira ACS, Lima CX, Levi JE, Granato C, Ferreira GM, Francisco RS, Granja F,
Garcia MT, Moretti ML, Perroud MW, Castiñeiras TMPP, Lazari CS, et al. 2020. Evolution and
epidemic spread of SARS-CoV-2 in Brazil. Science 369:1255-1260.

84.   Ellinghaus D, Degenhardt F, Bujanda L, Buti M, Albillos A, Invernizzi P, Fernández J, Prati D,
Baselli G, Asselta R, Grimsrud MM, Milani C, Aziz F, Kässens J, May S, Wendorff M,
Wienbrandt L, Uellendahl-Werth F, Zheng T, Yi X, de Pablo R, Chercoles AG, Palom A, Garcia-
Fernandez AE, Rodriguez-Frias F, Zanella A, Bandera A, Protti A, Aghemo A, Lleo A, Biondi A,
Caballero-Garralda A, Gori A, Tanck A, Carreras Nolla A, Latiano A, Fracanzani AL, Peschuck
A, Julià A, Pesenti A, Voza A, Jiménez D, Mateos B, Nafria Jimenez B, Quereda C, Paccapelo C,

Gassner C, Angelini C, Cea C, Solier A, et al. 2020. Genomewide Association Study of Severe
Covid-19 with Respiratory Failure. N Engl J Med doi:10.1056/NEJMoa2020283.

85.   2020. The COVID-19 Host Genetics Initiative, a global initiative to elucidate the role of host
genetic factors in susceptibility and severity of the SARS-CoV-2 virus pandemic. Eur J Hum
Genet 28:715-718.

86.   Weisblum Y, Schmidt F, Zhang F, DaSilva J, Poston D, Lorenzi JCC, Muecksch F, Rutkowska M,
Hoffmann H-H, Michailidis E, Gaebler C, Agudelo M, Cho A, Wang Z, Gazumyan A, Cipolla M,
Luchsinger L, Hillyer CD, Caskey M, Robbiani DF, Rice CM, Nussenzweig MC, Hatziioannou T,
Bieniasz PD. 2020. Escape from neutralizing antibodies by SARS-CoV-2 spike protein variants.
bioRxiv doi:10.1101/2020.07.21.214759:2020.07.21.214759.

87.   Li T, Han X, Wang Y, Gu C, Wang J, Hu C, Li S, Wang K, Luo F, Huang J, Long Y, Song S,
Wang W, Hu J, Wu R, Mu S, Hao Y, Chen Q, Gao F, Shen M, Long S, Gong F, Li L, Wu Y, Xu
W, Cai X, Qu D, Yuan Z, Gao Q, Zhang G, He C, Nai Y, Deng K, Du L, Tang N, Xie Y, Huang
A, Jin A. 2020. A key linear epitope for a potent neutralizing antibody to SARS-CoV-2 S-RBD.
bioRxiv doi:10.1101/2020.09.11.292631:2020.09.11.292631.

88.   Hansen J, Baum A, Pascal KE, Russo V, Giordano S, Wloga E, Fulton BO, Yan Y, Koon K, Patel
K, Chung KM, Hermann A, Ullman E, Cruz J, Rafique A, Huang T, Fairhurst J, Libertiny C,
Malbec M, Lee W-y, Welsh R, Farr G, Pennington S, Deshpande D, Cheng J, Watty A, Bouffard
P, Babb R, Levenkova N, Chen C, Zhang B, Romero Hernandez A, Saotome K, Zhou Y, Franklin
M, Sivapalasingam S, Lye DC, Weston S, Logue J, Haupt R, Frieman M, Chen G, Olson W,
Murphy AJ, Stahl N, Yancopoulos GD, Kyratsous CA. 2020. Studies in humanized mice and
convalescent humans yield a SARS-CoV-2 antibody cocktail. Science 369:1010-1014.

89.   Long SW, Olsen RJ, Eagar TN, Beres SB, Zhao P, Davis JJ, Brettin T, Xia F, Musser JM. 2017.
Population Genomic Analysis of 1,777 Extended-Spectrum Beta-Lactamase-Producing
Klebsiella pneumoniae Isolates, Houston, Texas: Unexpected Abundance of Clonal
Group 307. mBio 8:e00489-17.

90.   Stucker KM, Schobel SA, Olsen RJ, Hodges HL, Lin X, Halpin RA, Fedorova N, Stockwell TB,
Tovchigrechko A, Das SR, Wentworth DE, Musser JM. 2015. Haemagglutinin mutations and
glycosylation changes shaped the 2012/13 influenza A(H3N2) epidemic, Houston, Texas. Euro
Surveill 20.

91.   Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol Biol Evol 30:772-80.

92.   Waterhouse AM, Procter JB, Martin DM, Clamp M, Barton GJ. 2009. Jalview Version 2--a
multiple sequence alignment editor and analysis workbench. Bioinformatics 25:1189-91.

93.   Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-likelihood trees for
large alignments. PLoS One 5:e9490.

94.   Chen T, Guestrin C. 2016. XGBoost: A Scalable Tree Boosting System, abstr Proceedings of the
22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining, San
Francisco, California, USA, Association for Computing Machinery,

95.   Nguyen M, Brettin T, Long SW, Musser JM, Olsen RJ, Olson R, Shukla M, Stevens RL, Xia F,
Yoo H. 2018. Developing an in silico minimum inhibitory concentration panel test for *Klebsiella*
pneumoniae. Scientific reports 8:421.

96.   Nguyen M, Long SW, McDermott PF, Olsen RJ, Olson R, Stevens RL, Tyson GH, Zhao S, Davis
JJ. 2019. Using machine learning to predict antimicrobial MICs and associated genomic features
for nontyphoidal Salmonella. Journal of Clinical Microbiology 57:e01260-18.

97.   Pedregosa F, Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M.,
Prettenhofer, P., Weiss, R., Dubourg, V., Vanderplas, J., Passos, A., Cournapeau, D., Brucher, M.,
Perrot, M., and Duchesnay, E. 2011. Scikit-learn: Machine Learning in Python. Journal of
Machine Learning Research 12 (2011) 2825-2830.

98.   Cai Y, Zhang J, Xiao T, Peng H, Sterling SM, Walsh RM, Jr., Rawson S, Rits-Volloch S, Chen B.
2020. Distinct conformational states of SARS-CoV-2 spike protein. Science
doi:10.1126/science.abd4251.

1329

A

B

1330

FIG 1 (A) Confirmed COVID-19 cases in the Greater Houston Metropolitan region.

Cumulative number of COVID-19 patients over time through July 7, 2020. Counties

include Austin, Brazoria, Chambers, Fort Bend, Galveston, Harris, Liberty, Montgomery,

and Waller. The shaded area represents the time period during which virus genomes

characterized in this study were recovered from COVID-19 patients. The red line

represents the number of COVID-19 patients diagnosed in the Houston Methodist

Hospital Molecular Diagnostic Laboratory. (B) Distribution of strains with either the

Asp614 or Gly614 amino acid variant in spike protein among the two waves of COVID-

19 patients diagnosed in the Houston Methodist Hospital Molecular Diagnostic

Laboratory. The large inset shows major clade frequency for the time frame studied.

1341

1342

FIG 2 Sequential time-series heatmaps for all COVID-19 Houston Methodist patients during the study period. Geospatial distribution of COVID-19 patients is based on zip code. Panel A (left) shows geospatial distribution of sequenced SARS-CoV-2 strains in wave 1 and panel B (right) shows wave 2 distribution. The collection dates are shown at the bottom of each panel. The insets refer to numbers of strains in the color spectrum used. Note difference in numbers of strains used in panel A and panel B insets.

1349

1350

1351 **FIG 3** Location of amino acid replacements in RNA-dependent RNA polymerase

1352 (RdRp/Nsp12) among the 5,085 genomes of SARS-CoV-2 sequenced. The various

1353 RdRp domains are color-coded. The numbers refer to amino acid site. Note that several

1354 amino acid sites have multiple variants identified.

1355

1356

FIG 4 Amino acid changes identified in Nsp12 (RdRp) in this study that may influence

interaction with remdesivir. The schematic at the top shows the domain architecture of

Nsp12. (Left) Ribbon representation of the crystal structure of Nsp12-remdesivir

monophosphate-RNA complex (PDB code: 7BV2). The structure in the right panel

shows a magnified view of the boxed area in the left panel. The Nsp12 domains are

colored as in the schematic at the top. The catalytic site in Nsp12 is marked by a black

circle in the right panel. The side chains of amino acids comprising the catalytic site of

RdRp (Ser758, Asp759, and Asp760) are shown as balls and stick and colored yellow.

The nucleotide binding site is boxed in the right panel. The side chains of amino acids

participating in nucleotide binding (Lys544, Arg552, and Arg554) are shown as balls and

sticks and colored light blue. Remdesivir molecule incorporated into the nascent RNA is

shown as balls and sticks and colored light pink. The RNA is shown as a blue cartoon

and bases are shown as sticks. The positions of Cα atoms of amino acids identified in

this study are shown as red and green spheres and labeled. The amino acids that are

shown as red spheres are located above the nucleotide binding site, whereas Cys812

located at the catalytic site is shown as a green sphere. The side chain of active site

residue Ser758 is shown as ball and sticks and colored yellow. The location of Cα atoms

of remdesivir resistance conferring amino acid Val556 is shown as blue sphere and

labeled.

1376

1377

FIG 5 Location of amino acid replacements in spike protein among the 5,085 genomes of SARS-CoV-2 sequenced. The various spike protein domains are color-coded. The numbers refer to amino acid site. Note that many amino acid sites have multiple variants identified.

1382

1383

FIG 6 Location of amino acid substitutions mapped on the SARS-CoV-2 spike protein.

Model of the SARS-CoV-2 spike protein with one protomer shown as ribbons and the

other two protomers shown as a molecular surface. The Cα atom of residues found to be

substituted in one or more virus isolates identified in this study is shown as a sphere on

the ribbon representation. Residues found to be substituted in 1–9 isolates are colored

tan, 10–99 isolates yellow, 100–999 isolates colored red (H49Y and F1052L), and >1000

isolates purple (D614G). The surface of the aminoterminal domain (NTD) that is distal to

the trimeric axis has a high density of substituted residues. RBD, receptor binding

domain.

1393

1394 **FIG 7** Cycle threshold (Ct) for every SARS-CoV-2 patient sample tested using the

1395 Hologic Panther assay. Data are presented as mean +/- standard error of the mean for

1396 strains with an aspartate (D614, n=102 strains, blue) or glycine (G614, n=812 strains,

1397 red) at amino acid 614 of the spike protein. Mann-Whitney test, *P<0.0001.

1398

FIG 8 Biochemical characterization of spike RBD variants. (A) Size-exclusion

chromatography (SEC) traces of the indicated spike-RBD variants. Dashed line indicates

the elution peak of spike-6P. (B) The relative expression of all RBD variants as

determined by the area under the SEC traces. All expression levels are normalized

relative to spike-6P. (C) Thermostability analysis of RBD variants by differential scanning

fluorimetry. Each sample had three replicates and only mean values were plotted. Black

vertical dashed line indicates the first melting temperature of 6P-D614G and orange

vertical dashed line indicates the first melting temperature of the least stable variant

(spike-G446V). (D) First apparent melting temperature of all RBD variants. (E) ELISA-

based binding affinities for ACE2 and (F) the neutralizing antibody CR3022 to the

indicated RBD variants. (G) Summary of EC50s for all measured RBD variants.

1410

1411

1412   **Table 1.** Nonsynonymous SNPs of SARS-CoV-2 *nsp12.*

Genomic Locus	Gene Locus	Amino Acid Change	Domain	Wave 1 (n=1026)	Wave 2 (n=4059)	Total (n=5085)
13446	C3T	A1V	N-terminus		2	2
13448	G5A	D2N	N-terminus	1		1
13487	C44T	A15V	N-terminus		138	138
13501	C58T	P20S	N-terminus		1	1
13514	G71A	G24D	N-terminus		3	3
13517	C74T	T25I	N-terminus		4	4
13520	G77A	S26N	N-terminus		1	1
13523	C80T	T27I	N-terminus		1	1
13526	A83C	D28A	N-terminus		1	1
13564	G121A	V41I	B hairpin		1	1
13568	C125T	A42V	B hairpin	1		1
13571	G128T	G43V	B hairpin	1		1
13576	G133T	A45S	B hairpin		12	12
13617	G174T	K58N	NiRAN		1	1
13618	G175T	D59Y	NiRAN		24	24
13620	C177G	D59E	NiRAN		1	1
13627	G184T	D62Y	NiRAN	1		1
13661	G218A	R73K	NiRAN		1	1
13667	C224T	T75I	NiRAN		2	2
13694	C251T	T84I	NiRAN		1	1
13712	A269G	K90R	NiRAN		1	1
13726	G283A	V95I	NiRAN		1	1
13730	C287T	A96V	NiRAN	2	2	4
13762	G319C	G107R	NiRAN		1	1
13774	C331A	P111T	NiRAN		1	1
13774	C331T	P111S	NiRAN		15	15
13777	C334T	H112Y	NiRAN		1	1
13790	A347G	Q116R	NiRAN		2	2
13835	G392T	R131M	NiRAN		1	1
13858	G415T	D139Y	NiRAN		3	3
13862	C419T	T140I	NiRAN	1	5	6
13868	A425G	K142R	NiRAN	1		1
13897	G454T	D152Y	NiRAN		4	4
13901	A458G	D153G	NiRAN		2	2
13957	C514T	R172C	NiRAN		2	2
13963	T520C	Y174H	NiRAN		1	1
13966	G523A	A175T	NiRAN		1	1
13975	G532T	G178C	NiRAN		4	4

Genomic Locus	Gene Locus	Amino Acid Change	Domain	Wave 1 (n=1026)	Wave 2 (n=4059)	Total (n=5085)
13984	G541A	V181I	NiRAN		1	1
13994	C551T	A184V	NiRAN		8	8
14104	T661C	F221L	NiRAN	2		2
14109	A666G	I222M	NiRAN	1		1
14120	C677T	P226L	NiRAN		2	2
14185	A742G	R248G	NiRAN		1	1
14187	G744T	R248S	NiRAN		1	1
14188	G745A	A249T	NiRAN		1	1
14225	C782A	T261K	Interface		4	4
14230	C787T	P263S	Interface		1	1
14233	T790C	Y264H	Interface		1	1
14241	G798T	K266N	Interface		1	1
14290	G847T	D283Y	Interface		1	1
14335	G892T	V298F	Interface		8	8
14362	C919A	L307M	Interface		2	2
14371	G928C	A310P	Interface	1		1
14396	C953T	T318I	Interface		1	1
14398	G955T	V319L	Interface		1	1
14407	C964T	P322S	Interface		2	2
14408	C965T	P322L	Interface	843	4050	4893
14500	G1057T	V353L	Interface		5	5
14536	C1093T	L365F	Interface		1	1
14557	G1114T	V372L	Fingers		4	4
14584	G1141T	A381S			1	1
14585	C1142T	A381V	Fingers		10	10
14593	G1150A	G384S	Fingers	1		1
14657	C1214T	A405V	Fingers		1	1
14708	C1265T	A422V			1	1
14747	A1304G	E435G	Fingers		2	2
14768	C1325T	A442V	Fingers		21	21
14786	C1343T	A448V	Fingers	3	6	9
14821	C1378T	P460S			1	1
14829	G1386T	M462I	Fingers		59	59
14831	G1388T	C463F	Fingers		3	3
14857	G1414T	V472F			1	1
14870	A1427G	D476G	Fingers		5	5
14874	G1431T	K477N	Fingers	1		1
14912	A1469G	N490S	Fingers	1	1	2
14923	G1480A	V494I	Fingers		2	2

Genomic Locus	Gene Locus	Amino Acid Change	Domain	Wave 1 (n=1026)	Wave 2 (n=4059)	Total (n=5085)
14980	C1537T	L513F	Fingers	1	1	2
14990	A1547G	D516G			1	1
15006	G1563C	E521D	Fingers	2	3	5
15016	G1573T	A525S	Fingers		3	3
15026	C1583T	A528V	Fingers	5	1	6
15037	C1594T	R532C	Fingers		1	1
15100	G1657C	A553P	Fingers		1	1
15101	C1658T	A553V	Fingers		1	1
15124	A1681G	I561V	Fingers		2	2
15202	G1759C	V587L	Palm		7	7
15211	A1768G	T590A			1	1
15226	G1783A	G595S	Palm		1	1
15243	G1800T	M600I	Palm	71	4	75
15251	C1808G	T603S	Palm		1	1
15257	A1814G	Y605C			1	1
15260	G1817A	S606N	Palm		1	1
15327	G1884T	M628I	Fingers	3	1	4
15328	C1885T	L629F	Fingers	1		1
15334	A1891G	I631V	Fingers		1	1
15341	C1898T	A633V	Fingers		1	1
15352	C1909T	L637F	Fingers		1	1
15358	C1915T	R639C	Fingers		1	1
15362	A1919G	K640R	Fingers		1	1
15364	C1921G	H641D	Fingers	1		1
15368	C1925T	T642I	Fingers	1		1
15380	G1937T	S646I	Fingers	1		1
15386	C1943T	S648L	Fingers		2	2
15391	C1948T	R650C	Fingers		1	1
15406	G1963T	A655S	Fingers		3	3
15407	C1964T	A655V	Fingers	1		1
15436	A1993G	M665V	Fingers		2	2
15438	G1995T	M665I	Fingers		24	24
15452	G2009T	G670V	Fingers		28	28
15487	G2044C	G682R	Palm		1	1
15497	C2054A	T685K	Palm		1	1
15572	A2129G	D710G	Palm		1	1
15596	A2153G	Y718S	Palm	2		2
15619	C2176T	L726F	Palm	1		1
15638	G2195A	R732K	Palm	1		1

Genomic Locus	Gene Locus	Amino Acid Change	Domain	Wave 1 (n=1026)	Wave 2 (n=4059)	Total (n=5085)
15640	A2197G	N733D	Palm	1		1
15640	A2197T	N733Y	Palm	1		1
15655	A2212G	T738A	Palm		2	2
15656	C2213T	T738I	Palm		2	2
15658	G2215A	D739N	Palm		2	2
15664	G2221A	V741M	Palm		1	1
15715	T2272C	S758P	Palm		1	1
15760	G2317A	G773S	Palm	1		1
15761	G2318A	G773D	Palm		1	1
15827	A2384G	E795G	Palm	1		1
15848	C2405T	T802I	Palm		1	1
15850	G2407T	D803Y	Palm		1	1
15853	C2410T	L804F	Palm		2	2
15878	G2435T	C812F	Palm		1	1
15886	C2443T	H815Y	Palm		1	1
15906	G2463T	Q821H	Thumb	1	1	2
15908	G2465T	G822V	Thumb		1	1
15979	A2536G	I846V	Thumb	4		4
16045	C2602T	L868F	Thumb		1	1
16084	C2641T	H881Y	Thumb		1	1
16148	A2705G	Y902C	Thumb		1	1
16163	C2720T	T907I	Thumb		45	45
16178	C2735T	S912L	Thumb		2	2
16192	C2749T	P917S	Thumb		80	80

1413
1414

Table 2. Nonsynonymous SNPs in SARS-CoV-2 spike protein.

Genomic Locus	Gene Locus	Amino Acid Change	Domain	Wave 1 (n=1026)	Wave 2 (n=4059)	Total (n=5085)
21575	C13T	L5F	S1	11	25	36
21578	G16T	V6F	S1		1	1
21587	C25T	P9S	S1	2		2
21588	C26T	P9L	S1	1	1	2
21594	T32C	V11A	S1		1	1
21597	C35T	S12F	S1		6	6
21604	G42T	Q14H	S1		1	1
21614	C52T	L18F	S1 - NTD	1	11	12
21618	C56T	T19I	S1 - NTD	1	1	2
21621	C59T	T20I	S1 - NTD		1	1
21624	G62T	R21I	S1 - NTD		6	6
21624	G62A	R21K	S1 - NTD		1	1
21624	G62C	R21T	S1 - NTD		3	3
21627	C65T	T22I	S1 - NTD	2	4	6
21638	C76T	P26S	S1 - NTD		17	17
21641	G79T	A27S	S1 - NTD	1	1	2
21641	G79A	A27T	S1 - NTD	1		1
21642	C80T	A27V	S1 - NTD		1	1
21648	C86T	T29I	S1 - NTD	1	4	5
21707	C145T	H49Y	S1 - NTD		142	142
21713	A151G	T51A	S1 - NTD		1	1
21724	G162T	L54F	S1 - NTD		11	11
21754	G192T	W64C	S1 - NTD		1	1
21767	C205T	H69Y	S1 - NTD	1	7	8
21770	G208A	V70I	S1 - NTD		1	1
21770	G208T	V70F	S1 - NTD	1		1
21774	C212T	S71F	S1 - NTD		1	1
21784	T222A	N74K	S1 - NTD	1		1
21785	G223C	G75R	S1 - NTD		1	1
21793	G231T	K77N	S1 - NTD		1	1
21824	G262A	D88N	S1 - NTD		1	1
21834	A272T	Y91F	S1 - NTD		1	1
21846	C284T	T95I	S1 - NTD	1	10	11
21852	A290G	K97R	S1 - NTD		1	1
21855	C293T	S98F	S1 - NTD	1	2	3
21861	T299C	I100T	S1 - NTD		2	2

Genomic Locus	Gene Locus	Amino Acid Change	Domain	Wave 1 (n=1026)	Wave 2 (n=4059)	Total (n=5085)
21918	T356C	I119T	S1 - NTD	1		1
21930	C368T	A123V	S1 - NTD		1	1
21941	G379T	V127F	S1 - NTD		1	1
21942	T380C	V127A	S1 - NTD		4	4
21974	G412T	D138Y	S1 - NTD	2		2
21985	G423T	L141F	S1 - NTD		1	1
21986	G424A	G142S	S1 - NTD		2	2
21993	A431G	Y144C	S1 - NTD	1		1
21995	T433C	Y145H	S1 - NTD	2		2
21998	C436T	H146Y	S1 - NTD	1	2	3
22014	G452A	S151N	S1 - NTD		1	1
22014	G452T	S151I	S1 - NTD		2	2
22017	G455T	W152L	S1 - NTD	1	1	2
22021	G459T	M153I	S1 - NTD		1	1
22021	G459A	M153I	S1 - NTD		1	1
22022	G460A	E154K	S1 - NTD		1	1
22028	G466C	E156Q	S1 - NTD	2		2
22037	G475A	V159I	S1 - NTD	1		1
22097	C535T	L179F	S1 - NTD		1	1
22104	G542T	G181V	S1 - NTD		1	1
22107	A545G	K182R	S1 - NTD		1	1
22135	A573T	E191D	S1 - NTD		1	1
22139	G577T	V193L	S1 - NTD		1	1
22150	T588G	N196K	S1 - NTD	1		1
22175	T613G	S205A	S1 - NTD		1	1
22205	G643T	D215Y	S1 - NTD		1	1
22206	A644G	D215G	S1 - NTD		2	2
22214	C652G	Q218E	S1 - NTD		1	1
22227	C665T	A222V	S1 - NTD		1	1
22241	G679A	V227I	S1 - NTD		2	2
22242	T680C	V227A	S1 - NTD	1		1
22244	G682C	D228H	S1 - NTD		2	2
22245	A683G	D228G	S1 - NTD	1		1
22246	T684G	D228E	S1 - NTD	2		2
22248	T686G	L229W	S1 - NTD	1		1
22250	C688A	P230T	S1 - NTD	1		1
22253	A691G	I231V	S1 - NTD	1		1

Genomic Locus	Gene Locus	Amino Acid Change	Domain	Wave 1 (n=1026)	Wave 2 (n=4059)	Total (n=5085)
22254	T692C	I231T	S1 - NTD	1		1
22259	A697G	I233V	S1 - NTD	1		1
22260	T698C	I233T	S1 - NTD	1		1
22262	A700G	N234D	S1 - NTD	1		1
22266	T704C	I235T	S1 - NTD	1		1
22281	C719T	T240I	S1 - NTD		5	5
22286	C724T	L242F	S1 - NTD		1	1
22295	C733T	H245Y	S1 - NTD		2	2
22304	T742C	Y248H	S1 - NTD		3	3
22311	C749T	T250I	S1 - NTD	1	4	5
22313	C751T	P251S	S1 - NTD		2	2
22320	A758G	D253G	S1 - NTD		2	2
22320	A758C	D253A	S1 - NTD	1		1
22323	C761T	S254F	S1 - NTD		3	3
22329	C767T	S256L	S1 - NTD	1		1
22335	G773T	W258L	S1 - NTD	1		1
22344	G782T	G261V	S1 - NTD	3		3
22346	G784T	A262S	S1 - NTD		4	4
22350	C788T	A263V	S1 - NTD	1		1
22382	A820G	T274A	S1 - NTD		1	1
22398	A836T	Y279F	S1 - NTD	1		1
22408	T846G	N282K	S1 - NTD		1	1
22425	C863T	A288V	S1 - NTD		1	1
22430	G868T	D290Y	S1 - NTD	1		1
22484	G922T	V308L	S1		3	3
22487	G925C	E309Q	S1		1	1
22532	G970C	E324Q	S1		1	1
22533	A971T	E324V	S1		1	1
22535	T973C	S325P	S1		1	1
22536	C974T	S325F	S1		1	1
22550	C988T	P330S	S1 - RBD		2	2
22574	T1012C	F338L	S1 - RBD	1		1
22608	C1046T	S349F	S1 - RBD		1	1
22616	G1054T	A352S	S1 - RBD		7	7
22661	G1099T	V367F	S1 - RBD		1	1
22673	T1111C	S371P	S1 - RBD		3	3
22679	T1117C	S373P	S1 - RBD		1	1

Genomic Locus	Gene Locus	Amino Acid Change	Domain	Wave 1 (n=1026)	Wave 2 (n=4059)	Total (n=5085)
22712	C1150T	P384S	S1 - RBD		1	1
22716	C1154T	T385I	S1 - RBD	3		3
22785	G1223C	R408T	S1 - RBD		1	1
22793	G1231T	A411S	S1 - RBD		1	1
22818	C1256T	A419V	S1 - RBD	1		1
22895	G1333T	V445F	S1 - RBD		1	1
22899	G1337T	G446V	S1 - RBD	2		2
22928	T1366C	F456L	S1 - RBD	1		1
23001	G1439T	C480F	S1 - RBD		1	1
23012	G1450C	E484Q	S1 - RBD	1		1
23046	A1484C	Y495S	S1 - RBD		1	1
23111	C1549T	L517F	S1 - RBD		1	1
23120	G1558T	A520S	S1 - RBD	1	6	7
23121	C1559T	A520V	S1 - RBD		1	1
23127	C1565T	A522V	S1 - RBD	1	1	2
23145	A1583G	K528R	S1 - RBD		2	2
23149	G1587T	K529N	S1		1	1
23170	C1608A	N536K	S1		1	1
23202	C1640A	T547K	S1		2	2
23202	C1640T	T547I	S1		1	1
23223	A1661T	E554V	S1		2	2
23224	G1662T	E554D	S1	4	31	35
23270	G1708T	A570S	S1		3	3
23277	C1715T	T572I	S1	5	5	10
23282	G1720T	D574Y	S1		1	1
23292	G1730T	R577L	S1	1		1
23311	G1749T	E583D	S1		6	6
23312	A1750G	I584V	S1		1	1
23315	C1753T	L585F	S1	1	7	8
23349	G1787A	S596N	S1		1	1
23373	C1811T	T604I	S1		2	2
23380	C1818A	N606K	S1		2	2
23403	A1841G	D614G	S1	841	4054	4895
23426	G1864T	V622F	S1		2	2
23426	G1864C	V622L	S1		2	2
23435	C1873T	H625Y	S1		1	1
23439	C1877T	A626V	S1		1	1

Genomic Locus	Gene Locus	Amino Acid Change	Domain	Wave 1 (n=1026)	Wave 2 (n=4059)	Total (n=5085)
23444	C1882G	Q628E	S1		7	7
23453	C1891T	P631S	S1	1		1
23457	C1895T	T632I	S1		1	1
23481	C1919T	S640F	S1	1	42	43
23486	G1924T	V642F	S1		1	1
23502	C1940T	A647V	S1		1	1
23536	C1974A	N658K	S1		4	4
23564	G2002T	A668S	S1		1	1
23586	A2024G	Q675R	S1		14	14
23587	G2025C	Q675H	S1		1	1
23587	G2025T	Q675H	S1		4	4
23589	C2027T	T676I	S1	1	2	3
23593	G2031T	Q677H	S1	1	1	2
23595	C2033T	T678I	S1	1		1
23624	G2062T	A688S	S2		4	4
23625	C2063T	A688V	S2		16	16
23655	C2093T	S698L	S2		1	1
23664	C2102T	A701V	S2		21	21
23670	A2108G	N703S	S2		1	1
23679	C2117T	A706V	S2		1	1
23684	T2122C	S708P	S2		1	1
23709	C2147T	T716I	S2		1	1
23718	C2156T	T719I	S2		1	1
23745	C2183T	P728L	S2	1		1
23755	G2193T	M731I	S2	3	1	4
23798	T2236C	S746P	S2		1	1
23802	C2240T	T747I	S2		1	1
23804	G2242A	E748K	S2		1	1
23832	G2270T	G757V	S2		1	1
23856	G2294T	R765L	S2		1	1
23868	G2306T	G769V	S2		3	3
23873	G2311T	A771S	S2		8	8
23877	T2315C	V772A	S2		1	1
23895	C2333T	T778I	S2		1	1
23900	G2338C	E780Q	S2		1	1
23936	C2374T	P792S	S2		1	1

Genomic Locus	Gene Locus	Amino Acid Change	Domain	Wave 1 (n=1026)	Wave 2 (n=4059)	Total (n=5085)
23948	G2386T	D796Y	S2		2	2
23955	G2393T	G798V	S2	1		1
23987	C2425T	P809S	S2		2	2
23988	C2426T	P809L	S2		1	1
23997	C2435T	P812L	S2		1	1
24003	A2441G	K814R	S2		1	1
24014	A2452G	I818V	S2 - FP		5	5
24026	C2464T	L822F	S2 - FP		97	97
24041	A2479T	T827S	S2 - FP		4	4
24077	G2515T	D839Y	S2	2		2
24089	G2527A	D843N	S2	1	1	2
24095	G2533T	A845S	S2		5	5
24099	C2537T	A846V	S2		1	1
24129	A2567G	N856S	S2		7	7
24138	C2576T	T859I	S2		5	5
24141	T2579C	V860A	S2		1	1
24170	A2608G	I870V	S2		3	3
24188	G2626T	A876S	S2		1	1
24197	G2635T	A879S	S2		31	31
24198	C2636T	A879V	S2		1	1
24212	T2650G	S884A	S2		11	11
24237	C2675T	A892V	S2		1	1
24240	C2678T	A893V	S2	1		1
24268	G2706T	M902I	S2		1	1
24287	A2725G	I909V	S2 - HR1		2	2
24314	G2752C	E918Q	S2 - HR1	1		1
24328	G2766C	L922F	S2 - HR1		2	2
24348	G2786T	S929I	S2 - HR1		1	1
24356	G2794T	G932C	S2 - HR1		1	1
24357	G2795T	G932V	S2 - HR1		1	1
24368	G2806A	D936N	S2 - HR1		3	3
24368	G2806C	D936H	S2 - HR1		1	1
24368	G2806T	D936Y	S2 - HR1	3	4	7
24374	C2812T	L938F	S2 - HR1		3	3
24378	C2816T	S939F	S2 - HR1		4	4
24380	T2818G	S940A	S2 - HR1		5	5
24389	A2827G	S943G	S2 - HR1		6	6

Genomic Locus	Gene Locus	Amino Acid Change	Domain	Wave 1 (n=1026)	Wave 2 (n=4059)	Total (n=5085)
24463	C2901A	S967R	S2 - HR1	2		2
24507	C2945T	S982L	S2 - HR1		1	1
24579	C3017T	T1006I	S2 - CH		1	1
24588	C3026G	T1009S	S2 - CH		1	1
24621	C3059T	A1020V	S2 - CH	1		1
24638	G3076T	A1026S	S2 - CH		2	2
24642	C3080T	T1027I	S2 - CH	5		5
24649	G3087T	M1029I	S2 - CH		1	1
24710	A3148T	M1050L	S2		1	1
24710	A3148G	M1050V	S2	1	1	2
24712	G3150T	M1050I	S2		2	2
24718	C3156A	F1052L	S2	1	166	167
24770	G3208T	A1070S	S2		2	2
24794	G3232T	A1078S	S2 - CD	3	2	5
24812	G3250T	D1084Y	S2 - CD	1	29	30
24834	G3272T	R1091L	S2 - CD	1		1
24867	G3305T	W1102L	S2 - CD		1	1
24872	G3310T	V1104L	S2 - CD		1	1
24893	G3331C	E1111Q	S2 - CD		2	2
24897	C3335T	P1112L	S2 - CD	2	2	4
24912	C3350T	T1117I	S2 - CD		1	1
24923	T3361C	F1121L	S2 - CD		2	2
24933	G3371T	G1124V	S2 - CD	1	2	3
24959	G3397T	V1133F	S2 - CD		1	1
24977	G3415T	D1139Y	S2 - CD		1	1
24986	C3424A	Q1142K	S2	1		1
24998	G3436T	D1146Y	S2		4	4
24998	G3436C	D1146H	S2		13	13
25019	G3457T	D1153Y	S2		11	11
25032	A3470T	K1157M	S2	1		1
25046	C3484T	P1162S	S2		5	5
25047	C3485T	P1162L	S2		3	3
25050	A3488T	D1163V	S2		2	2
25088	G3526T	V1176F	S2		18	18
25101	A3539G	Q1180R	S2		1	1
25104	A3542G	K1181R	S2		4	4
25116	G3554A	R1185H	S2		2	2

Genomic Locus	Gene Locus	Amino Acid Change	Domain	Wave 1 (n=1026)	Wave 2 (n=4059)	Total (n=5085)
25121	A3559T	N1187Y	S2		1	1
25135	G3573T	K1191N	S2		1	1
25137	A3575C	N1192T	S2	1		1
25158	A3596C	D1199A	S2		1	1
25160	C3598T	L1200F	S2		1	1
25163	C3601A	Q1201K	S2		1	1
25169	C3607T	L1203F	S2	1		1
25183	G3621T	E1207D	S2		1	1
25186	G3624T	Q1208H	S2	1		1
25217	G3655T	G1219C	S2	1	3	4
25234	G3672T	L1224F	S2		1	1
25241	A3679G	I1227V	S2	1		1
25244	G3682T	V1228L	S2		2	2
25249	G3687T	M1229I	S2		1	1
25249	G3687C	M1229I	S2		2	2
25250	G3688A	V1230M	S2		1	1
25266	G3704T	C1235F	S2		4	4
25273	G3711T	M1237I	S2		2	2
25284	G3722T	C1241F	S2		1	1
25287	G3725T	S1242I	S2		4	4
25297	G3735T	K1245N	S2		1	1
25301	T3739G	C1247G	S2		1	1
25302	G3740T	C1247F	S2		4	4
25305	G3743T	C1248F	S2		2	2
25317	C3755T	S1252F	S2		1	1
25340	G3778T	D1260Y	S2		2	2
25350	C3788T	P1263L	S2	1	2	3
25352	G3790T	V1264L	S2		1	1
25365	T3803C	V1268A	S2		1	1

1416

1417 The domain region of RBD is based on structural information found in Cai et al.
1418 2020 (98).

1419

1420

Forty-nine of these amino acid replacements (V11A, T51A, W64C, I119T,
E156Q, S205A, D228G, L229W, P230T, N234D, I235T, T274A, A288V, E324Q,
E324V, S325P, S349F, S371P, S373P, T385I, A419V, C480F, Y495S, L517F,
K528R, Q628E, T632I, S708P, T719I, P728L, S746P, E748K, G757V, V772A,

K814R, D843N, S884A, M902I, I909V, E918Q, S982L, M1029I, Q1142K,
K1157M, Q1180R, D1199A, C1241F, C1247G, and V1268A) were not
represented in a publicly available database (34) as of August 19, 2020

1428