Evidence for and level of herd immunity against SARS-CoV-2 infection: the ten-community study

Andrew Jeremijenko MD,[1] Hiam Chemaitelly MSc,[2,3] Houssein H. Ayoub PhD,[4] Moza Abdellatif Hassan Abdulla PhD, [1] Abdul-Badi Abou-Samra MD PhD,[1] Jameela Ali A.A. Al Ajmi MD,[1] Nasser Ali Asad Al Ansari PhD,[1] Zaina Al Kanaani PhD,[1] Abdullatif Al Khal MD,[1] Einas Al Kuwari MD,[1] Ahmed Al-Mohammed MD,[1] Naema Hassan Abdulla Al Molawi PhD,[1] Huda Mohamad Al Naomi MD,[1] Adeel A. Butt MBBS MS,[1] Peter Coyle PhD,[1] Reham Awni El Kahlout MSc,[1] Imtiaz Gillani MSc[1] Anvar Hassan Kaleeckal MSc,[1] Naseer Ahmad Masoodi MD,[1] Anil George Thomas MHA,[1] Hanaa Nafady-Hego MD PhD,[1,5] Ali Nizar Latif MD,[1] Riyazuddin Mohammad Shaik MSc,[1] Nourah B M Younes PhD,[1] Hanan F. Abdul Rahim PhD,[6] Hadi M. Yassine PhD,[7,8] Mohamed G. Al Kuwari MD,[9] Hamad Eid Al Romaihi MD,[10] Sheikh Mohammad Al Thani MD,[10] Roberto Bertollini MD[10] and Laith J. Abu-Raddad PhD,[2,3,11]

[1]*Hamad Medical Corporation, Doha, Qatar*
[2]*Infectious Disease Epidemiology Group, Weill Cornell Medicine-Qatar, Cornell University, Doha, Qatar*
[3]*World Health Organization Collaborating Centre for Disease Epidemiology Analytics on HIV/AIDS, Sexually Transmitted Infections, and Viral Hepatitis, Weill Cornell Medicine–Qatar, Cornell University, Qatar Foundation – Education City, Doha, Qatar*
[4]*Department of Mathematics, Statistics, and Physics, Qatar University, Doha, Qatar*
[5]*Microbiology and Immunology Department, Faculty of Medicine, Assiut University, Assiut, Egypt*
[6]*College of Health Sciences, QU Health, Qatar University, Doha, Qatar*
[7]*Biomedical Research Center, Qatar University, Doha, Qatar*
[8]*Department of Biomedical Science, College of Health Sciences, Member of QU Health, Qatar University, Doha, Qatar*
[9]*Primary Health Care Corporation, Doha, Qatar*
[10]*Ministry of Public Health, Doha, Qatar*
[11]*Department of Population Health Sciences, Weill Cornell Medicine, Cornell University, New York, New York, USA*

[*]Address reprints requests or correspondence to

Dr. Andrew Jeremijenko, Hamad Medical Corporation, P.O. Box 3050, Doha, Qatar. E-mail: AJeremijenko@hamad.qa.

Professor Laith J. Abu-Raddad, Infectious Disease Epidemiology Group, World Health Organization Collaborating Centre for Disease Epidemiology Analytics on HIV/AIDS, Sexually Transmitted Infections, and Viral Hepatitis, Weill Cornell Medicine - Qatar, Qatar Foundation - Education City, P.O. Box 24144, Doha, Qatar. Telephone: +(974) 4492-8321. Fax: +(974) 4492-8333. E-mail: lja2002@qatar-med.cornell.edu.

ABSTRACT

Background: Qatar experienced a large severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) epidemic that disproportionately affected the craft and manual workers (CMWs) who constitute 60% of the population. This study aimed to investigate level of immunity in communities within this population as well as infection exposure required to achieve herd immunity.

Methods: Anti-SARS-CoV-2 seropositivity was assessed in ten CMW communities between June 21 and September 9, 2020. PCR positivity, infection positivity (antibody and/or PCR positive), and infection severity rate were also estimated. Associations with anti-SARS-CoV-2 positivity were investigated using regression analyses.

Results: Study included 4,970 CMWs who were mostly men (95.0%) and <40 years of age (71.5%). Seropositivity ranged from 54.9% (95% CI: 50.2-59.4%) to 83.8% (95% CI: 79.1-87.7%) in the different CMW communities. Pooled mean seropositivity across all communities was 66.1% (95% CI: 61.5-70.6%). PCR positivity ranged from 0.0% to 10.5% (95% CI: 7.4-14.8%) in the different CMW communities. Pooled mean PCR positivity was 3.9% (95% CI: 1.6-6.9%). Median cycle threshold (Ct) value was 34.0 (range: 15.8-37.4)—majority (79.5%) of PCR-positive individuals had Ct value >30 indicative of earlier rather than recent infection. Infection positivity (antibody and/or PCR positive) ranged from 62.5% (95% CI: 58.3-66.7%) to 83.8% (95% CI: 79.1-87.7%) in the different CMW communities. Pooled mean infection positivity was 69.5% (95% CI: 62.8-75.9%). Only five infections were ever severe and one was ever critical—an infection severity rate of 0.2% (95% CI: 0.1-0.4%).

Conclusions: Based on an extended range of epidemiological measures, active infection is rare in these communities with limited if any sustainable infection transmission for clusters to occur.

At least some CMW communities in Qatar have reached or nearly reached herd immunity for SARS-CoV-2 infection at a proportion of ever infection of 65-70%.

Keywords: SARS-CoV-2; epidemiology; COVID-19; infection; seroprevalence; immunity

Introduction

Since the start of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) epidemic, millions of infections have been laboratory-confirmed globally [1], and millions others must have gone undocumented [2]. Two key questions remain unanswered: has any community reached herd immunity to render infection transmission chains unsustainable? What level of exposure to the infection (attack rate) is needed to reach herd immunity?

Qatar, a peninsula in the Arabian Gulf with a diverse population of 2.8 million [3], experienced a large-scale SARS-CoV-2 epidemic [4, 5]. By August 27, 2020, the rate of laboratory-confirmed infections in Qatar was at 50,324 per million population, one of the highest worldwide [6, 7]. The epidemic, currently in an advanced stage [4], seems to have followed a classic susceptible-infected-recovered "SIR" pattern with an epidemic peak around May 20 followed by a steady decline for the next four months [4].

The most affected subpopulation by this epidemic was that of the expatriate craft and manual workers (CMWs) among whom community transmission was first identified [4]. These workers constitute about 60% of the Qatar population and are typically single men aged 20-49 years [8]. CMWs at a given workplace or company not only work together during the day, but also live together as a *community* in large dormitories or housing complexes where they share rooms, bathrooms, and cafeteria-style meals [4, 9, 10]. These communities stay mostly in contact with their own community members and infrequently mingle with other communities, creating a geographic "bubble" that proved important for the pattern of infection transmission [4]. With reduced options for effective social and physical distancing, SARS-CoV-2 transmission in these CMW communities resembled that of influenza outbreaks in schools [4, 11, 12], and especially boarding schools [12].

Given the large number of diagnosed infections in CMWs [4], the large proportion of infections that were asymptomatic [4], the high polymerase chain reaction (PCR)-positivity rates in the random testing campaigns conducted around the epidemic peak in different CMW communities [4], and the observed "SIR" epidemic curve with rapid declines in incidence for over four months despite easing of the social and physical distancing restrictions [4], all pose a question as to whether herd immunity may have been reached in at least some of these communities.

Our aim was to assess ever exposure to the SARS-CoV-2 infection and attainment of herd immunity in several CMW communities by assessing the level of detectable antibodies. Operationally, we defined herd immunity as the proportion of the population ever infected ("attack rate") beyond which infection transmission/circulation becomes unsustainable in this population with limited if any new infections occurring. The study was conducted to inform the national response and preparedness for potential future infection waves.

Methods

Data sources

Testing for detectable SARS-CoV-2-specific antibodies on blood specimens was conducted in ten CMW communities between June 21 and September 9, 2020 as part of a priori designed study combined with a testing and surveillance program led by the Ministry of Public Health (MOPH) and Hamad Medical Corporation (HMC), the main public healthcare provider and the nationally-designated provider for all COVID-19 healthcare needs in Qatar. The goal of this program was to assess the level of infection exposure in different subpopulations and economic sectors.

Study design was opportunistic utilizing the MOPH-HMC program and the need for rapid data collection to inform the national response. The ten CMW communities were selected for feasibility and/or given earlier random PCR testing campaigns or contact tracing that suggested substantial infection levels. For instance, CMW Community 1 was part of a random PCR testing campaign that identified a high positivity rate of 59% in late April. In six select communities, PCR testing was also simultaneously conducted to assess active infection using nasopharyngeal swabs, and in one select community (CMW Community 1), an interview schedule (based on World Health Organization (WHO) suggested questionnaire [13]) was administered to collect data on socio-demographics and history of exposure and symptoms.

The population size of each of these communities ranged from few hundreds to few thousands who live in shared accommodations provided by the employers. The companies that employ these workers belonged to the service or industrial sectors, but the bulk of the employees, even in the industrial companies, worked on providing services such as catering, cleaning and other janitorial services, warehousing, security, and port workers.

Employers were contacted and those agreeing to participate were asked to advertise the availability and location of testing sites to their employees. Individuals' participation was voluntary. Employees interested in being tested and in knowing their status were provided with transportation to HMC testing sites. Informed consent and questionnaire were provided and collected in nine languages (Arabic, Bengali, English, Hindi, Urdu, Nepali, Sinhala, Tagalog, and Tamil) to cater to the main language groups in these CMW communities. National guidelines and standard of care were applied to all identified PCR positive cases. No action was mandated by the national guidelines to those found antibody positive, and thus no action was taken apart from notifying individuals of their sero-status.

Results of the serological testing were subsequently linked to the HMC centralized and standardized database comprising all SARS-CoV-2 PCR testing conducted in Qatar since the start of the epidemic [4, 6]. The database also includes data on hospitalization and on the WHO severity classification [14] for each laboratory-confirmed infection.

The study was approved by HMC and Weill Cornell Medicine-Qatar Institutional Review Boards.

Laboratory methods

Testing for SARS-CoV-2-specific antibodies in the serological samples was performed using an electrochemiluminescence immunoassay, the Roche Elecsys® Anti-SARS-CoV-2 (Roche, Switzerland). Results' interpretation was per manufacturer's instructions: reactive for cutoff index ≥1.0 and non-reactive for cutoff index <1.0 [15].

PCR testing was performed on aliquots of Universal Transport Medium (UTM) used for nasopharyngeal swabs' collection (Huachenyang Technology, China). Aliquots were: extracted on the QIAsymphony platform (QIAGEN, USA) and tested with real-time reverse-transcription PCR (RT-qPCR) using the TaqPath™ COVID-19 Combo Kit (Thermo Fisher Scientific, USA) on a ABI 7500 FAST (ThermoFisher, USA); extracted using a custom protocol [16] on a Hamilton Microlab STAR (Hamilton, USA) and tested using the AccuPower SARS-CoV-2 Real-Time RT-PCR Kit (Bioneer, Korea) on a ABI 7500 FAST; or loaded directly to a Roche cobas® 6800 system and assayed with the cobas® SARS-CoV-2 Test (Roche, Switzerland).

All laboratory testing was conducted at HMC Central Laboratory following standardized protocols.

Statistical analysis

Frequency distributions were used to describe CMWs' characteristics and to estimate different SARS-CoV-2 epidemiological measures. Anti-SARS-CoV-2 pooled mean prevalence across CMW communities was estimated using meta-analysis. Here, a DerSimonian-Laird random-effects model [17] was applied to pool seroprevalence measures that were weighted using the inverse-variance method [18, 19].

Chi-square tests and univariable logistic regressions were implemented to explore associations with anti-SARS-CoV-2 positivity. Odds ratios (ORs), 95% confidence intervals (CIs), and p-values were generated. Covariates with p-value ≤0.2 in univariable regression analysis were included in the multivariable logistic regression model where applicable. Covariates with p-value ≤0.05 in the multivariable model were considered as showing evidence for an association with the outcome. The distribution of PCR cycle threshold (Ct) values among persons testing PCR positive was further generated, and summary statistics reported.

Results

A total of 4,970 CMWs from the ten CMW communities participated in this study (Table 1). Participants were mostly men (95.0%), below 40 years of age (71.5%), and of Nepalese (43.0%), Indian (33.1%), or Bangladeshi (11.6%) origin. Regression analyses identified each of sex, nationality, and CMW Community to be independently associated with seropositivity.

Women had 87% lower odds of being seropositive compared to men (adjusted odds ratio (AOR): 0.13; 95% CI: 0.09-0.19; Table 1). Compared to all other nationalities, AOR was 6.78 (95% CI: 4.31-10.66) for Bangladeshis, 4.93 (95% CI: 3.27-7.42) for Nepalese, 3.60 (95% CI: 2.40-5.41) for Indians, 3.43 (95% CI: 1.99-5.90) for Kenyans, 2.81 (95% CI: 1.66-4.76) for Sri Lankans,

and 2.23 (95% CI: 1.32-3.75) for Filipinos. Some differences in seropositivity by CMW Community were noted. No significant differences in seropositivity by age group were found.

Table 2 shows the characteristics and associations with anti-SARS-CoV-2 positivity for only CMW Community 1 where a specific interview schedule was administered and collected specific socio-demographic data and history of exposure and symptoms. Close to 40% of participants had intermediate or low educational attainment, and a third had higher schooling levels or vocational training. University education was associated with 75% (OR: 0.25; 95% CI: 0.09-0.67) lower odds of seropositivity compared to intermediate or lower educational attainment. No statistically-significant associations with seropositivity were found for contact with an infected person, presence of symptoms, or symptoms requiring medical attention. Appendix Table S1 shows also the characteristics and associations with anti-SARS-CoV-2 positivity for CMW Communities 2-10. For each of these communities, associations were found for sex and nationality, but no notable associations were found for age group.

Figure 1 illustrates key SARS-CoV-2 epidemiological measures in the different CMW communities. Out of a total of 4,970 anti-SARS-CoV-2 test results for these CMWs, 3,199 (64.4%; 95% CI: 63.0-65.7%) were seropositive. Seropositivity ranged from 54.9% (95% CI: 50.2-59.4%) in CMW Community 5 to 83.8% (95% CI: 79.1-87.7%) in CMW Community 3 (Figure 1A). The pooled mean anti-SARS-CoV-2 positivity across the ten CMW communities was 66.1% (95% CI: 61.5-70.6%).

Out of a total of 2,016 PCR test results for these CMWs, 112 (5.6%; 95% CI: 4.6-6.6%) were positive. PCR positivity ranged from 0.0% (95% CI: 0.0-3.9%) in CMW Community 1 and 0.0% (95% CI: 0.0-9.0%) in CMW Community 8 to 10.5% (95% CI: 7.4-14.8%) in CMW Community 3 (Figure 1B). Pooled mean PCR positivity across the six CMW communities where PCR testing

was conducted was 3.9% (95% CI: 1.6-6.9%). Ct values ranged from 15.8-37.4 with a median of 34.0 (Figure 2). The vast majority (79.5%) of PCR-positive individuals had a Ct value >30 suggestive of no active infection [20, 21]. Significant differences in PCR positivity were found by nationality and CMW Community (Appendix Table S2).

Infection positivity (antibody and/or PCR positive) ranged from 62.5% (95% CI: 58.3-66.7%) in CMW Community 4 to 83.8% (95% CI: 79.1-87.7%) in CMW Community 3 (Figure 1C). Pooled mean infection positivity across the six CMW communities with antibody and PCR results was 69.5% (95% CI: 62.8-75.9%).

Data were linked to the national SARS-CoV-2 PCR testing and hospitalization database. Of the 3,199 antibody positive CMWs, 1,012 (31.6%; 95% CI: 30.0-33.3%) were *previously* diagnosed with SARS-CoV-2 infection (had a laboratory-confirmed PCR positive result *before* this study). For the CMW communities that were previously part of broad PCR testing because of a case identification and/or a random testing campaign, the diagnosis rate ranged from 28.0% (95% CI: 19.1-38.2%) in CMW Community 8 to 82.9% (95% CI: 76.8-87.9%) in CMW Community 1. Meanwhile, where no such broad PCR testing was conducted, the diagnosis rate was only 13.2% (95% CI: 10.7-16.1%) in CMW Community 10, 7.4% (95% CI: 4.7-11.2%) in CMW Community 2, and 0.4% (95% CI: 0.0-2.3%) in CMW Community 3. Only a very small fraction of antibody *negative* persons, 14 out of 1,771 (0.8%; 95% CI: 0.4-1.3%), had been previously diagnosed as PCR positive (Appendix Table S3).

Of the total sample, 21 individuals had a hospitalization record associated with a SARS-CoV-2 infection diagnosis, of whom, infection severity per WHO classification was mild for five, moderate for ten, severe for five, and critical for one. All 21 individuals eventually cleared their infection and were discharged from the hospital. All these individuals also tested anti-SARS-

CoV-2 positive. Accordingly, the proportion of those with a confirmed *severe* or *critical* infection out of 3,233 who had a laboratory-confirmed infection (antibody and/or PCR positive result) was 0.2% (95% CI: 0.1-0.4%).

Discussion

Above results support that herd immunity has been reached (or at least nearly reached) in these CMW communities, and that the level of herd immunity needed for SARS-CoV-2 infection is an attack rate (proportion ever infected) of about 65-70%.

This conclusion is supported by: i) these CMW communities had comparable seroprevalence of about 65-70%; ii) PCR positivity was low and the vast majority of those PCR positive had high Ct value suggestive of an earlier rather than recent infection [20, 21]; iii) only few persons had active infection (Ct value <25) and no significant infection cluster was identified in any of these CMW communities during this study (suggestive of isolated infections and unsustainable infection transmission for clusters to occur); iv) the level of 65-70% infection exposure is in concordance with that predicted using the "classical" formula for herd immunity of $\sim 1 - 1/R_0$ [22, 23], with R_0, the basic reproduction number, being at 2.5-4 [24, 25].

Notably, although large clusters of infection were very common in such CMW communities before and around the epidemic peak towards end of May, no major cluster has been identified in any CMW community in Qatar for several weeks now, despite the progressive easing of the social and physical distancing restrictions since June 15, 2020.

These findings indicate that reaching herd immunity in such largely homogenous communities requires high exposure level of about 65-70%. However, it is possible that *true* herd immunity may have been reached even at a lower attack rate. Mathematical modeling indicates that

infection exposure for a novel infection (especially in the first cycle) can considerably

"overshoot" the "classical" herd immunity level of $\sim 1 - 1/R_0$, more so if the social contact rate

within this community is homogeneous (illustration is shown in Appendix Figure S1).

Meanwhile, heterogeneity in social contact rate can reduce the final attack rate (Appendix Figure

S1) [23, 26].

This study had other notable findings. Severity rate for SARS-CoV-2 infection was low (0.2%),

possibly because of the young age of the CMWs. No COVID-19 deaths were reported in these

CMW communities. Remarkably, in the communities where no prior broad PCR testing was

conducted, <15% of the antibody positive subjects had ever been diagnosed as PCR positive

prior to this study. There was a large difference in infection exposure between women and men

(Table 1). This difference, with the variable proportion of women across these communities,

explains also part of the variation seen in seroprevalence across these communities (Figure 1

versus Appendix Figure S2). This finding may be attributed to women and men living in

different housing accommodations and having different work roles. Women, a very small

minority in these CMW communities, live in small shared accommodations as opposed to the

large ones hosting men.

There were differences by nationality (Table 1), but these are explained by nearly all

Bangladeshis and Nepalese and most Indians being the workers in these communities, while a

proportion of Indians and much of the other nationalities holding administrative or managerial

positions with lower social contact rates and possibly living in a different kind of

accommodations compared to the bulk of the workers. No major differences in infection

exposure by age were found, though there was some tendency for those >40 years of age to have

lower infection exposure (Appendix Table S1), possibly due to different occupations within these communities.

This study has limitations. Testing was conducted in select CMW communities and therefore findings may not be generalizable to the wider CMW population in Qatar. Response rate could not be precisely ascertained given uncertainty around the number of CMWs who were aware of the invitation to participate, but based on employer-reported counts of the size of each community, the response rate is >50% and participants expressed high interest in knowing their antibody status. The validity of study outcomes is contingent on the sensitivity and specificity of the used assays. However, the laboratory methods were based on high-quality commercial platforms, and each diagnostic method was validated in the laboratory before its use. Notably, the antibody assay had high specificity reported at 99.8% [15] by the manufacturer and at 100% by a validation study by Public Health England [27].

In conclusion, at least some of the CMW communities in Qatar, who constitute about 60% of the total population [8], have reached or nearly reached herd immunity for SARS-CoV-2 infection, providing to our knowledge the first empirical evidence for herd immunity worldwide. While achieving herd immunity at a national level is difficult within few months [28], herd immunity could be achieved in specific communities within few months. In such relatively homogenous communities, reaching herd immunity required infection of 65-70% of the members of the community. These findings suggest that the SARS-CoV-2 epidemic in a homogenous population is unlikely to be unsustainable before as much as two-thirds of the population become infected. This also suggests that a SARS-CoV-2 vaccine needs at least 65-70% efficacy at universal coverage for herd immunity to be achieved in a population naïve to SARS-CoV-2 infection [29].

Table 1. Characteristics of the craft and manual workers (CMWs) and associations with anti-SARS-CoV-2 positivity.

Characteristics	Tested	Anti-SARS-CoV-2 positive		Univariable regression analysis		Multivariable regression analysis	
	N (%)	N (%)	p-value	OR (95% CI)	p-value[*]	OR (95% CI)	p-value[†]
Sex							
Men	4,721 (95.0)	3,153 (66.8)	<0.001	1.00		1.00	
Women	249 (5.0)	46 (18.5)		0.11 (0.08-0.16)	<0.001	0.13 (0.09-0.19)	<0.001
Age (years)							
<29	1,579 (31.8)	1,031 (65.3)	<0.001	1.00		1.00	
30-39	1,973 (39.7)	1,226 (62.1)		0.87 (0.76-1.00)	0.052	0.90 (0.78-1.05)	0.178
40-49	1,040 (20.9)	680 (65.4)		1.00 (0.85-1.18)	0.962	1.12 (0.93-1.35)	0.216
50+	339 (6.8)	225 (66.4)		1.05 (0.82-1.34)	0.705	1.21 (0.92-1.59)	0.170
Missing	39 (0.8)	37 (94.9)		9.83 (2.36-40.95)	0.002	9.57 (2.22-41.32)	0.002
Nationality							
All other nationalities[‡]	125 (2.5)	40 (32.0)		1.00		1.00	
Filipino	186 (3.7)	68 (36.6)		1.22 (0.76-1.98)	0.408	2.23 (1.32-3.75)	0.003
Sri Lankan	147 (3.0)	77 (52.4)		2.34 (1.42-3.84)	0.001	2.81 (1.66-4.76)	<0.001
Kenyan	152 (3.1)	77 (50.7)		2.18 (1.33-3.57)	0.002	3.43 (1.99-5.90)	<0.001
Indian	1,647 (33.1)	1,035 (62.8)	<0.001	3.59 (2.44-5.30)	<0.001	3.60 (2.40-5.41)	<0.001
Nepalese	2,136 (43.0)	1,468 (68.7)		4.67 (3.17-6.88)	<0.001	4.93 (3.27-7.42)	<0.001
Bangladeshi	577 (11.6)	434 (75.2)		6.45 (4.23-9.82)	<0.001	6.78 (4.31-10.66)	<0.001
CMW community							
CMW Community 6[§]	1,505 (30.3)	946 (62.9)	<0.001	1.00[§]		1.00[§]	
CMW Community 5	443 (8.9)	243 (54.9)		0.72 (0.58-0.89)	0.002	0.75 (0.60-0.95)	0.015
CMW Community 4	534 (10.7)	330 (61.8)		0.96 (0.78-1.17)	0.664	0.85 (0.66-1.09)	0.193
CMW Community 10	957 (19.3)	620 (64.8)		1.09 (0.92-1.29)	0.332	0.98 (0.82-1.17)	0.812
CMW Community 7	188 (3.8)	122 (64.9)		1.09 (0.80-1.50)	0.585	0.99 (0.71-1.37)	0.934
CMW Community 2	456 (9.2)	282 (61.8)		0.96 (0.77-1.19)	0.695	1.10 (0.85-1.41)	0.462
CMW Community 9	202 (4.1)	126 (62.4)		0.98 (0.72-1.33)	0.894	1.29 (0.92-1.81)	0.142
CMW Community 8	139 (2.8)	93 (66.9)		1.19 (0.83-1.73)	0.344	1.45 (0.98-2.15)	0.064
CMW Community 1	255 (5.1)	193 (75.7)		1.84 (1.36-2.49)	<0.001	1.90 (1.37-2.64)	<0.001
CMW Community 3	291 (5.9)	244 (83.8)		3.07 (2.21-4.26)	<0.001	2.64 (1.89-3.68)	<0.001

CI, confidence interval; OR, odds ratio.

[*]Covariates with p-value ≤0.2 in the univariable analysis were included in the multivariable analysis.

[†]Covariates with p-value ≤0.05 in the multivariable analysis were considered predictors of SARS-CoV-2 antibody positivity.

[‡]Includes all other nationalities that contributed <10% of the sample in each community.

[§]CMW Community 6 which had the largest sample size was chosen as a reference.

Table 2. Characteristics of the Craft and Manual Worker (CMW) Community 1 and associations with anti-SARS-CoV-2 positivity including socio-demographics and history of exposure and symptoms.

Characteristics	Tested	Anti-SARS-CoV-2 positive		Univariable regression analysis[*]	
	N (%[†])	N (%[‡])	p-value	OR (95% CI)	p-value[§]
Sex					
Men	240 (94.1)	189 (78.8)	<0.001	1.00	
Women	15 (5.9)	4 (26.7)		0.10 (0.03-0.32)	<0.001
Age (years)					
<29	105 (41.2)	84 (80.0)	0.322	1.00	
30-39	111 (43.5)	83 (74.8)		0.74 (0.39-1.41)	0.360
40-49	27 (10.6)	19 (70.4)		0.59 (0.23-1.54)	0.284
50+	12 (4.7)	7 (58.3)		0.35 (0.10-1.21)	0.098
Nationality					
All other nationalities[¶]	48 (18.8)	23 (47.9)	<0.001	1.00	
Indian	32 (12.5)	20 (62.5)		1.81 (0.73-4.51)	0.202
Nepalese	157 (61.6)	132 (84.1)		5.74 (2.82-11.67)	<0.001
Bangladeshi	18 (7.1)	18 (100.0)		omitted	--
Educational attainment					
Intermediate or lower	101 (39.6)	88 (87.1)	<0.001	1.00	
Secondary/High school/Vocational	80 (31.4)	69 (86.3)		0.93 (0.39-2.20)	0.863
University	27 (10.6)	17 (63.0)		0.25 (0.09-0.67)	0.005
Missing	47 (18.4)	19 (40.4)		0.10 (0.04-0.23)	<0.001
Contact with infected person					
No	124 (48.6)	93 (75.0)	0.303	1.00	
Yes	14 (5.5)	13 (92.9)		4.33 (0.54-34.48)	0.166
Unknown/Missing	117 (45.9)	87 (74.4)		0.97 (0.54-1.73)	0.909
Presence of symptoms in the past two weeks					
Asymptomatic	184 (72.2)	148 (80.4)	<0.001	1.00	
One symptom	16 (6.3)	16 (100.0)		omitted	--
Two or more symptoms	12 (4.7)	12 (100.0)		omitted	--
Missing	43 (16.9)	17 (39.5)		0.16 (0.08-0.32)	<0.001
Symptoms required medical attention					
No	210 (82.4)	174 (82.9)	<0.001	1.00	
Yes	3 (1.2)	3 (100.0)		omitted	--
Unknown/Missing	42 (16.5)	16 (38.1)		0.13 (0.06-0.26)	<0.001

CI, confidence interval; OR, odds ratio.
[*]No multivariable regression was performed for lack of relevance.
[†]Percentage of the total sample.
[‡]Percentage positive out of those tested.
[§]Covariates with p-value ≤0.05 in the univariable analysis were considered as showing evidence for an association with anti-SARS-CoV-2 positivity.
[¶]Includes all other nationalities that contributed <10% of the sample.

Figure 1. Measures of SARS-CoV-2 A) antibody positivity, B) PCR positivity, C) infection positivity (antibody and/or PCR positive), and D) diagnosis rate, across the craft and manual worker (CMW) communities. Of note that PCR testing was done in only six communities.

Figure 2. Distribution of PCR cycle threshold (Ct) values among craft and manual workers (CMWs) identified as SARS-CoV-2 PCR positive during the study period.

Author contributions

AJ and LJA co-conceived and co-designed the study. AJ led data collection. HC performed the data analyses and wrote the first draft of the article. HHA contributed to analysis of data. All authors contributed to data acquisition, database development, testing, program development, discussion and interpretation of the results, and to the writing of the manuscript. All authors have read and approved the final manuscript.

Competing interests

We declare no competing interests.

Acknowledgement

We would like to thank Her Excellency Dr. Hanan Al Kuwari, the Minister of Public Health, for her vision, guidance, leadership, and support. We also would like to thank Dr. Saad Al Kaabi, Chair of the System Wide Incident Command and Control (SWICC) Committee for the COVID-19 national healthcare response, for his leadership, analytical insights, and for his instrumental role in enacting the data information systems that made these studies possible. We further extend our appreciation to the SWICC Committee and the Scientific Reference and Research Taskforce (SRRT) members for their informative input, scientific technical advice, and enriching discussions. We would also like to thank Dr. Mariam Abdulmalik, the CEO of the Primary Health Care Corporation and the Chairperson of the Tactical Community Command Group on COVID-19, as well as members of this committee, for providing support to the teams that worked on the field surveillance. We also would like to acknowledge the dedicated efforts of the Clinical Coding Team and the COVID-19 Mortality Review Team, both at Hamad Medical

Corporation, and the Surveillance Team at the Ministry of Public Health. We further thank all companies that have facilitated the participation of their employees in this study.

Funding

The authors are grateful for support provided by the Ministry of Public Health, Hamad Medical Corporation, and the Biomedical Research Program and the Biostatistics, Epidemiology, and Biomathematics Research Core, both at Weill Cornell Medicine-Qatar. The statements made herein are solely the responsibility of the authors.

References

1. *COVID-19 Outbreak Live Update. Available from: https://www.worldometers.info/coronavirus/. Accessed on September 06, 2020.* 2020.
2. Ioannidis, J., *The infection fatality rate of COVID-19 inferred from seroprevalence data.* medRxiv, 2020: p. 2020.05.13.20101253.
3. Planning and Statistics Authority- State of Qatar, *Qatar Monthly Statistics. Available from: https://www.psa.gov.qa/en/pages/default.aspx. Accessed on: may 26,2020.* 2020.
4. Abu-Raddad, L.J., et al., *Characterizing the Qatar advanced-phase SARS-CoV-2 epidemic.* medRxiv, 2020: p. 2020.07.16.20155317v2.
5. Al Kuwari, H.M., et al., *Characterization of the SARS-CoV-2 outbreak in the State of Qatar, February 28-April 18, 2020.* BMJ open (accepted for publication), 2020.
6. Hamad Medical Corporation, *SARS-CoV-2 hospitalizations and care.* 2020.
7. Ministry of Public Health-State of Qatar, *Coronavirus Disease 2019 (COVID-19). Available from: https://covid19.moph.gov.qa/EN/Pages/default.aspx. Accessed on: May 25, 2020.* 2020.
8. Planning and Statistics Authority- State of Qatar, *Labor force sample survey. Available from: https://www.psa.gov.qa/en/statistics/Statistical%20Releases/Social/LaborForce/2017/statistical_analysis_labor_force_2017_En.pdf. Accessed on: May 01, 2020.* 2017.
9. Ministry of Interior-State of Qatar, *Population distribution by sex, age, and nationality: results of Kashef database.* 2020.
10. De Bel-Air, F., *Demography, Migration, and Labour Market in Qatar. Available from: https://www.researchgate.net/publication/323129801_Demography_Migration_and_Labour_Market_in_Qatar-_UPDATED_June_2017. Accessed on May 01, 2020.* 2018, Gulf Labour Markets and Migration.
11. Jackson, C., et al., *School closures and influenza: systematic review of epidemiological studies.* BMJ Open, 2013. **3**(2).
12. Glatman-Freedman, A., et al., *Attack rates assessment of the 2009 pandemic H1N1 influenza A in children and their contacts: a systematic review and meta-analysis.* PLoS One, 2012. **7**(11): p. e50228.
13. World Health Organization, *Population-based age-stratified seroepidemiological investigation protocol for COVID-19 virus infection. Available from: file:///C:/Users/hsc2001/Downloads/covid-19-seroepidemiological-investigation-protocol-v3.pdf. Accessed on: April 15, 2020.* 2020.
14. World Health Organization, *Clinical management of COVID-19. Available from: https://www.who.int/publications-detail/clinical-management-of-covid-19. Accessed on: May 31st 2020.* 2020.
15. The Roche Group, *Roche's COVID-19 antibody test receives FDA Emergency Use Authorization and is available in markets accepting the CE mark. Available from: https://www.roche.com/media/releases/med-cor-2020-05-03.htm. Accessed on: June 5, 2020.* 2020.
16. Kalikiri, M.K.R., et al., *High-throughput extraction of SARS-CoV-2 RNA from nasopharyngeal swabs using solid-phase reverse immobilization beads.* medRxiv, 2020: p. 2020.04.08.20055731.
17. DerSimonian, R. and N. Laird, *Meta-analysis in clinical trials.* Control Clin Trials, 1986. **7**(3): p. 177-88.
18. Miller, J.J., *The Inverse of the Freeman – Tukey Double Arcsine Transformation.* The American Statistician, 1978. **32**(4): p. 138-138.
19. Barendregt, J.J., et al., *Meta-analysis of prevalence.* J Epidemiol Community Health, 2013. **67**(11): p. 974-8.

20.	Sethuraman, N., S.S. Jeremiah, and A. Ryo, *Interpreting Diagnostic Tests for SARS-CoV-2.* JAMA, 2020.

21.	Wajnberg, A., et al., *Humoral immune response and prolonged PCR positivity in a cohort of 1343 SARS-CoV 2 patients in the New York City region.* medRxiv, 2020: p. 2020.04.30.20085613.

22.	Anderson, R.M., et al., *How will country-based mitigation measures influence the course of the COVID-19 epidemic?* Lancet, 2020. **395**(10228): p. 931-934.

23.	Britton, T., F. Ball, and P. Trapman, *A mathematical model reveals the influence of population heterogeneity on herd immunity to SARS-CoV-2.* Science, 2020. **369**(6505): p. 846-849.

24.	He, W., G.Y. Yi, and Y. Zhu, *Estimation of the basic reproduction number, average incubation time, asymptomatic infection rate, and case fatality rate for COVID-19: Meta-analysis and sensitivity analysis.* J Med Virol, 2020.

25.	MIDAS Online COVID-19 Portal, *COVID-19 parameter estimates: basic reproduction number. Available from: https://github.com/midas-network/COVID-19/tree/master/parameter_estimates/2019_novel_coronavirus. Accessed on: MAy 19, 2020.* 2020.

26.	Aguas, R., et al., *Herd immunity thresholds for SARS-CoV-2 estimated from unfolding epidemics.* medRxiv, 2020: p. 2020.07.23.20160762.

27.	Public Health England, *Evaluation of Roche Elecsys AntiSARS-CoV-2 serology assay for the detection of anti-SARS-CoV-2 antibodies. Available from: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/891598/Evaluation_of_Roche_Elecsys_anti_SARS_CoV_2_PHE_200610_v8.1_FINAL.pdf. Accessed on June 5, 2020.* 2020.

28.	Pollan, M., et al., *Prevalence of SARS-CoV-2 in Spain (ENE-COVID): a nationwide, population-based seroepidemiological study.* Lancet, 2020. **396**(10250): p. 535-544.

29.	Makhoul, M., et al., *Epidemiological impact of SARS-CoV-2 vaccination: mathematical modeling analyses.* medRxiv, 2020: p. 2020.04.19.20070805.

Appendix

Evidence for and level of herd immunity against SARS-CoV-2 infection: the ten-community study

Andrew Jeremijenko MD,[1] Hiam Chemaitelly MSc,[2,3] Houssein H. Ayoub PhD,[4] Moza Abdellatif Hassan Abdulla PhD, [1] Abdul Badi Abou Samra MD PhD,[1] Jameela Ali A.A. Al Ajmi MD,[1] Nasser Ali Asad Al Ansari PhD,[1] Zaina Al Kanaani PhD,[1] Abdullatif Al Khal MD,[1] Einas Al Kuwari MD,[1] Ahmed Al-Mohammed MD,[1] Naema Hassan Abdulla Al Molawi PhD,[1] Huda Mohamad Al Naomi MD,[1] Adeel A. Butt MBBS MS,[1] Peter Coyle PhD,[1] Reham Awni El Kahlout MSc,[1] Imtiaz Gillani MSc[1] Anvar Hassan Kaleeckal MSc,[1] Naseer Ahmad Masoodi MD,[1] Anil George Thomas MHA,[1] Hanaa Nafady-Hego MD PhD,[1,5] Ali Nizar Latif MD,[1] Riyazuddin Mohammad Shaik MSc,[1] Nourah B M Younes PhD,[1] Hanan F. Abdul Rahim PhD,[6] Hadi M. Yassine PhD,[7,8] Mohamed G. Al Kuwari MD,[9] Hamad Eid Al Romaihi MD,[10] Sheikh Mohammad Al Thani MD,[10] Roberto Bertollini MD[10] and Laith J. Abu-Raddad PhD,[2,3,11]

[1]*Hamad Medical Corporation, Doha, Qatar*

[2]*Infectious Disease Epidemiology Group, Weill Cornell Medicine-Qatar, Cornell University, Doha, Qatar*

[3]*World Health Organization Collaborating Centre for Disease Epidemiology Analytics on HIV/AIDS, Sexually Transmitted Infections, and Viral Hepatitis, Weill Cornell Medicine–Qatar, Cornell University, Qatar Foundation – Education City, Doha, Qatar*

[4]*Department of Mathematics, Statistics, and Physics, Qatar University, Doha, Qatar*

[5]*Microbiology and Immunology Department, Faculty of Medicine, Assiut University, Assiut, Egypt*

[6]*College of Health Sciences, QU Health, Qatar University, Doha, Qatar*

[7]*Biomedical Research Center, Qatar University, Doha, Qatar*

[8]*Department of Biomedical Science, College of Health Sciences, Member of QU Health, Qatar University, Doha, Qatar*

[9]*Primary Health Care Corporation, Doha, Qatar*

[10]*Ministry of Public Health, Doha, Qatar*

[11]*Department of Population Health Sciences, Weill Cornell Medicine, Cornell University, New York, New York, USA*

Table S1. Characteristics of the Craft and Manual Worker (CMW) Communities 2-10 and associations with anti-SARS-CoV-2 positivity.

Characteristics	Tested	Anti-SARS-CoV-2 positive		Univariable regression analysis	
	N (%)	N (%)	p-value	OR (95% CI)	p-value[*]
CMW Community 2					
Sex					
Men	385 (84.4)	270 (70.1)	<0.001	1.00	
Women	71 (15.6)	12 (16.9)		0.09 (0.04-0.17)	<0.001
Age (years)					
≤29	301 (66.0)	180 (59.8)	0.005	1.00	
30-39	143 (31.4)	99 (69.2)		1.51 (0.99-2.31)	0.055
40-49	12 (2.6)	3 (25.0)		0.22 (0.06-0.84)	0.027
Nationality					
All other nationalities	47 (10.3)	29 (61.7)	<0.001	1.00	
Kenyan	60 (13.2)	27 (45.0)		0.51 (0.23-1.11)	0.088
Nepalese	296 (64.9)	181 (61.1)		0.98 (0.52-1.84)	0.942
Bangladeshi	53 (11.6)	45 (84.9)		3.49 (1.34-9.07)	0.010
CMW Community 3					
Sex					
Men	291 (100.0)	244 (83.8)	--	--	--
Women	--		--	--	--
Age (years)					
≤29	81 (27.8)	69 (85.2)	0.837	1.00	
30-39	109 (37.5)	89 (81.7)		0.77 (0.35-1.69)	0.520
40-49	79 (27.1)	68 (86.1)		1.08 (0.44-2.60)	0.872
50+	22 (7.6)	18 (81.8)		0.78 (0.23-2.72)	0.700
Nationality					
All other nationalities	5 (1.7)	4 (80.0)	0.972	1.00	
Indian	99 (34.0)	83 (83.8)		1.30 (0.14-12.37)	0.821
Nepalese	187 (64.3)	157 (84)		1.31 (0.14-12.12)	0.813
CMW Community 4					
Sex					
Men	460 (86.1)	309 (67.2)	<0.001	1.00	
Women	74 (13.9)	21 (28.4)		0.19 (0.11-0.33)	<0.001
Age (years)					
≤29	257 (48.1)	168 (65.4)	<0.001	1.00	
30-39	200 (37.5)	108 (54.0)		0.62 (0.43-0.91)	0.014
40-49	38 (7.1)	17 (44.7)		0.43 (0.22-0.85)	0.016
50+	3 (0.6)	2 (66.7)		1.06 (0.09-11.85)	0.963
Missing	36 (6.7)	35 (97.2)		18.54 (2.50-137.60)	0.004
Nationality					
All other nationalities	49 (9.2)	11 (22.4)	<0.001	1.00	
Indian	62 (11.6)	33 (53.2)		3.93 (1.70-9.07)	0.001
Nepalese	155 (29.0)	89 (57.4)		4.66 (2.22-9.79)	<0.001
Bangladeshi	268 (50.2)	197 (73.5)		9.59 (4.65-19.77)	<0.001
CMW Community 5					
Sex					
Male	437 (98.7)	243 (55.6)	0.007	1.00	
Women	6 (1.4)	0 (0.0)		Omitted	--
Age (years)					
≤29	117 (26.4)	67 (57.3)	0.542	1.00	
30-39	193 (43.6)	106 (54.9)		0.91 (0.57-1.44)	0.687
40-49	108 (24.4)	54 (50.0)		0.75 (0.44-1.26)	0.275
50+	25 (5.6)	16 (64.0)		1.33 (0.54-3.25)	0.536

Characteristics	Tested	Anti-SARS-CoV-2 positive		Univariable regression analysis	
	N (%)	N (%)	p-value	OR (95% CI)	p-value[*]
Nationality					
Other nationalities	46 (10.4)	17 (37.0)	<0.001	1.00	
Indian	160 (36.1)	73 (45.6)		1.43 (0.73-2.81)	0.297
Sri Lankan	63 (14.2)	35 (55.6)		2.13 (0.98-4.64)	0.056
Nepalese	174 (39.3)	118 (67.8)		3.59 (1.82-7.08)	<0.001
CMW Community 6					
Sex					
Men	1465 (97.3)	939 (64.1)	<0.001	1.00	
Women	40 (2.7)	7 (17.5)		0.12 (0.05-0.27)	<0.001
Age (years)					
≤29	389 (25.8)	258 (66.3)	0.114	1.00	
30-39	644 (42.8)	387 (60.1)		0.76 (0.59-0.99)	0.045
40-49	374 (24.9)	244 (65.2)		0.95 (0.71-1.29)	0.753
50+	98 (6.5)	57 (58.2)		0.71 (0.45-1.11)	0.132
Nationality					
All other nationalities	173 (11.5)	63 (36.4)	<0.001	1.00	
Indian	649 (43.1)	408 (62.9)		2.96 (2.09-4.19)	<0.001
Nepalese	547 (36.3)	366 (66.9)		3.53 (2.47-5.05)	<0.001
Bangladeshi	136 (9.0)	109 (80.1)		7.05 (4.18-11.89)	<0.001
CMW Community 7					
Sex					
Men	188 (100.0)	122 (64.9)	--	--	--
Women	--	--	--	--	--
Age (years)					
≤29	91 (48.4)	61 (67.0)	0.703	1.00	
30-39	54 (28.7)	32 (59.3)		0.72 (0.36-1.44)	0.346
40-49	29 (15.4)	21 (72.4)		1.29 (0.51-3.25)	0.588
50+	11 (5.9)	6 (54.5)		0.59 (0.17-2.09)	0.414
Missing	3 (1.6)	2 (66.7)		0.98 (0.09-11.28)	0.989
Nationality					
All other nationalities	24 (12.8)	6 (25.0)	<0.001	1.00	
Indian	42 (22.3)	2 (4.8)		0.15 (2.76-81.64)	0.028
Nepalese	122 (64.9)	114 (93.4)		42.75 (13.28-137.66)	<0.001
CMW Community 8					
Sex					
Men	134 (96.4)	92 (68.7)	0.023	1.00	
Women	5 (3.6)	1 (20.0)		0.11 (0.01-1.05)	0.056
Age (years)					
≤29	26 (18.7)	16 (61.5)	0.330	1.00	
30-39	70 (50.4)	48 (68.6)		1.36 (0.53-3.48)	0.517
40-49	33 (23.7)	20 (60.6)		0.96 (0.33-2.76)	0.942
50+	10 (7.2)	9 (90.0)		5.63 (0.62-51.37)	0.126
Nationality					
All other nationalities	26 (18.7)	7 (26.9)	<0.001	1.00	
Indian	64 (46.0)	46 (71.9)		6.94 (2.49-19.31)	<0.001
Bangladeshi	25 (18.0)	18 (72.0)		6.98 (2.04-23.88)	0.002
Nepalese	24 (17.3)	22 (91.7)		29.86 (5.53-161.34)	<0.001
CMW Community 9					
Sex					
Men	164 (81.2)	125 (76.2)	<0.001	1.00	
Women	38 (18.8)	1 (2.6)		0.01 (0.00-0.06)	<0.001
Age (years)					

Characteristics	Tested	Anti-SARS-CoV-2 positive		Univariable regression analysis	
	N (%)	N (%)	p-value	OR (95% CI)	p-value[*]
≤29	30 (14.9)	16 (53.3)	0.201	1.00	
30-39	58 (28.7)	33 (56.9)		1.16 (0.48-2.80)	0.750
40-49	61 (30.2)	38 (62.3)		1.45 (0.60-3.50)	0.414
50+	53 (26.2)	39 (73.6)		2.44 (0.95-6.25)	0.064
Nationality					
All other nationalities	28 (13.9)	17 (60.7)	0.766	1.00	
Nepalese	85 (42.1)	51 (60.0)		0.97 (0.41-2.33)	0.947
Indian	89 (44.1)	58 (65.2)		1.21 (0.50-2.90)	0.668
CMW Community 10					
Sex					
Men	957 (100.0)	620 (64.8)	--	--	--
Women	--	--	--	--	--
Age (years)					
≤29	189 (19.7)	116 (61.4)	0.119	1.00	
30-39	392 (41.0)	243 (62.0)		1.03 (0.72-1.47)	0.887
40-49	273 (28.5)	190 (69.6)		1.44 (0.98-2.13)	0.067
50+	103 (10.8)	71 (68.9)		1.40 (0.84-2.32)	0.199
Nationality					
All other nationalities	69 (7.2)	42 (60.9)	0.061	1.00	
Bangladeshi	62 (6.5)	37 (59.7)		0.95 (0.47-1.92)	0.889
Nepalese	389 (40.6)	238 (61.2)		1.01 (0.60-1.71)	0.961
Indian	437 (45.7)	303 (69.3)		1.45 (0.86-2.46)	0.162

CI, confidence interval; OR, odds ratio.

[*]Covariates with p-value ≤0.05 in the univariable analysis were considered as showing evidence for an association with anti-SARS-CoV-2 positivity.

Table S2. Characteristics of the craft and manual workers (CMWs) and associations with SARS-CoV-2 polymerase chain reaction (PCR) positivity.

Characteristics	Tested	SARS-CoV-2 positive		Univariable regression analysis		Multivariable regression analysis	
	N (%)	N (%)	p-value	OR (95% CI)	p-value[*]	OR (95% CI)	p-value[†]
Sex							
Men	1,844 (91.5)	100 (5.4)	0.395	1.00		--	--
Women	172 (8.5)	12 (7.0)		1.31 (0.70-2.43)	0.396	--	--
Age (years)							
<29	785 (38.9)	39 (5.0)	0.486	1.00		--	--
30-39	787 (39.0)	44 (5.6)		1.13 (0.73-1.76)	0.581	--	--
40-49	351 (17.4)	25 (7.1)		1.47 (0.87-2.46)	0.148	--	--
50+	93 (4.6)	4 (4.3)		0.86 (0.30-2.46)	0.778	--	--
Nationality							
All other nationalities[‡]	149 (7.4)	15 (10.1)	0.002	1.00		1.00	
Nepalese	918 (45.5)	46 (5.0)		0.47 (0.26-0.87)	0.016	0.43 (0.22-0.82)	0.011
Bangladeshi	265 (13.1)	9 (3.4)		0.31 (0.13-0.74)	0.008	0.44 (0.17-1.12)	0.085
Filipino	112 (5.6)	4 (3.6)		0.33 (0.11-1.03)	0.055	0.49 (0.15-1.63)	0.242
Indian	572 (28.4)	38 (6.6)		0.64 (0.34-1.19)	0.157	0.69 (0.33-1.45)	0.329
CMW community							
CMW Community 6[§]	832 (41.3)	42 (5.0)	<0.001	1.00[§]		1.00[§]	
CMW Community 1	92 (4.6)	0 (0.0)		omitted	--	omitted	--
CMW Community 8	39 (1.9)	0 (0.0)		omitted	--	omitted	--
CMW Community 4	363 (18)	10 (2.8)		0.53 (0.26-1.07)	0.078	0.63 (0.30-1.36)	0.240
CMW Community 2	424 (21.0)	32 (7.5)		1.54 (0.95-2.47)	0.077	1.60 (0.91-2.79)	0.099
CMW Community 3	266 (13.2)	28 (10.5)		2.21 (1.34-3.65)	0.002	2.44 (1.45-4.08)	0.001

CI, confidence interval; OR, odds ratio.

[*]Covariates with p-value ≤0.2 in the univariable analysis were included in the multivariable analysis.

[†]Covariates with p-value ≤0.05 in the multivariable analysis were considered predictors of SARS-CoV-2 antibody positivity.

[‡]Includes all other nationalities that contributed <5% of the sample in each community.

[§]CMW Community 6 which had the largest sample size was chosen as a reference.

Table S3. Characteristics of anti-SARS-CoV-2 negative persons who had tested positive for SARS-CoV-2 infection using polymerase chain reaction (PCR) at some point prior to conduct of this study.

Person number	Age	Sex	PCR test date	Ct value	Antibody test date	Symptoms
1	36	Man	13 May	35.5	02 August	Not recorded
2	28	Man	04 May	34.1	03 August	Not recorded
3	25	Man	04 May	35.1	03 August	Not recorded
4	26	Man	28 April	22.0	08 August	Not recorded
5	40	Man	06 June	35.5	23 July	Not recorded
6	33	Man	09 June	23.7	27 July	Not recorded
7	27	Man	12 May	35.8	28 July	Not recorded
8	34	Man	13 May	Unknown	28 July	Not recorded
9	31	Man	23 May	27.4	20 August	Not recorded
10	31	Man	06 May	19.6	20 August	Not recorded
11	33	Man	07 May	18.0	20 August	Not recorded
12	39	Man	11 July	32.9	20 August	Not recorded
13	44	Man	05 August	35.2	20 August	Asymptomatic
14	34	Man	08 May	32.0	24 August	Not recorded

Figure S1. Herd immunity and heterogeneity in risk of exposure to the infection. SARS-CoV-2 active infection prevalence (A) and attack rate (proportion of the population that has ever been infected) (B) in a community where the risk of exposure is homogeneous versus in a community where the risk of exposure is heterogeneous. In both of these scenarios, the basic reproduction number R_0 was assumed equal to 3 [1, 2]. These simulations were generated using a classic age-structured susceptible-exposed-infectious-recovered "SEIR" mathematical model [3]. Heterogeneity in the second modeled scenario was introduced through variable exposure risk by age.

Figure S2. Measures of SARS-CoV-2 A) antibody positivity, B) PCR positivity, C) infection positivity (antibody and/or PCR positive), and D) diagnosis rate, among *only men* craft and manual workers (CMW) across the CMW communities.

References

1.   He, W., G.Y. Yi, and Y. Zhu, *Estimation of the basic reproduction number, average incubation time, asymptomatic infection rate, and case fatality rate for COVID-19: Meta-analysis and sensitivity analysis.* J Med Virol, 2020.
2.   MIDAS Online COVID-19 Portal, *COVID-19 parameter estimates: basic reproduction number. Available from:* https://github.com/midas-network/COVID-19/tree/master/parameter_estimates/2019_novel_coronavirus*. Accessed on: MAy 19, 2020.* 2020.
3.   Abu-Raddad, L.J., et al., *Characterizing the Qatar advanced-phase SARS-CoV-2 epidemic.* medRxiv, 2020: p. 2020.07.16.20155317v2.