Title
The Relationship Between Cannabis Legalization, Suicide and Mental Health Remains
Unchanged After Recreational Access

Jeffrey A. Singer, MD[1], Jacob James Rich, MA[2], Michael Schemenaur, MA[3], and Robert
Capodilupo, MPhil[4]

1. Health Policy Studies, Cato Institute, Washington, DC

2. Drug Policy Project, Reason Foundation, Washington, DC

3. Department of Economics, Eller College of Management, University of Arizona, Tucson, AZ

4. Yale Law School, New Haven, CT

Abstract:
Background. We extend the model introduced by Anderson et al. [M.D. Anderson, D.I. Rees, J.J.
Sabia, *American Journal of Public Health* 104, 2369-2376] to evaluate the public mental health
implications of new developments in marijuana policy, such as recreational marijuana access and
additional years of data on suicide mortality.

Methods. We obtained state-level suicide data from the National Vital Statistics System's
Mortality Detail Files for 1999-2018. We used panel regression analysis to examine the
association between suicides per 100 000 population and both medical and recreational
marijuana access. Three specifications of the equation were employed as a robustness test.

Results. After adjusting for economic conditions, alcohol taxes and consumption, time effects,
state fixed effects, and state-specific linear time trends, the legalization of recreational marijuana
was associated with a 2.9% (95% confidence interval [CI] = -5.6, -.03) reduction in the suicide
rate for the most rigorous specification, but didn't pass the robustness check at the $P < .05$ level
for the other two specifications. However, recreational marijuana legalization was associated
with a 5.4% reduction (95% CI = -9.4%, -1.4%) in suicide rates for males in the 40 to 49 age
group, passing the robustness check for all specifications.

Conclusion. Suicide rates among men in the 40 to 49 age group fell after recreational marijuana
legalization compared with those in states that did not legalize, which is consistent with the
results found by Anderson et al. for medical marijuana when that same population was younger.
There was no consistent relationship between medical marijuana legalization and suicide rates
for any population. Additionally, mental health indicators at the state level were unaffected by
either recreational or medical marijuana legalization. We confirm the findings of Anderson et al.
(2014), observing no consistent relationship between marijuana legalization and observable
mental health outcomes. However, suicide rates increase at lower rates on average relative to
other states following recreational marijuana legalization.

**The Relationship Between Cannabis Legalization, Suicide and Mental Health Remains
Unchanged After Recreational Access**

As of September 1 2020, 11 states and the District of Columbia have voted to legalize marijuana
for recreational use, while medical marijuana is available in 33 states. A November 2019 survey
from the Pew Research Center found that 67 percent of adults support national legalization of
marijuana, including 63 percent of the Baby Boomer generation, 76 percent of Millennials, and
65 percent of Gen X. (Daniller, 2019) In October 2018 Canada became the first G7 country to
federally legalize marijuana for recreational use, and the same month Mexico's Supreme Court
ruled marijuana prohibition unconstitutional, setting the stage for legalization in that country.
(Hudak, 2020) In the not-too-distant future marijuana use in the United States might be federally
decriminalized and the matter devolved to the states. Legislation to that effect was introduced in
the 116[th] Congress. (Nadler, 2020) These developments reflect the increasing acceptance of the
recreational use of marijuana, which trends suggest will further increase in days ahead.

Policy makers considering marijuana legalization should consider the public health
implications of increasing access to a psychoactive substance that could lead to substance use
disorder. In a 2009 Norwegian study, Pedersen concluded "exposure to marijuana by itself does
not lead to depression but that it may be associated with later suicidal thoughts and attempts."
(Pederson, 2008) A 2014 systematic review and meta-analysis of longitudinal studies by
Canadian researchers found an increased risk for developing depressive disorders may be
associated with heavy marijuana use. (Lev-Ran, Roerecke, George, McKenzie, & Rehm, 2014)
Agrawal et al., in a 2017 retrospective study, found that the correlation between marijuana use,
suicidal thoughts, and major depressive disorders in discordant twins was not solely attributable
to predisposing factors. (Agrawal, et al., 2017) This observation confirmed similar findings
reported by researchers at the same institution ten years earlier. (Lynskey, et al., 2004) A 2015
study using community-based samples from the Australian Twin Registry by Delforterie et al.
found a "modest association" of marijuana use with suicidal thoughts and attempts, concluding
that the association of suicidal thoughts and behaviour with marijuana use disorder requires
further study. (Delforterie, et al., 2015) In a 2020 study in *JAMA Network Open,* Gorfinkel et al.
found that people reporting depression tend to use marijuana at a higher rate. (Gorfinkel, Stohl,
& Hasin, 2020)

However, much of the literature finds no relationship between marijuana use and mental

health outcomes. In particular a 2014 regression analysis of state-level data from the National

Vital Statistics System (NVSS) Mortality Detail Files from 1990-2007 by Anderson et al. found

no general association between medical marijuana legalization and suicide rates, but found

10.8% and 9.4% reductions in the suicide rates in males age 20 to 29 and 30 to 39 in the U.S.,

respectively. (Anderson, Rees, & Sabia, 2014) Additionally, a 2018 Canadian study found no

correlation between suicidal behaviour and heavy marijuana use in male or female patients with

psychiatric disorders. (Naji, et al., 2018) With the mental outcomes of marijuana use contested

by much of the literature, policymakers would benefit from research investigating the population

effects on mental health after marijuana legalization.

Marijuana use is tracked by the Substance Abuse and Mental Health Services

Administration (SAMHSA) with the National Survey on Drug Use and Health (NSDUH). As of

2018, about 43.5 million people (or 15.9 percent of the U.S. population above the age of 12)

consumed marijuana at least once that year—a historic high. Of those marijuana users, 3.1

million used marijuana for the first time, which has been a relatively consistent finding every

year since Colorado became the first state to legalize marijuana in 2014. That year 12.9 percent

of the U.S. population used marijuana in the past year and the recent trend of increase is

predicted to continue into the near future. (SAMHSA, 2015) Marijuana use reported in the past

month has also steadily increased in the U.S. since 2007, after a subtler increase starting after an

all-time low in 1992. (Golub, Johnson, & Dunlap, 2005)

Reports of major depressive episodes have increased since 2010, while both serious

thoughts of suicide and severe mental illness have increased each year since the inception of the

survey questions in 2008. (SAMHSA, 2019) Suicide rates in the U.S. have been steadily

increasing since an all-time low in 1999 and were at an all-time high in 2018. Respondents who

reported past month marijuana use in 2017 were much more likely to experience major

depression (2.29 times), suicidal thoughts (3.2 times), and slightly more prone to any/severe

mental illness (1.16/1.64 times) than nonusers. (SAMHSA, 2018) This observation has garnered

much interest in the relationship between marijuana use and possible mental health deterioration.

But with increasing evidence of marijuana's medicinal benefits, and with its growing popularity

as a recreational drug, it is necessary to definitively answer the questions surrounding heavy

marijuana use and risk of depression and suicide.

In the years since the Anderson et al. review of data from 1990-2007, the advent of

recreational marijuana laws has provided unprecedented access to marijuana. Significant gaps

remain in the knowledge of the potential harms and benefits of marijuana use, and the effects

these laws have on various facets of mental illness are a fertile topic for new research. (National

Academies of Sciences, Engineering, and Medicine, 2017)

Methods

Employing most of the same techniques used by Anderson et al. (2014), we use population-

weighted ordinary least squares (OLS) regressions for the observed groups to conduct a state-

level panel analysis for various time periods of available data to estimate the effect recreational

and medical marijuana laws have had on various proxies for mental health: suicide rates per 100

000 people for the general population, males, females, and both sexes separately for the age

groups 15-19, 20-29, 30-39, 40-49, 50-59, and 60 and above (1999-2018); the percentages of the

population that reported "serious thoughts of suicide in the past year," "any mental illness in the

past year," and "serious mental illness in the past year" for the age groups 18-25, 18 and up, and

26 and up (2009-2018); and the percentages of the population that reported they "had at least one

major depressive episode in the past year" for the age groups 12-17, 18-25, 18 and up, and 26

and up (2006-2018).

Similar to Anderson et al. (2014), we control for observable state-level characteristics the

literature considers risk factors for suicide: unemployment rates, GDP per capita, beer taxes, and

ethanol consumption per capita for the population age 21 and older. Also following Anderson et

al. (2014), we attempt to control for unobserved factors with time effects, fixed effects, and state-

specific linear time trends. Since previous studies suggest stricter alcohol policies can reduce

suicides, (Markowitz, Chatterji, & Kaestner, 2003; Birckmayer & Hemenway, 1999; Jones,

Pieper, & Robertson, 1992; Robertson, 2014) Anderson et al. (2014) controlled for whether a

0.08 blood-alcohol-content (BAC) law was in effect, whether a zero-tolerance drunk-driving law

was in effect, and for the beer excise tax. We removed the controls for BAC laws because every

state had implemented a 0.08 BAC limit for our entire period, except for Utah, which

implemented a .05 limit December 25 2018. Additionally, we removed indicators for zero-

tolerance drunk-driving laws because every state had implemented such restrictions by 2001.

(Carpenter, Heavy alcohol use and youth suicide: Evidence from tougher drunk driving laws,

2004; Carpenter, How do Zero Tolerance Drunk Driving Laws work?, 2004) We argue those

variables ultimately attempted to capture state government interventions that reduce alcohol use,

which is why we added a measurement for per capita ethanol consumption.

To test the robustness of our findings, we follow Anderson et al. (2014) and employ a

sensitivity analysis involving three separate specifications for each dependent variable: first only

adjusting for state and year effects; then adjusting for state effects, year effects, and covariates;

and finally adjusting for state effects, year effects, covariates, and state-specific linear time

trends in the most rigorous model. Robust standard errors were employed and clustered at the

state level for all specifications. Additionally, we took the natural log transform of all dependent

variables, and because the log of zero is undefined, we dropped all observations of zero suicides.

If the coefficients for the marijuana law indicators remain relatively stable and significant for all

regression specifications, there is evidence that a predictive relationship exists between

marijuana legalization and the mental health variables of interest. (Miron & Tetelbaum, 2009;

Thornton, 2011)

Suicide mortality (X60-X84, Y87.0, and U03 intentional self-harm) data and population

estimates are sourced from the Centers for Disease Control and Prevention (CDC) Wide-ranging

Online Data for Epidemiologic Research (WONDER) NVSS database, reports of mental illness

from the SAMHSA NSDUH, unemployment rates and GDP per capita from the Bureau of Labor

Statistics, ethanol consumption from the National Institute on Alcohol Abuse and Alcoholism,

and beer excise tax figures from the Tax Policy Center at the Urban Institute and Brookings

Institution. Marijuana policy indicators were calculated referencing reports by the Marijuana

Policy Project and the National Organization for the Reform of Marijuana Laws (NORML).

These indicators consider the dates laws were enacted and for the first year assume a value

between 0 and 1 equal to the proportion of days the policies were effective. The year recreational

marijuana laws became effective were used to graph unadjusted suicides per 100 000 population

over the year of law change to compare rates with states that did not legalize recreational

marijuana anytime during the period.

Results

Table 1 presents the dates that various marijuana laws went into effect. Average suicide rates per

100 000 population over the year of law change are displayed in Figure 1, graphing average

suicides rates for the five jurisdictions that have legalized recreational marijuana for at least three

years (Alaska, Colorado, District of Columbia, Oregon, and Washington) next to the 40 states

that didn't legalize marijuana for the entire period of interest centred at 2014, the average year of

implementation for the 5 legalizing states. Reviewing the slopes of both groups before policies

went into effect, states that would eventually legalize recreational marijuana had a higher

average annual increase in suicide rates (0.708 per 100 000 people) relative to states that never

legalized (0.374 per 100 000 people). However, after legalization policies went into effect, states

permitting access to recreational marijuana reduced their average annual increase to a level

(0.339 per 100 000) below both groups before legalization and the non-legalizing group after

legalization (0.411 per 100 000).

137   Table 2 presents the regression results measuring the effect marijuana legalization had on

the mental health outcomes. The only relationship that passed each of the robustness check

specifications was between suicide rates for males in the 40 to 49 age group and recreational

marijuana legalization, where legalization is followed by a 5.4% reduction (95% confidence

interval [CI] = -9.4%, -1.4%) in suicides per capita in the most rigorous model. Interestingly,

Anderson et al. found that medical marijuana legalization led to 10.8% (95% CI =$\square$−17.1%,

−3.7%) and 9.4% (95% CI$\square$=$\square$−16.1%, −2.4%) drops in the 20 to 29 and 30 to 39 age groups,

respectively, many of whom would now be in the 40 to 49 age group. This population is now

older because the youngest data from the previous study were published 13 years ago.

146   The only other relationship that was significant for the most rigorous specification was a

2.9% reduction (95% CI = -5.6, -.03) in the total suicide rate following recreational marijuana

legalization. The relationship between recreational marijuana legalization was also weakly

associated reductions in the total suicide rate for the other two specifications, but the estimates

didn't quite achieve significance at the standard $P < .05$ level.

Discussion

Despite suicide rates reaching unprecedented levels across the country, suicides in states with

recreational marijuana access have increased at slower rates after their policies went into effect

relative to their rates before legalization and relative to other states across the country. Our

analysis finds strong evidence that recreational marijuana laws reduce suicide rates for middle

age men and weak evidence recreational marijuana laws reduce suicide rates for the population

as a whole. In line with the conclusions of Anderson et al. (2014), we find that marijuana laws in

general don't have consistent effects on suicide rates or reports of mental illness, despite

correlations between marijuana use and reports of depression.

While 24% of past year marijuana users report depression, 25.1% of people who report

depression admit to using marijuana in the past year. (SAMHSA, 2018) The correlations

concerning policy makers may be due to simultaneity, or "reverse causality," which could be the

result of self-medication. Regardless, there is no evidence that the liberalization of marijuana in

the U.S. is leading to reductions in public mental health.

Limitations

Unfortunately, due to the limitations of public SAMHSA data because of privacy

concerns, it is not possible to utilize geographical controls for individual data. Our analysis

suffers from the same limitations that plague observational studies in general. (Koopman &

Lynch, 1999) To improve on future public research independent of the government's efforts,

SAMHSA should add state-level identifiers to its public individual-response data for all years

after 2002. Since NSDUH state-level samples are too large to identify the individuals who took

the survey, privacy will not be a concern if state of residence information is released to the public

for each individual response. Permitting state-level identifiers will allow for the employment of

accepted metric methods that use geographic identifiers to control for environmental and

reporting differences. The consequences of drug use in general are of the utmost interests to

public health officials, and the literature would be best advanced by allowing public access to

currently restricted data.

There have also been legitimate challenges to the accuracy of the NSDUH and suicide

rates. Critics claim that because the NSDUH requires the surveyed individuals to have a

residence, much of the population (especially homeless people) are not represented by the

estimates. (Hedden, et al., 2012) Additionally, it is widely believed suicide rates are

underreported. (Warshauer & Monk, 1978) These are legitimate criticisms, but the regression

analyses are much more concerned with the changes in rates, as opposed to the absolute accuracy

of the rates themselves. Additionally, there's no reason to believe states with recreational

marijuana access would be more likely to underreport suicides. Since consistent methods for

collecting these data were employed for the entire period of interest for all dependent variables,

we are confident that the variable regression coefficients reflect the relationships between

evolving trends of substance availability and the mental health identifiers of interest.

Public Health Implications and Conclusion

The use of any drug, psychoactive or otherwise, entails certain risks along with any benefits.

Marijuana is no exception. Critics of marijuana legalization point to studies showing correlations

between heavy cannabis use and suicide, depression, and mental health disorders. However, such

studies demonstrating correlation have yet to confirm causation, which should be determined by

a model's ability to predict. (Arrow, 1963) Although those reporting depression to SAMHSA

have increasingly used marijuana since states began increasing access to regulated cannabis,

(Gorfinkel, Stohl, & Hasin, 2020) we observe no evidence that these laws have predictive

relationships with reports of any mental illness. Additionally, recreational marijuana access is

followed by reductions in suicide for some age groups. Anderson et al. (2014) had found similar

relationships between medical marijuana access and suicide rates, and the lack of an association

between mental health and medical marijuana access in our study is likely because recreational

marijuana access is a superior substitute requiring fewer barriers to purchase cannabis.

We propose as medicinal and recreational use of marijuana becomes more widespread

and mainstream, concerns about the correlation between marijuana use and depression shouldn't

interfere with state or federal efforts to decriminalize or legalize marijuana. In fact, legalization

will have the salutary effect of allowing more rigorous research—now inhibited by federal

prohibition—into the further benefits, as well as any other potential harms, from the long-term

use of marijuana, and promote safer use.

References

Agrawal, A., Nelson, E. C., Bucholz, K. K., Tillman, R., Grucza, R. A., Statham, D. J., . . .

Lynskey, M. T. (2017, September 1). Major depressive disorder, suicidal thoughts and

218   behaviours, and cannabis involvement in discordant twins: a retrospective cohort study.
219   *Lancet Psychiatry, 4*(9), 704-714.
220 Anderson, M. D., Rees, D. I., & Sabia, J. J. (2014, December). Medical Marijuana Laws and
221   Suicides by Gender and Age. *American Journal of Public Health, 104*(12), 2369-2376.
222 Arrow, K. J. (1963, December). Uncertainty and the Welfare Economics of Medical Care. *The*
223   *American Economic Review, 53*(5), 941-973.
224 Birckmayer, J., & Hemenway, D. (1999). Minimum-age drinking laws and youth suicide, 1970-
225   1990. *American Journal of Public Health, 89*(9), 1365-1368.
226 Bureau of Labor Statistics. (2020). Databases, Tables & Calculators by Subject. Washington,
227   District of Columbia, United States.
228 Carpenter, C. (2004, May). Heavy alcohol use and youth suicide: Evidence from tougher drunk
229   driving laws. *Journal of Policy Analysis and Management, 23*(4), 831-842.
230 Carpenter, C. (2004, January). How do Zero Tolerance Drunk Driving Laws work? *Journal of*
231   *Health Economics, 23*(1), 61-83.
232 Centers for Disease Control and Prevention. (2020). Wide-ranging Online Data for
233   Epidemiologic Research (WONDER). Atlanta, Georgia, United States.
234 Daniller, A. (2019, November 14). Two-thirds of Americans support marijuana legalization.
235   *News In The Numbers*.
236 Delforterie, M. J., Lynskey, M. T., Huizink, A. C., Creemers, H. E., Grant, J. D., Few, L. R., . . .
237   Agrawal, A. (2015, May 1). The relationship between cannabis involvement and suicidal
238   thoughts and behaviors. *Drug and Alcohol Dependence, 150*, 98-104.
239 Golub, A., Johnson, B. D., & Dunlap, E. (2005). The Growth in Marijuana Use Among
240   American Youths During the 1990s and the Extent of Blunt Smoking. *Journal of*
241   *Ethnicity in Substance Abuse, 4*(3-4), 121.
242 Gorfinkel, L. R., Stohl, M., & Hasin, D. (2020, August 18). Association of Depression With
243   Past-Month Cannabis Use Among US Adults Aged 20 to 59 Years, 2005 to 2016. *JAMA*
244   *Network Open, 3*(8).
245 Hedden, S., Gfroerer, J., Barker, P., Smith, S., Pemberton, M. R., Saavedra, L. M., . . . Novak, S.
246   P. (2012). *Comparison of NSDUH Mental Health Data and Methods with Other Data*
247   *Sources.* Rockville, MD: Department of Health and Human Services.
248 Hudak, J. (2020). *Marijuana: A Short History.* Washington, D.C.: Brookings Institution.
249 Jones, N. E., Pieper, C. F., & Robertson, L. S. (1992, January). The effect of legal drinking age
250   on fatal injuries of adolescents and young adults. *American Journal of Public Health,*
251   *82*(1), 112-115.
252 Koopman, J. S., & Lynch, J. W. (1999, October 7). Individual causal models and population
253   system models in epidemiology. *American Journal of Public Health, 89*(8), 1170-1174.
254 Lev-Ran, S., Roerecke, M., George, T. P., McKenzie, K., & Rehm, J. (2014, March). The
255   association between cannabis use and depression: a systematic review and meta-analysis
256   of longitudinal studies. *Psychological Medicine, 44*(4), 797-810.
257 Lynskey, M. R., Glowinski, A. L., Todorov, A. A., Bucholz, K. K., Madden, P. A., Nelson, E.
258   C., . . . Heath, A. C. (2004). Major Depressive Disorder, Suicidal Ideation, and Suicide
259   Attempt inTwins Discordant for Cannabis Dependence and Early-Onset Cannabis Use.
260   *JAMA Psychiatry, 61*(10), 1016-1032.
261 Markowitz, S., Chatterji, P., & Kaestner, R. (2003, March). Estimating the impact of alcohol
262   policies on youth suicides. *Journal of Mental Health Policy Economics, 6*(1), 37-46.

Miron, J. A., & Tetelbaum, E. (2009, April). Does the Minimum Legal Drinking Age Save
Lives? *Economic Inquiry, 47*(2), 317-336.
Nadler, J. (2020). H.R.3884 - Marijuana Opportunity Reinvestment and Expungement Act of
2019. Retrieved from https://www.congress.gov/bill/116th-congress/house-bill/3884
Naji, L., Rosic, T., Dennis, B., Bhatt, M., Sanger, N., Hudson, J., . . . Samaan, Z. (2018, June
11). The association between cannabis use and suicidal behavior in patients with
psychiatric disorders: an analysis of sex differences. *Biology of Sex Differences, 9*(22).
National Academies of Sciences, Engineering, and Medicine. (2017). *The Health Effects of
Cannabis and Cannabinoids: The Current State of Evidence and Recommendations for
Research.* Washington: National Academies Press.
National Institute on Alcohol Abuse and Alcoholism. (2020, April). Surveillance Report #115.
Washington, District of Columbia, United States.
Pederson, W. (2008, November). Does cannabis use lead to depression and suicidal behaviours?
A population-based longitudinal study. *Acta Psychiatrica Scandinavica, 118*(5), 395-403.
Robertson, L. S. (2014, December). Science and public health principles used to reduce road
deaths. *American Journal of Public Health, 104*(12), 2256-2258.
SAMHSA. (2015). *2013-2014 NSDUH State Estimates Of Substance Use And Mental Disorders.*
Rockville, MD.
SAMHSA. (2018). National Survey on Drug Use and Health (NSDUH-2017). Rockville,
Maryland, United States.
SAMHSA. (2019). *Key Substance Use and Mental Health Indicators in the United States:
Results from the 2018 National Survey on Drug Use and Health.* Rockville, MD:
Substance Abuse and Mental Health Services.
Tax Policy Center. (2020). State Alcohol Excise Taxes. Washington, District of Columbia,
United States.
Thornton, J. A. (2011, October 14). Does More Medical Care Improve Population Health? New
Evidence for an Old Controversy. *Applied Economics, 34*(1), 3325-3336.
Warshauer, M., & Monk, M. (1978, April). Problems in suicide statistics for whites and blacks.
American Journal of Public Health, 68(4), 383-388.

292
293

304 **Tables and Figures**

306 Table 1 — Effective Dates of Marijuana Legalization

State	FIPS	Recreational Effective	Medical Effective	Decriminalized Effective
Alabama	1			
Alaska	2	2/24/15	1999	1975-1990, 8/29/2003-6/14/2006, 2/24/2015
Arizona	4		3/14/11	
Arkansas	5		4/8/17	
California	6	11/9/16		1976
Colorado	8	1/3/13	6/1/01	1975
Connecticut	9		3/5/12	6/6/11
Delaware	10		5/13/11	12/18/15
District of Columbia	11	2/26/15	3/29/13	7/17/14
Florida	12		1/3/17	
Georgia	13			
Hawaii	15	1/11/20	8/8/17	1/11/20
Idaho	16			
Illinois	17	1/1/20	1/1/14	
Indiana	18			
Iowa	19			
Kansas	20			
Kentucky	21			
Louisiana	22		8/6/19	
Maine	23	12/17/16	5/10/10	5/1/76
Maryland	24		1/1/17	10/1/14
Massachusetts	25	12/15/16	1/1/13	1/2/09
Michigan	26	12/6/19	10/25/18	
Minnesota	27		7/1/15	1976
Mississippi	28			
Missouri	29		12/6/18	1/1/17
Montana	30		11/2/04	
Nebraska	31			1/1/79
Nevada	32	1/1/17	10/1/01	10/1/01
New Hampshire	33		7/23/13	9/16/17
New Jersey	34		1/1/11	
New Mexico	35		7/1/07	7/1/19
New York	36		7/7/14	7/29/77
North Carolina	37			7/1/77
North Dakota	38		1/28/17	5/2/19

Ohio	39		9/6/16	8/22/75
Oklahoma	40		8/30/18	
Oregon	41	7/1/15	12/3/98	10/1/73
Pennsylvania	42		5/17/16	
Rhode Island	44		1/3/06	4/1/13
South Carolina	45			
South Dakota	46			
Tennessee	47			
Texas	48			
Utah	49			
Vermont	50	7/1/18	5/19/04	7/1/13
Virginia	51			7/1/20
Washington	53	12/2/12	1998	12/2/12
West Virginia	54			
Wisconsin	55			
Wyoming	56			

308   Figure 1

Year of Law Change	Centred Years	Legal Slope	Illegal Slope
-3 to -1	2011-2013	0.7084	0.3744
0 to 3	2014-2017	0.3397	0.4116

Table 2 — Regression Results

Dependent Variable	Policies	Coefficients Controls, TE, FE, ST	[95% Conf.	Interval]	Coefficients Controls, TE, FE	[95% Conf.	Interval]	Coefficients TE, FE	[95% Conf.	Interval]
Total Suicide Rate [a]	Recreational Marijuana	-0.0299 *	-0.0567	-0.0030	-0.0444	-0.0918	0.0030	-0.0340	-0.0852	0.0171
	Medical Marijuana	-0.0176	-0.0696	0.0344	0.0134	-0.0182	0.0450	0.0272	-0.0045	0.0590
Male Suicide Rate [a]	Recreational Marijuana	0.0749	-0.1508	0.3006	0.0783	-0.0963	0.2529	0.0838	-0.0917	0.2593
	Medical Marijuana	-0.0587	-0.1683	0.0510	-0.0073	-0.0761	0.0614	-0.0070	-0.0729	0.0590
Female Suicide Rate [a]	Recreational Marijuana	-0.0373	-0.0938	0.0192	-0.0561	-0.1330	0.0207	-0.0432	-0.1275	0.0410
	Medical Marijuana	0.0233	-0.0500	0.0967	0.0546 *	0.0091	0.1001	0.0766 **	0.0278	0.1253
Suicide Rate Male 15 to 19 [b]	Recreational Marijuana	-0.0843	-0.2344	0.0657	0.0319	-0.1021	0.1660	0.0387	-0.0756	0.1530
	Medical Marijuana	0.0058	-0.0730	0.0846	-0.0263	-0.1093	0.0566	-0.0242	-0.1050	0.0566
Suicide Rate Male 20 to 29 [a]	Recreational Marijuana	-0.0131	-0.0645	0.0382	-0.0018	-0.0589	0.0553	0.0142	-0.0368	0.0652
	Medical Marijuana	-0.0084	-0.0755	0.0587	-0.0136	-0.0805	0.0534	-0.0200	-0.0852	0.0451
Suicide Rate Male 30 to 39 [a]	Recreational Marijuana	-0.0026	-0.0422	0.0370	-0.0184	-0.0822	0.0454	-0.0214	-0.0834	0.0406
	Medical Marijuana	-0.0213	-0.0739	0.0313	-0.0263	-0.0674	0.0148	-0.0252	-0.0627	0.0124
Suicide Rate Male 40 to 49 [a]	Recreational Marijuana	-0.0543 **	-0.0942	-0.0145	-0.0873 **	-0.1400	-0.0346	-0.0824 **	-0.1402	-0.0245
	Medical Marijuana	-0.0339	-0.1186	0.0507	0.0110	-0.0406	0.0626	0.0307	-0.0231	0.0845
Suicide Rate Male 50 to 59 [a]	Recreational Marijuana	-0.0259	-0.0629	0.0110	-0.0973 **	-0.1535	-0.0412	-0.0827 *	-0.1567	-0.0086
	Medical Marijuana	-0.0357	-0.0840	0.0126	0.0195	-0.0213	0.0603	0.0439	-0.0059	0.0938
Suicide Rate Male 60 and Up [a]	Recreational Marijuana	0.0023	-0.0555	0.0602	-0.0545	-0.1126	0.0036	-0.0469	-0.1187	0.0250
	Medical Marijuana	-0.0216	-0.0826	0.0394	0.0219	-0.0133	0.0570	0.0340	-0.0043	0.0723
Suicide Rate Female 15 to 19 [c]	Recreational Marijuana	-0.1013	-0.2976	0.0950	-0.1706	-0.3420	0.0007	-0.1431	-0.3040	0.0178
	Medical Marijuana	-0.1753	-0.3846	0.0340	-0.0761	-0.2200	0.0678	-0.0839	-0.2142	0.0465
Suicide Rate Female 20 to 29 [d]	Recreational Marijuana	0.0057	-0.0920	0.1034	-0.0470	-0.1198	0.0257	-0.0281	-0.1033	0.0471
	Medical Marijuana	-0.0484	-0.1596	0.0627	-0.0167	-0.0909	0.0576	0.0056	-0.0674	0.0785
Suicide Rate Female 30 to 39 [e]	Recreational Marijuana	0.0121	-0.0686	0.0928	-0.0354	-0.1182	0.0474	-0.0185	-0.0950	0.0579
	Medical Marijuana	0.0900	-0.0005	0.1806	0.1128 **	0.0450	0.1807	0.1276 **	0.0554	0.1998

Suicide Rate Female 40 to 49 [f]	Recreational Marijuana	-0.0867	-0.1762	0.0028	-0.1547	**	-0.2558	-0.0537	-0.1234	*	-0.2249	-0.0219
	Medical Marijuana	0.0001	-0.0854	0.0856	0.0620		-0.0209	0.1450	0.0919		-0.0021	0.1859
Suicide Rate Female 50 to 59 [b]	Recreational Marijuana	-0.0532	-0.1900	0.0836	-0.0548		-0.2322	0.1227	-0.0767		-0.2684	0.1151
	Medical Marijuana	0.0750	-0.0257	0.1757	0.0808		-0.0079	0.1695	0.1019	*	0.0098	0.1940
Suicide Rate Female 60 and Up [g]	Recreational Marijuana	-0.0009	-0.1391	0.1373	0.0031		-0.1335	0.1396	-0.0051		-0.1464	0.1363
	Medical Marijuana	0.0388	-0.0764	0.1540	0.0396		-0.0363	0.1156	0.0580		-0.0157	0.1317
Reports of Major Depression 12 to 17 [h]	Recreational Marijuana	-0.0401	-0.0864	0.0063	0.0171		-0.0241	0.0583	0.0278		-0.0048	0.0604
	Medical Marijuana	0.0107	-0.0443	0.0657	0.0073		-0.0263	0.0409	0.0141		-0.0193	0.0476
Reports of Major Depression 18 to 25 [h]	Recreational Marijuana	0.0614	-0.0103	0.1331	0.0908	**	0.0515	0.1300	0.0927	**	0.0650	0.1205
	Medical Marijuana	0.0004	-0.0461	0.0468	0.0125		-0.0229	0.0478	0.0192		-0.0126	0.0510
Reports of Major Depression 26 and Up [h]	Recreational Marijuana	-0.0508	-0.1227	0.0210	0.0115		-0.0329	0.0559	0.0138		-0.0236	0.0512
	Medical Marijuana	-0.0084	-0.0409	0.0241	-0.0078		-0.0437	0.0281	-0.0027		-0.0348	0.0294
Reports of Major Depression 18 and Up [h]	Recreational Marijuana	-0.0220	-0.0852	0.0411	0.0315		-0.0071	0.0701	0.0357		0.0061	0.0654
	Medical Marijuana	-0.0066	-0.0380	0.0248	-0.0031		-0.0341	0.0278	0.0026		-0.0247	0.0299
Serious Mental Illness 18 to 25 [i]	Recreational Marijuana	0.0717	-0.0438	0.1872	0.0837	**	0.0375	0.1299	0.0603	**	0.0223	0.0983
	Medical Marijuana	-0.0186	-0.0802	0.0431	-0.0042		-0.0416	0.0333	-0.0035		-0.0445	0.0376
Serious Mental Illness 26 and Up [i]	Recreational Marijuana	-0.0658	-0.1630	0.0314	-0.0060		-0.0639	0.0518	-0.0023		-0.0486	0.0440
	Medical Marijuana	0.0454	-0.0379	0.1286	0.0024		-0.0370	0.0418	-0.0063		-0.0483	0.0357
Serious Mental Illness 18 and Up [i]	Recreational Marijuana	-0.0375	-0.1300	0.0550	0.0096		-0.0440	0.0632	0.0128		-0.0304	0.0561
	Medical Marijuana	0.0335	-0.0419	0.1090	0.0029		-0.0301	0.0358	-0.0032		-0.0388	0.0324
Serious Thoughts of Suicide 18 to 25 [i]	Recreational Marijuana	-0.0124	-0.0554	0.0306	0.0333		-0.0322	0.0988	0.0192		-0.0545	0.0929
	Medical Marijuana	0.0039	-0.0702	0.0780	-0.0240		-0.0648	0.0168	-0.0205		-0.0579	0.0168
Serious Thoughts of Suicide 26 and Up [i]	Recreational Marijuana	-0.0038	-0.0980	0.0905	0.0603		-0.0075	0.1280	0.0470		-0.0042	0.0983
	Medical Marijuana	0.0489	-0.0112	0.1091	-0.0281		-0.0790	0.0229	-0.0295		-0.0899	0.0310
Serious Thoughts of Suicide 18	Recreational Marijuana	-0.0034	-0.0789	0.0721	0.0464		-0.0148	0.1076	0.0342		-0.0157	0.0841

and Up [i]										
	Medical Marijuana	0.0330	-0.0157	0.0817	-0.0291	-0.0630	0.0048	-0.0285	-0.0719	0.0149
Any Mental Illness 18 to 25 [i]	Recreational Marijuana	-0.0052	-0.0452	0.0348	0.0260	-0.0019	0.0538	0.0341 **	0.0090	0.0591
	Medical Marijuana	0.0012	-0.0429	0.0454	0.0101	-0.0168	0.0369	0.0092	-0.0170	0.0354
Any Mental Illness 26 and Up [i]	Recreational Marijuana	-0.0277	-0.0898	0.0345	0.0295	-0.0135	0.0725	0.0230	-0.0127	0.0588
	Medical Marijuana	-0.0146	-0.0598	0.0307	-0.0248	-0.0710	0.0213	-0.0269	-0.0738	0.0199
Any Mental Illness 18 and Up [i]	Recreational Marijuana	-0.0230	-0.0776	0.0315	0.0286	-0.0087	0.0659	0.0258	-0.0049	0.0566
	Medical Marijuana	-0.0119	-0.0519	0.0280	-0.0191	-0.0590	0.0209	-0.0204	-0.0605	0.0198

*P < .05

**P < .01

[a] n = 1 020

[b] n = 1 012

[c] n = 951

[d] n = 1 005

[e] n = 1 008

[f] n = 1 014

[g] n = 1 009

[h] n = 663

[i] n = 510