INFERRED RESOLUTION THROUGH HERD IMMMUNITY OF FIRST COVID-19 WAVE IN MANAUS, BRAZILIAN AMAZON

*Thomas A. A. Prowse, School of Mathematical Sciences, The University of Adelaide, Adelaide, South Australia 5005, Australia

*Tara Purcell, Victorian Infectious Diseases Reference Laboratory Epidemiology Unit at The Peter Doherty Institute for Infection and Immunity, The University of Melbourne and Royal Melbourne Hospital, Melbourne, Victoria 3000, Australia

Djane Clarys Baía-da-Silva, Fundação de Medicina Tropical Dr Heitor Vieira Dourado, Manaus, Brazil

Vanderson Sampaio, Fundação de Vigilância em Saúde; Fundação de Medicina Tropical Dr Heitor Vieira Dourado, Manaus, Brazil

Wuelton Marcelo Monteiro, Universidade do Estado do Amazonas; Fundação de Medicina Tropical Dr Heitor Vieira Dourado, Manaus, Brazil

James Wood, School of Population Health, UNSW Sydney, Australia

Ivo Mueller, Population Health and Immunity Division, Walter + Eliza Hall Institute, Parkville, Victoria, Australia; Department of Medical Biology, FMDHS, Univ. of Melbourne, Parkville; Department of Parasites and Insect Vectors, Institut Pasteur, Paris, France

Jodie McVernon, Victorian Infectious Diseases Reference Laboratory Epidemiology Unit at The Peter Doherty Institute for Infection and Immunity, The University of Melbourne and Royal Melbourne Hospital, Melbourne, Victoria 3000, Australia; Centre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, The University of Melbourne; Infection and Immunity Theme, Murdoch Childrens Research Institute, Parkville.

#Marcus V. G. Lacerda, Instituto Leônidas & Maria Deane, Fiocruz; Fundação de Medicina Tropical Dr Heitor Vieira Dourado, Manaus, Brazil

#Joshua V. Ross, School of Mathematical Sciences, The University of Adelaide, Adelaide, South Australia 5005, Australia

* equal first authors

equal last authors

INTRODUCTORY PARAGRAPH (189 words)

As in many other settings, peak excess mortality preceded the officially reported 'first wave' peak of the COVID-19 epidemic in Manaus, Brazil, reflecting delayed case recognition and limited initial access to diagnostic testing. To avoid early information bias, we used detailed age and gender stratified death certificate and hospitalisation data to evaluate the epidemic's trajectory and infer the cause of its decline using a stochastic model. Our results are consistent with heterogenous transmission reducing over time due to the development of herd immunity. Relative to a baseline model that assumed homogenous mixing across Manaus, a model that permitted a small, self-isolated population fraction raised the estimated herd-immunity threshold from 28% to 30% and reduced the final attack rate from 86% to 65%. In the latter scenario, a substantial proportion of vulnerable, older individuals remained susceptible to infection. Given uncertainties regarding the distancing behaviours of population subgroups with different social and economic characteristics, and the duration of sterilising or transmission-modifying immunity in exposed individuals, we conclude that the potential for epidemic outbreaks remains, but that future waves of infection are likely to be much less pronounced than that already experienced.

LETTER TEXT (1,495 words)

Background/Rationale (420 words)

Globally, marked differences have been observed in morbidity and mortality due to SARS-CoV-2 infection. This variability mostly reflects the extent and timeliness of spontaneous and imposed changes in social mixing and the sensitivity of surveillance systems to detect cases and deaths[1]. Many countries that successfully constrained the initial epidemic through distancing measures are now experiencing second waves in still-susceptible populations[2].

Estimates of R_0 for SARS-CoV-2 in the range 2 to 6[3,4] suggest that population immunity of approximately 50-80% is required to achieve herd protection. However, heterogeneous behaviour, infectivity and immunity within subpopulations could plausibly decrease this threshold to 10-20%[5]. Controversy remains regarding the extent of population exposure required to achieve such constraint[6].

Comparison of population attack rates to inform this question is made challenging by

imperfect case ascertainment, compounded by limited diagnostics and overwhelmed health

systems, particularly in high incidence settings. Assessment of epidemic activity therefore

requires the use of less biased metrics than confirmed case reports. Excess mortality is an

objective measure which, with cause-of-death certification, can be used as an indicator of

direct and indirect COVID-19 associated mortality[7]. With hospitalisations data, it can inform

retrospective estimation of cumulative cases and deaths[8].

Brazil experienced a severe first wave of COVID-19 disease, with mass mortality reported in

many states, mainly in the north where seasonality of respiratory infections contributed to

higher vulnerability. A socialized health system provided free and global access to tertiary care

hospitals, but inequalities might explain different mortality rates in the population[9]. In Manaus,

the highly urbanised capital of Amazonas state, the first case of COVID-19 was reported on 13

March 2020[10]. By 11 August 2020, 37,597 cases and 2,051 deaths were reported[11].

However, burial and death records indicate far higher mortality than official reports, suggesting

late recognition of importation and underreporting. Previous studies have assumed that the first

wave in Manaus was significantly mitigated by non-pharmaceutical interventions (NPIs)[12].

While these restrictions may have partly constrained early transmission, local reports indicate

that implementation was highly variable[13]. Moreover, a possible role for immunity is suggested

by the observation of declining cases and deaths over a period in which restrictions were

officially eased.

We use death certificate and hospitalisation records to parameterise an epidemiological model

of the COVID-19 epidemic in Manaus. The model allows inference of age- and gender-

stratified infection-fatality ratios to explore evidence for development of herd immunity as a

driver of local epidemic resolution, with implications for the ongoing risk posed by SARS-

CoV-2 to this population.

Main results (388 words)

We calculated 3,457 excess deaths in Manaus, Brazil, between 19 March and 24 June 2020

(Supplementary Table 1) representing 0.16% of the city's population. Males 30 years and over

experienced greater excess mortality than females; individuals aged 75 years or more

accounted for 39% of the excess (Figure 1a). During this period, 7% of the 75+ male population

in Manaus died (Figure 1b). COVID-19 deaths were first reported from 26 March and increased

weekly thereafter in keeping with improved access to diagnostics and/or increasing prevalence

(Figure 2), comprising 53% of the total excess. Other reported causes of death included

respiratory diseases, unattended and unknown causes of mortality, cardiovascular, endocrine

and cancer-related mortality, the majority of which were compatible[14] with a clinical diagnosis

of COVID-19 or are known comorbidities associated with severe outcomes (Figure 2).

Stochastic transmission models captured the observed peak in excess deaths in late April,

including age and gender variation (Figure 3, Supplementary Figures S1, S3). The models also

captured synchronous peaks in hospitalisations (Figure 3, Supplementary Figures S2, S4),

although model fit to these data was poorer, potentially reflecting delayed recognition of

COVID-19 cases and/or capacity exceedance during the epidemic peak.

Despite their different assumptions about the proportion of the Manaus population at risk, the

two model scenarios yielded equivalent fits to the data (Figure 3A, B). The baseline model

estimated heterogenous transmission (k=0.065 [0.047, 0.093], mean [95 % credible intervals]),

a mean time-to-death of 13 days (Supplementary Figures S5), and inferred a population-wide

attack rate of 85.7 [84.6, 86.7]% by 20 July. Attack rates were lowest in younger age classes

(Figure 3A(i)), reflecting lower susceptibility assumed for people under 15 years.

Infection fatality rate (IFR) estimates for the baseline model ranged from almost zero for both

genders aged 30-35 years, to 3.0 [2.8, 3.3]% and 7.4 [6.8, 8.0]% for females and males over 75

years, respectively (Figure 4). At a population level, reduced mobility resulted in variable R_t,

while estimates of R_{eff} fell below 1 after 10 April (Supplementary Figure S6), when the

estimated attack rate was 52.6 [51.1, 54.1]%.

Outcomes were similar for the second model which estimated 24.5 [23.0, 25.1]% of the Manaus

population was effectively removed from the susceptible pool. Relative to the baseline model,

this second model raised the calculated herd-immunity threshold from 28% to 30% and reduced

the final attack rate to 65.0 [63.7, 65.6]%.

Implications (687 words)

Early time-series modelling of the COVID-19 epidemic in Brazilian states up to 6 May 2020

used an unstructured model informed by state-level mobility indicators and reported deaths[12].

The model inferred population attack rates ranging from 0.13% to 10.6%, for Minas Gerais

and Amazonas, respectively, and predicted ongoing epidemic growth throughout May. Other

modelled R_0 estimates based on reported COVID-19 cases from Amazonas state have similarly

anticipated continued growth in cases through the month of June[13]. These models could neither

explain nor accurately forecast observations of epidemic decline in Manaus.

Our analysis of excess deaths and hospitalisations in Manaus identifies rapid epidemic growth

from late March, peaking in late April, with a return to baseline mortality in early June. These

age- and gender-stratified models inferred heterogeneous transmission consistent with previous

studies of SARS-CoV-1[15] and SARS-CoV-2[16], and a herd-immunity threshold of around 30%.

However, we estimated that this threshold was substantially overshot in Manaus, with resulting

attack rates inferred to be 86% and 65% for models that assumed the risk of infection applied

to 100% or *c.* 75% of the population, respectively.

Although both models estimated declining SARS-CoV-2 transmission over time due to herd

immunity, their implications for the possibility of subsequent waves are qualitatively different.

Although there are currently no data to support the assertion that some fraction of the Manaus

population self-isolated, we consider the second model is likely more realistic because: (1) the

estimated infection-fatality ratios are higher than reported from high-income countries[17] which

we anticipate to be plausible; and (2) a population attack rate of 65% is more consistent with

recent estimates based on a Manaus blood bank serosurvey[18] which reported the largest

increase in seropositivity over the month of April.

We focus on Manaus as the epicentre of the COVID-19 epidemic in Amazonas, avoiding

conflation of case numbers in this dense city of more than 2 million people with slower

growing rural outbreaks across the state[6]. Given known information bias associated with

limited early testing capacity, we used excess deaths and hospitalisations as the most

objective indicators of epidemic activity presently available. Detailed cause-of-death data

support the hypothesis that the majority of excess mortality over this time was COVID-19

related.

An important caveat is that estimated attack rates accumulate all infections and are agnostic to
the presence or degree of symptoms. Severity of the clinical course of COVID-19 is associated
with magnitude and persistence of the host immune response[19]. In consequence, our estimates
of 'exposure' cannot be directly related to predicted antibody seroprevalence at the end of the
first wave[20]. It is reasonably anticipated that population immunity will wane over time,
requiring robust memory responses[21] to prevent reinfection or modify the clinical course[22].

Our findings support emerging evidence that population heterogeneity of SARS-CoV-2
transmission, attributable to a range of biological and sociological factors, reduces the herd
immunity threshold[16,23,24]. Reduction of superspreading events through constraints on mixing
group sizes is concordant with genomic studies showing successive extinction of imported
strains under the influence of social and mobility restrictions[25].

While future infection clusters and outbreaks remain possible in Manaus due to unexposed
subgroups, population-level immunity will likely constrain widespread transmission unless
immunity wanes. As in other settings, underlying vulnerability of older age groups may be
further exacerbated over time by reduced health seeking behaviours of individuals with pre-
existing and new medical conditions[26]. Social measures should be informed by understanding
of key enablers of superspreading and amplification, awareness of at-risk populations and
tailored to context depending on population experience of the first wave.

Excess mortality data are not sufficiently timely to support real-time decision making, but in
areas with limited testing may be a more reliable indicator of past epidemic activity than
confirmed cases. To accurately support response to and assess the impact of the COVID-19
and other public health emergencies, improved access to diagnostics, and strengthening of
reporting systems are needed in low- and middle-income settings. Cross-sectional and
longitudinal seroprevalence studies are essential to understand markers and maintenance of
immunity to inform prediction of long-term epidemiologic trends and bridging to likely vaccine
impacts[18,27].

Acknowledgements

TAAP is supported by the NHMRC Centre for Research Excellence in Policy Relevant
Infectious Disease Simulation and Mathematical Modelling. JM is supported by an NHMRC

Principal Research Fellowship GNT1117140. IM is supported by an NHMRC Principal

Research Fellowship GNT1155075.

METHODS (ONLINE ONLY)

Excess mortality and reported causes

Death certificate data from January 2015 to June 2020 were sourced from the Brazilian Ministry of Health Mortality Information System (SIM). Information on age, gender, and cause and date-of-death was recorded. Cause of death was reported using the World Health Organization's (WHO) International Classification of Diseases 10th revision (ICD-10). The 2010 population census and the 2019 municipal population estimates were accessed from the Brazilian Institute of Geography and Statistics (IBGE). This project was approved by the *Fundação de Medicina Tropical Dr Heitor Vieira Dourado* Ethics Review Board (Approval 4.033.218).

Background mortality was calculated by averaging the number of deaths observed per week, commencing on January 1 each year, for the previous 5 years (2015-2019). Background mortality was further stratified by gender and into 5-year age bands, culminating at 75+. For each subgroup, we subtracted this five-year average from deaths observed for the period between March 19 and June 24 of 2020 to determine excess deaths. This period was selected as excess mortality was first observed in the week beginning on March 19 and the data returned to baseline background mortality in the week beginning on June 25.

The age and gender structure reported in the 2010 population census was used to estimate the 2019 population size for each age and gender class. Population estimates were aggregated into 5-year age bands. The 5-year population estimates for 2019 were used to determine excess mortality, as a proportion of the population.

The ICD-10 codes assigned to each death certificate were aggregated into 7 categories: diseases of the respiratory system (J00-J99), circulatory system (I00-I99), endocrine system (E00-E99), unattended (R98) and unknown cause of death (R99), coronavirus infection (B34.2) and all remaining codes were categorised as 'other'. All analyses were performed using R software.

Hospitalisation data

Hospitalisation data for patients admitted to all hospitals (private and public) in Manaus from January 2020 to July 2020 was accessed from the Influenza Surveillance Information System (SIVEP-Gripe). Data entry in this system is compulsory by law. A deidentified line list

included patient demographics, comorbidities, clinical symptoms, investigations, clinical
management and outcome data.

Daily admissions to both hospital and intensive care units (ICU) were aggregated each week
in 2020 and stratified into 3 age bands (0-29, 30-64 and 65+). For each week, the proportion
of hospital and ICU admissions within each age group was determined. Between April and
July, the proportional distribution of hospital and ICU admissions across the three categories
remained fairly stable, with no evidence of rationing of service access on the basis of age.

Epidemiological model

We developed a stochastic, discrete-time, susceptible-infected-recovered epidemiological
model for Manaus which we calibrated using daily time-series data from the death-certificate
and hospitalisation records. The model assumed an initial population size equal to that of
Manaus in 2019 (2,182,761), which was split into $N=36$ age/gender groups based on
proportions reported in the 2010 census (i.e., 5-year age classes up to an age of 75 years, and
then a pooled age class for all individuals aged 75 years and over). To allow inference on the
background mortality rate in each age/gender group, the model was initiated on January 1,
2020, approximately two months prior to the introduction of COVID-19. The model was
terminated after $n=202$ days on July 20, 2020, to ensure complete reporting of both death
certificates and hospitalisations over the modelled period. The model detailed below was fitted
within a Bayesian framework using *JAGS* (v. 4.3.0) software[28] and a mixture of informative
and uninformative priors (for full details of the model code, including details of all prior
distributions, see Supplementary Appendix S1).

COVID-19 introduction and virus transmission. We initialised the model by allowing
importation of COVID-19 cases into Manaus over the week beginning March 5, 2020 (one
week before the first reported case), and inferred an importation model such that the number
of introduced cases into each group in each day of this first week arose from a Poisson process
with a common mean inferred from the data. To model community transmission, we inferred
a reproduction number on day t that was modified by an index of human mobility

$$R_t = e^{(\log(R_0) + aX_t)}$$

where R_0 is the basic reproduction number, X_t is the mobility covariate and a is the inferred coefficient. We derived the time-series X_t by first averaging daily data for five separate indices of community mobility (available as the mobility change relative to baseline for retail/recreation, grocery/pharmacy, parks, transit stations and workplaces, accessed from www.google.com/covid19/mobility), and then smoothed the series with a 7-day moving-average smoother. We then assumed the effective reproduction number $R_{eff_{(i,j,t)}}$ in group i due to mixing with group j on day t is

$$R_{eff_{(i,j,t)}} = R_t \, Q_i \, M_{i,j} \, S_{i,t}$$

where Q_i is the susceptibility of group i to infection, $M_{i,j}$ is the rate of mixing between the two groups, and $S_{i,t}$ is the time-varying proportion of susceptible (previously uninfected) individuals remaining in the focal group. Based on previous studies[29,30], we used prior means for the susceptibilities of 0-15, 15-60, and 60+ year old age classes of 0.5, 1.0, and 1.3, respectively.

Mixing between groups was governed by an age-structured mixing matrix reported previously for Brazil [31], which we corrected to be symmetrical (by averaging the upper and lower triangles), scaled to a mean of $1/N$, and applied equally to both genders. The expected number of new cases in each group each day was then calculated as

$$C_{i,t} = \sum_{j=1}^{N} R_{eff_{(i,j,t)}} \left(c_{j,1:(t-1)} \cdot g_{(t-1):1} \right)$$

where $c_{j,1:(t-1)}$ is the case-history vector for group j and $g_{(t-1):1}$ is the portion of the generation interval distribution relevant to those cases. We used the same generation interval distribution as Mellan et al.[12] which concentrated >99% of an individual's infectivity within the first 3 weeks of infection (median generation interval=5 days).

To account for heterogeneity in transmission, we assumed the offspring distribution in group i due to mixing with infectious individuals from group j on day t was governed by a negative binomial distribution with mean equal to $C_{i,t}$ and variance equal to $C_{i,t}(1 + C_{i,t}/\emptyset_{i,t})$, where

$$\emptyset_{i,t} = kC_{i,t}/R_t$$

and k is the overdispersion parameter, such that smaller values of k represent greater

transmission heterogeneity (i.e., the more transmission is due to a small number of people,

including by so-called "superspreaders"). Given no previous study has documented strong

evidence of different susceptibility between genders, we first generated the expected number

of new cases in each age class (regardless of gender), and then assumed gender-specific cases

arose from a binomial distribution with probability equal to the proportion of the total

susceptible individuals for that age class attributable to each gender.

Background and COVID-induced mortality. We developed a model for the expected number

of deaths per day which was comprised of three components. First, we modelled the

background (pre-COVID-19) death rate d_i estimated separately for each group. Second,

available data on the time from symptom onset to death for confirmed COVID-19 cases (from

the Manaus hospitalisation records) were used to infer a 4-parameter $(y_0, u_{\min}, u_0, b)$ mortality

distribution (m) as a function of the time x since symptom onset ($x \in \{-5, …, N\text{-}5\}$) of the form

242
$$m_{x+a} = \frac{y_0 e^{u_{\min}(x+a) \ + \ ((u_0 - u_{\min})/b)(1 - e^{-b(x+a)})}}{\sum_{j=1}^{N} m_j}$$

243 where a is the average time from infection to symptoms, which was fixed at 5 days[3,32,33].

244 Finally, we modelled the infection-fatality ratio in each group i (IFR_i) as a log-linear function

245 of age and gender, with prior distributions on this components' parameters based on a recent

246 meta-analysis of COVID-19-induced mortality[34]. Together, these components yielded the

247 following model for the expected number of deaths

248
$$D_{i,t} = d_i + IFR_i \sum_{j=1}^{N} \left(c_{j,1:(t-1)} \cdot m_{(t-1):1} \right)$$

and the observed number of deaths each day was assumed to arise from a Poisson distribution

with mean equal to $D_{i,t}$.

Hospitalisations due to COVID-19. Hospitalisations were modelled in a similar way to

COVID-induced deaths, in that available data on the time from symptom onset to

hospitalisation were used to infer a 4-parameter hospitalisation distribution (h) of the same

functional form as that used for the mortality distribution above. We modelled the infection-

hospitalisation ratio in each group i (IHR_i) as a logistic function of age and gender. Daily
hospitalisations were assumed to arise from a Poisson distribution with mean equal to

$$H_{i,t} = IHR_i \sum_{j=1}^{N} \left(c_{j,1:(t-1)} \cdot h_{(t-1):1} \right) .$$

Model scenarios. Given no data were available on whether some portion of the Manaus
population has been self-isolating since March 2020, we initially fitted a 'baseline' model to
death-certificate and hospitalisation data which assumed homogeneous age-structured mixing
across the entire Manaus population. We compared these model outcomes to those from a
second model that permitted a self-quarantined proportion of the population which could not
be exposed to the SARS-CoV-2 virus. Communication with local experts including authors on
this paper suggested that more wealthy residents of Manaus had been able to greatly reduce
social interactions during the first epidemic wave and avoid exposure to the virus. We reviewed
detailed socio-demographic data on Manaus[35], but were not able to source quantitative
estimates of the relevant population fraction. However, we considered that this would not
exceed the upper 2 income quintiles (40%) of the population but was likely to be greater than
the wealthiest 10%, and accordingly, chose a non-informative uniform (0.1,0.4) prior on this
population fraction.

REFERENCES

1. Dye, C., Cheng, R., Dagpunar, J. & Williams, B. The scale and dynamics of COVID-19 epidemics across Europe. (https://www.medrxiv.org/content/10.1101/2020.06.26.20131144v2, 2020).
2. Klein, A. Australia looks to be finally beating its second wave of coronavirus. in *New Scientist* (https://www.newscientist.com/article/2252690-australia-looks-to-be-finally-beating-its-second-wave-of-coronavirus, 2020).
3. Li, Q*., et al.* Early Transmission Dynamics in Wuhan, China, of Novel Coronavirus–Infected Pneumonia. **382**, 1199-1207 (2020).
4. Sanche, S*., et al.* High Contagiousness and Rapid Spread of Severe Acute Respiratory Syndrome Coronavirus 2. *Emerging Infectious Disease journal* **26**, 1470 (2020).
5. Aguas, R*., et al.* Herd immunity thresholds for SARS-CoV-2 estimated from unfolding epidemics. 2020.2007.2023.20160762 (2020).
6. Ferrante, L*., et al.* Brazil's policies condemn Amazonia to a second wave of COVID-19. *Nature Medicine* (2020).
7. Felix-Cardoso, J., Vasconcelos, H., Rodrigues, P. & Cruz-Correia, R. Excess mortality during COVID-19 in five European countries and a critique of mortality analysis data. 2020.2004.2028.20083147 (2020).
8. Thompson, R.N*., et al.* Key questions for modelling COVID-19 exit strategies. *Proc Biol Sci* **287**, 20201405 (2020).
9. Croda, J*., et al.* COVID-19 in Brazil: advantages of a socialized unified health system and preparation to contain cases %J Revista da Sociedade Brasileira de Medicina Tropical. **53**(2020).
10. Amazonas Health Surveillance Foundation FVS. Amazonas confirms 1st case of Covid-19 and authorities guarantee that the assistance network is prepared for assistance. (2020).
11. Amazonas Health Surveillance Foundation FVS. COVID-19 Monitoring Panel. Manaus, Brazil. (2020).
12. Mellan, T.A., Hoeltgebaum, H.H. & Mishra, S. Estimating COVID-19 cases and reproduction number in Brazil. (Imperial College London, 2020).
13. de Souza, W.M*., et al.* Epidemiological and clinical characteristics of the COVID-19 epidemic in Brazil. *Nat Hum Behav* **4**, 856-865 (2020).
14. Excess deaths associated with COVID-19. Provisional death counts for coronavirus disease (COVID-19). National Center for Health Statistics, Centers for Disease Control and Prevention.
15. Lloyd-Smith, J.O., Schreiber, S.J., Kopp, P.E. & Getz, W.M. Superspreading and the effect of individual variation on disease emergence. *Nature* **438**, 355-359 (2005).
16. Endo, A., null, n., Abbott, S., Kucharski, A. & Funk, S. Estimating the overdispersion in COVID-19 transmission using outbreak sizes outside China [version 3; peer review: 2 approved]. **5**(2020).
17. Levin, A., Meyerowitz-Katz, G., Owusu-Boaitey, N., Cochran, K. & Walsh, S. Assessing the age specificity of infection fatality rates for COVID-19: systematic review, meta-analysis, and public policy implications. (https://www.medrxiv.org/content/10.1101/2020.07.23.20160895v4.full.pdf, 2020).
18. Buss, L*., et al.* COVID-19 herd immunity in the Brazilian Amazon. (https://www.medrxiv.org/content/10.1101/2020.09.16.20194787v1, 2020).
19. Long, Q.X*., et al.* Clinical and immunological assessment of asymptomatic SARS-CoV-2 infections. *Nat Med* **26**, 1200-1204 (2020).

320   20.   Hallal, P*., et al.* Remarkable variability in SARS-CoV-2 antibodies across Brazilian
321   regions: nationwide serological survey in 27 states.
322   (https://www.medrxiv.org/content/10.1101/2020.05.30.20117531v1, 2020).
323   21.   Rodda, L*., et al.* Functional SARS-CoV-2 specific immune memory persists after
324   mild COVID-19.
325   (https://www.medrxiv.org/content/10.1101/2020.08.11.20171843v2, 2020).
326   22.   Ledford, H. COVID-19 reinfection: three questions scientists are asking. *Nature* **585**,
327   168-169 (2020).
328   23.   Britton, T., Ball, F. & Trapman, P. A mathematical model reveals the influence of
329   population heterogeneity on herd immunity to SARS-CoV-2. **369**, 846-849 (2020).
330   24.   Wang, Y. & Teunis, P. Strongly Heterogeneous Transmission of COVID-19 in
331   Mainland China: Local and Regional Variation. **7**(2020).
332   25.   Pybus, O.G*., et al.* Preliminary analysis of SARS-CoV-2 importation and
333   establishment of UK transmission lineages. (https://virological.org/t/preliminary-
334   analysis-of-sars-cov-2-importation-establishment-of-uk-transmission-lineages/507,
335   2020).
336   26.   Kluge, H.H.P*., et al.* Prevention and control of non-communicable diseases in the
337   COVID-19 response. *Lancet* **395**, 1678-1680 (2020).
338   27.   Koopmans, M. & Haagmans, B. Assessing the extent of SARS-CoV-2 circulation
339   through serological studies. *Nat Med* **26**, 1171-1172 (2020).
340   28.   Plummer, M. JAGS: A Program for Analysis of Bayesian Graphical Models using
341   Gibbs Sampling. *3rd International Workshop on Distributed Statistical Computing
342   (DSC 2003); Vienna, Austria* **124**(2003).
343   29.   Zhang, J*., et al.* Changes in contact patterns shape the dynamics of the COVID-19
344   outbreak in China. **368**, 1481-1486 (2020).
345   30.   Davies, N.G*., et al.* Age-dependent effects in the transmission and control of COVID-
346   19 epidemics. *Nature Medicine* (2020).
347   31.   Prem, K., Cook, A.R. & Jit, M. Projecting social contact matrices in 152 countries
348   using contact surveys and demographic data. *PLOS Comput. Biol.* **13**, e1005697
349   (2017).
350   32.   Lauer, S.A*., et al.* The Incubation Period of Coronavirus Disease 2019 (COVID-19)
351   From Publicly Reported Confirmed Cases: Estimation and Application. *Annals of
352   Internal Medicine* **172**, 577-582 (2020).
353   33.   He, X*., et al.* Temporal dynamics in viral shedding and transmissibility of COVID-19.
354   *Nature Medicine* **26**, 672-675 (2020).
355   34.   Levin, A.T., Meyerowitz-Katz, G., Owusu-Boaitey, N., Cochran, K.B. & Walsh, S.P.
356   Assessing the age specificity of infection fatality rates for COVID-19: systematic
357   review, meta-analysis, and public policy implications. 2020.2007.2023.20160895
358   (2020).
359   35.   United Nations Development Programme. Atlas of human development in Brazil
360   (http://www.atlasbrasil.org.br/2013/en/perfil_m/manaus_am/, accessed 16/09/2020).
361

364   **Figure 1.** (a) Total excess mortality by age and gender for the period between March 19 and

365   June 24 of 2020. Excess mortality is the 2020 weekly observed mortality less the expected

366   mortality summed for the period between March 19 and June 24 of 2020 for each subgroup.

367   (b) Excess mortality for the period between March 19 and June 24 of 2020 divided by the

368   estimated 2019 subgroup populations to obtain population proportions.

Figure 2. Observed weekly mortality for the period between January 01 and July 22 of 2020 aggregated by major ICD-10 categories of attributed causes of death.

Figure 3. Integrated model fits by gender and model, pooled at the level of three aggregated
age classes: (A) 0-30 years; (B) 30-60 years; and (C) 60+ years. For each age class, panels (i)
and (ii) show the fit of the model (lines) to the reported number of deaths and hospitalisations
per day (points), while panel (iii) shows inferred attack rates. Note that lines showing the
model fits overlap in panels (i) and (ii), and attack rates for males and females overlap in
panel (iii).

Figure 4. Model-based estimates (mean ± 95 % credible intervals) of age- and gender-
structured (A) infection-fatality ratios (IFRs) and (B) infection-hospitalisation ratios (IHRs).
Note that in panel (A), IFRs were fixed at zero for all age classes below 30 years.

383
Figure S1. Baseline model fits to daily deaths data for all 32 age and gender classes. Shown

are the model-inferred expected death rate from all causes (lines) ± 95% credible intervals

(ribbons), and the mortality data used for model fitting (points).

Figure S2. Baseline model fits to hospitalisations data for all 32 age and gender classes.

Shown are the model-inferred expected hospitalisation rate (lines) ± 95% credible intervals

(ribbons) and the hospitalisations data used for model fitting (points).

391
392 **Figure S3**. The fit of Model 2 to daily deaths data for all 32 age and gender classes. Shown

393 are the model-inferred expected death rate from all causes (lines) ± 95% credible intervals

394 (ribbons), and the mortality data used for model fitting (points).

395
Figure S4. The fit of Model 2 to hospitalisations data for all 32 age and gender classes.

Shown are the model-inferred expected hospitalisation rate (lines) ± 95% credible intervals

(ribbons) and the hospitalisations data used for model fitting (points).

Figure S5. Estimates from the baseline model (mean ± 95 % credible intervals) of the shape

of the (A) mortality distribution and (B) hospitalisation distribution, as a function of time

since infection. Vertical bars indicate empirical frequencies derived from hospitalisation

records for Manaus. In (A), the dashed line indicates the mortality distribution used by a

previous COVID-19 modelling study for the State of Amazonas, Brazil[12].

Figure S6. Estimates from the baseline model (mean ± 95 % credible intervals) of: (A) the

impact of personal mobility on the time-varying reproduction number (R_t); and (B) the

population-level effective reproduction number (R_{eff}) over time. In the latter, $R_{eff} < 1$ indicates

the switch to negative epidemic growth due to the development of herd immunity.

Table S1. Excess mortality in Manaus, Amazonas, during the COVID-19 epidemic (here defined as the period between March 19 and June 24, 2020).

Age	Females			Males			Total	
	Expected mortality	Observed mortality	Excess mortality	Expected mortality	Observed mortality	Excess mortality	Total Excess mortality	Excess mortality (%)
0-5	84.5	65	-19.5	109.6	83	-26.6	**-46.1**	**-23.7**
5-10.	13.2	3	-10.2	19.3	10	-9.3	**-19.5**	**-60**
10-15.	13.5	6	-7.5	19.3	11	-8.3	**-15.8**	**-48.2**
15-20	26.1	14	-12.1	64.3	45	-19.3	**-31.4**	**-34.7**
20-25	24.1	22	-2.1	116.4	78	-38.4	**-40.5**	**-28.8**
25-30	26.7	30	3.3	92.2	81	-11.2	**-7.9**	**-6.6**
30-35	30	35	5	81.2	88	6.8	**11.8**	**10.6**
35-40	38.3	53	14.7	79	118	39	**53.7**	**45.8**
40-45	44.6	88	43.5	72.4	154	81.7	**125.1**	**107**
45-50	53.3	110	56.7	79.8	184	104.2	**160.9**	**120.8**
50-55	59.2	114	54.8	99	225	126	**180.8**	**114.3**
55-60	73.2	180	106.8	121.2	310	188.8	**295.6**	**152.1**
60-65	95.6	215	119.4	139.2	436	296.8	**416.2**	**177.3**
65-70	101.6	283	181.4	148.6	510	361.4	**542.8**	**216.9**
70-75	110.2	276	165.8	143.6	456	312.4	**478.2**	**188.4**
75+	468.4	996	527.6	382.4	1208	825.6	**1353.2**	**159.1**
Total	**1262.4**	**2490**	**1227.6**	**1767.5**	**3997**	**2229.5**	**3457.2**	**114.1**