Impact of visual impairment on balance and visual processing functions in students with

special educational needs

Choi KY[1], Wong HY[1], Cheung HN[1], Tseng JK[2], Chen CC[2], Wu CL[2], Eng H[1], Woo GC[1,3],

Cheong AMY[1,3]

[1] School of Optometry, The Hong Kong Polytechnic University, Hong Kong

[2] Department of Optometry, Asia University, Taichung, Taiwan

[3] Center for Eye and Vision Research, HKSAR

Running head:

Keywords:

Correspondence's Address: Allen MY Cheong

School of Optometry, The Hong Kong Polytechnic University, Hung Hom, Hong Kong

Telephone: 852-2766-6108

Facsimile: 852-2764-6051

Electronic mail: allen.my.cheong@polyu.edu.hk

Word count: 3,660

Number of tables: 2

Number of figures: 2

Abstract

Introduction

Vision is a critical factor for children's development. However, prevalence of visual impairment

(VI) is high in students with special educational needs (SEN). Other than vision disability, this

group of students is prone to having functional deficits. It is unclear whether visual problems

relate to these compromised functional deficits. This study aimed to assess the impact of vision

on visual processing functions and balance performance in SEN students through a community

service in special schools.

Methods

A total of 104 (age 14.3 ± 4.3 years, 43 girls) SEN students in Taiwan were assessed, who were

classified as having normal vision (NV) or vision impairment (VI). Visual acuity (distance and

near) and contrast sensitivity (CS) were measured as the visual outcomes. Visual processing

function assessment included orientation recognition (mailbox game, MB) and facial expression

recognition [Heidi expression test, in terms of card matching] (HC), and examiner's facial

expression matching (HF)]. Dynamic balance was assessed with Timed Up and Go (TUG) test,

while static standing balance was assessed using a force plate to measure the postural sway in

double-legged feet-together and tandem stance with eyes open and closed conditions. Static

balance was presented in terms of the change in the centre of pressure in maximal medial-lateral

(ML) and antero-posterior (AP) sways, sway variability (V), and sway path length (L).

Results

Although visual acuity was significantly worse in VI than NV ($p < 0.001$), CS was similar in the

two groups ($p = 0.08$). MB, HC, and HF also did not differ significantly between groups ($p >$

0.05). NV performed better in the TUG than VI ($p = 0.03$). There was a significant interaction

between eye condition and the vision group ($p < 0.05$) for static balance. Pairwise comparisons

showed that NV swayed significantly less in ML than VI under tandem stance-open eye

condition (p = 0.04), but significantly more in closed eye condition (p = 0.03). Conversely, VI

had less V and shorter L than NV under tandem stance-close eye condition (p = 0.03).

Conclusion

This study is the first to our knowledge to examine the effect of vision on visual processing

functions and balance performance in SEN students. Vision did not appear to be the major reason

for impairment in visual processing. However, vision plays an important role in maintaining

dynamic and static balance in SEN students.

Introduction

Vision in children is crucial for their daily life, as it contributes greatly to their development of functional abilities and essential skills needed for schooling and learning such as reading comprehension and mathematical concepts. However, visual disabilities are common in children with special education needs (SEN).[1] Perinatal adversity is one of the major causes of vision loss,[2,3] including preterm birth, improper neonatal environment, and neurological damage, which can affect visual acuity, contrast sensitivity, and ocular alignment. The level of visual impairment has been shown to be dependent of the severity of the SEN (e.g. grading of cerebral palsy),[4] which was partially attributable to cerebral visual impairment.[5] In addition to neurological and anatomical damage, correctable refractive error is also highly prevalent in children with SEN,[6,7] so prescription of spectacles or other optical aids would be beneficial to improve their vision. However, current evidence shows inadequate eye care service for children with SEN,[8,9] leading to an inappropriate educational experience due to the misunderstanding of the individual's visual status.

Balance function is also compromised in children with SEN, in terms of static and dynamic balance. Children with more severe cerebral palsy (higher gross motor function classification system level) recorded a greater magnitude of postural sway during a static balance measure than children with normal development,[10] as well as dynamic balance function measured with the Timed Up and Go (TUG) test.[11] Children with Down syndrome tended to spend more time to execute in functional balance tasks, such as performing a standing reach[12] and the TUG test.[13] The postural control system was also found to be underdeveloped in autistic[14] and deaf children.[15] For children with other intellectual disabilities, it was controversial whether their balance function was inferior to their peers with normal development.[16,17] However, visual impairment was associated with reduced balance function.[17,18] It was also reported that postural sway for male participants with visual impairment, regardless whether their eyes were open or closed, was similar to that of sighted participants with their eyes closed.[19] This implies that vision has less impact on postural control despite compromised balance function in visually impaired patients. Balance function in children with mild disruption of binocular vision, due to strabismus or amblyopia, was significantly reduced.[20] However, postural control was significantly improved after corrective surgery in children with strabismus,[21] suggesting that

there is a possibility to improve balance function by correcting vision or improved visual

function. Children with visual impairment had poorer postural sway in both double-leg and

single-leg standing compared with their sighted peers.[22] However, reduced static balance

function was only found in eye-open but not eye-closed condition. Both dynamic functional

balance and coordination were also reported to be weaker in children who were visually

impaired.[23, 24] Despite the important role of vision on balance function, no studies have

considered the impact of visual impairment in children who are more prone to have

compromised balance function (e.g. children with SEN).

In view of the importance of vision in childhood development, thorough assessment of visual

function and related visual processing function can provide information on the limitations of the

SEN children.[25] Visuo-spatial and visuo-perceptual impairment, in terms of facial recognition

and line orientation judgement, are common in children with bilateral cerebral palsy[26] and

autism.[27, 28] In contrast, adolescents who were born extremely pre-term or with extremely low

birth weight and visual processing disorders had poorer academic performance independent of

other perinatal diversities.[29] However, based on the various capabilities of the SEN children, it

has been suggested that evaluation for visual processing function could be better delivered in

play situations,[30] especially for children who functioned normally in other visual tasks such as

visual acuity.[31] Despite the importance of vision on learning different perceptual skills, it is

unclear whether visual impairment further impairs visual processing performance in children

with SEN.

Previous studies have suggested poorer balance function, weaker visual processing function, and

higher prevalence of vision impairment in students with SEN. However, it is unclear whether

compromised vision further impairs these SEN students' balance and visual processing

performance. To our knowledge, this is the first study aiming to investigate the impact of vision

on visual processing function and balance performance (dynamic and static) in students attending

special-care schools in Taiwan.

Methods

Study population

This study analysed the cross-sectional data collected as part of a community service project to provide eye care services for students with special needs attending ten special schools in Taiwan. All tests were conducted by optometrists and trained university students. A total of 157 students agreed to participate in the study, of whom 127 students had measurable visual acuity and at least one functional measurement. Students with solely visual impairment, but otherwise normal development (i.e. absence of other non-visual disabilities, n = 23) were excluded. The remaining 104 students (age 14.3 ± 4.3 years, range 4 – 19 years, 43 girls) were included in the analysis to assess the impact of vision on students with different disabilities, including cerebral palsy, autism, and Down syndrome. Written consent and verbal assent (if feasible) were obtained from the guardians and the students, respectively. All study procedures followed the tenets of the Declaration of Helsinki and were approved by The Hong Kong Polytechnic University Human Subjects Ethics Subcommittee.

Data collection

Demographics and information on the disabilities were obtained by a structured questionnaire completed by either the guardians or school teachers. The questionnaires included birth history, types of disabilities, self-report vision status, and visual problems.

Distance visual acuity was measured monocularly by Lea symbols at 3 m (or 1.5 m if vision was poor) using matching toys (The Good-Lite Company, USA). If the students were intellectually incapable of performing the test, tests with lower cognitive requirement were used, sequentially Cardiff acuity test at 1 m by pointing at the direction, Cardiff acuity test at 1 m by preferential looking, and Lea gratings at 57 cm by preferential looking. Habitual visual acuity of the better-seeing eye was converted into LogMAR acuity, allowing subjects to be classified into two groups: normal vision (NV, LogMAR < 0.50) and visual impairment (VI, LogMAR $\geq$ 0.50). Habitual visual acuity was used for grouping because the functional performances were assessed with visual correction aids. Near visual acuity was measured binocularly by LEA near vision card (The Good-Lite Company, USA). Binocular contrast sensitivity (CS) was measured using the letter version of Mars (The Mars Perceptrix Corp., US), or Lea low-contrast symbols flip-chart test / Hiding Heidi test (The Good-Lite Company, USA) at 40 cm if students were intellectually incapable, then LogCS was recorded. External and internal ocular health was

assessed using slit lamp biomicroscopy and direct ophthalmoscope (or binocular indirect

ophthalmoscope with pupil dilation upon parent's consent), respectively.

Visual processing function was evaluated by two paediatric tests (The Good-Lite Company,

USA). Visual orientation recognition was measured using Lea mailbox game and the average

time for five trials was recorded (MB). Facial expression recognition was measured using the

Heidi expression test. The test was divided into two parts: (1) expression recognition with the

Heidi cards matching (HC), (2) expression recognition with the examiner's facial expression

(HF). Both parts were timed and the average time needed for the five trials was recorded.

Dynamic balance function was measured by the Timed Up and Go (TUG) test.[32] The students

were asked to rise from a chair, walk three meters on a straight line, turn around, return to the

chair, and sit down. The chair height depended on the height of the students using the common

chairs provided by the schools, as the age of the sample covered a wide range. The test was

repeated three times, with the time needed for each trial being averaged and recorded. Static

balance function, in terms of postural sway, was measured using a force plate (BP400600, AMTI,

US) with double-leg feet-together standing and tandem stance. Double-legged feet-together

standing was chosen to mimic the natural standing position in daily life, while tandem stance

condition was chosen over one-leg standing to mimic the postural stability during adverse

conditions because of the limited capability of the SEN students in the current study. Postural

sway was measured at each condition while standing steadily for 20 s under each of the four

conditions: (1) feet together + eyes open (FO); (2) tandem stance + eyes open (TO); (3) feet

together + eyes closed (FC), and (4) tandem stance + eye closed (TC). Students were asked to

fixate at a distant target at 3 m under eye open condition (Figure 1). The force plate measured the

subjects' position at the centre of pressure (COP), at which several variables were generated and

included in the analysis: maximum medial-lateral and antero-posterior sway (ML and AP – i.e.

maximal amplitude of COP in the ML and AP dimensions), sway variability (V – i.e. variability

of COP around its mean value computed by the root mean square of the COP displacement in

ML and AP), and sway path length (L – i.e. total distance of COP travelled).

Figure 1. Measurement of postural sway while standing on a force platform for different

conditions. Students were asked to fixate at a distant target at 3 m under open eye condition (left

panel). Upper condition of the right panel was double-leg standing with feet together, while the

lower condition was tandem stance.

Statistical analysis

As the distribution of some of the outcome variables was significantly different from normal,

non-parametric tests were used in the analysis. Mann-Whitney U test was used to compare visual

functions, visual processing functions, and dynamic balance function between the two groups.

For static balance, the results were transformed to achieve normality using percentile ranking

followed by inverse-normal transformation into normally distributed Z-scores.[33] Two-way

repeated measures ANOVA was used to analyse the main effects and the interaction effect

between the grouping (NV and IV), eye conditions, and feet conditions on the static balance.

Greenhouse-Geisser test was used whenever sphericity could not be assumed, and Bonferroni

adjustment was used in the post-hoc comparisons. Inability to perform the test was regarded as

missing data, which was treated with pairwise deletion. Significance level was set as $p < 0.05$.

All statistical procedures were performed using SPSS v22 (IBM Inc, US).

Results

Among the 104 students, 62 (59.6%) were classified as having visual impairment (i.e. visual acuity of the better eye was ≥ 0.50 LogMAR), in which the major causes of the reduced vision were optic nerve related (18.8%, including optic atrophy, optic nerve hypoplasia, and glaucoma), retinal impairment (17.6%, including retinopathy of prematurity, retinal dystrophy, and macular anomalies), uncorrected/under-corrected refractive error (14.1%), ocular media opacity (12.9%), and oculomotor anomaly (11.8%). However, a large proportion of VI (24.8%) was probably not due to ocular problems, but other neurological causes (e.g. cerebral visual impairment due to cerebral palsy). After subjective refraction or retinoscopy, 32 students (12 NV and 20 VI) were found to have uncorrected / under-corrected refractive errors and benefited from prescription of updated optical aids to improve their vision thereafter, with a mean improvement of visual acuity of LogMAR 0.33 ± 0.17. Twelve out of 20 VI students, who were prescribed with an updated optical aid, had best-corrected visual acuity better than LogMAR 0.50. Notably, among the 62 VI students, only 25 guardians/ teachers (40.3%) reported their children/ students having visual problems in the questionnaire, revealing an insufficient awareness of visual problems encountered in the special needs population. Given the different capabilities of the students, it was expected that some students could not complete all the tasks. The distribution of missing data in visual processing and balance functions were independent of the grouping ($p > 0.05$), indicating a similar capability in two groups.

The demographics, visual functions, visual processing functions, and balance performances of both groups of students are listed in Table 1. The age distribution was similar in the two groups ($p = 0.44$) and the LogCS did not differ significantly between VI and NV groups (Mann-Whitney $U = 886.00$, $p = 0.08$). VI and NV groups also had similar performances for the visual processing functions, in terms of MB, HC, and HF ($p > 0.05$). With respect to dynamic balance, NV students performed significantly better than VI students (Mann-Whitney $U = 1040.00$, $p = 0.03$), by a median difference of 3.61 s to complete the TUG task.

Table 1. Comparison of demographic, visual and functional performance between disability groups

[median (IQR) or percentage]

	Normal vision (NV, n = 42)	Visual impairment (VI, n = 62)	p-value
Demographic:			
Age (in years)	16.0 (5.3)	15.0 (5.0)	0.32
Gender			0.88
Female	17 (40.5%)	26 (41.9%)	
Gestation age			0.78
Full-term (≥ 37 weeks)	28 (66.7%)	42 (67.7%)	
Pre-term (< 37 weeks)	6 (14.3%)	11 (17.8%)	
Unknown	8 (19.0%)	9 (14.5%)	
Presence of self-reported non-visual disabilities (e.g. cerebral palsy, Down syndrome, autism)			0.47
(i) 1	(i) 66.7 %	(i) 54.8 %	
(ii) 2	(ii) 16.7 %	(ii) 21.0 %	
(iii) 3+	(iii) 16.7 %	(iii) 24.2 %	
Self-reported visual disabilities	6 (14.3%)	25 (40.3%)	**< 0.01**
Vision measures:			
Distance acuity of the better eye (LogMAR)	0.30 (0.26)	0.87 (0.55)	**< 0.001**
Near acuity (LogMAR)	0.21 (0.35)	0.65 (0.89)	**< 0.001**
Contrast sensitivity (LogCS)	1.64 (0.30)	1.60 (0.86)	0.08
Functional measures:			
Mailbox (MB) – average completion time (s)	1.68 (1.37)	1.78 (1.42)	0.71
Heidi cards (HC) – average completion time (s)	11.00 (13.25)	15.34 (19.90)	0.14
Heidi faces (HF) – average completion time (s)	15.00 (25.33)	18.00 (36.30)	0.52
Time-up-go (TUG) – average completion time (s)	10.60 (4.52)	13.61 (9.05)	**0.03**
Static balance (medial-lateral sway, ML) (cm)*			
(0) Feet-together with eye open (FO)	1.62 (0.93)	1.87 (1.05)	0.37
(ii) Tandem stance with eye open (TO)	1.92 (0.68)	2.45 (0.91)	**0.02**
(iii) Feet-together with eye close (FC)	1.47 (1.23)	1.81 (1.50)	0.24
(iv) Tandem stance with eye close (TC)	2.69 (0.97)	2.23 (1.39)	0.39

* Only 20 (47.6%) and 32 (51.6%) students from the NV and VI groups, respectively, could complete the different conditions in the static balance.

The results for different balance variables are shown in Figure 2. In general, SEN students' static

balance function varied substantially when measured in different conditions. Results from the

two-way repeated measures ANOVA showed that the within-subject main effects for eye

condition (i.e. eye open vs. eye close) and feet condition (i.e. feet together vs. tandem stance)

were insignificant ($F_{1,50} < 1.15$, all $p > 0.05$) in all the outcomes, as was between-subject effect

for vision group (i.e. NV vs. VI) ($F_{1,50} < 0.92$, all $p > 0.05$). However, the interaction between the

eye condition and vision group was significant in V ($F_{1,50} = 4.24$, $p = 0.05$) and L ($F1,50 = 4.66$,

$p = 0.04$), revealing that the absence of visual input (i.e. eye close condition) had stronger impact

on balance function in students with NV than VI. Further interaction among all three factors (eye

condition, feet condition, and vision group) was significant in the ML ($F_{1,50} = 7.44$, $p = 0.01$), but

not in other outcomes ($p > 0.05$). Detailed statistical results are listed in Table 2.

In post-hoc pairwise comparisons, NV students swayed significantly less in ML than VI students

under TO condition ($p = 0.04$), but ML sway became significantly greater under closed eye

condition ($p = 0.02$). Similar findings were observed for V and L with NV students having

significantly less sway under closed eye than open eye condition ($p = 0.03$). In contrast, VI

students swayed significantly more than NV student in V and L under eye closed and tandem

stance condition ($p = 0.03$). The findings demonstrated that when visual input was provided (i.e.

open eye condition), students with normal vision had better static balance performance. However,

when visual input was deprived (i.e. close eye condition), students with vision impairment

swayed significantly less.

Figure 2. Static balance sway parameters in different eye and feet conditions of normal (NV) and

vision impairment (VI) groups. (A) Medial-lateral sway (ML); (B) Antero-posterior sway (AP);

(C) Sway variability (V); (D) Sway path length (L). FO: Feet together standing with eye open;

TO: Tandem stance with eye open; FC: Feet together standing with eye close; TC: Tandem

stance with eye close. The graph is plotted with transformed data, and the error bars represent the

standard error of mean.

259   Table 2. Results of two-way mixed repeated measures ANOVA for static balance measures.

	$F_{1,50}$	p-value		$F_{1,50}$	p-value
Medial-lateral sway (ML)			Antero-posterior sway (AP)		
Group	0.92	0.34	Group	0.18	0.67
Eye condition	< 0.001	> 0.99	Eye condition	0.08	0.77
Feet condition	0.001	0.97	Feet condition	1.15	0.29
Eye * Group	2.92	0.09	Eye * Group	0.34	0.57
Feet * Group	0.33	0.57	Feet * Group	3.37	0.07
Eye * Feet * Group	**7.44****	**0.01**	Eye * Feet * Group	1.85	0.18
Sway variability (V)			Sway path length (L)		
Group	0.61	0.44	Group	0.50	0.48
Eye condition	0.21	0.65	Eye condition	0.09	0.77
Feet condition	0.06	0.81	Feet condition	0.02	0.89
Eye * Group	**4.24**	**0.05***	**Eye * Group**	**4.66**	**0.04***
Feet * Group	1.25	0.27	Feet * Group	1.46	0.23
Eye * Feet * Group	2.22	0.14	Eye * Feet * Group	2.14	0.15

260   * indicates p < 0.05, ** indicates p < 0.01

Discussion

This study is the first to our knowledge to examine the effect of vision on two essential daily

tasks – visual processing and balance functions in children with SEN attending special care

schools. In line with previous findings, prevalence of reduced visual acuity was high in this study.

Our findings showed that vision did not play a major role in the poor performance in functional

tests, such as orientation recognition and facial recognition – two important visual processing

tasks. In contrast, vision was significantly associated with the balance function, both dynamic

and static. SEN students with visual impairment had poorer performance in TUG test and sway

parameters under open eye condition. However, when the visual input was deprived (i.e. eye

close condition), students with normal vision performed significantly worse than those with

visual impairment.

Previous studies have reported an increased risk, as high as 75%, of visual impairment in

disabled children,[34] whereas the current study found 59.6% of students attending the special

schools had reduced vision, although this figure did not include those SEN students with solely

visual impairment (n = 23). In the US, cortical visual impairment was the leading cause of visual

impairment and blindness in children[35, 36] and constituted 19% of the visually challenged

children. Our study, echoes the previous studies in finding that the causes of reduced vision for

24.8% of the SEN students were likely to be neurological, while the remainder were due to

different types of ocular anomalies (e.g. optic atrophy, glaucoma). Although the types of

disabilities were self-reported by guardians or teachers rather than full medical records (because

of compliance with patients' privacy),[37] it was speculated that cerebral visual impairment was

likely to be the major cause of vision loss in students attending the special schools.

Vision did not appear to be a major obstacle for visual processing functions. As in a previous

study, visuo-spatial and visuo-perceptual abilities were found to be impaired in 90% and 60% of

subjects with cerebral palsy, respectively,[26] in terms of orientation judgement and facial

recognition, which was independent of their visual acuity. Facial details processing deficits are

also common in autistic spectrum disorder,[27, 28] even if autistic individuals have normal vision,

leading to problems such as unsustainable eye contact and switching focus for social function.

Inhibition of visual input caused by visual impairment might have limited impact of such social

information. In the current study, orientation judgement and facial expression recognition were

similar in both groups of students – normal vision and visual impairment. Similarly, age did not

show a significant association with the performance in visual processing functions (Spearman's

test, p > 0.05). In line with previous studies, our results indicated that non-visual disabilities, e.g.

intellectual disability, might attribute for the impaired visual processing functions instead of

vision itself. Further studies on functional assessments, such as joint attention and imitation, are

warranted.

The balance functions of SEN students are reduced, regardless of the type of disability. In

children with normal development, the dynamic balance function measured by TUG ranged from

approximately 4 to 7 s.[13, 38] The time needed to complete the TUG task increased from a median

of 7.5 s in gross motor function classification system level I, to 17.8 s in level II, and finally 50.7

s in level III for children with cerebral palsy,[11] while was approximately 9 s in adolescents with

Down syndrome.[13] Our sample recorded a median of 10.60 s and 13.61 s in SEN students with

normal vision and vision impairment, respectively. The correlation analysis showed a significant

but weak association between habitual visual acuity and TUG (Spearman's $\rho = 0.23$, p = 0.04).

This indicated that visual and non-visual disabilities compromise SEN students' dynamic

balance function in different ways and might have a composite effect, as dynamic balance

function was further reduced in students with both visual and non-visual disabilities. Several

studies have compared static balance function in sighted and visually impaired children. In

summary, they concluded that visually impaired individuals had greater postural sway than

sighted individuals under open eye condition, especially in single-leg-standing positions:

adults,[19, 39] adolescent,[17] and children.[18, 22] However, no significant difference were observed

between the sighted and visually impaired individuals under closed eye condition. In the current

study, a similar outcome was found in SEN students. Those students with normal vision had

better static balance performance in open eye condition than those with visual impairment under

tandem stance, while no significant difference was observed in feet together standing. When

visual input was deprived (i.e. eye closed condition), their static balance function became

significantly poorer. However, students with both visual and non-visual disabilities (i.e. VI group)

swayed similarly under eye open and eye closed conditions. It is speculated that students with

visual impairment may rely more on their somatosensory and/or vestibular inputs rather than

vision to maintain their postural control, while NV students rely more on the visual input.

Coincidently, it was observed that similar balance performance was found in older adults with

VI.[40] They swayed significantly more than NV subjects in static standing under eye open, but

displayed no difference with sighted aged-match subjects when eyes were closed. The reliance

on somatosensory input was shown when subjects with VI stood under a sway-reference support

surface with eye open swayed significantly more than the sighted subjects.[40] Further studies

examining the contribution of multi-sensory inputs to postural control, and functional activities,

such as obstacle avoidance and crossing, are needed. Such knowledge could facilitate the design

of balance training protocols for students with SEN.

Taiwan initiated a national-wide registry policy for children with disabilities in 1980's to better

understand and provide support for the SEN. The combined prevalence of SEN increased from

1.0% to 1.5% over three decades,[41] affecting more boys, especially in rural areas.[42]

Institutionalised care, e.g. special schools, has been the mainstream for the special care service,[43]

providing educational, occupational, and vocational training for registrants. Although resources

had been pooled for rehabilitation services, the prevalence of the beneficiaries were still low,[44] in

which the service recipients only accounted for 24.5% of an 957-subject SEN sample within a 7-

month period. As well as the SEN themselves, the primary family carer may also experience

challenges, which the carers' health status and quality of life can be significantly lower.[45] In the

current study, despite the students attending special care schools, awareness of visual impairment

in the guardians or teachers was still low, in that only 40.3% reported their children or students

to have visual impairment. In addition, near one third of the SEN students (n = 32) were found to

have uncorrected / under-corrected refractive error, whose visual acuity was significantly

improved by an average of LogMAR 0.33 by merely updating their optical aids. This

phenomenon indicates the necessity of regular eye examinations in this special population, as

suggested in previous studies.[8, 9, 46]

There were several limitations in the current study. Firstly, the self-reporting disability status was

less reliable than reviewing a full medical history. However, due to the compliance with patient

privacy, this information could not be retrieved from the guardians or teachers. Secondly, the

severity of non-visual disabilities was not assessed in the current study, which is speculated to

affect the capability of SEN student to perform different tasks. Most of our tests required

students to have a certain degree of cognitive ability to participate. It is not surprising that almost

half of the participating students could not complete the various conditions in balance

measurement, in particular the tandem stance in eye closed condition. Finally, this study lacked a

longitudinal follow-up to observe the effect of improved vision after updated optical corrections

on visual processing functions and balance functions in SEN students.

To conclude, visual impairment in SEN students is common. However, despite the high

prevalence of visual impairment in this population, our findings suggested that some SEN

students' visual function could be improved by prescribing appropriate and updated refractive

correction. Although reduced vision is not a major limiting factor explaining the deficits of

visual processing function in these SEN students, vision remains an important integral for

children's all-round development. Adequate visual input was also found to be critical for SEN

students to maintain their dynamic and static balance functions. Hence, regular checking and

preserving the vision for SEN students is of high importance.

Acknowledgment

This study was supported by The Hong Kong Polytechnic University Community Service Fund (83CT) and Chinese Mainland Affairs Office Fund. We thank the financial support from Lions Clubs International MD300 Taiwan Eye Care Network Committee, Taiwan. We thank Dr Pei-Chang Wu and other ophthalmologists from Chang Gung Hospital for providing part of the ocular health examination during the trip. In addition, we appreciated the assistance provided by all volunteering students from The Hong Kong Polytechnic University and Asia University, volunteers from Taiwan Lions Club and teachers from the participating schools. We also thank Prof. William Tsang and Ms. Eva Chan from The Open University of Hong Kong to provide professional advice in preparation of this manuscript. We acknowledge Dr. Maureen Boost for proofreading this manuscript.

Reference

1. Salt A, Sargent J. Common visual problems in children with disability. *Arch Dis Child* 2014; 99(12): 1163-1168.

2. O'Connor AR, Fielder AR, editors. Visual outcomes and perinatal adversity. Seminars in Fetal and Neonatal Medicine; 2007: Elsevier.

3. Fazzi E, Galli J, Micheletti S. Visual impairment: a common sequela of preterm birth. *NeoReviews* 2012; 13(9): e542-e550.

4. Ghasia F, Brunstrom J, Gordon M, Tychsen L. Frequency and severity of visual sensory and motor deficits in children with cerebral palsy: gross motor function classification scale. *Invest Ophthalmol Vis Sci* 2008; 49(2): 572-580.

5. Good WV, Hou C, Norcia AM. Spatial contrast sensitivity vision loss in children with cortical visual impairment. *Invest Ophthalmol Vis Sci* 2012; 53(12): 7730-7734.

6. Sobrado P, Suá rez J, García□Sá FA, Usón E. Refractive errors in children with cerebral palsy, psychomotor retardation, and other non□cerebral palsy neuromotor disabilities. *Dev Med Child Neurol* 1999; 41(6): 396-403.

7. Thankappan B, Sidhan N, Aparna K. Ocular disorders in children with autism in special schools. *J Med Sci Clinc Res* 2017; 5: 25199-25203.

8. Woodhouse JM, Davies N, McAvinchey A, Ryan B. Ocular and visual status among children in special schools in Wales: the burden of unrecognised visual impairment. *Arch Dis Child* 2014; 99(6): 500-504.

9. McKerr L, McConnell EL, Black SA, et al. Meeting vision needs of children with special educational needs: Case studies of the impact on behaviour and academic achievement. *Br J Learn Disabil* 2020; 48(1): 45-58.

10. Bickley C, Linton J, Sullivan E, Mitchell K, Slota G, Barnes D. Comparison of simultaneous static standing balance data on a pressure mat and force plate in typical children and in children with cerebral palsy. *Gait Posture* 2019; 67: 91-98.

11. Carey H, Martin K, Combs-Miller S, Heathcock JC. Reliability and responsiveness of the timed up and go test in children with cerebral palsy. *Pediatr Phys Ther* 2016; 28(4): 401-408.

12. Chen H-L, Yeh C-F, Howe T-H. Postural control during standing reach in children with Down syndrome. *Res Dev Disabil* 2015; 38: 345-351.

13. Nicolini□Panisson RDA, Donadio MV. Normative values for the Timed 'Up and Go'test in children and adolescents and validation for individuals with Down syndrome. *Dev Med Child Neurol* 2014; 56(5): 490-497.

14. Minshew NJ, Sung K, Jones BL, Furman JM. Underdevelopment of the postural control system in autism. *Neurology* 2004; 63(11): 2056-2061.

15. de Sousa AMM, de França Barros J, de Sousa Neto BM. Postural control in children with typical development and children with profound hearing loss. *Int J Gen Med* 2012; 5: 433.

16. Blomqvist S, Olsson J, Wallin L, Wester A, Rehn B. Adolescents with intellectual disability have reduced postural balance and muscle performance in trunk and lower limbs compared to peers without intellectual disability. *Res Dev Disabil* 2013; 34(1): 198-206.

17. Klavina A, Jekabsone I. Static balance of persons with intellectual disabilities, visual impairment, and without disabilities. *Eur J Adapt Phys Act* 2014; 7(1): 50-57.

18. Klavina A, Zusa-Rodke A, Galeja Z. The assessment of static balance in children with hearing, visual and intellectual disabilities. *Acta Gymnica* 2017; 47(3): 105-111.

19. Alghadir AH, Alotaibi AZ, Iqbal ZA. Postural stability in people with visual impairment. *Brain and Behavior* 2019; 9(11): e01436.

20. Zipori AB, Colpa L, Wong AM, Cushing SL, Gordon KA. Postural stability and visual impairment: Assessing balance in children with strabismus and amblyopia. *PLoS One* 2018; 13(10): e0205857.

21.   Legrand A, Quoc EB, Vacher SW, et al. Postural control in children with strabismus: effect of eye surgery. *Neurosci Lett* 2011; 501(2): 96-101.
22.   Müürsepp I, Arjokesse R, Ereline J, Pääsuke M, Gapeyeva H. Impact of visual impairment on static and dynamic postural control and habitual physical activity in children aged 10–16 years. *Br J Vis Impair* 2018; 36(3): 227-237.
23.   Zylka J, Lach U, Rutkowska I. Functional balance assessment with pediatric balance scale in girls with visual impairment. *Pediatr Phys Ther* 2013; 25(4): 460-466.
24.   Rutkowska I, Lieberman LJ, Bednarczuk G, et al. Bilateral coordination of children who are blind. *Percept Mot Skills* 2016; 122(2): 595-609.
25.   Hyvärinen L. Assessment of visual processing disorders in children with other disabilities. *Neuroophthalmology* 2009; 33(3): 158-161.
26.   Pueyo R, Junqué C, Vendrell P, Narberhaus A, Segarra D. Neuropsychologic impairment in bilateral cerebral palsy. *Pediatr Neurol* 2009; 40(1): 19-26.
27.   Guillon Q, Rogé B, Afzali MH, Baduel S, Kruck J, Hadjikhani N. Intact perception but abnormal orientation towards face-like objects in young children with ASD. *Sci Rep* 2016; 6: 22119.
28.   Walsh JA, Creighton SE, Rutherford M. Emotion perception or social cognitive complexity: what drives face processing deficits in autism spectrum disorder? *J Autism Dev Disord* 2016; 46(2): 615-623.
29.   Molloy CS, Di Battista AM, Anderson VA, et al. The contribution of visual processing to academic achievement in adolescents born extremely preterm or extremely low birth weight. *Child Neuropsychol* 2017; 23(3): 361-379.
30.   Hyvärinen L. Considerations in evaluation and treatment of the child with low vision. *Am J Occup Ther* 1995; 49(9): 891-897.
31.   Hyvärinen L, Walthes R, Jacob N, Lawrence L, Chaplin K. Delayed visual development: development of vision and visual delays. American Academy of Ophthalmology, 2017.
32.   Podsiadlo D, Richardson S. The timed "Up & Go": a test of basic functional mobility for frail elderly persons. *J Am Geriatr Soc* 1991; 39(2): 142-148.
33.   Templeton GF. A two-step approach for transforming continuous variables to normal: implications and recommendations for IS research. *Commun Assoc Inf Syst* 2011; 28(1).
34.   Rahi JS, Cable N, Group BCVIS. Severe visual impairment and blindness in children in the UK. *The Lancet* 2003; 362(9393): 1359-1365.
35.   Steinkuller PG, Du L, Gilbert C, Foster A, Collins ML, Coats DK. Childhood blindness. *J AAPOS* 1999; 3(1): 26-32.
36.   Kong L, Fry M, Al-Samarraie M, Gilbert C, Steinkuller PG. An update on progress and the changing epidemiology of causes of childhood blindness worldwide. *J AAPOS* 2012; 16(6): 501-507.
37.   Ministry of Health and Welfare. Physicians Act. Taiwan: Laws & Regulations Database of The Republic of China, 2020.
38.   Itzkowitz A, Kaplan S, Doyle M, et al. Timed up and go: Reference data for children who are school age. *Pediatr Phys Ther* 2016; 28(2): 239-246.
39.   Giagazoglou P, Amiridis IG, Zafeiridis A, Thimara M, Kouvelioti V, Kellis E. Static balance control and lower limb strength in blind and sighted women. *Eur J Appl Physiol* 2009; 107(5): 571-579.
40.   Chen EW, Fu AS, Chan K, Tsang WW. Balance control in very old adults with and without visual impairment. *Eur J Appl Physiol* 2012; 112(5): 1631-1636.
41.   Lai DC, Tseng YC, Guo HR. Trends in the prevalence of childhood disability: Analysis of data from the national disability registry of Taiwan, 2000–2011. *Res Dev Disabil* 2013; 34(11): 3766-3772.
42.   Lai DC, Tseng YC, Hou YM, Guo HR. Gender and geographic differences in the prevalence of intellectual disability in children: Analysis of data from the national disability registry of Taiwan. *Res Dev Disabil* 2012; 33(6): 2301-2307.
43.   Lin LP, Lin JD. Perspectives on intellectual disability in Taiwan: epidemiology, policy and services for children and adults. *Curr Opin Psychiatry* 2011; 24(5): 413-418.

44.   Lin JD, Yen CF, Loh CH, Li CW, Wu JL. Rehabilitation service utilization and determinants
among people with an intellectual disability: Preliminary findings in Taiwan. *Disabil Rehabil* 2006;
28(23): 1499-1506.
45.   Chou YC, Chiao C, Fu LY. Health status, social support, and quality of life among family carers
of adults with profound intellectual and multiple disabilities (PIMD) in Taiwan. *J Intellect Dev Disabil*
2011; 36(1): 73-79.
46.   Das M, Spowart K, Crossley S, Dutton GN. Evidence that children with special needs all require
visual assessment. *Arch Dis Child* 2010; 95(11): 888-892.