A steady trickle-down from metro districts and improving epidemic-parameters characterize the increasing COVID-19 cases in India

Santosh Ansumali[a,e], Aloke Kumar[b], Samarth Agarwal[a], H. J. Shashank[a],
Meher K. Prakash[1c,e]

[a]*Engineering Mechanics Unit, Jawaharlal Nehru Centre for Advanced Scientific Research, Jakkur, Bangalore, India*
[b]*Department of Mechanical Engineering, Indian Institute of Science, Bangalore, India*
[c]*Theoretical Sciences Unit, Jawaharlal Nehru Centre for Advanced Scientific Research, Jakkur, Bangalore, India*
[d]*Sankhya Sutra Labs, Manyata Embassy Business Park, Bengaluru, Karnataka, India*
[e]*VNIR Biotechnologies Pvt Ltd, Bangalore Bioinnovation Center, Helix Biotech Park, Electronic City Phase I, Bangalore, India*

Abstract

Background **By mid-September of** 2020**, the number of daily new infections in India have crossed** $95,000$**. To facilitate an intuition for the spatio-temporal development of the pandemic and to help resource deployment planning, we analyze and describe how the disease burden almost-predictably shifted from large metropolitan districts to suburban districts.**

Methods **We gathered the publicly available granular data from** 186 **different districts (equivalent of counties) on their COVID-19 infections and deaths during the 15 April to 31 August** 2020 **period. These districts presented an active case burden of** $559,566$ **and a cumulative** $2,715,656$ **infections as of August 31. The epidemiological data of these districts was fit to a susceptible-asymptomatic-infected-recovered-dead (SAIRD) model and the underlying epidemic parameters for each of these districts during the course of 4 months was estimated. We validated these parameters against known epidemiological characteristic distributions**

---

[1]Corresponding Author: meher@jncasr.ac.in

and analyzed them to understand their changes in space-time during the pandemic.

Findings The center of the burden of the current-active infections which on May 15 was in the large metro districts with most international access shifted continuously and smoothly shifted towards districts which could be accessed by domestic airports and by trains. A linear trend-analysis showed a continuous improvement in most epidemic parameters consistently across the districts with four categories of accessibility from an international travel perspective - large metro, metro, urban and sub-urban districts. The reproduction numbers improved from 1.77 ± 0.58 on May 15 to 1.07 ± 0.13 on August 31 in large metro districts (p-Value of trend 0.0001053); and from 1.58 ± 0.39 on May 15 to 0.94 ± 0.11 on August 31 in sub-urban districts (p-Value of trend 0.0067). The recovery rate per infected person per day improved from 0.0581 ± 0.009 on May 15 to 0.091 ± 0.010 on August 31 in large metro districts (p-Value of trend 0.26×10^{-12}); and from 0.059 ± 0.011 on May 15 to 0.100 ± 0.010 on August 31 in sub-urban districts (p-Value of trend 0.12×10^{-16}). The death rate of symptomatic individuals which includes the case-fatality-rate as well as the time from symptoms to death, consistently decreased from 0.0025 ± 0.0014 on May 15 to 0.0013 ± 0.0003 on August 31 in large metro districts (p-Value of trend **0.0010**); and from 0.0018 ± 0.0008 on May 15 to 0.0014 ± 0.0003 on August 31 in sub-urban districts (p-Value of trend 0.2789).

Interpretation As the daily infections continue to rise at a national level, it is important to notice a 'local-flattening' in larger metro districts, and a shift of the pandemic-burden towards smaller sized districts in a clear hierarchical fashion of accessibility from an international travel perspective. The pandemic burden shifting towards remotely accessible regions, with possibly lesser health care facilities, is a call for attention to the re-organization of resources.

Funding None

Keywords: Case-burden, Epidemic diffusion, Hotspots, granular, time-varying epidemic-parameters, COVID-19, India

Introduction

In the COVID-19 pandemic that began in December 2019, each month witnessed the critical rise of infections in completely different countries. The initial declaration of national emergency which was a "one-size fits all" was quickly irrelevant in many countries as each state or county perceived the need for stringency differently. However, now the focus is it identify the next hotspot[1, 2]. The differences arose mainly because even within the first wave of the pandemic, the solidarity extended through inter-state loans of ventilators demonstrated how at any given time the geographical regions in different phases of the pandemic, rising, stable or declining, perceived the threat differently from others. As the rise of infections in different geographical regions is asynchronous, the critical care burden shifts dynamically with each region attempting to achieve its 'local-flattening' of the peak at a different time.

Epidemiological models have focused on making predictions of the rise of infections, and a peak of critical care burden on a country or a state basis. There have been longitudinal studies[3, 4] demonstrating how the reproduction number varied over time in the different states or countries and possibly correlating it to the effects of lockdowns. Further, a few cross-sectional studies correlated the hotspots in the spread of infections in the initial phase to the traffic flow out of Wuhan.[5] These studies mostly focused on one region at a time, either on estimating parameters, or making projections. In this work, we build a summarizing perspective of the geographical development of the pandemic, by analyzing a cumulative 2.71 million infections and 52 thousand deaths of COVID-19 from India over a period of four and a half months.

To examine how the geographic spread of COVID-19, we identified the districts of interest based on case-burden, estimated the time-varying epidemiological parameters for these regions, categorized these regions according to the ease of travel access in the hierarchical order of availability of international flights, domestic flights, trains or other quaternary modes of transport, and analyzed the trends in parameters over time in each of these categories. The data shows predictable shifts to remote regions. The infections spreading to newer and under-catered locations presents both an opportunity and a challenge, an asynchronous peak of critical requirements which can be addressed both by manufacturing and re-organizing the resources on an *ad hoc* basis.

Research in context

Evidence before this study

We searched *pubmed* and *medRxiv* up to 20 September 2020 with the keywords "COVID-19", "time variation or longitudinal", "cantons or states", "hotspots", "diffusion" "active-cases". The published research thus far focused mainly on reporting either the time variation of reproduction number at the state-level, or the predictions of rise of infections at the canton level. However, from these constantly updating predictions with new hotspots appearing, there is no continuity of knowledge and possibility to obtain a comprehensive picture of the development of the pandemic. A study correlating the number of infections in different regions in China to the traffic-flow out of Wuhan, and another describing the development of hotspots along the highways in Brazil describe some aspects of this geographical spread. Firstly, a similar analysis for with granular Indian data has been missing. Further, the questions of how the center of the active-case burden diffused and the spatio-temporal trends in epidemiological parameters underlying the large number of infections have not been the focus of these earlier studies.

Added value of this study

Our study begins with the understanding that the different geographic regions such as districts (or counties) may be in different phases of the pandemic, and mixing of population within them happens more likely than from outside. Considering this geographical heterogeneity, and the ease of access to these different districts, we develop a summary how the pandemic evolved in the regions of different accessibility. Predictable trends in the case-burden diffusion are identified.

Implications of all the available evidence

Authorities should understand the shift in the dynamics of the critical-care requirements, not by geographically contiguous regions, but by ease of travel or by common economic interests which guided such ease of travel. Strategies should be driven by knowing where the pandemic burden is likely to move, before it happens. The trend of the case-burden predictably shifting towards suburban districts can be useful in re-organizing the available resources as per the needs.

Methods

COVID-19 Data

The district-wise COVID-19 data was gathered from a publicly available data repository: *http://www.covid19india.org.* 186 districts with at least 20 daily new infections on 31 August 2020 were used in the analysis. These 186 districts accounted for $2,715,656$ of the $3,687,939$ cumulative infections in India, and had an active case burden of $559,566$ of the $785,127$ total active infections in India. These 186 districts out of the 739 districts in India, span across 22 different states which are currently governed by many different political parties and account for a 696 million population in 2020, estimated from the 2001 and 2011 census data. The daily new infections, daily active cases, daily deaths data from these 186 districts was used in all of our analyses.

Model

An important feature of COVID-19 is the role played by asymptomatics in the spread of the disease.[6] Classical SIR models ignores this important aspect of the disease and thus one should work with models where asymptomatic fraction is explicitly included in the model. A susceptible-asymptomatic-infected-recovered-dead (SAIRD) framework, an extension of an SAIR model we developed earlier[7] by including the deaths, was used to model the spread of the pandemic. The compartmental nature of the model is shown schematically in Fig.(1A) and the governing equations are given in Fig.(1B). The asymptomatics in this framework are assumed to include both the pre-symptomatics who get converted and those who will continue to remain asymptomatic until their recovery. The rates of recovery and transmission of infections of the asymptomatic and the symptomatic individuals are assumed to be the same. The model thus relies on four different epidemiological parameters - infection rate (β), recovery rate (γ), death rate (γ_D), asymptomatic to symptomatic conversion (δ), which we extracted from the data as follows.

Parameter Estimation

Even in the simplest scenario of the exponential growth with of infections I with time, we found that choosing a linear fit between dI/dt and I led to a better parameter estimation than an exponential fit between I and t, although both are mathematically equivalent.[8] The parameter estimation

Figure 1: *A.* A schematic of the SAIRD model and its *B.* modelling equations used in the present analysis. The SAIR model on which this scheme is based is described in Ref. [7]. The parameters used in these equations are assumed to be constant over different segments of the pandemic trajectory (piece-wise continuous) and were estimated by fitting the predictions for infections, recoveries, and deaths over time.

scheme we evolved[7] for all the parameters thus relies on rearranging the underlying equations in a way that allowed us to provide most reliable estimates despite the uncertainties and noise in the data. The time-varying epidemiological parameters were estimated for the different districts. The infection spread-rate (β), recovery-rate (γ) and the death-rate (γ_D) were estimated in a time-dependent way, identifying time periods over which a single parameter best describes the observations. The (pre- or) asymptomatic to symptomatic conversion rate (δ) was assumed to be a constant for each district.

$\gamma(t)$ **estimation.** To estimate γ we note that, $dR_I/dt = \gamma I$, which can be easily integrated to obtain $R_I(t_2) - R_I(t_1) = \gamma \left(\int_{t_1}^{t_2} I dt \right)$. However, it is well understood that γ changes over time owing to changes in policies and behavioral patterns. We note that the plot for R_I vs. $\int I dt$ is piecewise linear and the value of γ for different periods can be calculated using simple linear regression and to estimate the slope and therefore the γ.

$\gamma_D(t)$ **estimation.** The estimation for γ_D then follows in a similar manner. $dDdt = \gamma_D I$, which is integrated to obtain $D(t_2) - D(t_1) = \gamma_D \left(\int_{t_1}^{t_2} I dt \right)$.

$\beta(t)$ **estimation.** To estimate β we employ $\log I(t) \sim (\beta - \gamma) \, t$.

δ **estimation.** The values of the hidden asymptomatics and δ are determined using a grid search and then choosing the values which provide the best fit. The estimates of δ for the 186 districts were 0.34 ± 0.02, which corresponds

to a time-scale of 4.23 ± 0.48 days. This is in line with numbers provided in Ref[9].

R_t **estimation.** The reproduction ratio, R_t, defined as the ratio, $\beta(t)/\gamma(t)$, is an effective measure of the current status of the spread of the infection and its containment.

Travel accessibility of districts

The degree of travel-accessibility of the districts was defined using *pre-pandemic* data. The international and domestic flight traffic to the different airports was gathered from the Airports Authority of India website (*https://www.aai.aero/sites/default/files/traffic-news/Mar2k20Annex2.pdf*), and was assigned to the districts. The train station with the highest through-traffic in each districts was noted using the data from the ClearTrip website (*https://www.cleartrip.com/trains/stations/list*). The districts were ranked 1 to 8 based on their accessibility through flights or trains and grouped into four categories - *Large metro districts:* International airports with an *annual* traffic of more than $25,000$ flights (1), more than 5,000 flights (2), with any direct international flights (3), *Metro districts:*more than 5000 *annual* domestic flights (4), any domestic flights (5), *Urban districts:* more than 50 trains stopping in the district (6), more than 20 trains (7), and *Sub-urban districts:* all districts with other quaternary means of reaching (8).

Results

Hotspots are not contiguous

In the period of study April 15-August 31, the number of infections continued to rise in India from around $1,000$ daily new infections to around 80,000 daily new infections. The initial epicenters were in the large metropolitan cities of Delhi (from the state of Delhi) and Mumbai (from the state of Maharashtra). With lockdowns and non-pharmaceutical interventions, the case burden in these cities reduced, but with a concomitant rise of COVID-19 cases in many smaller districts. The active-case burden thus diffused out of large metro districts. However, mapping the hotspot districts in the geographic map of the India certainly does not immediately reveal how the infections increased or spread. An example of the development of hotspots in the state of Maharashtra is shown in Figure 2. One may immediately notice two exceptions - that newer hotspot districts need not be contiguous

128 with the existing hotspots and further some of the contiguous districts did
129 not develop into hotspots. Thus it is apparent that a plain spatial diffusion
model is not sufficient.

Figure 2: Evolution in time of the pandemic over the districts as seen in the geographic map
of Maharashtra. The active case burden is approximated to the nearest cluster indicated
in the color-bar. One can see that the the spread of the infections is at times continuous,
and on some other occasions begins in districts disconnected from the current hotspots.

Correlating hotspots with international travel accessibility

Since COVID-19 infections started with import of cases, we asked if the down-
stream development of newer hotspots is also related to the ease of accessing
a district from an international perspective by various means of transport.
We ranked the districts 1 to 8 depending upon the level of accessibility, and
further categorized the districts as large metro, metro, urban and sub-urban
districts using a hierarchical accessibility by international airports, domestic
airports, train hubs and other quaternary means of transport, respectively.
In Figure 3, we show the number of active COVID-19 cases during the period

May 15 - August 31 across the districts in four broad accessibility categories. The figure clarifies that the hotspots in each state, represented by a sector, spread radially outwards suggesting a spread away from the districts with international airports towards districts which are only accessible only via trains. The states where the most accessible regions have not yet recorded a lot of infections, did not also a downstream spread of infections into their remote districts. The graphic highlights the significant role of importation of the cases, and a weak or poor-mixing across different states initially due to lockdowns, followed by quarantines upon inter-state travels during many stages of lockdown release.

Figure 3: Evolution in time of the pandemic over the districts. Each sector represents the state denoted by its 2 letter code, such as DE for Delhi, MH for Maharashtra. The 8 different layers from inside to outside are the ones in the decreasing order of accessibility. These 8 layers are further clustered as those having international airports, having only domestic airports, connected only by trains and have no train or airplane connections, and are separated by the red-circles. In each state, the hotspots continue to move outwards as the time progresses. Two segments labeled OT are meant other states, which are not included in the analysis.

The pandemic-burden is shifting away from larger metros

As the infections gain a greater geographic spread, it is important to know where the current center of the disease burden is, and if there is a predictable

shift in its position. By weighing the number of currently active cases with the accessibility level (1 to 8) of the district, we defined the current center of active burden as $(\sum_{districts} I_{district} \times \text{accessibility})/(\sum_{districts} I_{district})$. The result shown in Figure 4 suggests a continuous and smooth drift of this center of active cases away from the large metro districts towards least accessible districts. The districts which are not easily accessible, are also usually the ones with lesser health care resources. The result underscores the need for deployment of the resources towards regions which do not have adequate critical health care support. We analyze the general trends in each of the different regions, to understand if a re-organization of the resources or a creation of newer resources will be a solution to address these critical needs.

Figure 4: The current burden of the active cases is calculated by weighing the active cases in each district by its accessibility code (1-internationally accessible large metro; 8-least accessible district, as indicated in Figure 3). One can see a clear shift of the case burden away from the large cities. The accessibility of the districts through international, domestic airports and trains are indicated.

Infections continue to rise, but with lower Reproduction numbers.

As a country India has not yet reached the peak of COVID-19 infections and the number of newly reported cases as well as the number of currently active cases continue to rise (Figure 2A). However, one can notice from the slopes that the rate of infection rise continues to decrease in districts of all levels of

accessibility. The time varying reproduction number (R_t) estimates from our analysis over the May 15-August 31 period which witnessed phased release of lockdowns suggest values which are much less than the basic reproduction number estimates between 2.5 to 4. The average reproduction number across the districts of different accessibility showed a decreasing trend (Figure 2B). A linear trend analysis suggested the large metro and sub-urban districts have a reduction in R_t with p-Values of 0.0001053 and 0.0067 respectively. While the average reproduction numbers appear to reduce even in metro and urban districts, the trends were not significant (p-Values of 0.4634 and 0.4618 respectively).

In order to distinguish if the slow down of infections occurred because of a herd immunity or because of the changes in behavioural patterns, we examined the infected percentage (=cumulative infections/population × 100%) of the population at which the R_t dropped below one. The infected percentages vary significantly, for example in the two contiguous districts of Mumbai and Pune, these values were 0.25% symptomatic and 1.08% including asymptomatics in Mumbai and 0.7% symptomatic and 2.21% including asymptomatics in Pune, which suggests that the slowdown of infections is achieved because of changes in behavioural patterns and an exercise of caution is still required to maintain the pandemic in control.

Recovery and death rates improved

The time to recovery from the time of appearance of symptoms for the symptomatic patients ($1/\gamma$) estimated for the 186 districts as 16.18±1.01 on May 15 was consistent with the 24.7 (22.9-28.1) that was reported in the early days of infections from China.[10]. However, by August 31, the recovery time showed an improvement as (γ) decreased to 10.08±0.69 on August 31. The time variation of recovery rates (γ) is shown in Figure 2C, and the trend analysis suggested an increase in the recovery rates with p-values $0.2667 \times 10^{-12}, 0.5504 \times 10^{-3}, 0.9180 \times 10^{-37}, 0.1250 \times 10^{-16}$ in the large metro, metro, urban and sub-urban districts. Similarly death rates γ_D which is a product of the case fatality rate (reported as $\approx 2\%$ in India, and consistent with an ≈ 10 days time to death) with the rate of death also showed an improvement over the time with p-Values of $0.0010, 0.0122, 0.6272 \times 10^{-3}$, 0.2789 respectively in the large metro, metro, urban and sub-urban districts.

Figure 5: The trends in different epidemiological parameters *A*. Number of daily new infections, *B*. Reproduction number, *C*. Recovery rate (γ) *D*. Death-rate (γ_D) are shown for the districts categorized by their accessibility is shown. The trends across the different districts groups are consistent. The reproduction rates, recovery and death rates are all improving, despite the increase in the new infections.

Comparison with other studies

In an analysis of the spread of SARS across countries,[11] it was realized that a smooth spatial diffusion of pandemic which was sufficient to describe Black Death[12] could not address a quicker reach of the epidemic to many cities. It was demonstrated that the time taken for the epidemic arrival time to the different cities was correlated not to the geographic distance from the epicenter, but rather to an effective distance defined by the degree of connectivity between the different international airports. The work thus introduced the concepts of development of epidemics that go beyond simple spatial diffusion.[12] In a cross-sectional analysis, the development of case-load of the infections in the different hotspots in China was correlated to the outflow of traffic from Wuhan in the period January 1 to January 24.[5] In another recent study on COVID-19 in Brazil, the connectivity between different cities via highways was attributed to be one of the reasons for super-spreading across major cities.[13] However, the work did not establish any patterns for the spread of the active-case burden or its flattening,[14] which is extremely important in an epidemic or a pandemic from the public health point-of-view.

While the present work is in line with all three works noted above in term of travel connectivity, there are several unique aspects we discovered in our study of the COVID-19 data from India. Possibly the global spread of pandemic from Wuhan to different major international destinations may have followed the similar pattern of effective-distance. [11] However, the emphasis, from the perspective of incidence within a country and critical resource planning is on the granular data within the country. Further, unlike the travel outflow which was obtained by tracking mobiles,[5] the accessibility we use via airplane and train-connectivity data from pre-pandemic period is believed to be a surrogate for the degree of connectivity and mixing among the populations. In the *network science* parlance, this is an emphasis on the centrality-measures which define the relative importance of the different cities that are being connected, by-passing the actual connectivity data which is harder to obtain, especially because our study was not limited to a single phase such as the January 1 to January 24.[5] Finally, by developing the hierarchical accessibility network, we could show trends such as the radial spreading out towards remote areas in each state and a smooth variation of the current-case-burden in Figure 4. Figures 3 and 4, in our opinion go beyond the immediate requirements of how the infections are rising in a specific

region and serve to provide a wholistic perspective on the development of the pandemic as well to help planning for allocating or re-organizing critical care or test resources towards less catered cities.

Conclusions

In summary, we show predictable trends of improvement of disease parameters, and spread of COVID-19 infection burden away from easily accessible metros with large international airports. The continuing number of cases, and the predictable drift suggests the need for planning or even reorganizing resources to districts which are down-stream in international access. A lack of correlation between the slow-down of infections and the extent of current infections suggests that the caution and change in behavioral patterns need to continue if the COVID-19 has to be kept in check.

Acknowledgements

We acknowledge discussions with P. Sunthar. MKP and SA acknowledge Aman Sharma and Fauzia Javed for help in collecting the airport and train traffic data.

Conflict of interests

The authors declare no conflicts of interest.

References

[1] J. Cohen, Scientists are racing to model the next moves of a coronavirus that's still hard to predict, Science 7 (2020).

[2] J. Budd, B. S. Miller, E. M. Manning, V. Lampos, M. Zhuang, M. Edelstein, G. Rees, V. C. Emery, M. M. Stevens, N. Keegan, et al., Digital technologies in the public-health response to covid-19, Nature medicine (2020) 1–10.

[3] S. Flaxman, S. Mishra, A. Gandy, H. J. T. Unwin, T. A. Mellan, H. Coupland, C. Whittaker, H. Zhu, T. Berah, J. W. Eaton, et al., Estimating the effects of non-pharmaceutical interventions on covid-19 in europe, Nature 584 (2020) 257–261.

[4] M. Chinazzi, J. T. Davis, M. Ajelli, C. Gioannini, M. Litvinova, S. Merler, A. P. y Piontti, K. Mu, L. Rossi, K. Sun, et al., The effect of travel restrictions on the spread of the 2019 novel coronavirus (covid-19) outbreak, Science 368 (2020) 395–400.

[5] J. S. Jia, X. Lu, Y. Yuan, G. Xu, J. Jia, N. A. Christakis, Population flow drives spatio-temporal distribution of covid-19 in china, Nature (2020) 1–5.

[6] R. Li, S. Pei, B. Chen, Y. Song, T. Zhang, W. Yang, J. Shaman, Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (sars-cov2), Science (2020).

[7] S. Kaushal, A. S. Rajput, S. Bhattacharya, M. Vidyasagar, A. Kumar, M. K. Prakash, S. Ansumali, Estimating hidden asymptomatics, herd immunity threshold and lockdown effects using a covid-19 specific model, arXiv preprint arXiv:2006.00045 (2020).

[8] M. K. Prakash, S. Kaushal, S. Bhattacharya, A. Chandran, A. Kumar, S. Ansumali, Minimal and adaptive numerical strategy for critical resource planning in a pandemic, Physical Review E 102 (2020) 021301.

[9] S. A. Lauer, K. H. Grantz, Q. Bi, F. K. Jones, Q. Zheng, H. R. Meredith, A. S. Azman, N. G. Reich, J. Lessler, The incubation period of coronavirus disease 2019 (covid-19) from publicly reported confirmed cases: estimation and application, Annals of internal medicine 172 (2020) 577–582.

[10] R. Verity, L. C. Okell, I. Dorigatti, P. Winskill, C. Whittaker, N. Imai, G. Cuomo-Dannenburg, H. Thompson, P. G. Walker, H. Fu, et al., Estimates of the severity of coronavirus disease 2019: a model-based analysis, The Lancet infectious diseases (2020).

[11] D. Brockmann, D. Helbing, The hidden geometry of complex, network-driven contagion phenomena, science 342 (2013) 1337–1342.

[12] J. V. Noble, Geographic and temporal development of plagues, Nature 250 (1974) 726–729.

[13] M. A. L. Nicolelis, R. L. G. Raimundo, P. S. Peixoto, C. S. de Andreazzi, How super-spreader cities, highways, hospital bed availability,

301 and dengue fever influenced the covid-19 epidemic in brazil, medRxiv
302 (2020).

303 [14] R. M. Anderson, H. Heesterbeek, D. Klinkenberg, T. D. Hollingsworth,
304 How will country-based mitigation measures influence the course of the
305 covid-19 epidemic?, The Lancet 395 (2020) 931–934.