Multi-trait genome-wide association study identifies new loci associated with sleep apnoea risk

Adrian I. Campos[1,2], Nathan Ingold[1,3], Yunru Huang[4,¶], Brittany L. Mitchell[1,3], Pik-Fang Kho[5], Xikun Han[6], Luis M. García-Marín[1], Jue-Sheng Ong[1], 23andMe Research Team[4], Matthew H. Law[1,3], Jennifer S. Yokoyama[7,8,9], Nicholas G. Martin[1], Xianjun Dong[10,11], Gabriel Cuellar-Partida[4], Stuart MacGregor[1], Stella Aslibekyan[4] and Miguel E. Rentería[1,2,3*]

1. Department of Genetics & Computational Biology, QIMR Berghofer Medical Research Institute, Brisbane, QLD, Australia.
2. School of Biomedical Sciences, Faculty of Medicine, The University of Queensland, Brisbane, QLD, Australia.
3. School of Biomedical Sciences, Faculty of Health, Queensland University of Technology, Brisbane, Queensland, Australia.
4. 23andMe, Inc., Sunnyvale, CA, USA.
5. Division of Cardiovascular Medicine, Department of Medicine, Stanford University School of Medicine, Stanford, CA, USA.
6. Program in Genetic Epidemiology and Statistical Genetics, Harvard University T.H. Chan School of Public Health, Boston, MA, USA.
7. Memory and Aging Center, University of California, San Francisco, San Francisco, CA, USA.
8. Weill Institute of Neurosciences, Department of Neurology, University of California, San Francisco, San Francisco.
9. Global Brain Health Institute, University of California, San Francisco, San Francisco, CA, USA
10. Genomics and Bioinformatics Hub, Brigham and Women's Hospital, Boston, MA, USA.
11. Department of Neurology, Brigham & Women's Hospital and Harvard Medical School, Boston, MA, USA.
¶. Current address: Genentech Inc., South San Francisco, CA, USA.
* **Correspondence:** Dr Miguel E. Rentería, Department of Genetics and Computational Biology, QIMR Berghofer Medical Research Institute, Locked Bag 2000, Royal Brisbane Hospital, Herston, QLD 4029, Australia. Email: miguel.renteria@qimrberghofer.edu.au

ABSTRACT

Background: Sleep apnoea is a disorder characterised by periods of halted breathing during sleep. Despite its association with severe health conditions such as cardiovascular disease, liver problems and metabolic syndrome, the aetiology of sleep apnoea remain understudied, and previous genetic analyses have not identified many robustly associated genetic risk variants.

Methods: We performed a genome-wide association study (GWAS) meta-analysis of sleep apnoea across five cohorts ($N_{Total}=523,366$) and multi-trait analysis of GWAS (MTAG) to boost statistical power, leveraging the high genetic correlation between sleep apnoea and snoring. GWAS and MTAG results were adjusted to control for the effects of body mass index (BMI) using the multi-trait-based conditional & joint analysis (mtCOJO) method. Lead hit replication was conducted in an independent sample GWAS from 23andMe, Inc. that included BMI as a covariate ($N_{Total}=1,477,352$; $N_{cases}=175,522$). Lastly, we explored genetic correlations with other complex traits and performed a phenome-wide screen for causally associated phenotypes using the latent causal variable method.

Results: Our MTAG with snoring uncovered a total of 49 independent significant sleep apnoea loci; adjusting for BMI showed that approximately half of those loci act via their effects on BMI/obesity. Twenty-nine variants replicated in the 23andMe cohort. Several complex traits, including multisite chronic pain, diabetes, eye disorders, high blood pressure, osteoarthritis, chronic

obstructive pulmonary disease, and BMI-related traits were genetically correlated with sleep apnoea.

Conclusions: Our study identifies multiple robust genetic markers for sleep apnoea, increasing our understanding of the aetiology of this condition and its relationship with other complex traits.

Keywords: Sleep apnoea, genetics, GWAS.

INTRODUCTION

Sleep apnoea is a disorder characterised by episodes of halted breathing during sleep, which leads to frequent arousal and intermittent hypoxia[1]. The most common type of sleep apnoea is obstructive sleep apnoea (OSA), which may affect up to 47% of adults and ~1% of children[2,3]. Central sleep apnoea (CSA) can affect ~0.9% of adults and is also characterised by a lack of breathing and oxygenation during sleep, but without the characteristic respiratory efforts observed in OSA[3,4]. Remarkably, the prevalence of sleep apnoea tends to be underestimated because those affected cannot gain awareness about their condition or may confuse it with habitual snoring.

OSA is caused by a reduced function of pharyngeal dilator muscles, brought about by the onset of sleep, causing the collapse of the upper airways and subsequent hypopnea or apnoea[5,6]. CSA can be caused by a lower respiratory drive, opioid use, obesity, lesions to the central nervous system, and impaired reactivity to carbon dioxide (CO_2), among others[4]. A high degree of overlap in characteristics between the sleep apnoea subtypes makes them hard to distinguish. Notably, sleep apnoea also leads to mental and physical fatigue, which has been shown to increase the risk of motor accidents[7], decrease mental wellbeing and overall quality of life[8]. In addition to these direct harmful effects, sleep apnoea has also been associated with an increased risk of obesity, hypertension[9], stroke[10] and increased levels of reactive oxygen species in blood, which increases oxidative stress in the body[11,12].

The proportion of sleep apnoea risk attributed to genetic factors (i.e., heritability) has been estimated between 35 and 75%[13,14]. Familial aggregation seems to be partially independent of weight[15]. Increased understanding of the genetic architecture of sleep apnoea could help generate risk prediction models and provide the necessary groundwork for developing interventions and therapies. In addition, having information on the effect of genetic variants on sleep apnoea risk could enable us to infer its causal relationship with other conditions using methods such as Mendelian randomisation[16]. Although some candidate gene studies for sleep apnoea have yielded some putatively associated genes[17,18], genome-wide association studies (GWAS) have failed to replicate those associations[19–21]. In fact, GWAS have identified very few genome-wide significant loci robustly associated (i.e., with evidence for replication in an independent cohort) with sleep apnoea to date.

Sleep apnoea is likely a highly polygenic trait, with many variants of small effect size contributing to the genetic liability of developing this condition. Thus, most studies with modest sample sizes will be underpowered to identify most of these risk variants and are susceptible to false-positive associations. Furthermore, the number of diagnosed cases of sleep apnoea within existing large population cohorts is low. In a sample of 500,000 individuals, the expected number of sleep apnoea cases (assuming a conservative prevalence of ~5%) would be ~25,000. However, in the UK

Biobank (~500,000 individuals), only ~8,000 sleep apnoea cases have been recorded. This is likely explained by the fact that sleep apnoea is recognised as an underdiagnosed condition[22,23]. Underdiagnosis reduces power as several cases would be considered unaffected controls in a formal analysis. Thus, combining large samples through meta-analysis and replicating findings in large, independent studies are essential to achieving reliable results.

We have conducted a GWAS meta-analysis of sleep apnoea across five cohorts. Also, we employed Multi-trait Analysis of Genome-Wide Association Summary Statistics (MTAG) to combine our sleep apnoea meta-analysis with a snoring meta-analysis across five cohorts to boost power by leveraging the high genetic correlations between sleep apnoea and snoring[24]. We also performed additional sensitivity analyses to control for the effect of BMI and identify loci associated with sleep apnoea independently from BMI. We characterised the genetic architecture of sleep apnoea by performing gene-based tests and genetic correlation analyses. Finally, the different GWAS results were validated by polygenic prediction using a leave-one cohort out cross-validation (LOO) framework and by examining evidence for replication in a GWAS for self-reported sleep apnoea diagnosis in an independent sample from 23andMe, Inc. We refer to our results broadly as sleep apnoea, although they likely represent a closer proxy to OSA given its higher prevalence than CSA.

METHODS

Sample information and phenotype ascertainment

This study analysed GWAS data from five cohorts from the UK (UK Biobank; UKB), Canada (Canadian Longitudinal Study of Aging; CLSA)[25,26], Australia (Australian Genetics of Depression Study; AGDS), the USA (Mass General Brigham Biobank), and Finland (FinnGen). Sample sizes and the number of cases and controls are listed in **Table 1**. Sleep apnoea cases were defined for each cohort using either self-report or diagnostic codes available in electronic health records (See **Supplementary Methods** for individual cohort details). Self-reported snoring cases were excluded from the analyses for the sleep apnoea GWAS across the UKB, CLSA and AGDS cohorts. An overview of the analysis pipeline used for sleep apnoea discovery analysis is available in **Supplementary Figure S1**.

GWAS

All GWAS studies included the following covariates: age, sex, batch (where relevant), and genetic ancestry principal components derived from genotype data. Standard quality control filters were applied at both the sample and variant levels. Variants were excluded from the analyses if they had a low minor allele frequency (MAF<0.01) or low imputation quality score (INFO<0.6). Individuals were excluded based on excess missingness, heterozygosity, or evidence from a deviation from European ancestry based on genetic principal components. For each cohort, a GWAS was performed using logistic regression models and including random effects to account for cryptic relatedness where relevant (**Supplementary Methods**). For the UKB snoring GWAS, we used the summary statistics from our previously published GWAS for snoring[27]. FinnGen GWAS results for sleep apnoea and snoring were obtained from the open-access FinnGen resource (http://r3.finngen.fi/).

GWAS meta-analyses

Sample-size weighted (P-value-based) meta-analyses for sleep apnoea and snoring were performed (separately for each phenotype) across the five cohorts described above using METAL (v2020-05-05).[28] Studies were weighted according to their effective sample size as described by the formula: $\square\square\square\square = 4/(1/\square\square\square\square\square\square + 1/\square\square\square\square\square\square\square\square\square)$ as recommended for studies with different levels of case-control imbalance (**Supplementary Methods**).

Multi-trait GWAS analyses

Multi-trait Analysis of Genome-Wide Association Summary Statistics (MTAG) was used to boost power for discovering sleep apnoea associated loci. MTAG performs a generalized meta-analysis of GWAS summary statistics for different but genetically correlated traits and accounts for sample overlap[24]. We performed MTAG analyses combining our sleep apnoea and snoring meta-analyses for this study. This is motivated by 1) the high genetic correlation between these traits (rG ~ 0.8)[27] and 2) the observation that snoring is one of the primary symptoms of obstructive sleep apnoea, the most common type of sleep apnoea[6].

BMI adjustment

Given the relationship between sleep apnoea, snoring, and BMI, we performed a secondary analysis adjusting our GWAS results (both the meta-analysis and the MTAG) for the effect of BMI. To adjust for BMI while avoiding biases due to collider bias, that is, the emergence of a spurious

association between a pair of variables when a common outcome is modelled as a covariate[29], we used multi-trait-based conditional and joint analysis (mtCOJO)[30,31].

23andMe replication GWAS

We sought to replicate variants identified in the discovery phase in an independent sample of participants from the 23andMe cohort (N=1,477,352). Cases were ascertained based on the question "*Have you ever been diagnosed with, or treated for any of the following conditions*?" with one of the choices being "*Sleep apnoea*" (Yes = 175,522; No = 1,301,830). Methods and results from this GWAS have been presented at the 2018 American Society for Human Genetics annual conference[32]. Briefly, a logistic regression GWAS was performed using sleep apnoea as the dependent variable while adjusting for sex, age, BMI, genetic principal components and genotype platform. Participants provided informed consent and participated in the research online, under a protocol approved by the external AAHRPP-accredited IRB, Ethical & Independent Review Services (E&I Review). Only unrelated participants of European ancestry who provided consent were included in the analysis. The full GWAS summary statistics for the 23andMe discovery data set can be made available through 23andMe to qualified researchers under an agreement with 23andMe that protects the privacy of the 23andMe participants. Please visit research.23andme.com/collaborate/#publication for more information and to apply to access the data. We defined evidence of replication after correcting for the number of significant variants with data available for replication per GWAS analysis. That is p<0.01 for the sleep apnoea meta-analysis, p<0.0016 for the sleep apnoea plus snoring MTAG and p<0.002 for the sleep apnoea plus snoring MTAG adjusted for BMI.

Gene-based association tests and eQTL colocalization

We used the "set-based association analysis for human complex traits" fastBAT method, which performs a set-based enrichment analysis using GWAS summary statistics while accounting for linkage disequilibrium (LD) between SNPs[33]. Statistical significance was defined using the Bonferroni method for multiple testing correction (p<2.07e-6). Genes identified as statistically significant were further assessed for expression quantitative trait loci (eQTL) colocalization using the *COLOC*[34] package in R. Briefly, we integrated our GWAS summary data with *cis*-eQTL data from whole blood, oesophagus, adipose and lung tissue from GTEx V8[35] to estimate the posterior probability that GWAS signals co-occur with eQTL signals while accounting for LD structure. This method estimates the posterior probabilities (PP) for five different scenarios. The scenario of interest is colocalization due to associations with both traits through the same SNPs (PP4). A threshold of PP4>=0.8 was considered as evidence for colocalization of GWAS signals and eQTL signals at the region of interest (**Supplementary Methods**).

S-MultiXcan based eQTL integration

Integration of eQTL with GWAS results interrogates whether the observed associations are consistent with gene-expression changes mediating the trait under study. This study integrated our GWAS results with eQTL data from GTEx using S-MultiXcan [36] as implemented in the Complex Traits Genetics Virtual Lab (CTG-VL). Briefly, this method employs a multiple regression of the phenotype on the predicted expression of a gene across multiple tissues based on eQTL data. When using only GWAS summary statistics, single-tissue associations are performed using S-PrediXcan and joint-effects from the single-tissue results are estimated using an approximation similar to that

of the conditional and joint multiple-SNP analysis[37]. Contrary to the eQTL colocalization described above, this analysis employs the whole GWAS summary statistics and is not restricted only to genes identified using fastBAT or other gene-based tests.

Heritability and genetic correlations
We used LD score regression to estimate the SNP-based heritability (h_{SNP}^2) for the sleep apnoea meta-analysis. Given that samples were not explicitly ascertained for sleep apnoea, we assumed the overall sample and population prevalence for sleep apnoea to be the prevalence estimated across cohorts (0.05), consistent with reported epidemiological estimates[2]. Genetic correlations (rG) between sleep apnoea and 1,522 phenotypes (with available GWAS summary statistics) were estimated using bivariate LD score regression in CTG-VL[38] based on a standard set of HapMap3 variants. The Benjamini-Hochberg FDR at 5% was used to define statistical significance.

Polygenic risk scoring
To assess the external validity of the GWAS, we performed polygenic-based prediction on a target sample of 9,221 unrelated Australian adults from the Australian Genetics of Depression Study[39] (AGDS) with complete data. Briefly, the meta- and MTAG analyses were repeated, leaving out the AGDS cohort to avoid sample overlap. We employed a recently developed method, SBayesR, to obtain the conditional effects of the studied variants, thus avoiding inflation due to correlated SNPs in LD[40]. Sleep apnoea polygenic risk scores (PRS; SA_{PRS}) were calculated using PLINK 1.9. The association between PRS and sleep apnoea in AGDS was assessed using a logistic regression model (python *statsmodels*). SA_{PRS} was the predictive variable of interest, with age, sex and the first ten genetic principal components included as covariates. Finally, binary classifiers based on a logistic regression were built, including either age and sex (base model) or age, sex and the PRS of interest (SA_{PRS} or $SAmtagSnoring_{PRS}$). These classifiers were used to further assess the polygenic predictive ability. The sample was divided randomly into training and testing datasets of equal sizes. Then, the classifier's ability to predict SA was assessed using the area under the receiver operating characteristic (ROC) curve. To avoid potential biases from the random division of training and testing datasets, the procedure was repeated 100 times to estimate a mean area under the curve (**Supplementary Methods**).

Latent Causal Variable analysis
The latent causal variable (LCV) method leverages GWAS summary statistics to estimate whether a causal association can explain a genetic correlation between traits rather than horizontal pleiotropy (i.e., shared genetic pathways)[41–44]. LCV conceptually relies on a latent variable L, assumed to be the causal factor underlying the genetic correlation between the traits [41–44]. LCV estimates the genetic causality proportion (GCP). A higher absolute GCP value indicates more evidence of a causal association among a pair of genetically correlated phenotypes. In contrast, a GCP value of zero would imply that horizontal pleiotropy underlies the genetic correlation between the phenotypes. However, the LCV method will be biased towards the null (a GCP value of 0) if a bi-directional association exists between traits. An absolute value for GCP < 0.60 indicates only partial genetic causality. Multiple testing correction was applied using Benjamini-Hochberg's False Discovery Rate (FDR < 5%). We performed a phenome-wide hypothesis-free LCV analysis to identify traits causally associated with sleep apnoea. Given the limitations of the LCV method (see **Discussion**), we consider this a hypothesis-generating approach. These hypotheses should be tested

in follow-up studies that include relevant Mendelian randomisation analyses and a synthesis of the available literature on the association between sleep apnoea and the trait of interest.

RESULTS

GWAS meta-analysis

The prevalence of both SA and snoring showed some variation across the five different cohorts included in this study (**Table 1** and **Supplementary Material**). Nonetheless, the genetic correlation estimates were high, albeit with large standard errors (**Supplementary Table S1**). These results suggest that the individual cohort GWAS converge onto a similar phenotype. Our meta-analysis identified five independent (LD $r^2<0.05$) genome-wide significant (p<5e-8) loci associated with sleep apnoea (**Figure 1a**). The signals spanned chromosomes 5, 11, 12 and 16 near genes *ANKRD31*, *STK33*, *BDNF*, *KDM2B* and *PRIM1* (**Supplementary Figure S2**). The LD-score regression SNP-based heritability on the liability scale was 13% (S.E.=0.087%). LD-score regression intercept suggested most inflation (λ_{GC}=1.21) was due to polygenic signal (intercept= 1.012, S.E.=0.009) rather than population stratification. Upon adjusting for BMI effects (see **Methods**), one new genome-wide hit on chromosome 15 was identified, but the evidence of association for all other loci was reduced below genome-wide significance (**Figure 1a**). The significant hit after adjusting for BMI was located near genes *HDGFL3*, *TM6SF1* and *BNC1* (**Supplementary Figure S2**).

MTAG

We used MTAG to boost statistical power and increase loci discovery by leveraging the genetic correlation between sleep apnoea and snoring. This analysis identified 43 independent genome-wide significant loci associated with sleep apnoea (**Figure 1b**). The direction and effect sizes of the independent hits were highly consistent across the sleep apnoea meta-analysis and the MTAG analysis with snoring (R^2>0.95 **Supplementary Figure S3**). After adjusting for BMI, 23 hits were genome-wide significant; most overlapped with the unadjusted results (**Figure 1b**).

Independent sample replication

We sought to replicate our GWAS results in an independent sample (N= 1,477,352) from 23andMe. Notably, the 23andMe sleep apnoea replication GWAS was adjusted for BMI (**see Methods**). Overall, ten of the independent variants identified by our analyses showed association beyond the genome-wide significance threshold (**Supplementary Table S2**) in the replication. Three out of the five loci were replicated after multiple testing corrections for the sleep apnoea meta-analysis. Furthermore, the variant that only became significant after adjusting for BMI was also replicated. For the sleep apnoea plus snoring MTAG, 30 out of 43 variants available in the 23andMe dataset were replicated. Finally, 22 out of 25 variants from the sleep apnoea plus snoring MTAG adjusted for BMI analysis were also replicated. This higher replication rate was expected since the 23andMe GWAS had been adjusted for BMI (see **Discussion**). Overall, 29 significant independent loci with evidence for replication were identified (**Table 2**). Furthermore, there was a large concordance in the direction and magnitude of effect sizes between our analyses and the 23andMe replication results (**Supplementary Figure S4**). Due to power, replication rates, and the interest in studying the aetiology of sleep apnoea beyond BMI effects, we focus below on the meta-analysis, the MTAG analysis and the MTAG analysis adjusted for BMI.

Gene-based tests and colocalization

The gene-based association analyses identified 22, 132 and 74 genes beyond the significance threshold (p<2.07e-6) for the sleep apnoea meta-analysis, the sleep apnoea plus snoring MTAG,

and the sleep apnoea plus snoring MTAG adjusted for BMI, respectively. As expected, many of these genes overlapped. Identified genes included *DLEU1, DLEU7, MSRB3, CTSF* and *SCAPER* (**Supplementary Figure S5** and **Supplementary Tables S3-S5**). Some of these genes were located within the same locus and in high LD. Thus, to identify genes linked to sleep apnoea through potential changes in gene expression, we performed eQTL colocalization analyses for any of the genes mentioned above. Of the 151 genes with available eQTL data, only 18 showed strong evidence of eQTL colocalization with either sleep apnoea, sleep apnoea plus snoring or sleep apnoea plus snoring adjusted for BMI (**Supplementary Tables S6-S8**).

eQTL integration
We used S-MultiXcan to integrate our GWAS summary statistics with eQTL data and identify genes associated with sleep apnoea through changes in predicted gene expression. These analyses identified 5 and 65 genes (**Supplementary Table S9**), for which evidence of association with sleep apnoea meta-analysis or sleep apnoea plus snoring MTAG reached statistical significance. Some of these genes included *DLEU7, PRIM1, COPZ2, SKAP1, DNAJB7, ACTBP13* and *ZBTB6,* among others. Although the results of S-MultiXcan partially overlapped those of the gene-based positional analysis, this approach identified four and 33 new genes which are likely associated with sleep apnoea through changes in gene expression. Genes with convergent evidence through gene-based association and S-MultiXcan include *FTO, STK33, ETFA, SKAP1, MAPT, BAZ2A, DCAF16, MACF1, NSF, COPZ2, SP6, LACTB2, LRRC4* and *HOXB3,* among others (**Supplementary Figure S6**).

Genetic correlations
Bivariate LD score regression was used to assess the genetic correlation between sleep apnoea and other complex traits. The trait with the highest genetic correlation (rg = 0.92) with sleep apnoea was a sleep apnoea GWAS performed on the UK-Biobank from a public GWAS repository (http://www.nealelab.is/uk-biobank/); this is essentially a subset of the UK-Biobank GWAS used in our meta-analysis. Other genetically correlated traits included respiratory diseases, type 2 diabetes, obesity, eye disorders, stroke, depression, alcohol addiction and smoking history, among others (**Supplementary Table S10-S12**). The sleep apnoea meta-analysis and the sleep apnoea plus snoring analyses showed a highly concordant pattern of genetic correlations. While also showing overall agreement, the sleep apnoea plus snoring adjusted for BMI results showed lower genetic correlations with BMI-related traits such as obesity, diabetes, and stroke (**Figure 2**).

Polygenic risk scoring
PRS based on either of our results were significantly associated with sleep apnoea in a leave one out polygenic prediction analysis. Odds ratios (OR) per standard deviation of PRS increased with the number of hits. For example, the meta-analysis based PRS (SA_{PRS}) showed an OR = 1.15 (1.08-1.21), whereas the PRS based on the sleep apnoea plus snoring showed an OR=1.21 (1.14-1.28). A similar pattern was observed for variance explained and significance (**Table 3**). These PRS were significantly associated with sleep apnoea even after adjusting for BMI measures in the AGDS cohort (**Table 3**), suggesting that signals independent of BMI contribute to polygenic prediction. Participants on the highest PRS decile showed between 50 to 87% higher odds of reporting sleep apnoea relative to the lowest decile (**Figure 3a**). Classifier models based on PRS showed prediction

ability higher than a random guess for the meta-analysis, and the MTAG showed an even higher predictive ability than the meta-analysis (**Figure 3b** and **3c** and **Supplementary Figure S7**).

Predicting traits causally associated with sleep apnoea
We used LCV to perform a hypothesis-free screening to assess whether a causal relationship can explain the potential genetic overlap between sleep apnoea and >400 traits and diseases. To this end, we employed the results of the MTAG GWAS with snoring, given its increased statistical power. We did not identify any traits with a statistically significant causal relationship on sleep apnoea risk. We found several traits for which evidence of a causal association with sleep apnoea was significant. Based on our analysis, traits that purportedly increase the risk for sleep apnoea included hypertension, asthma, lung cancer, obesity, having a period of mania, and hernia. Conversely, we found evidence for levels of vitamin D and sex hormone-binding globulin (SHBG) (from either a male- or female-only GWAS) to potentially reduce the risk for sleep apnoea (**Figure 4**).

DISCUSSION

This study aimed to increase our understanding of the genetic aetiology of sleep apnoea risk, an area that has stagnated due to the difficulty in achieving the required sample sizes for GWAS studies. Combining data across different cohorts and integrating our results with gene expression datasets, our sleep apnoea GWAS meta-analysis identified five independent loci (**Supplementary Figure 6**) associated with sleep apnoea meta-analysis. The evidence of association for these loci decreased below statistical significance upon adjustment for BMI, and new loci on chromosome 15 near *HDGFL3* reached significance. While this manuscript was under review, another study describing a GWAS for obstructive sleep apnoea (OSA) in FinnGen and the UK Biobank was published[45]. That study identified five genome-wide significant loci associated with OSA and BMI's clear, strong causal component. This is consistent with our observation of genome-wide hits showing weaker evidence of association upon a statistical adjustment for BMI effects[37]. In this study, we further used MTAG to boost power and identify additional loci likely conferring sleep apnoea risk by combining our sleep apnoea meta-analysis with a snoring meta-analysis and identified several variants linked to sleep apnoea over and above the effect of BMI. We also sought replication in an independent sample from 23andMe. The 23andMe GWAS adjusted for BMI, and we could replicate 29 loci associated with sleep apnoea, suggesting our results are robust signals linked to other sleep apnoea pathways.

We employed gene-based tests and identified several genes associated with sleep apnoea, including *DLEU1, DLEU7 CTSF, MSRB3, FTO*, and *TRIM66*. FTO association is likely due to this loci's strong effect on BMI and adiposity[46]. Loss-of-function of *MSRB3*, which encodes a methionine sulfoxide reductase, has been associated with human deafness. This finding is consistent with reported associations between hearing impairment and sleep apnoea[47]. *CTSF* has been linked to the airway wall area (Pi10) as measured quantitatively using CT chest images[48]. This is consistent with the fact that small airway dimensions have been linked to sleep apnoea measures in a COPD comorbid sample[49], and with the fact that obesity is believed to increase OSA risk by an increase in the fat levels of upper airway structures and the compression of airway walls[50]. *DLEU1* and *DLEU7* are both located within a region associated with leukaemia. While *DLEU7* is a protein-coding gene, *DLEU1* has been recently discovered to be part of a bigger gene, *BCMS,* that has a potential tumour suppressing function[51]. Although this locus has been linked to snoring[27], its role in the pathogenicity for sleep apnoea remains to be clarified.

Genes with evidence from positional gene mapping and gene-expression integration included *SKAP1, MAPT, STK33* and *ETFA. SKAP1, STK33* and *MAPT* are genes related to the MAPK signalling pathway. *MAPT* is genetically and neuropathologically associated with neurodegenerative disorders, including Alzheimer's disease and frontotemporal dementia[52]. Furthermore, *ETFA* expression has changed in an Alzheimer's disease mouse model in response to aducanumab, an amyloid-beta antibody[53]. There is a known link between sleep apnoea and Alzheimer's disease[54]. Recent studies with mouse models suggest that intermittent hypoxia induces cholinergic forebrain degeneration[55]. Furthermore, other observations suggest OSA severity might be linked to increased amyloid-beta plaques[56]. Although informative, these studies cannot still distinguish whether a true causal association underlies sleep apnoea and Alzheimer's disease in humans. Our results should enable the exploration of this question by enabling causal inference studies using instrumental variable analysis.

We did not replicate previously reported candidate gene associations such as *TNFA*, *APOE*, *PTGER3* and *LPAR1*[17]. This could be explained by differences between our analysis and those identifying the candidate genes. For example, the *LPAR1* association was observed in participants of African ancestry[57]. Nonetheless, studies assessing the support for candidate gene associations using GWAS have found poor consistency[58]. Our results suggest a similar trend for candidate gene studies of sleep apnoea. Our study should be powered to detect previously reported candidate-gene effect sizes; for instance, polymorphisms within *TNFA* were reported to show an odds ratio of 2.01 for sleep apnoea[59]. Future studies should perform a systematic evaluation of candidate gene studies and GWAS concordance in sleep apnoea, an objective outside the scope of the current study.

As a proof-of-principle of the utility of having well-powered GWAS summary statistics, we performed a hypothesis-free inference of causal associations between >400 traits and our sleep apnoea MTAG. Consistent with previous findings[45], our approach inferred obesity to likely increase the risk for sleep apnoea. Similar results were found for asthma, lung cancer, hernia, hypertension, ever having a period of mania and stroke. The extent to which hypertension, hernia, and stroke are associated with sleep apnoea above and beyond obesity as a shared causal component is unclear; given adequate data, future studies could explore this via multivariable MR analyses. Conversely, we found that SHBG levels derived from both male-only, female-only, and combined-sex GWAS decreased the risk for sleep apnoea. A similar finding was observed for endogenous testosterone levels derived from a male-only GWAS. This is consistent with observations of SHBG and testosterone levels negatively correlating with OSA severity[60]. However, continuous positive airway pressure therapy does not seem to reverse these abnormal changes[61,62], which would be consistent with the direction of causality predicted through LCV (from hormone level to phenotype). LCV also identified vitamin D levels as causal determinants of sleep apnoea risk. This is consistent with reports linking vitamin D with OSA[63]. Nonetheless, it is also possible that BMI explains this result. Given that vitamin D levels are increased with sun exposure[64], and sun exposure increases with physical activity, the well-documented inverse relationship between obesity (or BMI) and vitamin D concentrations might better explain the observed association[65,66]. Overall, our LCV analysis identified a set of testable hypotheses, which can be further explored through MR analyses contrasting the observational associations with sleep apnoea with genetically derived effect sizes.

This study was performed using cohorts of European ancestry. Thus, generalisations and comparisons with other ancestry groups should be performed with caution. In order to maximize sample size, we included cohorts with different definitions of sleep apnoea, including self-reported diagnosis and self-reported diagnostic items. While this is warranted to achieve the required sample sizes for GWAS, heterogeneity might be introduced by minimal phenotyping, but our genetic correlation analyses suggested these definitions have a comparable genetic architecture. Ideally, any sleep apnoea study would ascertain cases employing a robust measure such as the apnoea-hypopnea index or oxygen saturation; GWAS of complex traits require very large sample sizes, making such an approach challenging. Although MTAG has proven successful for boosting the discovery of loci associations, combining traits with extreme power differences might inflate signals related to the most powered phenotype[24]. In our study, adjusting for BMI seemed to affect the pattern of genetic correlations, particularly increasing the correlations with BMI and related

traits such as stroke and obesity. Replication of the sleep apnoea plus snoring adjusted for BMI results was higher than the other analyses. This result is expected for two reasons: First, it benefited from the increased power of combining GWAS for apnoea and snoring through MTAG whilst it also adjusted for BMI using mtCOJO. Second, the GWAS performed by 23andMe included BMI as a covariate. As such, it resembles a phenotype that is in line with those for which the sleep apnoea plus snoring adjusted for BMI is boosting power. Finally, some limitations of the approach used for causal inference need to be acknowledged. LCV is still dependent on the power of the original GWAS for both traits. Traits with a potential causal association with sleep apnoea might have not been included in the set of tested traits. Finally, this method assumes no bi-directional causality and will likely be biased towards the null in such cases. Thus, a null finding in our study does not reflect a lack of association, especially if bidirectional relationships are suspected.

In summary, here, we performed a GWAS meta-analysis of sleep apnoea across five European-ancestry cohorts and identified five independent genome-wide significant loci. Conditional analyses suggested a large contribution of BMI to sleep apnoea, and most of the discovered genome-wide hits were, in fact explained by BMI. After adjusting for BMI, the meta-analysis identified one genome-wide significant locus. MTAG of sleep apnoea with snoring identified 43 independent hits and 23 after conditioning on BMI. Overall, 29 significant independent hits were replicated in an independent sleep apnoea GWAS from 23andMe. All analyses showed a significant polygenic prediction of sleep apnoea in a leave-one-out PRS analysis. Our results largely confirm the previously observed overlap with BMI and highlight genetic overlap with traits such as stroke, asthma, hypertension, glaucoma and cataracts. We further found evidence indicating a potential causal role of SHBG and vitamin D levels decreasing the risk for sleep apnoea. If multivariable MR and interventional studies confirm, new treatments based on modifying these risk factors might be used for OSA treatment or early intervention. This general framework can be used in a hypothesis-free manner to generate testable hypotheses of risk factors for complex traits[43]. The associations identified here can be used as instrumental variables in targeted MR studies to understand the relationship between sleep apnoea and hypothesised causally related traits. Identifying robust loci associated with sleep apnoea is an essential step towards a deeper biological understanding, translating into novel treatments and risk assessment strategies.

Acknowledgements

The opinions expressed in this manuscript are the author's own and do not reflect the views of the Canadian Longitudinal Study on Aging or any affiliated institution. This project used data from the UK-Biobank under application number 25331. We want to acknowledge the Mass General Brigham Biobank for providing genomic and health information data. We want to acknowledge the participants and investigators of the FinnGen study. This research was made possible using the data/biospecimens collected by the Canadian Longitudinal Study on Aging (CLSA). Funding for the Canadian Longitudinal Study on Aging (CLSA) is provided by the Government of Canada through the Canadian Institutes of Health Research (CIHR) under grant reference: LSA 94473 and the Canada Foundation for Innovation. This research has been conducted using the CLSA dataset [Baseline Comprehensive Dataset version 4.0, Follow-up 1 Comprehensive Dataset version 1.0], under Application Number 190225. The CLSA is led by Drs. Parminder Raina, Christina Wolfson and Susan Kirkland. Data collection for the Australian Genetics of Depression Study was possible, thanks to funding from the Australian National Health & Medical Research Council (NHMRC) to N.G.M. (GNT1086683). A.I.C. is supported by UQ Research Training Scholarships from The University of Queensland (UQ). P.F.K. is supported by an Australian Government Research Training Program Scholarship from Queensland University of Technology (QUT). M.E.R. thanks the support of the NHMRC and Australian Research Council (GNT1102821). SM is supported by a research fellowship from the Australian National Health and Medical Research Council (NHMRC). This work was supported by an NHMRC project grant [1123248]. We would like to thank the research participants and employees of 23andMe for making this work possible. The following members of the 23andMe Research Team contributed to this study: Michelle Agee, Stella Aslibekyan, Adam Auton, Elizabeth Babalola, Robert K. Bell, Jessica Bielenberg, Katarzyna Bryc, Emily Bullis, Briana Cameron, Daniella Coker, Devika Dhamija, Sayantan Das, Sarah L. Elson, Teresa Filshtein, Kipper Fletez-Brant, Pierre Fontanillas, Will Freyman, Pooja M. Gandhi, Karl Heilbron, Barry Hicks, David A. Hinds, Karen E. Huber, Ethan M. Jewett, Yunxuan Jiang, Aaron Kleinman, Katelyn Kukar, Keng-Han Lin, Maya Lowe, Marie K. Luff, Jennifer C. McCreight, Matthew H. McIntyre, Kimberly F. McManus, Steven J. Micheletti, Meghan E. Moreno, Joanna L. Mountain, Sahar V. Mozaffari, Priyanka Nandakumar, Elizabeth S. Noblin, Jared O'Connell, Aaron A. Petrakovitz, G. David Poznik, Anjali J. Shastri, Janie F. Shelton, Jingchunzi Shi, Suyash Shringarpure, Chao Tian, Vinh Tran, Joyce Y. Tung, Xin Wang, Wei Wang, Catherine H. Weldon, Peter Wilton.

Author Contributions

MER and SM conceived and directed the study. AIC performed the AGDS GWAS, the meta and MTAG analyses, as well as post-GWAS gene-based tests, genetic correlation, PRS analyses and LCV with help from LMGM and BLM. NI performed sleep apnoea and Snoring GWAS for UK-B and CLSA. PFK and GCP performed the COLOC expression analyses. XD performed the Mass General Brigham Biobank GWAS. XH, MHL, JSO, NGM, GCP and JSY provided technical expertise and helped in the interpretation of results. YH and SA performed the replication GWAS. All co-authors contributed to manuscript drafting and provided relevant intellectual input for this manuscript.

Conflict of Interest

Stella Aslibekyan, Yunru Huang and Gabriel Cuéllar-Partida are current or former employees of 23andMe, Inc. and may hold stock or stock options. All other authors declare no conflicts of interest.

Role of the funding sources
The funding sources had no input or involvement in the design, execution or reporting of the present study.

Ethics statements
This study was performed under the oversight of the QIMR Berghofer Human Research Ethics Committee. All participants provided informed consent.

UK Biobank: The UK Biobank study was approved by the National Health Service National Research Ethics Service (ref. 11/NW/0382) and all participants provided written informed consent to participate in the UK Biobank study. Information about ethics oversight in the UK Biobank can be found at https://www.ukbiobank.ac.uk/ethics/.

CLSA: All participants provided informed consent. The CLSA abides by the requirements of the Canadian Institutes of Health Research (CIHR). The protocol of the CLSA has been reviewed and approved by 13 research ethics boards across Canada. A complete and detailed list is available at: https://www.clsa-elcv.ca/participants/privacy/who-ensures-high-ethical-standards/research-ethics-boards.

FinnGen: The Ethical Review Board of the Hospital District of Helsinki and Uusimaa approved the FinnGen study protocol (HUS/990/2017)

Mass General Brigham Biobank: All participants provided informed consent and the study protocol was approved by the Mass General Brigham Ethics Committee.

AGDS: all participants provided informed consent prior to participating in the study. This study and all questionnaires used for AGDS were approved by the QIMR Berghofer Human Research Ethics Committee.

23andMe Inc: Participants provided informed consent and participated in the research online, under a protocol approved by the external AAHRPP-accredited IRB, Ethical & Independent Review Services (E&I Review).

Data availability
The full GWAS summary statistics for this study will be made available through the NHGRI-EBI GWAS Catalogue (https://www.ebi.ac.uk/gwas/downloads/summary-statistics). Data are available from the Canadian Longitudinal Study on Aging (www.clsa-elcv.ca) for researchers who meet the criteria for access to de-identified CLSA data. UK Biobank and FinnGen data are also accessible through their respective application procedures.

TABLES

Table 1 Cohort and prevalence overview			
Cohort	Total sample size	Apnoea cases	Snoring cases
UK-Biobank	408,317	7,902	152,303
Finngen	66,216	9,096	4,270
Mass General Brigham Biobank	20,047	3,102	4,175
CLSA	18,427	3,391	6,852
AGDS	10,359	1,517	4,450
Total	523,366	25,008	172,050

Table 2. Independent hits associated with sleep apnoea and replicated in 23andMe

SNP	CHR	BP	A1	A2	P_23&me	BETA*	SE*	P_META	Signal Source
rs1537818	1	39647038	G	A	2.76E-05	-0.01755	0.004182	1.31E-09	MTAG
rs633715	1	177852580	T	C	4.60E-07	-0.02466	0.004898	3.49E-08	MTAG_BMIadj
rs72902175	2	157013035	T	C	9.30E-10	0.035999	0.005866	3.67E-14	MTAG
rs1403848	3	77609655	C	A	7.51E-05	-0.01569	0.003962	9.30E-09	MTAG
rs4076077	5	170863509	T	C	3.70E-06	-0.01797	0.003882	4.26E-09	MTAG
rs1428381	5	122693901	G	A	0.000369	0.015265	0.004283	4.83E-09	MTAG
rs2715039	7	84094964	C	A	4.61E-05	-0.01611	0.003952	2.04E-08	MTAG
rs7005777	8	78233600	T	G	5.18E-05	0.017513	0.00433	1.12E-08	MTAG
rs8176749	9	136131188	T	C	1.47E-05	-0.03212	0.007433	3.78E-09	MTAG
rs10756798	9	16739763	T	C	3.70E-09	-0.02425	0.004115	3.28E-08	MTAG_BMIadj
rs1444789	10	9064361	T	C	2.40E-13	-0.03701	0.005042	1.10E-09	MTAG
rs6265	11	27679916	T	C	1.12E-05	-0.02198	0.00501	1.79E-14	MTAG
rs1815739	11	66328095	T	C	1.19E-06	0.018979	0.003906	2.10E-08	MTAG
rs4923536	11	28422496	G	A	1.52E-10	0.025071	0.003915	7.51E-11	MTAG_BMIadj
rs28758996	12	121960480	G	A	0.00122	-0.01282	0.003963	1.21E-08	META
rs1389799	12	65824846	G	A	3.57E-25	0.04184	0.004032	1.38E-18	MTAG_BMIadj
rs4554968	12	4372609	G	A	0.000854	0.013381	0.004011	4.47E-08	MTAG_BMIadj
rs592333	13	51340315	G	A	9.04E-23	-0.03997	0.004068	1.69E-14	MTAG
rs11852496	15	83817559	T	C	3.17E-05	-0.01918	0.004605	1.71E-06	META
rs11634019	15	76634680	T	C	4.44E-10	0.027288	0.00438	1.84E-09	MTAG
rs11075985	16	53805207	C	A	1.13E-05	0.017161	0.003907	5.41E-20	META
rs8045335	16	60607116	G	A	1.41E-09	-0.02374	0.003922	1.24E-08	MTAG
rs9933881	16	1740691	T	C	3.68E-07	-0.03664	0.007183	2.54E-08	MTAG
rs12603115	17	46248994	T	C	3.95E-06	-0.01812	0.003926	8.14E-10	MTAG
rs227731	17	54773238	T	G	2.40E-11	-0.02603	0.003896	3.96E-09	MTAG
rs4987719	18	60960310	T	C	1.28E-08	0.061095	0.01068	4.72E-09	MTAG
rs35445111	19	32172047	G	A	2.62E-12	0.04751	0.006817	1.62E-11	MTAG
rs6113592	20	22229505	G	A	6.42E-07	0.019892	0.003998	7.82E-11	MTAG
rs6038517	20	6458205	G	A	0.000217	-0.01703	0.0046	2.19E-08	MTAG

GWAS and replication significant hits. Results are shown for variants with genome-wide evidence of association (p<5e-8) in at least one of the main analyses, and evidence of replication in 23andMe. *Information based on the 23andMe GWAS. SA - sleep apnoea meta-analysis; MTAG - sleep apnoea plus snoring MTAG.

MTAG_BMIadj - sleep apnoea plus snoring MTAG adjusting for BMI using mtCOJO

Table 3. Sleep apnoea polygenic prediction				
Model	OR (95%C.I.)	P	Nagelkerke R2 (%)	Variance explained (%)
SAprs	1.15 (1.08-1.21)	**5.6e-06**	0.4	0.45
SAprs (adjusted for BMI)	1.09 (1.03-1.16)	**4.2e-03**	0.17	0.45
SAmtagSnoringprs	1.21 (1.14-1.28)	**3.9e-10**	0.77	0.87
SAmtagSnoringprs (adjusting for BMI)	1.14 (1.07-1.21)	**2.4e-05**	0.37	0.87

Results of PRS derived using our GWAS (leaving out the AGDS cohort) predicting sleep apnoea in the AGDS sample with and without accounting for BMI measures in AGDS. Note significant prediction even after adjusting for BMI measures. SA - sleep apnoea meta-analysis; SAmtagSnoring - sleep apnoea plus snoring MTAG.

Figures

Figure 1 Discovery sleep apnoea genetic associations
Miami plots depict the results of the meta-analysis for sleep apnoea before and after adjusting for BMI (a) or MTAG for sleep apnoea increasing power using the snoring meta-analysis before and after adjusting for BMI (b). Each dot represents a genetic variant. The x-axis represents the variant genomic position (chromosome labels are shown). The y-axis depicts the significance of the association with sleep apnoea. In the BMI adjusted analyses, highlighted variants show the genome-wide hits of the unadjusted GWAS.

Figure 2 SA is genetically correlated with psychiatric, behavioural and cardiorespiratory traits

Forest plots showing the genetic correlations from CTG-VL[38] between SA meta-analysis, MTAG between SA meta-analysis and snoring meta-analysis (SAmtagSnoring) and MTAG between SA meta-analysis and snoring adjusted for BMI (SAmtagSnoringBMIadj). Markers depict rg whereas lines represent 95% confidence intervals derived from the rg standard error. Not all traits with a significant association (FDR < 0.05) are shown, see supplementary data.

Figure 3 Sleep Apnoea polygenic prediction
a) Plot showing the odds ratio (OR) per change in polygenic risk score (PRS) decile. Error bars depict the 95% confidence intervals b) Example of a receiver operating characteristic (ROC) curve derived from assessing the ability of a logistic regression to predict SA using either a base model (covariates only), or the base model plus the PRS of interest. The higher the area under the curve, the higher the predictive power of the model. c) Average area under ROC curve after 100 iterations of leave out validation randomly assigning training and testing subsamples. Error bars depict the standard deviation of the mean. Full results (100 ROC curves per model) are available in **Supplementary Figure 5**. SA - sleep apnoea meta-analysis; SAmtagSnoring - sleep apnoea plus snoring MTAG.

Figure 4 Predicting traits causally associated with sleep apnoea

Volcano plot showing the results of a hypothesis-free latent causal variable analysis looking for traits causally associated with sleep apnoea. Each point represents a trait of interest from a dataset of 413 GWAS for traits and diseases. The x axis represents the genetic causal proportion which ranges from -1 (trait X likely causes sleep apnoea) to 1 (trait X is likely caused by sleep apnoea). The y-axis is the absolute Z-score of the GCP which is a measure of the statistical strength of the GCP estimation. Points are coloured by their genetic correlation, which indicates, if causality exists, whether the association is of increased or decreased risk. As a reference, hypertension would be predicted to increase the risk for sleep apnoea, whereas SHBG levels would be predicted to reduce the risk for sleep apnoea.

References

1. Lévy, P. *et al.* Obstructive sleep apnoea syndrome. *Nat Rev Dis Primers* **1**, 15015 (2015).

2. Veasey, S. C. *Encyclopedia of Neuroscience*. vol. 2009 975–981 (Elsevier, 2009).

3. Donovan, L. M. & Kapur, V. K. Prevalence and Characteristics of Central Compared to Obstructive Sleep Apnea: Analyses from the Sleep Heart Health Study Cohort. *Sleep* **39**, 1353–1359 (2016).

4. Eckert, D. J., Jordan, A. S., Merchia, P. & Malhotra, A. Central sleep apnea: Pathophysiology and treatment. *Chest* **131**, 595–607 (2007).

5. Mezzanotte, W. S., Tangel, D. J. & White, D. P. Influence of sleep onset on upper-airway muscle activity in apnea patients versus normal controls. *Am. J. Respir. Crit. Care Med.* **153**, 1880–1887 (1996).

6. White, D. P. Sleep Apnea. *Proceedings of the American Thoracic Society* **3**, 124–128 (2006).

7. Teran-Santos, J., Jimenez-Gomez, A., Cordero-Guevara, J. & Burgos--Santander, C. G. The association between sleep apnea and the risk of traffic accidents. *N. Engl. J. Med.* **340**, 847–851 (1999).

8. Engleman, H. M. & Douglas, N. J. Sleep\textperiodcentered 4: Sleepiness, cognitive function, and quality of life in obstructive sleep apnoea/hypopnoea syndrome. *Thorax* **59**, 618–622 (2004).

9. Lavie, P., Herer, P. & Hoffstein, V. Obstructive sleep apnoea syndrome as a risk factor for hypertension: population study. *BMJ* **320**, 479–482 (2000).

10. Munoz, R. *et al.* Severe sleep apnea and risk of ischemic stroke in the elderly. *Stroke* **37**, 2317–2321 (2006).

11. Simiakakis, M. *et al.* Lack of effect of sleep apnea on oxidative stress in obstructive sleep apnea syndrome (OSAS) patients. *PLoS One* **7**, e39172 (2012).

12. Alonso-Fernández, A. *et al.* Effects of CPAP on oxidative stress and nitrate efficiency in sleep apnoea: a randomised trial. *Thorax* **64**, 581–586 (2009).

13. Szily, M. *et al.* Genetic influences on the onset of obstructive sleep apnoea and daytime sleepiness: a twin study. *Respir. Res.* **20**, 125 (2019).

14. Casale, M. *et al.* Obstructive sleep apnea syndrome: from phenotype to genetic basis. *Curr. Genomics* **10**, 119–126 (2009).

15. Redline, S., Tosteson, T., Tishler, P. V., Carskadon, M. A. & Millman, R. P. Studies in the genetics of obstructive sleep apnea. Familial aggregation of symptoms associated with sleep-related breathing disturbances. *Am. Rev. Respir. Dis.* **145**, 440–444 (1992).

16. Burgess, S., Butterworth, A. & Thompson, S. G. Mendelian randomization analysis with multiple genetic variants using summarized data. *Genet. Epidemiol.* **37**, 658–665 (2013).

17. Mukherjee, S., Saxena, R. & Palmer, L. J. The genetics of obstructive sleep apnoea. *Respirology* **23**, 18–27 (2018).

18. Kalra, M. *et al.* Association of ApoE genetic variants with obstructive sleep apnea in children. *Sleep Med.* **9**, 260–265 (2008).

19. Farias Tempaku, P. *et al.* Genome-wide association study reveals two novel risk alleles for incident obstructive sleep apnea in the EPISONO cohort. *Sleep Med.* **66**, 24–32 (2020).

20. Chen, H. *et al.* Multiethnic Meta-Analysis Identifies RAI1 as a Possible Obstructive Sleep Apnea-related Quantitative Trait Locus in Men. *Am. J. Respir. Cell Mol. Biol.* **58**, 391–401 (2018).

21. Cade, B. E. *et al.* Genetic Associations with Obstructive Sleep Apnea Traits in Hispanic/Latino Americans. *Am. J. Respir. Crit. Care Med.* **194**, 886–897 (2016).

22. Fuhrman, C., Fleury, B., Nguyên, X.-L. & Delmas, M.-C. Symptoms of sleep apnea syndrome: high prevalence and underdiagnosis in the French population. *Sleep Med.* **13**, 852–858 (2012).

23. Costa, L. E. *et al.* Potential underdiagnosis of obstructive sleep apnoea in the cardiology outpatient setting. *Heart* **101**, 1288–1292 (2015).

24. Turley, P. *et al.* Multi-trait analysis of genome-wide association summary statistics using

MTAG. *Nat. Genet.* **50**, 229–237 (2018).

25. Raina, P. S. *et al.* The Canadian longitudinal study on aging (CLSA). *Can. J. Aging* **28**, 221–229 (2009).

26. Raina, P. *et al.* Cohort Profile: The Canadian Longitudinal Study on Aging (CLSA). *Int. J. Epidemiol.* **48**, 1752–1753j (2019).

27. Campos, A. I. *et al.* Insights into the aetiology of snoring from observational and genetic investigations in the UK Biobank. *Nat. Commun.* **11**, 817 (2020).

28. Willer, C. J., Li, Y. & Abecasis, G. R. METAL: fast and efficient meta-analysis of genomewide association scans. *Bioinformatics* **26**, 2190–2191 (2010).

29. Aschard, H., Vilhjálmsson, B. J., Joshi, A. D., Price, A. L. & Kraft, P. Adjusting for heritable covariates can bias effect estimates in genome-wide association studies. *Am. J. Hum. Genet.* **96**, 329–339 (2015).

30. Zhu, Z. *et al.* Causal associations between risk factors and common diseases inferred from GWAS summary data. *Nat. Commun.* **9**, 1–12 (2018).

31. Byrne, E. M. *et al.* Conditional GWAS analysis to identify disorder-specific SNPs for psychiatric disorders. *Mol. Psychiatry* 1–12 (2020).

32. Yunru Huang, The 23andMe Research Team, Robert Gentleman. Genetic Association with Sleep Apnea in European Population. in (ASHG).

33. Bakshi, A. *et al.* Fast set-based association analysis using summary data from GWAS identifies novel gene loci for human complex traits. *Sci. Rep.* **6**, 32894 (2016).

34. Giambartolomei, C. *et al.* Bayesian test for colocalisation between pairs of genetic association studies using summary statistics. *PLoS Genet.* **10**, e1004383 (2014).

35. Aguet, F. *et al.* The GTEx Consortium atlas of genetic regulatory effects across human tissues. *Genetics* 1860 (2019).

36. Barbeira, A. N. *et al.* Integrating predicted transcriptome from multiple tissues improves association detection. *PLoS Genet.* **15**, e1007889 (2019).

37. Yang, J. *et al.* Conditional and joint multiple-SNP analysis of GWAS summary statistics identifies additional variants influencing complex traits. *Nat. Genet.* **44**, 369–75, S1–3 (2012).

38. Cuellar-Partida, G. *et al.* Complex-Traits Genetics Virtual Lab: A community-driven web platform for post-GWAS analyses. *Genetics* 194 (2019).

39. Byrne, E. M. *et al.* Cohort profile: the Australian genetics of depression study. *BMJ Open* **10**, e032580 (2020).

40. Lloyd-Jones, L. R. *et al.* Improved polygenic prediction by Bayesian multiple regression on summary statistics. *Nat. Commun.* **10**, 5086 (2019).

41. O'Connor, L. J. & Price, A. L. Distinguishing genetic correlation from causation across 52 diseases and complex traits. *Nat. Genet.* **50**, 1728–1734 (2018).

42. Haworth, S. *et al.* Assessment and visualization of phenome-wide causal relationships using genetic data: an application to dental caries and periodontitis. *Eur. J. Hum. Genet.* **29**, 300–308 (2021).

43. García-Marín, L. M., Campos, A. I., Martin, N. G., Cuéllar-Partida, G. & Rentería, M. E. Inference of causal relationships between sleep-related traits and 1,527 phenotypes using genetic data. *Sleep* **44**, (2021).

44. García-Marín, L. M. *et al.* Phenome-wide screening of GWAS data reveals the complex causal architecture of obesity. *bioRxiv* (2020) doi:10.1101/2020.10.20.20216598.

45. Strausz, S. *et al.* Genetic analysis of obstructive sleep apnoea discovers a strong association with cardiometabolic health. doi:10.1101/2020.08.04.235994.

46. Fawcett, K. A. & Barroso, I. The genetics of obesity: FTO leads the way. *Trends Genet.* **26**, 266–274 (2010).

47. Chopra, A. *et al.* Sleep Apnea Is Associated with Hearing Impairment: The Hispanic Community Health Study/Study of Latinos. *J. Clin. Sleep Med.* **12**, 719–726 (2016).

48. Cho, M. H. *et al.* A Genome-Wide Association Study of Emphysema and Airway Quantitative Imaging Phenotypes. *Am. J. Respir. Crit. Care Med.* **192**, 559–569 (2015).

49. Koch, A. L. *et al.* Obstructive Sleep Apnea and Airway Dimensions in Chronic Obstructive Pulmonary Disease. *Ann. Am. Thorac. Soc.* **17**, 116–118 (2020).

50. Dempsey, J. A., Veasey, S. C., Morgan, B. J. & O'Donnell, C. P. Pathophysiology of sleep apnea. *Physiol. Rev.* **90**, 47–112 (2010).

51. Wolf, S. *et al.* B-cell neoplasia associated gene with multiple splicing (BCMS): the candidate B-CLL gene on 13q14 comprises more than 560 kb covering all critical regions. *Hum. Mol. Genet.* **10**, 1275–1285 (2001).

52. Rademakers, R., Cruts, M. & van Broeckhoven, C. The role of tau (MAPT) in frontotemporal dementia and related tauopathies. *Hum. Mutat.* **24**, 277–295 (2004).

53. Bastrup, J. *et al.* Anti-Aβ Antibody Aducanumab Regulates the Proteome of Senile Plaques and Closely Surrounding Tissue in a Transgenic Mouse Model of Alzheimer's Disease. *J. Alzheimers. Dis.* **79**, 249–265 (2021).

54. Polsek, D. *et al.* Obstructive sleep apnoea and Alzheimer's disease: In search of shared pathomechanisms. *Neurosci. Biobehav. Rev.* **86**, 142–149 (2018).

55. Qian, L. *et al.* Cholinergic basal forebrain degeneration due to obstructive sleep apnoea increases Alzheimer's pathology in mice. *bioRxiv* (2020) doi:10.1101/2020.03.12.989848.

56. Owen, J. E. *et al.* Alzheimer's disease neuropathology in the hippocampus and brainstem of people with obstructive sleep apnea. *Sleep* **44**, (2021).

57. Patel, S. R. *et al.* Association of genetic loci with sleep apnea in European Americans and African-Americans: the Candidate Gene Association Resource (CARe). *PLoS One* **7**, e48836 (2012).

58. Farrell, M. S. *et al.* Evaluating historical candidate genes for schizophrenia. *Mol. Psychiatry* **20**, 555–562 (2015).

59. Zhong, A., Xiong, X., Xu, H. & Shi, M. An updated meta-analysis of the association between tumor necrosis factor-α -308G/A polymorphism and obstructive sleep apnea-hypopnea syndrome. *PLoS One* **9**, e106270 (2014).

60. Meston, N. *et al.* Endocrine effects of nasal continuous positive airway pressure in male patients with obstructive sleep apnoea. *J. Intern. Med.* **254**, 447–454 (2003).

61. Zhang, X.-B., Jiang, X.-T., Du, Y.-P., Yuan, Y.-T. & Chen, B. Efficacy of continuous positive airway pressure on testosterone in men with obstructive sleep apnea: a meta-analysis. *PLoS One* **9**, e115033 (2014).

62. Cignarelli, A. *et al.* Effects of CPAP on Testosterone Levels in Patients With Obstructive Sleep Apnea: A Meta-Analysis Study. *Front. Endocrinol.* **10**, 551 (2019).

63. Archontogeorgis, K., Nena, E., Papanas, N. & Steiropoulos, P. The role of vitamin D in obstructive sleep apnoea syndrome. *Breathe (Sheff)* **14**, 206–215 (2018).

64. Holick, M. F. Vitamin D Deficiency. *N. Engl. J. Med.* **357**, 266–281 (2007).

65. Hyppönen, E. & Boucher, B. J. Adiposity, vitamin D requirements, and clinical implications for obesity-related metabolic abnormalities. *Nutr. Rev.* **76**, 678–692 (2018).

66. Revez, J. A. *et al.* Genome-wide association study identifies 143 loci associated with 25 hydroxyvitamin D concentration. *Nat. Commun.* **11**, 1647 (2020).