The effect of maternal pre-/early-pregnancy BMI and pregnancy smoking and alcohol on congenital heart diseases: a parental negative control study

Short title: Pregnancy exposures and congenital heart diseases

Authors

Kurt Taylor[1,2*], Ahmed Elhakeem[1,2], Johanna Lucia Thorbjørnsrud Nader[3], Tiffany Yang[4], Elena Isaevska[5], Lorenzo Richiardi[5], Tanja Vrijkotte[6], Angela Pinot de Moira[7], Deirdre M Murray[8,9], Daragh Finn[8,9], Dan Mason[4], John Wright[4], Sam Oddie[10], Nel Roeleveld[11], Jennifer R Harris[12], Anne-Marie Nybo Andersen[7], Massimo Caputo[13,14], Deborah A. Lawlor[1,2,14]

Affiliations

[*] Corresponding author: kurt.taylor@bristol.ac.uk – Oakfield House, MRC Integrative Epidemiology Unit Bristol, Population Health Science Institute, University of Bristol, BS8 2BN.

[1] Population Health Science, Bristol Medical School, Bristol BS8 2BN, UK

[2] MRC Integrative Epidemiology Unit at the University of Bristol, Bristol BS8 2PS, UK

[3] Department of Genetics and Bioinformatics, Division of Health Data and Digitalisation, Norwegian Institute of Public Health, Oslo, Norway

[4] Bradford Institute for Health Research, Bradford Teaching Hospitals NHS Foundation Trust, Bradford BD9 6RJ, UK

[5] Cancer Epidemiology Unit, Department of Medical Sciences, University of Turin and CPO Piemonte, Turin, Italy

[6] Amsterdam UMC, University of Amsterdam, Department of Public and Occupational Health, Amsterdam Public Health research institute, Amsterdam, the Netherlands

[7] Section for Epidemiology, Department of Public Health, University of Copenhagen, Denmark

[8] The Irish Centre for Fetal and Neonatal Translational Research (INFANT), University College Cork, Cork, Ireland

[9] Department of Paediatrics and Child Health, University College Cork, Cork, Ireland

[10] Centre for Reviews and Dissemination, University of York, Heslington, York, UK

[11] Department for Health Evidence, Radboud Institute for Health Sciences, Radboud university medical center, Nijmegen 6500 HB, The Netherlands

[12] Division of Health Data and Digitalisation, and Centre for Fertility and Health, Norwegian Institute of Public Health, Oslo, Norway

[13] Translational Science, Bristol Medical School, Bristol BS2 8DZ, UK

[14] Bristol NIHR Biomedical Research Center, Bristol BS1 2NT, UK

Abstract

Background: Congenital heart diseases (CHDs) are the most common congenital anomaly. The causes of

CHDs are largely unknown. Higher prenatal body mass index (BMI), smoking and alcohol consumption are

associated with increased risk of CHDs. Whether these are causal is unclear.

Methods and Results: Seven European birth cohorts including 232,390 offspring (2,469 CHD cases [1.1%])

were included. We applied negative exposure paternal control analyses to explore the intrauterine effects

of maternal BMI, smoking and alcohol consumption during pregnancy, on offspring CHDs and CHD

severity. We used logistic regression and combined estimates using a fixed-effects meta-analysis. Analyses

of BMI categories resulted in similar increased odds of CHD in overweight (mothers OR: 1.15 (1.01, 1.31)

and fathers 1.10 (0.96, 1.27)) and obesity (mothers OR: 1.12 (0.93, 1.36) and fathers 1.16 (0.90, 1.50)).

The association of mean BMI with CHD was null. Maternal smoking was associated with increased odds of

CHD (OR: 1.11 (0.97, 1.25)) but paternal smoking was not (OR: 0.96 (0.85, 1.07)). The difference increased

when removing offspring with genetic/chromosomal defects (mothers OR: 1.15 (1.01, 1.32) and fathers

0.93 (0.83, 1.05)). The positive association with maternal pregnancy smoking appeared to be driven by

non-severe CHD cases (OR: 1.22 (1.04, 1.44)). Associations with maternal (OR: 1.16 (0.52, 2.58)) and

paternal (OR: 1.23 (0.74, 2.06)) moderate/heavy pregnancy alcohol consumption were similar.

Conclusions: We found evidence of an intrauterine effect for maternal smoking on offspring CHDs, but no

evidence for higher maternal BMI or alcohol consumption. Our findings provide further support for why

smoking cessation is important during pregnancy.

Key words: congenital heart disease; risk factors; negative control

1 Introduction

Congenital heart diseases (CHDs) are the most common congenital anomaly (CA), affecting 6-8

per 1000 live births and 10% of stillbirths, and are the leading cause of death from CAs [1]. Many CHD

patients present with sequela from surgical intervention and late complications related to the anomaly,

resulting in health problems that persist into adulthood [2,3]. The causes of CHDs are largely unknown, but

intrauterine mechanisms may play a role in their underlying pathophysiology [4]. Identifying modifiable risk

factors for CHDs is important for improving etiological understanding and developing preventive

interventions.

Several modifiable maternal characteristics have been found to be associated with increased risk

of CHDs, including maternal pre/early pregnancy body mass index (BMI) [5–7], smoking [8] and alcohol [9]

consumption in pregnancy. Whether these are causal is unclear. A recent systematic review and meta-

analysis of the association of BMI with CHDs found that risk of CHDs was higher in those whose mothers

were overweight or obese at the start of pregnancy, compared with those who were normal weight.

Results for underweight mothers were not reported [5], but a large cohort study consisting of >2,000,000

singletons found no clear association for maternal underweight status and CHDs [6]. These results from

conventional multivariable approaches may be explained by residual confounding due to incomplete

identification or adjustment for confounders. Maternal active smoking [8] and maternal exposure to alcohol

[9] were both associated with offspring CHDs in recent meta-analyses. However, 68% and 69% of the studies

within the meta-analyses (for maternal smoking and alcohol, respectively) did not adjust for confounders.

Therefore, those studies showing associations for smoking and alcohol cannot determine whether these

reflect the magnitude of causal effect or are biased by confounding.

Negative control studies are widely used in laboratory science and in recent years have become

increasingly used to explore causal effects in epidemiology [10]. The idea behind negative control studies is

that either the exposure or the outcome in the real experiment is substituted for a negative control

exposure (or outcome) that is not a plausible risk factor but would have similar sources of bias or

confounding as in the main experiment. In epidemiology this approach has been primarily used for

determining the extent to which hypothesized intrauterine and early life exposures might be associated

with outcomes as a result of residual confounding [10,11]. Negative parental exposure control studies are

used for this purpose. This involves comparing the confounder adjusted associations of maternal

pregnancy exposures with the offspring outcome of interest to similarly adjusted associations of the same

characteristics (negative controls) in the father. The assumptions of this approach are that: (i) measured

and unmeasured confounders influence the exposures in the same direction and with a similar magnitude

in mothers and fathers and (ii) there is no plausible reason why the exposure in the father would affect

the offspring outcome (or at a minimum the paternal association would be much weaker than in the

mother). In the present study we are assuming that paternal BMI, smoking and alcohol cannot causally

influence offspring CHDs through intrauterine mechanisms. Under these assumptions, if there is a causal

intrauterine effect of any of the maternal pregnancy exposures, we would expect to see a maternal-

specific association, with no (or a much weaker) association with the equivalent paternal exposure. Similar

associations in mothers and fathers would suggest that these are largely driven by residual confounding.

We aimed to explore the causal intrauterine effects of maternal pregnancy BMI, smoking and

alcohol on CHDs using data from the Horizon 2020 LifeCycle project [12]. As well as the negative parental

control study providing scope to explore residual confounding, the use of a large existing collaboration of

birth cohorts has considerable benefit. First, both offspring with and without CHDs are from the same

underlying population and have been selected for inclusion and assessed in identical ways. Related to this

most studies of risk factors for CHDs are case control studies which dominate meta-analysis results. These

have advantages in that they have large numbers of CHD cases and hence greater statistical power than

most cohorts, but they are prone to selection bias as response rates in controls are commonly low and

some controls are selected from hospitals or clinics. Furthermore, case control studies are susceptible to

information bias due to differential recall and reporting of the exposure between cases and controls.

Second, we have harmonized data on all exposures, confounders and outcomes. Third, we have large

numbers, with 232,390 participants in total and 2,469 CHD cases. Lastly, the ethos of the LifeCycle

collaboration is that all studies contribute to each research question unless they do not have data on

either exposure or outcome, meaning publication bias is minimized.

Methods

Inclusion criteria and participating cohorts

This study was part of the Horizon2020 LifeCycle Project. LifeCycle is a collaboration of largely

European birth cohorts that aims to determine the impact of early-life stressors on risk of developing

adverse cardio-vascular/-metabolic, respiratory, cognitive and mental health outcomes (http://lifecycle-

project.eu) [12]. A LifeCycle cohort was eligible for inclusion if it had information on CHD in the offspring

ascertained by any method and data on at least one of the following: i) mother's pre-/early-pregnancy

BMI, ii) maternal smoking during pregnancy iii) maternal alcohol consumption during pregnancy, iv)

exposures i-iii above measured in the father at a similar time to their pregnant partners. Eligible LifeCycle

cohorts could be from any geographical area and with participants from any ethnic background. In total,

seven cohorts were eligible and all participated: The Amsterdam Born Children and their Development

Study (ABCD) [13], Avon Longitudinal Study of Parents and Children (ALSPAC) [14,15], Cork SCOPE BASELINE

Study (BASELINE) [16], Born in Bradford (BiB) [17], Danish National Birth Cohort (DNBC) [18], Norwegian Mother,

Father and Child Cohort Study (MoBa) [19,20] and Nascita e INFanzia: gli Effetti dell'Ambiente (NINFEA) [21,22].

Individual cohort descriptions can be found in the Supplementary Material (Text S1). We excluded

multiple births from the study population since they differ from single births for CA outcomes [23,24]. Some

previous studies have excluded infants with any known chromosomal/genetic/teratogenic defects on the

assumption that modifiable risk factors are unlikely to contribute in the presence of known (genetic)

causes. We have not made these exclusions in our main analyses since they are often presented as

complex syndromes with variations in phenotype and severity which may be influenced by the modifiable

exposures we explore here. In additional analyses we explore whether their removal alters our main

results.

BMI, smoking and alcohol measurements

We used harmonized LifeCycle data for exposure and confounder data, with the exclusion of

paternal alcohol consumption which had not been harmonized by LifeCycle when we started this project.

ABCD and BASELINE are additional LifeCycle cohorts (all others were core). This means that at the time of

this study they were not part of the (phase 1) LifeCycle data harmonization. We harmonized the data for

these cohorts to resemble the harmonized LifeCycle variables. Cohort-specific information on methods of

data collection can be found in Supplementary Material (Text S2).

LifeCycle harmonized maternal BMI used measured or self-reported pre-/early-pregnancy weight

and height. Pre-pregnancy weight was prioritized and if not available the earliest pregnancy measures

were used. Paternal BMI was similarly reported (by the father or their pregnant partner) or measured and

we prioritized the timing to be pre- or as early as possible in their partners pregnancy. BMI was used as a

continuous variable for the main analyses. In cohorts that had >100 CHD cases, we also categorized BMI

as underweight (BMI <18.5 kg/m^2), normal weight (BMI 18.5 to <25 kg/m^2), overweight (BMI 25 to <30

kg/m^2) and obese (BMI ≥30 kg/m^2). ALSPAC, BiB, DNBC and MoBa contributed to these analyses.

We used two LifeCycle smoking variables for maternal and paternal smoking at the time of

pregnancy: (i) smoking in the first trimester (yes/no) where this was available, otherwise any smoking

during pregnancy (yes/no) and (ii) categorized into non-smokers, light (< 10 cigarettes smoked per day)

and heavy (≥ 10 cigarettes per day) throughout the entire pregnancy. Paternal smoking was categorised

as 'any smoking (yes/no)' at the time of partners pregnancy.

We used two LifeCycle variables for maternal alcohol consumption: (i) binary (yes/no), which like

smoking prioritized the first trimester if available but was otherwise any alcohol intake during pregnancy

and (ii) categorized into non-drinkers (none), light (>0 and <3 units per week) and moderate/heavy (≥3

units per week) drinkers during pregnancy. Two studies (ALSPAC and MoBa) had data on paternal alcohol

consumption in pregnancy and thus were able to harmonize variables relating to paternal alcohol for this

project. We generated one variable, categorized as: non-drinkers, light (>0 and <7units per week) or

moderate/heavy (≥7 units per week) drinkers (Text S3).

The rationale for prioritizing maternal pregnancy smoking and alcohol during the first trimester is

because fetal cardiac development starts early in pregnancy and much of the development occurs in the

first trimester [25]. 47% and 96% of mothers had measures specifically in the first trimester for smoking and

alcohol, respectively.

Congenital heart disease outcomes

Information on CHDs was retrieved from a variety of sources depending on the cohort. ALSPAC,

BiB, DNBC and NINFEA had International Classification of Diseases v10 (ICD-10) coded data. BASELINE had

individual CHD diagnoses assigned by a cardiologist based on echocardiography. For ABCD and MoBa, we

had a non-specific CHD diagnosis (yes/no). Data in ABCD, BASELINE, DNBC, and NINFEA were restricted to

liveborn infants, whereas other cohorts included stillbirths.

In the ABCD cohort, data on CHDs in liveborn children were obtained from three different sources:

(i) the infant questionnaire, which was filled out by the mother at an average infant age of 12.9 weeks, (ii)

the questionnaire filled out by the mother at an average infant age of 5.1 years, and (iii) clinical data of

the Youth Health Care Registration. In the ALSPAC cohort, cases were obtained from a range of data

sources, including health record linkage and questionnaire data up until age 25 following European

Surveillance of Congenital Anomalies (EUROCAT) guidelines [26]. In BASELINE, at 2 months, mothers were

asked of any medical problems and/or referrals. If a baby had been referred to a specialist, it was checked

by a cardiologist to see if they had results from an echocardiogram with exact diagnoses reported. Further

diagnoses up until age 12 were identified through records from the echocardiogram. In the BiB cohort,

there were two separate sources to identify CAs. Both sources were used in this study: (i) CAs up to 5

years of age, identified in GP records by Bishop et al [27] following EUROCAT guidelines. ICD-10 codes were

mapped to clinical term (CT)-V3 codes prior to extraction from GP records. (ii) Data extracted from the

Yorkshire and Humber CAs register database. Data were ICD-10 coded. All of these were confirmed

postnatally. In the DNBC, all diagnoses of congenital anomalies (according to EUROCAT guide 1.4 section

3.2 and 3.3) up until the age of 15 years were extracted from the Danish National Patient Register (DNPR)

which is linked to the cohort data[28,29]. Diagnoses were ICD-coded. These data were restricted to children

born alive. In MoBa, information on whether a child had a CHD or not was obtained though linkage to the

Medical Birth Registry of Norway (MBRN). All maternity units in Norway must notify births to the MBRN.

In the NINFEA cohort, CHDs were reported in the second questionnaire compiled 6 months after birth.

Mothers compiled a checklist that included pre-specified anomalies. If the child died or had any surgery

performed in the first 6 months, the cause of death and type of surgery were also checked to see if any

CA was reported. Data were coded using ICD-10 codes by an experienced pediatrician and were

reassessed by an independent MD. Further details of the sources of data for CHDs in each cohort are

provided in the Supplementary Material (Text S4).

In all studies, our main outcome was any CHD. Where data allowed (e.g. when we had full ICD-

codes), any CHD was defined according to EUROCAT, which excludes isolated patent ductus arteriosus

(PDA) and peripheral pulmonary artery stenosis in preterm births (gestational age <37 weeks) (Table S2).

We also categorized cases into severe CHD (heterotaxia, conotruncal defect, atrioventricular septal

defect, anomalous pulmonary venous return, left ventricle outflow tract obstruction, right ventricle

outflow tract obstruction, other complex defects) and the remainder as non-severe CHD (PDA [in full term

infants], valvular pulmonary stenosis, ventricular septal defect [VSD], atrial septum defects [ASD],

unspecified septal defects, isolated valve defects, other specified heart defects, unspecified heart defects)

[30,31] (Table S2).

Confounders

Analyses were adjusted for a number of confounders based on their known or plausible influence

on one or more of the maternal pregnancy exposures and on CHD: Maternal age (all exposures), parity

(all exposures), ethnicity (all exposures), socioeconomic position (SEP; all exposures), smoking (for BMI

and alcohol analyses), alcohol use (for BMI and smoking analyses). In the paternal negative control

analyses confounders were similar: fathers' age (all exposures), number of children (all exposures),

ethnicity (all exposures), SEP (all exposures) smoking (for BMI and alcohol), alcohol use (for BMI and

smoking). We also adjusted for offspring sex in all adjusted analyses. We used educational attainment for

both parents' measures of SEP. Full details of our selection and harmonization of confounders is provided

in the Supplementary Material (Text S5).

Statistical analysis

Analyses were conducted in either R (version 3.6.1) or Stata (version 16). An analysis plan was

written and published in October 2019, with any subsequent changes and their rationale documented in

the publication [32]. All associations between exposures and CHDs were performed within participating

studies using logistic regression (binary for main analyses and multinomial for CHD severity analyses). In

the two largest cohorts (DNBC and MoBa), we assessed deviation from linearity in our models in the BMI

analyses by running our main confounder adjusted model with BMI split into fifths. We ran regression

models with these fifths as four indicator variables (non-linear) and compared this model with one in

which the fifths were treated as a continuous (score) variable. We used a likelihood ratio comparison to

compare these two models. All analyses were run unadjusted and adjusted for maternal/paternal age,

SEP, parity (maternal) or number of children (paternal), ethnicity, smoking and/or alcohol (depending on

exposure) and offspring sex. In the adjusted models, studies were asked to adjust for as many of the

confounders as possible. All analyses were performed with maximal numbers (i.e. numbers included in

each model will vary due to missing data on exposure/outcome or confounders). In a sensitivity analysis,

we repeated our main analyses using complete-case data to assess whether missing data were influencing

the results.

For the main negative control analyses – i.e. where we directly compared maternal to paternal

exposure-CHD associations – we used multivariable logistic regression in which both maternal and

paternal exposures were adjusted for the other parent's exposure. This produces a maternal association

that adjusts for maternal confounders as well as the paternal exposure, and similarly a paternal

association adjusting for paternal confounders and the maternal exposure. The rationale for mutually

adjusting for the other parent's exposure is that parental BMI, smoking and alcohol may relate to each

other through assortative mating and/or convergence of behaviours that occurs overtime in couples [33].

Causal structural graphs together with simulated data show failure to undertake this mutual adjustment

will bias the negative control analysis results [34]. Also, paternal exposures may have some intrauterine

impact, for example via passive smoking or paternal support for the mother to reduce alcohol and have a

normal BMI during her pre-conceptual period or in pregnancy [35]. Mutual adjustment for maternal and

paternal confounders was necessary for ensuring both parental results were fully adjusted. Comparisons

between maternal and paternal associations from this model were assessed by visually comparing the

two results. In addition, statistical evidence of any differences was obtained by calculating differences in

log odds of CHD between the fathers' and mothers' associations and report the corresponding P-value

(P_{diff}), under the null hypothesis that there is no difference between the maternal and paternal estimate.

Analyses were conducted separately in each study and then meta-analysed using the *meta*

package in R [36]. All the data used in the present study originated from European birth cohorts, with broadly

similar methods and therefore, we assumed that they were each estimating an association from the same

underlying populations and used a fixed-effects meta-analysis. To explore this assumption, differences

between studies were assessed using I^2 and Cochrane Q P-values for heterogeneity [37].

Additional analyses

We repeated the main analyses after excluding infants with any known

chromosomal/genetic/teratogenic defects. Methods of data collection and definition of these variables

can be found in Supplementary Material (Table S3). Folic acid supplementation has been shown to lower

risk of birth defects and adverse pregnancy outcomes [38,39]. We repeated the adjusted maternal analyses

with additional adjustment for first trimester folic acid supplementation (yes/no).

Results

Participant characteristics

Figures S1-S7 in the Supplementary Material show flowcharts designating the assignment of

participants into analysis groups for each cohort. In total, 7 cohorts including 232,390 offspring with 2,469

CHD cases (1.1%) were included. The prevalence of CHD was close to 1% in most cohorts, with the lowest

being in ABCD (0.4%) and the highest in DNBC (1.4%) (**Table 1**). **Table 1** shows the distributions of maternal

and paternal characteristics for each cohort. Mean maternal age across the cohorts was broadly similar

(all late 20s to early 30s). Mean BMI was also similar across the cohorts but proportions in different

categories varied, with the lowest prevalence of pre-/early-pregnancy obesity seen in NINFEA (5%) and

the highest in BiB (21%). There was also variation in maternal smoking and alcohol consumption across

the cohorts, with notably high levels of both smoking (25% and 26%, respectively) and alcohol (55% and

45%, respectively) in ALSPAC and DNBC. Fathers were generally older than mothers and more likely to

smoke and drink alcohol, with the overall patterns of between study differences being similar to those for

the mothers. There were differing levels of missing data in each cohort (summarized in Table S4 and also

illustrated in cohort specific flow charts (Figures S1-S7). To check whether missing data influenced any of

our results, we report complete-case analysis results for our main analyses in the Supplementary Material.

Overall, complete-case results from meta-analyses were comparable (Tables S5-S8).

Table 1. Characteristics of the participating cohorts.

	Category	ABCD N = 8,131	ALSPAC N = 13,049	BASELINE N = 1,436	BiB N = 12,799	DNBC N = 89,107	MoBa N = 101,975	NINFEA N = 5,893
Country		Netherlands	UK	Ireland	UK	Denmark	Norway	Italy
Recruitment period		2003-2004	1991-1992	2008-2011	2007-2011	1996-2002	1999-2008	2005-2016
Offspring								
CHD	Any	34 (0.4)	103 (0.8)	10 (0.7)	145 (1.1)	1264 (1.4)	879 (0.9)	34 (0.6)
CHD severity in those with CHD	Non-severe	-	73/103 (70.9)	-	93/145 (64.1)	896/1264 (70.9)	-	27/34 (79.4)
	Severe	-	30/103 (29.1)	-	52/145 (35.9)	368/1264 (29.1)	-	7/34 (20.6)
Chromo/Genetic defects*		26 (0.3)	58 (0.4)	-	198 (1.5)	698 (0.8)	169 (0.2)	7 (0.1)
Maternal								
Age, years		30.7 (5.3)	28.9 (4.8)	30.7 (4.4)	26.0 (5.7)	29.9 (4.3)	30.2 (4.6)	33.1 (4.3)
BMI, kg/m^2		23.1 (4.1)	22.6 (4.4)	24.4 (4.1)	26.0 (5.7)	23.6 (4.3)	24.0 (4.3)	22.5 (3.8)
BMI categories	Underweight (<18.5)	360 (4.9)	1271 (11.6)	23 (1.6)	444 (4.4)	3861 (4.5)	3077 (3.2)	501 (8.5)
	Normal (18.5 to <25)	5270 (71.8)	7426 (67.7)	914 (63.6)	4586 (45.4)	57894 (67.8)	63706 (65.4)	4156 (70.5)
	Overweight (25 to <30)	1245 (17.0)	1537 (14.0)	345 (24.0)	2952 (29.2)	16578 (19.4)	21280 (21.8)	826 (14.0)
	Obese (≥30)	467 (6.4)	736 (6.7)	154 (10.7)	2127 (21.0)	7017 (8.2)	9337 (9.6)	286 (4.9)
Pregnancy smoking	Yes (any)	769 (9.5)	3147 (24.7)	357 (24.9)	1788 (16.4)	22514 (26.0)	9650 (9.6)	472 (8.1)
	Light	-	1684 (15.7)	-	1362 (12.5)	15777 (17.9)	7856 (7.7)	438 (7.5)
	Heavy	-	1096 (10.2)	-	426 (3.9)	7431 (8.5)	1587 (1.6)	30 (0.5)
Pregnancy alcohol	Yes (any)	1686 (20.8)	6894 (54.6)	527 (36.7)	-	38733 (44.7)	22799 (27.7)	1508 (25.8)
	Light	-	3044 (46.8)	-	-	46774 (52.9)	10461 (12.4)	1416 (24.4)
	Mod/Heavy	-	871 (13.4)	-	-	3717 (4.2)	509 (0.6)	230 (3.9)
Parity	Nulliparous	4500 (55.3)	5645 (45.0)	1436 (100)	4912 (39.8)	42203 (47.4)	46988 (46.9)	4070 (72.4)
Education	Low	3005 (37.3)	2374 (20.0)	-	5717 (56.9)	22225 (27.6)	2735 (2.9)	278 (4.8)
	Medium	2640 (32.8)	7985 (67.1)	208 (14.6)	1563 (15.6)	17756 (22.0)	31430 (33.1)	1892 (32.4)
	High	2403 (29.9)	1538 (12.9)	1219 (85.4)	2769 (27.6)	40675 (50.4)	60847 (64.0)	3677 (62.9)
Folic acid supp	Yes	5677 (70.7)	1070 (8.5)	-	-	56998 (69.0)	74466 (74.3)	4741 (82.5)
Paternal								
Age, years		35.1 (5.8)	30.9 (5.8)	32.2 (4.8)	30.4 (6.6)	32.2 (5.2)	32.7 (5.4)	36.2 (5.2)
BMI, kg/m^2		25.0 (3.5)	25.2 (3.3)	26.8 (3.6)	26.8 (4.7)	25.2 (3.2)	25.8 (3.3)	24.8 (3.2)
BMI categories	Underweight (<18.5)	28 (0.8)	41 (0.5)	2 (0.2)	53 (1.9)	271 (0.4)	242 (0.2)	43 (0.75)
	Normal (18.5 to <25)	1966 (54.8)	4308 (53.3)	345 (30.9)	953 (35.0)	33502 (53.5)	42952 (44.4)	3332 (58.4)
	Overweight (25 to <30)	1372 (38.2)	3111 (38.5)	594 (53.3)	1137 (41.7)	24529 (39.2)	43888 (45.3)	1977 (34.6)
	Obese (≥30)	223 (6.2)	616 (7.6)	174 (15.6)	582 (21.4)	4335 (6.9)	9759 (10.1)	355 (6.2)
Smoking	Yes	-	3459 (37.9)	277 (24.9)	1021 (32.0)	26242 (30.9)	27803 (27.3)	-
Alcohol	None	-	449 (5.5)	-	-	-	2963 (4.1)	-
	Light drinking	-	4251 (51.8)	-	-	-	59577 (82.3)	-
	Mod/heavy drinking	-	3505 (42.7)	-	-	-	9882 (13.6)	-
Education	Low	190 (8.5)	2959 (25.9)	-	4299 (52.9)	17069 (21.8)	4245 (4.4)	956 (16.6)
	Medium	398 (17.9)	6391 (55.9)	-	1115 (13.7)	28230 (36.0)	43576 (45.1)	2464 (42.8)
	High	1670 (73.9)	2079 (18.2)	-	2709 (33.3)	33118 (42.2)	48782 (50.5)	2335 (40.6)

Data are means ± SD or *n* (%). Study N's are based on singletons with data on at least one outcome and one exposure. '-' indicates data were not available. Light smoking, <10 cigarettes per day; heavy smoking, ≥10 cigarettes per day; maternal light drinking, >0 and <3 units per week during pregnancy; maternal moderate/heavy drinking, ≥3 units per week during pregnancy; paternal light drinking, >0 and <7 units per week; paternal moderate/heavy drinking, ≥7 units per week. Abbreviations: ABCD, The Amsterdam Born Children and their Development Study; ALSPAC, The Avon Longitudinal Study of Parents and Children; BiB, The Born in Bradford Study; DNBC, The Danish National Birth Cohort; MoBa, the Norwegian Mother, Father and Child Cohort Study; NINFEA, (Nascita e INFanzia: gli Effetti dell'Ambiente; Birth and Childhood: Effects of the Environment); BMI, body mass index; kg, kilogram; m, meters; mod, moderate; supp, supplementation; CHD, congenital heart disease; CA, congenital anomaly. * Chromosomal/genetic/teratogenic anomalies with a cause thought to be already known (see Table S3 for classifications).

BMI and CHDs

In confounder and other parent BMI adjusted analyses, there was no difference in the odds of

offspring CHD per 1kg/m^2 difference in maternal BMI (OR: 1.00, 95%CI: 0.99, 1.02) or paternal mean BMI

(OR: 1.01, 95%CI: 0.99, 1.03) (P$_{diff}$ = 0.43), with both being close to the null (**Figure 1A**). Results were similar

across studies (I^2 = 0% & 0%, P$_{heterogeneity}$ = 0.45 & 0.68 for maternal and paternal BMI respectively).

Unadjusted and confounder only adjusted results did not differ notably from those presented in **Figure 1**

(Figure S8). Figures S9 and S10 show the odds ratios of CHD by fifths of the BMI distribution for mothers

and fathers in DNBC and MoBa, respectively. Whilst there was statistical evidence for a linear trend in

DNBC mothers (p-value for per fifth increase = 0.05) the graph shows this was driven by increased risk

only in the highest fifth, with the 2nd, 3rd and 4th fifth (compared to the first) consistent with the null. In

MoBa mothers there was no clear pattern with some evidence that the 4th compared to the 1st fifth was

associated with lower risk with both other categories being consistent with the null (p-value for linear

trend in MoBa = 0.22). Whilst the p-values for the likelihood ratio comparing the linear model with the

category model (0.03 and 0.09, for DNBC and MoBa mothers, respectfully) provide statistical support for

the category model in each, this is based on just one of the fifths. Results for the fathers are broadly

consistent with those for the mothers, and overall these results are consistent with no association of

maternal or paternal mean BMI with offspring CHD risk.

In analyses of BMI categories, there were increased odds of offspring CHD in overweight and

obese mothers and fathers compared with those of a normal BMI, with similar magnitudes of association

in both parents (P$_{diff}$ overweight = 0.65 & P$_{diff}$ obese = 0.83) (**Figure 1B**). Underweight mothers had an

increased odds of offspring CHD, whereas underweight fathers had a decreased odds of offspring CHD.

Because of very small numbers of underweight parents, particularly fathers, however, results were

imprecise with wide confidence intervals and there was no statistical evidence for between parental

differences for underweight (P_{diff} underweight = 0.27). Individual study results for BMI categories are

shown in Figures S11-S13; there was no statistical evidence of heterogeneity across studies in the results

shown in Figure 1B (I^2 = 0%, Figure S13).

Analyses of continuously measured BMI with CHD cases separated into severe and non-severe

showed similar null associations for both mothers and fathers (P_{diff} = 1.00 for severe and 0.53 for non-

severe) (Figure S14) as well as after adjustment for folic acid supplementation in maternal analyses (Figure

S15). Results with both continuous and categories of BMI were unchanged when offspring with

chromosomal/genetic defects were removed from the study population (Figure S16 and Table S9).

A: Continuous BMI (per unit change)

B: BMI Categories

Figure 1. Associations between maternal and paternal pre/early pregnancy body mass index (BMI) and

offspring congenital heart disease (CHD). Figure 1A shows odds ratios of CHD for a one-unit (1kg/m^2)

difference in maternal BMI (top graph) and paternal BMI (bottom graph) in each study and pooled across

studies. Figure 1B shows the pooled (across ALSPAC, BiB, DNBC, MoBa) results for maternal (top) and

paternal (bottom) BMI categories. Results are odds ratios of CHD in comparison to normal BMI. The study

specific results for BMI categories are shown in supplementary material Figures S11-S13. 147,292 mothers

(1,430 with an offspring with CHD) and 133,620 fathers (1,325 with an offspring with CHD) were included

in the analyses presented in this figure. All results are adjusted for confounders (depending on cohort:

maternal and paternal age, education, ethnicity, smoking, alcohol, maternal parity and offspring sex) as

well as the other parents BMI.

Smoking and CHDs

In confounder and other parental smoking adjusted analyses any maternal smoking in pregnancy

was associated with increased odds of CHD (OR: 1.11, 95%CI: 0.97, 1.25), whereas paternal smoking at

the time of their partners pregnancy did not increase odds of offspring CHD (OR: 0.96, 95%CI: 0.85, 1.07)

(P_{diff} = 0.09) (**Figure 2A**). There was no statistical evidence of heterogeneity across studies for maternal or

paternal estimates (I^2 = 0% & 0%, $P_{heterogeneity}$ = 0.82 & 0.65 for maternal and paternal smoking,

respectively). Results for unadjusted analyses were consistent with the confounder and mutual parent

smoking adjusted result, whereas confounder only analyses were slightly attenuated for maternal

smoking (Figure S17). When removing offspring with a chromosomal/genetic defect, the magnitude of

the association for maternal smoking and CHDs increased slightly (OR: 1.15, 95%CI: 1.01, 1.32), and that

for paternal smoking decreased slightly (OR: 0.93, 95%CI: 0.83, 1.05), (P_{diff} = 0.02) (**Figure 2B** & Figure S18).

Adjusting for folic acid supplementation did not change results from main analyses (Figure S19).

The positive association between maternal smoking and offspring CHD appeared to be driven by

an association with non-severe CHD (OR: 1.22, 95%CI: 1.04, 1.44) with no increased risk of severe CHD

(OR: 0.95, 95%CI: 0.82, 1.11) (**Figures 2C & 2D**). When we analysed maternal smoking frequency

categories (i.e. none, light and heavy smoking), the results did not support an effect of heaviness over and

above what we saw with any (first trimester) smoking (Figure S20). Related to this, the maternal and

paternal associations for these categories were statistically consistent (P_{diff} = 0.25 & 0.38 for light and

heavy smoking, respectively) (Figure S20).

A: Main analyses

B: Genetic/Chromosomal defects removed from study population

C: Non−severe CHD

D: Severe CHD

Figure 2. Associations in each study and pooled across studies for maternal and paternal pregnancy

smoking and offspring congenital heart disease (CHD). Maternal first trimester smoking was prioritised

and used where possible. Figure 2A shows odds ratios of any CHD for any maternal smoking during

pregnancy (top graph) and paternal smoking (bottom graph). Figure 2B shows odds ratios of any CHD after

removing those with a chromosomal/genetic defect from the study population. 166,516 & 158,444

mothers (1,802 & 1,527 with an offspring with CHD) and 156,072 & 146,096 fathers (1,734 & 1,449 with

an offspring with CHD) were included in 2A and 2B respectively. Figures 2C and 2D show odds ratios of

1   non-severe CHD and severe CHD respectively. 89,250 mothers (828 non-severe CHD & 347 severe CHD)

2   and 86,258 fathers (813 non-severe CHD & 333 severe CHD) were included in the CHD severity analyses

3   shown (2C & 2D). All results are adjusted for confounders (depending on cohort: maternal and paternal

4   age, education, ethnicity, alcohol, maternal parity and offspring sex) as well as the other parents smoking.

Alcohol and CHDs

Due to lack of relevant paternal data, we were unable to undertake negative control analyses for

any first trimester alcohol consumption. Maternal only associations for that exposure are presented here

followed by the negative control analyses for levels of alcohol intake at any time in pregnancy. With

adjustment for all confounders, any maternal first trimester alcohol consumption was not associated with

odds of offspring CHD in meta-analyses from 5 cohorts (OR: 1.03, 95%CI: 0.94, 1.13). Results were

unchanged in unadjusted models (Figure S21) and with additional adjustment for folic acid

supplementation (Figure S22). There was a small increase in risk when restricting these analyses to non-

severe CHD (OR: 1.07, 95%CI: 0.93, 1.22) although confidence intervals included the null. Associations for

severe CHD were null (OR: 0.91, 95%CI: 0.73, 1.12) (Figure S23).

In confounder and other parental alcohol adjusted analyses, there was weak evidence of an

association between maternal light alcohol intake and CHDs (OR: 1.15, 95%CI: 0.90, 1.48), although this

did not statistically differ from paternal light intake (OR: 1.01, 95%CI: 0.63, 1.62) (P_{diff} = 0.63). Associations

for moderate/heavy intake were consistent for maternal and paternal alcohol use (P_{diff} = 0.75) with point

estimates showing weak positive associations, but with wide confidence intervals that included the null

(**Figure 3A and 3B**). We did not test associations between levels of alcohol intake and CHD severity due to

small numbers. Results for alcohol analyses were materially unchanged when removing offspring with a

chromosomal/genetic defect from the study population (Table S10). Due to the small number of cohorts

having paternal alcohol data, we also show confounder adjusted models (without mutual paternal

adjustment) for maternal alcohol intake (**Figure 3C**). The point estimate for maternal light drinking was

very close to the null and that for heavy drinking suggested it resulted in increased risk of offspring CHD.

However, both of these estimates had wide confidence intervals due to relatively few women reporting

drinking (particularly heavily) during pregnancy. Results in unadjusted analyses were unchanged (Figure

S24).

A: Parental light drinking negative control analyses

B: Parental mod/heavy drinking negative control analyses

C: Maternal alcohol (confounder adjusted)

Figure 3. Associations in each study and pooled across studies for maternal and paternal pregnancy alcohol intake and offspring CHDs. Figure 3A shows confounder and other parent's alcohol adjusted odds ratios of any CHD for maternal light drinking during pregnancy (top graph) and paternal light drinking (bottom graph). Figure 3B shows confounder and other parent's alcohol adjusted odds ratios of any CHD for maternal moderate/heavy drinking during pregnancy (top graph) and paternal moderate/heavy drinking (bottom graph). 64,156 mothers with 524 CHD cases and 64,163 fathers with 529 CHD cases were included in the alcohol negative control analyses shown (3A & 3B). Figure 3C shows confounder adjusted odds ratios of any CHD for maternal light drinking during pregnancy (top graph) and maternal mod/heavy drinking (bottom graph) (165,706 mothers with 1,823 CHD cases). Confounders (depending on cohort):

1   maternal and paternal age, education, ethnicity, smoking, maternal parity, offspring sex (and other

2   parental alcohol intake in panels A & B).

Discussion

In this large multi-cohort study, we found evidence that maternal pregnancy smoking increased offspring CHD risk via intrauterine mechanisms and that this appeared to be driven by a specific effect on non-severe CHDs. We did not find robust evidence to suggest a causal intrauterine effect of higher maternal pre-/early-pregnancy mean BMI or overweight or obesity on offspring CHD risk. Nor did we find evidence of an intrauterine effect of alcohol consumption on offspring CHD risk, although we acknowledge that for alcohol, we had less data and limited statistical power. To our knowledge, this is the first study to use a parental negative control method to explore whether maternal exposures have a causal intrauterine effect on offspring CHDs or whether associations are explained by residual confounding, which would generate a similar association for parental exposures.

We found increased odds of offspring CHD in mothers who were overweight and obese. This is consistent with the most recent systematic review and meta-analysis, which included 2,416,546 participants (57,172 with offspring CHD), from 19 studies and reported increased risk of any offspring CHD in women who were overweight or obese during pregnancy [5]. However, adjustment for confounders was poor, with 10 of the 19 included studies not providing information on confounder adjustment or not adjusting for any confounders. With more stringent confounder adjustment and the findings from a negative control study, our results suggest that the increased risk of offspring CHD in overweight and obese mothers is largely the result of residual confounding. We also found that mothers who were underweight at the start of pregnancy were at increased risk of having offspring with CHD, whereas underweight in fathers appeared to be protective of offspring CHD. There were 9,537 underweight mothers (4.4%) but only 680 underweight fathers (0.4%), making the fathers analyses imprecise and our negative control analyses lacking in power to reliably identify parental differences. The recent systematic review mentioned above did not report on associations of underweight with CHDs because too few

studies looked at this. A large Swedish linkage study of over 2 million singleton live born infants (born

between 1992 to 2012 with 28,628 CHD cases), has explored associations with maternal underweight, as

well as overweight and three grades of obesity [7]. It is difficult to compare the results from that study with

ours as they only present associations of maternal BMI with specific subtypes of CHDs, and not with any

CHD as in our main analyses. Risks of offspring CHD were similar in underweight compared to normal

weight women for all types of CHD except for mitral to tricuspid valve defects (14 cases), pulmonary valve

defects (24 cases) and right ventricular defects (5 cases), where there was some evidence of increased

prevalence with underweight. However, these estimates were imprecise, with confidence intervals

including the null. Whilst our findings suggest maternal underweight might increase offspring risk of any

CHD, we lacked power to rule out residual confounding in our negative control analyses and published

studies have limited power to explore any effects in women. The global obesity epidemic, which is

reflected in contemporary obstetric populations, might limit any potential concerns about maternal

underweight. However, as the prevalence of CHD in some low- and middle-income countries is high [40],

and these countries currently experience the double burden of under- and over-nutrition we would argue

further exploration of any possible impact of maternal underweight is warranted.

Consistent with our findings, a recent meta-analysis of >8 million participants (137,575 CHD cases)

from 125 studies reported positive associations between maternal pregnancy smoking and offspring CHDs

[8]. There was substantial heterogeneity (I^2 = 89%) in their pooled results and only 68% of the included

studies report adjustment for confounders. The authors also report positive associations between

maternal passive smoking and paternal active smoking with offspring CHDs, both of which (somewhat

unexpectedly) had stronger magnitudes of association than results from maternal active smoking. Our

results, including the negative control study, add to the previous research findings by providing more

robust evidence that these associations are unlikely to be explained by residual confounding and are

potentially causal. Other research has shown that pregnancy smoking is a risk factor for orofacial clefts [41].

The prevalence of CHD is around 1% in the general population, as shown in our study, yet in those with

orofacial clefts, CHD prevalence rates of up to 20% have been reported [42]. Both the heart and the palate

develop during early pregnancy around weeks 5 to 9. Therefore, it is plausible that smoking in early

pregnancy could disturb common biological pathways in these conditions. We found that the associations

for maternal smoking were largely driven by an effect in non-severe CHDs, with the association

strengthening when those with chromosomal or genetic defects were removed. Previous research has

reported positive associations between maternal smoking and septal defects, in particular for ASDs [43–45]

which are defined as non-severe according to the classification system used in our study.

In confounder adjusted analyses maternal alcohol consumption in the first trimester of pregnancy

was not associated with offspring CHD. There was some evidence that maternal moderate or heavy

alcohol consumption any time in pregnancy was associated with increased risk of offspring CHD. Whilst

associations between mothers and fathers light, moderate and heavy alcohol consumption, compared

with none, were statistically consistent, only 2 cohorts (80,627 participants, 703 with offspring CHD) had

alcohol information on fathers around the time of their partners pregnancy. Associations for fathers in

particular were imprecise with wide confidence intervals. Two recent meta-analyses found consistent

modest increases in risk of offspring CHD in mothers reporting alcohol consumption in pregnancy (OR:

1.11 (95%CI: 0.96, 1.29) [46] and 1.16 (1.05, 1.27) [9]. Although the first of these concluded 'no association' it

can be seen that the results for the two are consistent, and the larger sample size of the second has

increased precision. Of note, the second of these studies also explored paternal consumption and found

increased risk of offspring CHD related to fathers' alcohol consumption (1.44 (1.19, 1.74)) [9]. Although the

odds ratio for fathers' consumption suggests a stronger effect, the confidence intervals are wide, and the

result is statistically consistent with that for mothers' alcohol. As in our study there were fewer studies

with data on parental alcohol consumption around the time of their partners pregnancy. Taken together

with our findings these suggest that positive associations of maternal alcohol consumption with offspring

CHD may be due to residual confounding rather than a causal intrauterine effect.

The key strengths of this study are its large sample size, the use of a negative paternal exposures

control study and the pooling of results from several cohort studies that are less prone to selection bias

that can occur in case control studies and are not selected based on publication, but on being part of an

existing collaboration. The latter reduces the risk of publication bias as studies were included if they had

data and not on the basis of (published) results. This also allowed us to explore replication across studies

and the consistency of findings between studies in our main analyses adds confidence to our conclusions.

The use of harmonized data from LifeCycle is a strength that limits between study heterogeneity.

However, harmonizing data across several studies, as we have done in LifeCycle, can mean that some

variables lose detail. Here that is particularly relevant for the exposure and confounding variables. For

example, we were not able to explore pack weeks of smoking across the entire pregnancy. Simplified

confounder measurements, such as Western versus non-Western for ethnicity could result in residual

confounding if more specific ethnic groups have strong influences on exposure and outcome.

Furthermore, there were other confounders that we considered, including type-1 / existing diabetes and

physical activity, but had too few numbers (diabetes) across all cohorts or too few studies with data

(physical activity) to include. However, we aimed to address any form or residual confounding in our

paternal negative control analyses. Under the assumption that adjusted for but poorly measured (e.g.

ethnicity) or unadjusted for (e.g. physical activity) influence paternal exposure in the same direction and

to the same extent as in mothers, observing parental consistency of association this implies the maternal

association is influenced by residual confounding.

We were not able to fully harmonize outcome data with the key differences between studies

being the extent to which they only included cases that were diagnosed antenatally or at birth or whether

they included cases later in life. MoBa (N = 101,975 participants and N = 879 cases) only had cases

diagnosed antenatally or around the time of birth, with the remaining cohorts having diagnoses beyond

antenatal care, ranging from 6 months to 25 years. Many previous studies have only included cases

diagnosed at birth or early infancy. They, and the cohorts included here that only have these early life

cases, may be biased by outcome misclassification (i.e. the offspring who would have been diagnosed

later in life are treated as not having CHD). This is an important point for consideration because although

most CHDs are identified in utero or at birth, many are diagnosed after discharge from hospital during

childhood or even adulthood [47]. Therefore, It is reassuring therefore that our main results are largely

consistent across studies. In confounder and other parent adjusted smoking analyses, the weakest

association was found in MoBa (**Figure 2A**). It is likely that we missed some non-severe cases in MoBa

which were diagnosed later in life. Given that we demonstrate the smoking results were largely driven by

non-severe CHDs, this could have biased MoBa (and therefore meta-analysis) results towards the null.

The negative control analyses assume that factors that would confound the maternal exposure-

offspring CHD associations would have a similar magnitude and direction of confounding for the

equivalent paternal associations, irrespective of whether the confounders are measured or if measured

how accurately and precisely they are measured. This is likely to be true for paternal negative control

exposure studies, as used here [10,11]. It also assumes that there is no plausible intrauterine mechanism

through which the paternal exposure could impact offspring CHD. Whilst it has been argued that paternal

epigenetic preconception effects [48], and in the case of smoking, passive smoking could have an impact,

we would expect any such paternal exposure effect to be weaker than the maternal exposure effect. Heart

development occurs in utero (specifically in early pregnancy) and passive smoking would expose the fetus

to a much lower dose than active maternal smoking. As proof of principle, this approach confirms the

strong effect of smoking on birth weight, and fetal growth assessed by repeat ultrasound scan, with no

paternal association [49]. It is possible that potential differences in misreporting smoking and alcohol

consumption between mothers and fathers could produce spurious parental differences. Pregnant

women are likely to underreport whether they smoke or drink alcohol and the amount they smoke or

drink, because of the social stigma of these, particularly in recent decades. As the report of alcohol and

smoking in the LifeCycle cohorts was collected early in pregnancy it is likely to be random in relation to an

offspring CHD as the vast majority would not have been diagnosed. Hence, this underreporting would be

expected to attenuate any true effect of smoking/alcohol on CHD towards the null. This misclassification

is less likely in fathers. Thus, the specific positive association of maternal smoking on CHDs and its

difference to the paternal association may be underestimated.

In summary, we found evidence to support a causal intrauterine effect of maternal smoking on

any CHD, particularly with non-severe CHDs, but did not find robust evidence for a causal effect of

maternal BMI or alcohol on offspring CHD risk. Whilst everyone should be encouraged not to smoke, and

all clinical guidelines advocate not starting smoking, and if women do smoke, to quit before becoming

pregnant, there are still high rates of smoking in some groups, particularly those from deprived

backgrounds. In the studies included in this paper, two contemporary cohorts, BASELINE (Ireland), with

births occurring between 2008 and 2011 and BiB (UK), with births occurring between 2007 and 2011,

smoking prevalence rates were 25% and 16% respectively. The prevalence in BiB masks the high rate in

white British women (33%) who are from socioeconomically deprived backgrounds, as over 50% of births

in that cohort are to Pakistani women who have low rates of smoking (3%) [17]. It is possible that

emphasizing the potential adverse effect on CHDs in specific groups might help in supporting women of

reproductive age not to start smoking and women who are smoking at the start of pregnancy to be

encouraged to quit. Furthermore, understanding the specific mechanisms that link maternal smoking to

increased offspring CHD risk could identify targets for interventions for its prevention.

Acknowledgements

The authors would like to acknowledge everyone who has supported and contributed to each cohort

included in this study. Please see Text S6 (Supplementary Material) for a full list of acknowledgments.

Sources of funding

The LifeCycle project received funding from the 452 European Union's Horizon 2020 research and

innovation programme 453 (Grant Agreement No. 733206 LifeCycle). This study is also supported by the

British Heart Foundation (AA/18/7/34219), Bristol NIHR Biomedical Research Centre grant, European

Research Council (Advanced Grant; 669545) and UK National Institute of Health Research (R01 DK10324).

K.T is supported by a British Heart Foundation Doctoral Training Program (FS/17/60/33474). K.T, A.E and

D.A.L work in a unit that is supported by the University of Bristol and UK Medical Research Council

(MC_UU_00011/6). DAL is a NIHR Senior Investigator (NF-0616-10102). M.C is supported by the British

Heart Foundation Chair in Congenital Heart Disease (CH/1/32804). Work by J.R.H. is partly supported by

the Research Council of Norway through its Centers of Excellence funding scheme, project number

262700. Details of funding for each individual LifeCycle cohort that have contributed to this study are

provided below.

The Amsterdam Born Children and Their Development Study (ABCD) was supported by the Netherlands

Organization for Health Research and Development (grant 2100.0076) and the Electromagnetic Fields and

Health Research program (grants 85600004 and 85800001). Core funding for ALSPAC is provided by the

UK Medical Research Council and Wellcome (217065/Z/19/) and the University of Bristol. Very many

grants have supported different data collections, including for some of the data used in this publication

and a comprehensive list of grants funding is available on the ALSPAC website

(http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf). The Cork

BASELINE Birth Cohort Study was funded by the National Children's Research Centre (Project grant: Jan

2009). BiB has received core support from Wellcome Trust (WT101597MA) a joint grant from the UK

Medical Research Council (MRC) and UK Economic and Social Science Research Council (ESRC)

(MR/N024397/1), the British Heart Foundation (CS/16/4/32482) and the National Institute for Health

Research (NIHR) under its Collaboration for Applied Health Research and Care (CLAHRC) for Yorkshire and

Humber and the Clinical Research Network (CRN). The Danish National Birth Cohort Study (DNBC) was

supported by the Danish Epidemiology Science Centre, the Lundbeck Foundation (grant 195/04), the

Egmont Foundation, the March of Dimes Birth Defect Foundation, the Augustinus Foundation, and the

Medical Research Council (grant SSVF 0646). The Norwegian Mother, Father and Child Cohort Study is

supported by the Norwegian Ministry of Health and Care Services and the Ministry of Education and

Research. The NINFEA study was partially funded by the Compagnia San Paolo Foundation.

Disclosures

D.A.L reports support from Roche Diagnostics and Medtronic Ltd for research unrelated to that

presented here. All other authors declare no conflicts of interest.

References

1. Wang H, Naghavi M, Allen C, Barber RM, Bhutta ZA, Carter A, et al. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980–2015: a systematic analysis for the Global Burden of Disease Study 2015. The Lancet [Internet]. 2016 Oct [cited 2020 Feb 20];388(10053):1459–544. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0140673616310121

2. McCracken Courtney, Spector Logan G., Menk Jeremiah S., Knight Jessica H., Vinocur Jeffrey M., Thomas Amanda S., et al. Mortality Following Pediatric Congenital Heart Surgery: An Analysis of the Causes of Death Derived From the National Death Index. J Am Heart Assoc [Internet]. 2018 Nov 20 [cited 2020 May 7];7(22):e010624. Available from: https://www.ahajournals.org/doi/10.1161/JAHA.118.010624

3. van der Bom T, Zomer AC, Zwinderman AH, Meijboom FJ, Bouma BJ, Mulder BJM. The changing epidemiology of congenital heart disease. Nat Rev Cardiol [Internet]. 2011 Jan [cited 2020 May 7];8(1):50–60. Available from: https://www.nature.com/articles/nrcardio.2010.166

4. Jenkins KJ, Correa A, Feinstein JA, Botto L, Britt AE, Daniels SR, et al. Noninherited Risk Factors and Congenital Cardiovascular Defects: Current Knowledge: A Scientific Statement From the American Heart Association Council on Cardiovascular Disease in the Young: *Endorsed by the American Academy of Pediatrics*. Circulation [Internet]. 2007 Jun 12 [cited 2020 Feb 5];115(23):2995–3014. Available from: https://www.ahajournals.org/doi/10.1161/CIRCULATIONAHA.106.183216

5. Liu X, Ding G, Yang W, Feng X, Li Y, Liu H, et al. Maternal Body Mass Index and Risk of Congenital Heart Defects in Infants: A Dose-Response Meta-Analysis. BioMed Res Int. 2019;2019:1315796.

6. Persson M, Cnattingius S, Villamor E, Söderling J, Pasternak B, Stephansson O, et al. Risk of major congenital malformations in relation to maternal overweight and obesity severity: cohort study of 1.2 million singletons. BMJ [Internet]. 2017 Jun 14 [cited 2020 Feb 5];j2563. Available from: http://www.bmj.com/lookup/doi/10.1136/bmj.j2563

7. Persson M, Razaz N, Edstedt Bonamy A-K, Villamor E, Cnattingius S. Maternal Overweight and Obesity and Risk of Congenital Heart Defects. J Am Coll Cardiol. 2019 08;73(1):44–53.

8. Zhao L, Chen L, Yang T, Wang L, Wang T, Zhang S, et al. Parental smoking and the risk of congenital heart defects in offspring: An updated meta-analysis of observational studies. Eur J Prev Cardiol. 2019 Mar 23;2047487319831367.

9. Zhang S, Wang L, Yang T, Chen L, Zhao L, Wang T, et al. Parental alcohol consumption and the risk of congenital heart diseases in offspring: An updated systematic review and meta-analysis. Eur J Prev Cardiol [Internet]. 2020 Mar [cited 2020 Mar 20];27(4):410–21. Available from: http://journals.sagepub.com/doi/10.1177/2047487319874530

10. Smith GD. Negative Control Exposures in Epidemiologic Studies: Epidemiology [Internet]. 2012 Mar [cited 2020 Mar 20];23(2):350–1. Available from: http://journals.lww.com/00001648-201203000-00027

11. Sanderson E, Macdonald-Wallis C, Davey Smith G. Negative control exposure studies in the presence of measurement error: implications for attempted effect estimate calibration. Int J Epidemiol [Internet]. 2018 Apr [cited 2020 Jun 22];47(2):587–96. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5913619/

12. Jaddoe VWV, Felix JF, Andersen A-MN, Charles M-A, Chatzi L, Corpeleijn E, et al. The LifeCycle Project-EU Child Cohort Network: a federated analysis infrastructure and harmonized data of more than 250,000 children and parents. Eur J Epidemiol [Internet]. 2020 Jul 23 [cited 2020 Jul 27]; Available from: https://doi.org/10.1007/s10654-020-00662-z

13. van Eijsden M, Vrijkotte TG, Gemke RJ, van der Wal MF. Cohort Profile: The Amsterdam Born Children and their Development (ABCD) Study. Int J Epidemiol [Internet]. 2011 Oct 1 [cited 2020 May 12];40(5):1176–86. Available from: https://academic.oup.com/ije/article/40/5/1176/656509

14. Boyd A, Golding J, Macleod J, Lawlor DA, Fraser A, Henderson J, et al. Cohort Profile: The 'Children of the 90s'—the index offspring of the Avon Longitudinal Study of Parents and Children. Int J Epidemiol [Internet]. 2013 Feb [cited 2020 Mar 3];42(1):111–27. Available from: https://academic.oup.com/ije/article-lookup/doi/10.1093/ije/dys064

15. Fraser A, Macdonald-Wallis C, Tilling K, Boyd A, Golding J, Davey Smith G, et al. Cohort Profile: the Avon Longitudinal Study of Parents and Children: ALSPAC mothers cohort. Int J Epidemiol. 2013 Feb;42(1):97–110.

16. O'Donovan SM, Murray DM, Hourihane JO, Kenny LC, Irvine AD, Kiely M. Cohort profile: The Cork BASELINE Birth Cohort Study: Babies after SCOPE: Evaluating the Longitudinal Impact on Neurological and Nutritional Endpoints. Int J Epidemiol [Internet]. 2015 Jun 1 [cited 2020 Jun 9];44(3):764–75. Available from: https://academic.oup.com/ije/article/44/3/764/629555

17. Wright J, Small N, Raynor P, Tuffnell D, Bhopal R, Cameron N, et al. Cohort Profile: The Born in Bradford multi-ethnic family cohort study. Int J Epidemiol [Internet]. 2013 Aug 1 [cited 2020 Mar 23];42(4):978–91. Available from: https://academic.oup.com/ije/article-lookup/doi/10.1093/ije/dys112

18. Olsen J, Melbye M, Olsen SF, Sørensen TI, Aaby P, Andersen AM, et al. The Danish National Birth Cohort--its background, structure and aim. Scand J Public Health. 2001 Dec;29(4):300–7.

19. Magnus P, Irgens LM, Haug K, Nystad W, Skjaerven R, Stoltenberg C, et al. Cohort profile: the Norwegian Mother and Child Cohort Study (MoBa). Int J Epidemiol. 2006 Oct;35(5):1146–50.

20. Magnus P, Birke C, Vejrup K, Haugan A, Alsaker E, Daltveit AK, et al. Cohort Profile Update: The Norwegian Mother and Child Cohort Study (MoBa). Int J Epidemiol. 2016;45(2):382–8.

21. Firestone R, Cheng S, Pearce N, Douwes J, Merletti F, Pizzi C, et al. Internet-Based Birth-Cohort Studies: Is This the Future for Epidemiology? JMIR Res Protoc [Internet]. 2015 Jun 12 [cited 2020 May 12];4(2). Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4526937/

22. Richiardi L, Baussano I, Vizzini L, Douwes J, Pearce N, Merletti F, et al. Feasibility of recruiting a birth cohort through the Internet: the experience of the NINFEA cohort. Eur J Epidemiol. 2007;22(12):831–7.

23. Best KE, Rankin J. Increased risk of congenital heart disease in twins in the North of England between 1998 and 2010. Heart [Internet]. 2015 Nov 15 [cited 2020 Mar 20];101(22):1807–12. Available from: http://heart.bmj.com/lookup/doi/10.1136/heartjnl-2015-307826

24. Glinianaia SV, Rankin J, Wright C. Congenital anomalies in twins: a register-based study. Hum Reprod [Internet]. 2008 Apr 11 [cited 2020 Mar 20];23(6):1306–11. Available from: https://academic.oup.com/humrep/article-lookup/doi/10.1093/humrep/den104

25. Gittenberger-de Groot AC, Bartelings MM, Deruiter MC, Poelmann RE. Basics of Cardiac Development for the Understanding of Congenital Heart Malformations. Pediatr Res [Internet]. 2005 Feb [cited 2020 Sep 4];57(2):169–76. Available from: http://www.nature.com/doifinder/10.1203/01.PDR.0000148710.69159.61

26. Taylor K, Thomas R, Mumme M, Golding J, Boyd A, Northstone K, et al. Ascertaining and classifying cases of congenital anomalies in the ALSPAC birth cohort. Wellcome Open Res [Internet]. 2020 Oct 6 [cited 2020 Oct 13];5:231. Available from: https://wellcomeopenresearch.org/articles/5-231/v1

27. Bishop C, Small N, Mason D, Corry P, Wright J, Parslow RC, et al. Improving case ascertainment of congenital anomalies: findings from a prospective birth cohort with detailed primary care record linkage. BMJ Paediatr Open. 2017;

28. Lynge E, Sandegaard JL, Rebolj M. The Danish National Patient Register. Scand J Public Health. 2011 Jul 20;39(7_suppl):30–3.

29. Schmidt M, Schmidt SAJ, Sandegaard JL, Ehrenstein V, Pedersen L, Sørensen HT. The Danish National Patient Registry: a review of content, data quality, and research potential. Clin Epidemiol. 2015 Nov;449.

30. Øyen Nina, Olsen Sjurdur F., Basit Saima, Leirgul Elisabeth, Strøm Marin, Carstensen Lisbeth, et al. Association Between Maternal Folic Acid Supplementation and Congenital Heart Defects in Offspring in Birth Cohorts From Denmark and Norway. J Am Heart Assoc [Internet]. 2019 Mar 19 [cited 2020 May 18];8(6):e011615. Available from: https://www.ahajournals.org/doi/full/10.1161/JAHA.118.011615

31. Dale MTG, Magnus P, Leirgul E, Holmstrøm H, Gjessing HK, Brodwall K, et al. Intake of sucrose-sweetened soft beverages during pregnancy and risk of congenital heart defects (CHD) in offspring: a Norwegian pregnancy cohort study. Eur J Epidemiol [Internet]. 2019 Apr 1 [cited 2020 May 15];34(4):383–96. Available from: https://doi.org/10.1007/s10654-019-00480-y

32. Taylor, Kurt, Elhakeem, Ahmed. LifeCycle Project: The effect of maternal pre-/early-pregnancy BMI and pregnancy smoking and alcohol on congenital heart diseases: a parental negative control study. 2020 [cited 2020 May 14]; Available from: https://osf.io/e4t8c/

33. Di Castelnuovo A, Quacquaruccio G, Donati MB, de Gaetano G, Iacoviello L. Spousal Concordance for Major Coronary Risk Factors: A Systematic Review and Meta-Analysis. Am J Epidemiol [Internet]. 2008 Oct 8 [cited 2020 Sep 4];169(1):1–8. Available from: https://academic.oup.com/aje/article-lookup/doi/10.1093/aje/kwn234

34. Madley-Dowd P, Rai D, Zammit S, Heron J. Simulations and directed acyclic graphs explained why assortative mating biases the prenatal negative control design. J Clin Epidemiol [Internet]. 2020 Feb 1 [cited 2020 May 18];118:9–17. Available from: https://www.jclinepi.com/article/S0895-4356(19)30734-6/abstract

35. Sharp GC, Lawlor DA. Paternal impact on the life course development of obesity and type 2 diabetes in the offspring. Diabetologia [Internet]. 2019 Oct [cited 2020 Mar 20];62(10):1802–10. Available from: http://link.springer.com/10.1007/s00125-019-4919-9

36. Balduzzi S, Rücker G, Schwarzer G. How to perform a meta-analysis with R: a practical tutorial. Evid Based Ment Health. 2019 Nov;22(4):153–60.

37. Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. BMJ [Internet]. 2003 Sep 6 [cited 2020 May 22];327(7414):557–60. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC192859/

38. Xu A, Cao X, Lu Y, Li H, Zhu Q, Chen X, et al. A Meta-Analysis of the Relationship Between Maternal Folic Acid Supplementation and the Risk of Congenital Heart Defects. Int Heart J. 2016 Dec 2;57(6):725–8.

39. He Y, Pan A, Hu FB, Ma X. Folic acid supplementation, birth defects, and adverse pregnancy outcomes in Chinese women: a population-based mega-cohort study. The Lancet [Internet]. 2016 Oct [cited 2020 Mar 20];388:S91. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0140673616320189

40. Hoffman JIE. The global burden of congenital heart disease : review article. Cardiovasc J Afr [Internet]. 2013 Jun 21 [cited 2020 Sep 24];24(4):141–5. Available from: http://www.cvja.co.za/onlinejournal/vol24/vol24_issue4/#43/z

41. Leite M, Albieri V, Kjaer SK, Jensen A. Maternal smoking in pregnancy and risk for congenital malformations: results of a Danish register-based cohort study. Acta Obstet Gynecol Scand [Internet]. 2014 [cited 2020 Jul 1];93(8):825–34. Available from: https://obgyn.onlinelibrary.wiley.com/doi/abs/10.1111/aogs.12433

42. Genisca AE, Frías JL, Broussard CS, Honein MA, Lammer EJ, Moore CA, et al. Orofacial clefts in the National Birth Defects Prevention Study, 1997-2004. Am J Med Genet A. 2009 Jun;149A(6):1149–58.

43. Zhang D, Cui H, Zhang L, Huang Y, Zhu J, Li X. Is maternal smoking during pregnancy associated with an increased risk of congenital heart defects among offspring? A systematic review and meta-analysis of observational studies. J Matern Fetal Neonatal Med [Internet]. 2017 Mar 19 [cited 2020 Mar 20];30(6):645–57. Available from: https://www.tandfonline.com/doi/full/10.1080/14767058.2016.1183640

44. Malik S, Cleves MA, Honein MA, Romitti PA, Botto LD, Yang S, et al. Maternal Smoking and Congenital Heart Defects. Pediatrics [Internet]. 2008 Apr 1 [cited 2020 Jun 11];121(4):e810–6. Available from: https://pediatrics.aappublications.org/content/121/4/e810

45. Källén K. Maternal smoking and congenital heart defects. Eur J Epidemiol. 1999 Sep;15(8):731–7.

46. Wen Z, Yu D, Zhang W, Fan C, Hu L, Feng Y, et al. Association between alcohol consumption during pregnancy and risks of congenital heart defects in offspring: meta-analysis of epidemiological observational studies. Ital J Pediatr. 2016 Feb 3;42:12.

47. Massin MM, Dessy H. Delayed recognition of congenital heart disease. Postgrad Med J [Internet]. 2006 Jul [cited 2020 Jun 12];82(969):468–70. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2563766/

48. Day J, Savani S, Krempley BD, Nguyen M, Kitlinska JB. Influence of paternal preconception exposures on their offspring: through epigenetics to phenotype. Am J Stem Cells. 2016;5(1):11–8.

49. Brand JS, Gaillard R, West J, McEachan RRC, Wright J, Voerman E, et al. Associations of maternal quitting, reducing, and continuing smoking during pregnancy with longitudinal fetal growth: Findings from Mendelian randomization and parental negative control studies. PLoS Med [Internet]. 2019 Nov 13 [cited 2020 Jul 1];16(11). Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6853297/