Lineage-specific protection and immune imprinting shape the age distributions of influenza B cases

Marcos C. Vieira[1]*, Celeste M. Donato[2,3], Philip Arevalo[1], Guus F. Rimmelzwaan[4], Timothy Wood[5], Liza Lopez[5], Q. Sue Huang[5], Vijaykrishna Dhanasekaran[6], Katia Koelle[7], Sarah Cobey[1]*

For correspondence:
Marcos Vieira, mvieira@uchicago.edu ; Sarah Cobey, cobey@uchicago.edu

[1]Department of Ecology and Evolution, University of Chicago, Chicago, U.S.A.; [2]Enteric Diseases Group, Murdoch Children's Research Institute, Parkville, Australia; [3]Department of Paediatrics, The University of Melbourne, Parkville, Australia; [4]Research Center for Emerging Infections and Zoonoses, University of Veterinary Medicine Hannover, Hannover, Germany; [5]Institute of Environmental Science and Research, Wellington, New Zealand; [6]Department of Microbiology, Biomedicine Discovery Institute, Monash University, Melbourne, Australia; [7]Department of Biology, Emory University, Atlanta, Georgia, U.S.A.

Abstract

How a history of influenza virus infections contributes to protection is not fully understood, but such protection might explain the contrasting age distributions of cases of the two lineages of influenza B. Fitting a statistical model to those distributions using data from New Zealand, we found they could be explained by historical changes in lineage frequencies combined with cross-protection between strains of the same lineage. In addition to protection regardless of the order of infection, we found additional protection against B/Yamagata in people first infected with it, similar to the immune "imprinting" observed in influenza A. While the data were not informative about imprinting protection against B/Victoria, B/Yamagata imprinting could explain the fewer B/Yamagata than B/Victoria cases in cohorts born in the 1990s and the bimodal age distribution of B/Yamagata cases. Characterizing antibody specificity after exposure could illuminate the link between birth cohorts' infection histories and susceptibility to influenza B.

Introduction

The incidence of pathogens that induce long-lasting immunity, such as measles or mumps, typically peaks at a young age and subsequently decreases as hosts gain immune protection with age (Galbraith et al., 1984; Anderson et al., 1987; Grenfell and Anderson, 1985). However, many pathogens are able to infect the same host multiple times due to the circulation of antigenically distinct strains in the population. Changes in the prevalences of strains over time can lead to different infection histories in hosts born at different times, and such differences in infection history can lead to complicated distributions of infection (Gupta et al., 1998; Kucharski and Gog, 2012a,b). Influenza viruses, for instance, circulate as antigenically distinct variants, including the "types" A and B, "subtypes" of influenza A, "lineages" of influenza B, and clades within them. Changes in variant prevalence over time generate different infection histories that are correlated across birth cohorts (Kilbourne, 2006; Rota et al., 1990; Shaw et al., 2002), but how differences in infection history affect protection and shape the age distribution of influenza virus infections is not fully understood.

Early childhood infections with influenza A have long-lasting consequences for protection against influenza A subtypes, a phenomenon termed "immune imprinting" (Gostic et al., 2016, 2019; Arevalo et al., 2020). Subtypes are distinguished by their surface proteins, hemagglutinin (HA) and neuraminidase (NA), with HA subtypes clustered into the major phylogenetic groups 1 (including, among others, H1, H2 and H5) and 2 (including H3 and H7). Different subtypes of influenza A have circulated in the 20th century (Kilbourne, 2006), and protection against severe infection and death is higher against subtypes whose HAs are genetically similar to the HA of the subtype with which a person was likely first infected (Gostic et al., 2016,

2019; Arevalo et al., 2020). For instance, early infection with H1N1 or H2N2 is associated with lifelong protection against severe infections with H5N1 and H1N1, and early H3N2 infection protects against severe infection with H7N9 and H3N2. Subtypes H1N1 and H2N2 were the only subtypes circulating between 1918 and 1968, and H3N2 has been more common than H1N1 since 1968. Thus, the age distributions of clinical infections with H1N1 and H5N1 skew young, and the distributions of H7N9 and H3N2 infections skew old (Gostic et al., 2016, 2019; Arevalo et al., 2020), because of the lasting impact of childhood immunity to the first HA encountered.

Despite its durability, imprinting protection does not completely prevent re-infection with the same subtype, and models based on imprinting protection alone cannot completely recapitulate the age distribution of cases (Gostic et al., 2019; Arevalo et al., 2020). Longitudinal analyses of antibody titers, which reveal subclinical infections, suggest that protection after infection with influenza type A decreases significantly within 3.5-7 years (Kucharski et al., 2015; Ranjeva et al., 2019). Repeated clinical infections of the same subtype have been observed in the same person (Smith and Davies, 1976; Frank and Taber, 1983; Davies et al., 1984; Horby et al., 2012) and are likely enabled by antigenic evolution of HA and NA, which experience strong positive selection (Bush et al., 1999; Ferguson et al., 2003; Smith et al., 2004; Bedford et al., 2014). Thus, protection against infection with a subtype appears to depend not only on imprinting protection (early infection with a subtype) but also cross-protection from recent infections with that subtype. This cross-protection can be sensitive to the precise strains with which a person was infected, apparent as birth cohort effects and reproducible in animal models (Linderman et al., 2014; Petrie et al., 2016).

Like the different subtypes of influenza A, the two lineages of influenza type B have distinct age distributions of medically attended infections. The B/Victoria and B/Yamagata lineages diverged sometime in the 1970s to early 1980s (Rota et al., 1990; Dudas et al., 2015) and have circulated with varying frequencies since, causing 25% of global influenza cases detected in 2000-2018 (Caini et al., 2019). While the incidence of medically attended infections is highest in children for both lineages, B/Yamagata appears less common than B/Victoria in teenagers or young adults but more common in older age groups. This pattern has been observed in the 2000s and 2010s in Oceania (Vijaykrishna et al., 2015), East Asia (Tan et al., 2013), Europe (Sočan et al., 2014; Puzelli et al., 2019) and North America (Skowronski et al., 2017) and globally in sequence databases (Virk et al., 2020). Changes in the expression of sialic acid receptors with age have been proposed to explain the lower mean age of B/Victoria cases (Vijaykrishna et al., 2015), but this explanation does not account for the higher frequency of B/Yamagata cases in middle-aged people.

Alternatively or in addition to physiological changes, differences in cohorts' susceptibility to influenza B lineages might arise from differences in cohorts' infection histories. It is unclear if people have increased protection against their first infecting influenza B lineage. One hypothesized mechanism for immune imprinting in influenza A is antibodies to conserved epitopes (Gostic et al., 2016, 2019). Since B/Victoria and B/Yamagata diverged from each other more recently (Rota et al., 1990; Chen and Holmes, 2008; Dudas et al., 2015) and evolve antigenically much more slowly than the influenza A subtypes (Bedford et al., 2014, 2015; Vijaykrishna et al., 2015), conserved epitopes within a lineage might also be conserved between lineages, leading to strong cross-lineage protection by antibodies targeting those epitopes, regardless of which lineage was encountered first.

To investigate how protection might arise from infections with B/Victoria and B/Yamagata and contribute to differences in their age distributions, we fitted a statistical model to medically attended influenza B cases in New Zealand (Figure 1). The model used the estimated historical frequencies of the lineages to estimate the probabilities of different infection histories and the strength of within- and cross-lineage protection from previous infections. We found that differences in birth cohorts' susceptibility to each lineage are consistent with strong cross-protection between strains of the same lineage. In addition to within-lineage protection independent of the order of infection, we found evidence of an additional protection against B/Yamagata in people first infected with it, similar to the immune imprinting protection previously shown for influenza A. Imprinting protection against B/Yamagata can explain why cohorts born in the 1990s, when B/Yamagata appears to have been the only lineage circulating in New Zealand and other countries outside of East Asia (Shaw et al., 2002), have since had fewer B/Yamagata cases than B/Victoria cases and fewer B/Yamagata cases than older birth cohorts. The strength of imprinting protection against B/Victoria could not be estimated with confidence, suggesting it might have little impact on the age distributions of cases. Characterizing individuals' antibody specificity after influenza virus infection, and understanding how such specificity arises from past exposure, could help explain variation in infection risk over time and the unexpected age distributions of influenza variants.

Figure 1. Historical frequencies and age distributions of the influenza B lineages. A. Frequency of B/Yamagata estimated from sequences deposited on GISAID and the NCBI Influenza Virus Database. Frequencies were estimated for New Zealand and Australia combined to increase power. The circles are annotated with the number of isolates used to estimate frequencies in each season. In 2001 and 2003, when both lineages were known to be circulating in Australia and New Zealand but the number of isolates from those countries combined were small, we estimated lineage frequencies using isolates from all countries represented in the sequence databases. **B.** Distribution of medically attended influenza B cases in New Zealand in 2001-2019 by birth year of the infected person. The fraction of cases in each birth year was calculated relative to all cases observed for each lineage (including birth years before 1960, not shown in the figure).

Results

Statistical model of influenza B infections by birth year

To test how age, early infections, and protection within and between lineages might shape the distribution of influenza B cases, we fitted a statistical model to medically attended infections detected by influenza surveillance from 2001 to 2019 in New Zealand (Materials and Methods: "Case data"; Figure 1B, Figure 1-Supplement 1). Some of these data, from 2002-2013, were previously analyzed by Vijaykrishna et al. (2015), who also compiled cases observed in Australia. We focused on the New Zealand data because they include outpatient cases obtained by sentinel surveillance under a standardized case definition and sampling strategy. The data set also includes non-sentinel cases identified from hospital samples, mostly inpatient, that were obtained without consistent sampling criteria. Of 4,036 cases identified at the level of the influenza B lineage, approximately 60% (2,430) came from sentinel surveillance and 40% (1,606) from non-sentinel surveillance. We performed the main analyses on the entire New Zealand data set but assessed the effect of removing the non-sentinel data and fitting the model only to the more reliable sentinel surveillance data.

Following work on influenza A (Gostic et al., 2016, 2019; Arevalo et al., 2020), the model describes the probability that a case of a particular lineage in a particular season occurs in a person born in a particular birth year, and it bases these probabilities on the following factors (Materials and Methods: "Statistical model of influenza B susceptibility based on infection history"):

1. *Demography*. The fraction of the population born in a particular birth year gives a baseline probability that a case occurs in a person born in that year. We used demographic data from New Zealand to estimate those fractions (Materials and Methods: "Demographic data").
2. *Age-specific effects on infection risk*. Age can affect rates of contact with infectious people and susceptibility to infection (Mossong et al., 2008; Melegaro et al., 2011). Because of likely changes in the frequency and pattern of social contacts associated with school attendance, we assumed that preschoolers (0-5 years old), school-age children and teenagers (6-17 years old) and people 18 years and older had different baseline probabilities of infection, which we estimated

from the case data. We compared the estimates for the preschool and school-age attack rates with estimates obtained independently from seroprevalence data in children.

3. *Age-specific effects on reporting.* We assumed infections in children younger than 2 years old had a different probability of receiving medical attention and being reported than infections in the rest of the population (estimates of protection were similar when we assumed this effect applied to children younger than 5). We estimated this differential reporting from the case data.

4. *Infection history.* The probability of observing a case of a lineage in a birth cohort could be modified by the average susceptibility of people in that cohort to that lineage, based on the birth cohort's infection history. Susceptibility is defined as the probability of infection given exposure, relative to that of a naive person. We estimated the probabilities of different infection histories for each birth cohort using a discrete-time model in which the probability of becoming infected with either lineage depends on the lineages' historical frequencies (Materials and Methods: "Infection history probabilities"), which we estimated using sequence databases (Materials and Methods: "Historical frequencies of influenza B lineages"). Consistent with historical surveillance and with trends observed in other countries outside East Asia (Shaw et al., 2002), we found that B/Yamagata was the only lineage circulating in New Zealand in the 1990s. B/Victoria started circulating again in the early 2000s, and the two lineages have circulated with alternating frequencies since (Figure 1A, Figure 1-Supplement 2). Because the exclusive circulation of B/Yamagata in New Zealand is supported by scarce sequence data and antigenic surveillance that did not always include representative B/Victoria strains, we investigated the sensitivity of the results to the frequency of B/Yamagata in the 1990s by alternatively estimating lineage frequencies from all countries represented in sequence databases for that period.

We examined four effects of past infection (Table 1):

(a) *Within-lineage protection.* Any previous infection with B/Victoria or B/Yamagata decreases susceptibility to future infections with the same lineage by fractions χ_{VV} and χ_{YY}, respectively.

(b) *Cross-lineage protection* Cross-lineage protection was estimated as a fraction (γ) of the corresponding within-lineage protection against that lineage: $\chi_{YV} = \gamma_{YV} \times \chi_{YY}$ and $\chi_{VY} = \gamma_{VY} \times \chi_{VV}$, where χ_{YV} is the protection from B/Victoria against B/Yamagata and χ_{VY} is the protection from B/Yamagata against B/Victoria. We assumed that once a person was infected with a lineage, within-lineage protection superseded protection from previous infection with the other lineage.

(c) *Lineage-specific imprinting.* To represent increased protection against the lineage of first infection, susceptibility to B/Victoria and B/Yamagata could be further reduced by R_V or R_Y if a person was first infected with the corresponding lineage. Thus, susceptibility to B/Victoria in people first infected with B/Victoria was $(1 - \chi_V)(1 - R_V)$, and susceptibility to B/Yamagata in people first infected with B/Yamagata was $(1 - \chi_Y)(1 - R_Y)$.

(d) *Infection with influenza B strains before 1988.* Because sequence data were too scarce before 1988 to reliably estimate the frequencies of B/Victoria and B/Yamagata, we treated all infections before 1988 as infections with a separate "ancestral" lineage A and estimated protection from those infections against B/Victoria (χ_{VA}) and B/Yamagata (χ_{YA}). Those infections encompass the ancestral influenza B lineage before the split between B/Victoria and B/Yamagata and also strains circulating between the split and 1988. An infection with A, if it occurred, was necessarily a person's first infection. We modeled those infections to capture the middle part of the age distributions of cases, but because of the uncertain identity and antigenic phenotype of strains circulating in that period, we interpreted the associated parameters with caution.

We assumed that the protection conferred by an infection against future infections depends on the lineage but not on the time between infections. Since the lineages were identified in the late 1980s, sequence divergence within them has been small (about 7% at the amino acid level for the HA of both lineages in 2019) compared to the sequence divergence between them (about 14%; Figure 1-Supplement 3; Materials and Methods: "Sequence divergence analysis"). Assuming that cross-protection from past exposures does not decay appreciably over time allowed us to calculate infection history probabilities exactly without the need for dynamical simulations (Materials and Methods: "Infection history probabilities").

We fitted the model by maximum likelihood using the probabilities of observing an infection in each cohort as the parameters of a multinomial distribution. To limit model complexity, we fitted to cases from people born since 1959, i.e., to people 60 years old or younger at the time the cases were observed. Including older people (Figure 1- Supplement 1) would have required additional parameters to describe age-related changes in susceptibility, vaccination, healthcare-seeking

Table 1. Possible infection histories in terms of the lineage of first infection and lineages encountered since.

1st infection	Lineages encountered later	Symbol	Susceptibility to B/Vic	Susceptibility to B/Yam
None	None	P_0	1	1
V	None	$P_{V,0}$	$(1 - \chi_{VV})(1 - R_V)$	$(1 - \chi_{YV})$
V	Y	$P_{V,Y}$	$(1 - \chi_{VV})(1 - R_V)$	$(1 - \chi_{YY})$
Y	None	$P_{Y,0}$	$1 - \chi_{VY}$	$(1 - \chi_{YY})(1 - R_Y)$
Y	V	$P_{Y,V}$	$1 - \chi_{VV}$	$(1 - \chi_{YY})(1 - R_Y)$
A	None	$P_{A,0,0}$	$1 - \chi_{VA}$	$1 - \chi_{YA}$
A	V	$P_{A,V,0}$	$1 - \chi_{VV}$	$1 - \max(\chi_{YA}, \chi_{YV})$
A	Y	$P_{A,Y,0}$	$1 - \max(\chi_{VA}, \chi_{VY})$	$1 - \chi_{YY}$
A	V and Y (any order)	$P_{A,\{VY\}}$	$1 - \chi_{VV}$	$1 - \chi_{YY}$

V indicates B/Victoria, Y indicates B/Yamagata, and A indicates strains circulating before 1988. Susceptibility to each lineage depends on within-lineage (χ_{VV}, χ_{YY}) and cross-lineage (χ_{VY}, χ_{YV}, χ_{VA}, χ_{YA}) cross-protection from prior infections regardless of their order and on additional protection against the lineage first encountered (R_V and R_Y). Only the strongest cross-lineage protection term is assumed to affect susceptibility. Within-lineage protection is constrained to be stronger than cross-lineage protection against the same lineage.

166 behavior, and contact rates, and potentially multiple ancestral strains. We assessed the effect of moving the birth year cutoff
167 seven years in each direction (1952 and 1966). Influenza vaccine coverage was low in New Zealand during the study period in
168 the non-elderly (6% or less for ages 0-49 years and 13% for ages 50-64 years in 2016; Institute of Environmental Science and
169 Research, 2017; Materials and Methods: "Demographic data"), and we thus ignored protection by vaccination.

Evidence for within-lineage protection and immune imprinting

The distributions of B/Victoria and B/Yamagata are most consistent with within-lineage protection and imprinting protection
against B/Yamagata, with weak or unidentifiable protection across lineages (Table 2, Figure 2, Figure 2-Supplement 1). Any
previous B/Victoria infection decreased the probability of medically attended infection with B/Victoria by 89% ($\chi_{VV} = 0.89$,
95% CI 0.85-0.96), and any previous B/Yamagata infection decreased the probability of future medically attended B/Yamagata
infection by 28% ($\chi_{YY} = 0.28$, 95% CI 0.1-0.43). The data are most consistent with weak or nonexistent protection from
B/Yamagata infections against B/Victoria ($\gamma_{VY} = 0$, 95% CI 0-0.07, corresponding to a reduction in susceptibility of 0-4%
after re-scaling by χ_{VV}), whereas protection from B/Victoria against B/Yamagata was non-identifiable ($\gamma_{YV} = 0$, 95% CI 0-1).
People for whom B/Yamagata was the lineage of first infection had an additional 87-100% reduction in susceptibility to clinical
B/Yamagata infections compared with people infected with B/Yamagata after a primary infection with B/Victoria or with
strains circulating before 1988 ($R_Y = 0.96$, 95% CI 0.87-1). The strength of imprinting for B/Victoria was non-identifiable
($R_V = 0.97$, 95% CI 0-1).

The model suggests imprinting protection against B/Yamagata contributes to two major features of the birth year distribution
of cases: the low incidence of B/Yamagata compared to B/Victoria in people born in the late 1980s and early 1990s and the
lower incidence of B/Yamagata in those birth cohorts compared with older cohorts (Figure 2). Because B/Yamagata appears
to have been the only lineage circulating in New Zealand in the 1990s, many more people born in the late 1980s and early
1990s had been infected with B/Yamagata than with B/Victoria by the time the cases were recorded in the 2000s and 2010s
(Figure 3). Under the maximum likelihood estimates of infection history probabilities, 87% of people born between 1987 and
1993 had been infected with B/Yamagata by 2010 (the midpoint of the surveillance period), of which 96% had B/Yamagata as
their first influenza B infection (Figure 3). In contrast, only 55% of people in those birth cohorts had been previously infected
with B/Victoria by 2010. Thus, cohorts born in the late 1980s and early 1990s were more susceptible to B/Victoria (about
50% as susceptible as a fully naive cohort) than to B/Yamagata (20% as susceptible as a fully naive cohort).

Older birth cohorts, in contrast, were also likely infected with B/Yamagata in the 1990s and 2000s (74% of people born
until 1986 had been infected with B/Yamagata by 2010), but unlike younger cohorts, they lacked the imprinting protection
against B/Yamagata (Figure 3). The high proportion of B/Yamagata cases in those older birth cohorts in the 2000s and 2010s,
even after many people in them had already been infected with B/Yamagata, is consistent with only moderate within-lineage
protection against B/Yamagata in the absence of immune imprinting. The difference in the proportion of B/Yamagata cases
between cohorts born around 1990 and older birth cohorts also appears not due to differences in transmission related with age

Table 2. Parameter estimates for the model fitted to the distribution of general practice (sentinel) and hospital-associated (non-sentinel) cases in New Zealand.

Parameter	MLE (95% CI)	Definition
β_1	0.09 (0.07-0.10)	Annual exposure probability for preschoolers (0-5 years old)
β_2	0.16 (0.13-0.20)	Annual exposure probability for people 6-17 years old
β_3	0.12 (0.09-0.15)	Annual exposure probability for people 18+ years old
χ_{VV}	0.89 (0.85-0.96)	Protection against B/Vic from any prior B/Vic infection
χ_{YY}	0.28 (0.1-0.43)	Protection against B/Yam from any prior B/Yam infection
γ_{YV}	0.00 (0.00-1.00)	Protection from B/Vic infection against B/Yam (as a fraction of χ_{YY})[†]
γ_{VY}	0.00 (0.00-0.07)	Protection from B/Yam infection against B/Vic (as a fraction of χ_{VV})[†]
R_V	0.97 (0.00-1.00)	Additional B/Vic protection if 1st infection was with B/Vic
R_Y	0.96 (0.87-1.00)	Additional B/Yam protection if 1st infection was with B/Yam
γ_{VA}	1.00 (0.93-1.00)	Protection against B/Vic from pre-1988 infections (as a fraction of χ_{VV})[†]
γ_{YA}	1.00 (0.00-1.00)	Protection against B/Yam from pre-1988 infections (as a fraction of χ_{YY})[†]
ρ	1.97 (1.73-2.26)	Reporting factor for children under 2 years old.

[†]Cross-lineage protection does not apply when within-lineage protection is present.

Figure 2. Observed and predicted distributions of influenza B cases in New Zealand by birth year. The model was simultaneously fitted to the age distributions in each observation year from 2001 to 2019, accounting for uncertainty in the birth year of each reported case given the patient's age. For plotting, we pooled observed and predicted numbers of cases across observation years for each birth year, assuming the earliest possible birth year for each age (e.g., an age of 10 in 2000 was assumed to correspond to the birth year 1989). Red lines and dots show the predicted distribution under the model. Vertical bars are 95% bootstrap confidence intervals. In the bottom row, predicted and observed fractions of cases were normalized by dividing by the fraction of the population born in that birth year (i.e., the null expectation if all birth years were infected at the same rate).

Figure 3. Probabilities of different infection histories with influenza B in New Zealand for people born between 1959 and 2010 and observed in 2010. Infection histories consist of the lineage of first infection and lineages encountered later regardless of their order. Probabilities were estimated by fitting the model to case data from New Zealand. (A,0,0): First infection before 1988 and no subsequent infections with either B/Victoria or B/Yamagata. (A,Y,0) and (A,V,0): First infection before 1988 followed by B/Yamagata but not B/Victoria and by B/Victoria but not B/Yamagata, respectively. (A,{V,Y}): First infection before 1988 followed by infections with both B/Victoria and B/Yamagata in any order. (Y,V) and (Y,0): First infection with B/Yamagata, with and without a subsequent B/Victoria infection. (V,Y) and (V,0): First infection with B/Victoria, with and without a subsequent B/Yamagata infection. (0): fully naive to influenza B.

198 itself, since we estimated similar baseline annual infection rates for school-age children and adults (Table 2).

199 Weak or nonexistent protection from B/Yamagata against B/Victoria is consistent with a surge of B/Victoria cases in
200 children born in the late 1980s and early 1990s as B/Victoria reemerged in New Zealand in the early 2000s (Figure 2-
201 Supplement 2). By 2000, 66% of people born in 1987-1993 had been infected with B/Yamagata. The high incidence of
202 B/Victoria in those birth cohorts in the early 2000s suggests their history of B/Yamagata exposure did not protect them against
203 medically attended B/Victoria infections. This lack of protection from B/Yamagata against B/Victoria was distinguishable
204 from potential changes in transmission with age (Figure 3-Supplement 1). The non-identifiability of cross-lineage protection
205 in the other direction, from B/Victoria against B/Yamagata, might be due to the model estimating that relatively few people
206 were infected with B/Victoria alone (Figure 3). Our model assumes that protection from B/Yamagata infections themselves
207 supersedes the protection from B/Victoria infections against B/Yamagata in people infected with both lineages.

208 The estimated protection from strains circulating before 1988 against clinical infections suggests those early strains may
209 have been antigenically more similar to B/Victoria than to B/Yamagata. Protection against B/Yamagata from strains circulating
210 before 1988 (relative to protection from B/Yamagata itself) was non-identifiable ($\gamma_{YA} = 1.00$, 95% CI 0.00-1.00, corresponding
211 to a 0-28% protection). It is possible that protection from those early strains against B/Yamagata violates our assumption that
212 it cannot be stronger than protection from B/Yamagata itself, but the high proportion of B/Yamagata cases throughout the
213 2000s and 2010s in birth cohorts that were likely infected with those early strains suggests that such protection is limited at
214 best (Figure 2-Supplement 2). In contrast, the model estimated that infections with early strains were as protective against
215 B/Victoria as were B/Victoria infections themselves ($\gamma_{VA} = 1.00$, 95% CI 0.93-1.00). Thus, when B/Victoria began circulating
216 at high frequencies in the 2000s, birth cohorts with a history of influenza B infection prior to 1988 were strongly protected
217 against medically attended B/Victoria infections. Because we grouped infections before the lineages split with infections
218 that occurred between the split and 1988, it is possible that the strong protection from those early strains against B/Victoria
219 derives from a high incidence of B/Victoria during that period (which we are unable to detect). However, surveillance in the
220 United States found only two seasons with high influenza B circulation between 1983 (the most recent estimate of the lineage
221 split) and 1988 (Rota et al., 1990), suggesting that at least some of the protection comes from ancestral influenza B strains
222 prior to the lineage split.

223 Estimates of protection were similar if we excluded non-sentinel cases (Figure 2-Supplement 3, Supplement 4), if we
224 moved the minimum birth year to which we fitted the model (1959) by 7 years in each direction (Figure 2-Supplement 5,
225 Supplement 6), and if we assumed a separate reporting parameter for children under 5 years old instead of children under

2 years old (Figure 2-Supplement 7). Fitted to both sentinel and non-sentinel cases combined, the model underestimated
cases in the most recent birth years due to variation over time in the number of non-sentinel cases reported in the youngest
children (Figure 2-Supplement 2). Fitted to sentinel cases alone, however, the model predicted the number of cases in these
birth cohorts well (Figure 2-Supplement 4).
We also fitted the model to influenza B cases recorded in Australia and compiled by Vijaykrishna et al. (2015), mostly from
non-sentinel surveillance at hospitals. Those data are also consistent with lineage-specific protection against both lineages
and with imprinting protection against B/Yamagata (Figure 2-Supplement 8). However, unlike the New Zealand case data,
the age distribution of Australian cases is consistent with substantial cross-lineage protection. These differences are likely to
be at least partly due to differences in surveillance and sampling. For instance, the surge in B/Victoria cases during the early
2000s in children born in the 1990s in New Zealand, which is consistent with little to no protection from B/Yamagata against
B/Victoria, appears to not have been represented in the Australian case data (Figure 2-Supplement 9).

Sensitivity analyses corroborate the exclusive circulation of B/Yamagata in New Zealand in the 1990s

While previous work suggests B/Yamagata was the only lineage circulating widely outside of East Asia in the 1990s (Shaw
et al., 2002), the evidence for the exclusive circulation of B/Yamagata in New Zealand during that period is currently limited.
Few isolates collected in New Zealand or Australia during the 1990s were deposited in sequence databases (although they are
all B/Yamagata, Figure 1A, Figure 1-Supplement 2), and antigenic surveillance often did not test circulating strains against
representative B/Victoria strains. We re-fitted the model using lineage frequencies estimated from all countries represented
in sequence databases for the 1990s — resulting in an average B/Victoria frequency of about 25% in that period (Figure 1-
Supplement 4) — and found that the age distributions could not be adequately fitted using those alternative frequencies (a
decrease of 53 log-likelihood units compared to the model fitted with only B/Yamagata in the 1990s for the same number of
parameters). In particular, even with strong imprinting and cross-lineage protection from B/Victoria, the model could not
fully capture the low number of B/Yamagata cases in cohorts born around 1990 (Figure 2-Supplement 10). Thus, the age
distributions of medically attended influenza B cases in New Zealand are most consistent with the exclusive circulation of
B/Yamagata in the 1990s.

Estimated infection probabilities are consistent with independent serological data

To test if our model produced realistic estimates of infection risk, we compared estimates from the model fitted to the New
Zealand case data with estimates based on cross-sectional serology from children in the Netherlands (Bodewes et al., 2011;
Materials and Methods: "Model validation with independent serological data"). Both the infection probabilities estimated
from our model and those inferred from cross-sectional serology represent an average probability across influenza seasons.
Despite potential differences between New Zealand and the Netherlands in the intensity of influenza B circulation and in
lineage frequencies, we found similar estimates of infection risk. The annual probability of influenza B exposure (which,
for naive people, is equal to the probability of infection) was similar between our model (9% for preschoolers and 16% for
school-age children, 95% CIs 7-10% and 13-20%) and the estimates from Dutch seroprevalence data (12% for preschoolers
and 22% for school-age children, 95% CIs 10-14% and 16-27%). These estimates are also within the range of infection
probabilities estimated from longitudinal studies (Monto et al., 1985; Horby et al., 2012; Hayward et al., 2014; Huang et al.,
2019). The fraction of children with detectable antibodies against B/Victoria was close to the prediction from our model,
but more children in the Netherlands had detectable B/Yamagata antibodies than predicted by the model for New Zealand
(Figure 3-Supplement 2). In addition to differences in lineage frequencies between New Zealand and the Netherlands, this
discrepancy might be due to the presence of antibodies from B/Victoria infections that cross-react with B/Yamagata but not
vice-versa (Rota et al., 1990; Levandowski et al., 1991a,b; Skowronski et al., 2012; Laurie et al., 2018).

Discussion

Our model suggests that differences in the age distributions of B/Victoria and B/Yamagata cases can be explained by historical
changes in lineage frequencies combined with cross-protection between strains of the same lineage and an additional protection
against B/Yamagata in people first infected with it. In particular, two major features of the age distributions seem to arise from
the dominance of B/Yamagata in New Zealand in the 1990s, which appears to to also have occurred elsewhere outside of East
Asia (Shaw et al., 2002). First, people born in the late 1980s and early 1990s had fewer B/Yamagata cases than B/Victoria
cases in the 2000s and 2010s because they had accumulated immunity against B/Yamagata but not B/Victoria during the

1990s. Second, imprinting of people born in the late 1980s and early 1990s with B/Yamagata led to fewer B/Yamagata cases in those birth cohorts than in older ones, most of whom were likely first infected before (or shortly after) the lineages split.

While imprinting protection against B/Yamagata has a clear effect on the age distributons of cases, the non-identifiability of imprinting protection against B/Victoria might be due to the lack of an extended period when B/Victoria circulated at consistently higher frequencies than B/Yamagata. Our analysis is limited by the scarcity of data on the antigenic phenotype of influenza B strains circulating prior to the lineage split and on lineage frequencies shortly after. B/Victoria seems to have been more common in the seasons preceding the discovery of the existence of two lineages (Rota et al., 1990), but surveillance in the United States suggests there were only two major influenza B epidemics between the most recent estimate of the lineage split in 1983 and the rise of B/Yamagata around 1990 (Rota et al., 1990). Thus, even if B/Victoria was the dominant lineage between the lineage split and the early 1990s, there might have been limited opportunity for B/Victoria imprinting. The high proportion of B/Yamagata cases and low proportion of B/Victoria cases in birth cohorts that were likely infected with strains circulating before the lineages split suggest those strains were antigenically more similar to B/Victoria than to B/Yamagata. Consistent with this possibility, the study that first identified the existence of two lineages in the late 1980s found that antibodies induced by B/Victoria strains from that period were more cross-reactive than those induced by B/Yamagata strains against a strain from the early 1980s (Rota et al., 1990).

While previous work revealed immune imprinting at the level of the groups and subtypes of influenza type A (Gostic et al., 2016, 2019; Arevalo et al., 2020), our results suggest a similar phenomenon occurs between more recently diverged and thus antigenically closer branches of influenza. The major antigens HA and NA evolve more slowly within the influenza B lineages than within influenza A subtypes (Bedford et al., 2014, 2015; Vijaykrishna et al., 2015), and their amino acid sequence divergences are much lower between B/Victoria and B/Yamagata ($\approx$14%, 2% and 7% in 2019 for the HA head, the HA stalk, and NA, respectively) than between H1N1 and H3N2 ($\approx$66%, 50% and 60%, respectively) (Figure 3-Supplement 3). Still, epitopes conserved within the lineages but variable across them (or between the lineages and their ancestor) might be the basis for imprinting protection against B/Yamagata (and potentially B/Victoria), as has been hypothesized for influenza A subtypes (Gostic et al., 2019).

Although protection from B/Victoria against B/Yamagata was non-identifiable, the model suggests protection from B/Yamagata against medically attended B/Victoria infections is weak at most. This lack of substantial protection from B/Yamagata against B/Victoria is consistent with the observation that B/Yamagata exposure often induces low or undetectable levels of antibodies cross-reactive to B/Victoria (Rota et al., 1990; Levandowski et al., 1991a,b; Skowronski et al., 2012; Laurie et al., 2018). However, the same studies of strains from the late 1980s, 2000s and 2010s show that exposure to B/Victoria induces antibodies that inhibit hemagglutination by B/Yamagata. Thus, while not identifiable from the age distributions of cases, protection from B/Victoria against B/Yamagata might occur. Meta-analyses suggest lineage-mismatched trivalent seasonal vaccines are generally effective (Tricco et al., 2013), but few studies report lineage-specific estimates of vaccine effectiveness that could reveal asymmetric cross-lineage protection. A test-negative design study estimated that inactivated vaccines containing B/Victoria had over 50% effectiveness against B/Yamagata infections, and vice-versa, in a population of mostly adults (Skowronski et al., 2019). However, a meta-analysis of randomized controlled trials of live attenuated vaccines in children estimated that the efficacy of vaccines containing B/Yamagata in seasons dominated by B/Victoria was low to nonexistent (B/Yamagata seasons in which a B/Victoria vaccine was used were not available) (Belshe et al., 2010). Individual longitudinal studies of infection or vaccination might be able to test the lack of protection from B/Yamagata exposures against B/Victoria inferred by our model and potentially reveal protection in the other direction. Studies of children who were likely first exposed to influenza B via vaccination might also reveal if imprinting can occur via trivalent vaccines containing only one influenza lineage or quadrivalent vaccines containing both.

Our results also suggest that the acquisition of a B/Yamagata NA gene by B/Victoria strains via reassortment in 2000-2001 (Dudas et al., 2015; Langat et al., 2017) did not lead to substantial protection against the reassortant strains in people previously infected with B/Yamagata. While most studies of the antibody response to influenza focus on HA, serological evidence suggests immune responses targeting NA instead of HA are more common against influenza B than against influenza A, especially in children (Huang et al., 2019). Yet birth cohorts with a history of B/Yamagata infections had a high proportion of clinical B/Victoria infections in the 2000s, even though all B/Victoria cases in the data occurred after B/Victoria had acquired a B/Yamagata NA.

The difference in the estimated strength of within-lineage protection without imprinting for B Victoria (85-96%) and B/Yamagata (10-43%) might be due to differences in the times of infection for the birth cohorts informing each of the estimates and different rates of antigenic evolution between the lineages. Because most people born since the late 1980s and infected

with B/Yamagata were imprinted with it, our estimate of within-lineage protection against B/Yamagata in the absence of immune imprinting is mostly informed by cases in older birth cohorts lacking imprinting protection against B/Yamagata. Most of the B/Yamagata infections in those birth cohorts would have occurred in the 1990s, when B/Yamagata was dominant. In contrast, our estimate of within-lineage non-imprinting protection against B/Victoria was informed by more recent B/Victoria infections in younger birth cohorts, including people who were imprinted with B/Yamagata in the 1990s and later infected with B/Victoria in the 2000s. Because these B/Victoria infections were more recent, they were potentially antigenically closer to modern strains than were the B/Yamagata infections that inform estimated within-lineage protection against B/Yamagata, and thus less likely to violate our assumption of constant within-lineage protection over time. However, B/Victoria appears to undergo slightly faster antigenic evolution than B/Yamagata (Bedford et al., 2014; Vijaykrishna et al., 2015). Longitudinal studies of infection risk might reveal if within-lineage protection against B/Yamagata in the absence of immune imprinting is stronger in cohorts born recently than estimated by our model.

It remains unclear precisely how differences in people's antibody responses and past infections shape their susceptibility to influenza. Our model and others (Gostic et al., 2016, 2019; Arevalo et al., 2020) suggest using birth cohorts as a proxy for shared infection history can capture broad effects of infection history on susceptibility to clinical infections. Characterizing antibody specificity after infection and vaccination might reveal the origins these patterns in terms of the antigens and epitopes targeted by people with different infection histories. Antibody targeting of NA and of particular HA epitopes is known to change with age (Nachbagauer et al., 2016; Rajendran et al., 2017; Sun et al., 2019), but few studies have linked antibody specificity to the infection history of particular birth cohorts and their susceptibility to particular variants (Linderman et al., 2014; Andrews et al., 2015; Petrie et al., 2016). This resolution could improve understanding of influenza epidemiology and the response to influenza vaccination.

Materials and methods

Case data

Medically attended influenza B cases in New Zealand were identified from samples taken from patients with influenza-like illness (ILI) attended by a network of sentinel general practitioners (2,430 cases with an identified influenza B lineage) and from non-sentinel hospital samples (1,606 cases with an identified lineage) analyzed by regional diagnostic laboratories and by the WHO National Influenza Centre at the Institute for Environmental Science and Research (ESR). Briefly, sentinel surveillance operates from May to September, with participating practices collecting nasopharyngeal or throat swabs from the first ILI patient examined on each Monday, Tuesday and Wednesday. ILI is defined as an "acute respiratory tract infection characterised by an abrupt onset of at least two of the following: fever, chills, headache and myalgia" (Huang et al., 2008). A subset of the New Zealand data (cases from 2002-2013) was previously compiled by Vijaykrishna et al. (2015) along with cases from Australia reported to the World Health Organization (WHO) Collaborating Centre for Reference and Research on Influenza in Melbourne.

Statistical model of influenza B susceptibility based on infection history

For lineage V (B/Victoria), we modeled the number of cases in people born in birth year b observed in season y as a multinomial draw with probabilities given by:

$$\theta_V(b, y) = D(b, y)\beta(b, y)Z_V(b, y)\rho(b, y) \tag{1}$$

with an analogous equation defining the multinomial distribution $\theta_Y(b, y)$ for lineage Y (B/Yamagata). $D(b, y)$ is the fraction of the population that was born in year b as of observation season y. $Z(b, y)$ is the susceptibility to lineage V during season y of a person born in year b relative to that of an unexposed person. $\beta(b, y)$ is a baseline probability of infection with influenza B that captures differences in transmission associated with age (thus depending on b and y) and is equal to β_1 if people born in year b are in preschool during season y (0-5 years old), β_2 if they are school-age children or teenagers (6-17 years old), or β_3 if they are 18 or older. $\rho(b, y)$ is an age-specific reporting rate equal to a parameter ρ if people are less than 2 years old and 1 otherwise. While $\beta(b, y)$ represents true differences in infection probabilities between preschoolers and post-preschool individuals and thus affects the infection history probabilities in the calculation of $Z(b, y)$, $\rho(b, y)$ represents differential reporting in children and does not affect those probabilities. For each lineage and observation season, values of θ from Eq. (1) are normalized by their sum across birth years to make them proper multinomial probabilities.

We defined relative susceptibility to V, $Z_V(b, y)$, as an expectation over all possible immune histories in terms of the lineage of first infection and subsequent infection with the other lineage. We let susceptibility be 1 for a person never

³⁷¹ exposed to influenza B. Cross-immunity from any previously encountered strain of V or Y decreased susceptibility to the

³⁷² corresponding lineage by χ_{VV} and χ_{YY}, respectively. Susceptibility was further reduced by R_V or R_Y if the lineage of first

³⁷³ infection was V or Y. In the absence of a previous homologous infection, previous infection with Y decreased susceptibility

³⁷⁴ to V by χ_{VY}, and previous infection to V decreased susceptibility to Y by χ_{YV}. Protection due to homologous infections

³⁷⁵ superseded cross-lineage protection. Protection against V from a previous Y infection was constrained to be a fraction of

³⁷⁶ the within-lineage protection to V (and vice-versa): $\chi_{VY} = \chi_{VV} \cdot \gamma_{VY}$, $\chi_{YV} = \chi_{YY} \cdot \gamma_{YV}$, where $0 \le \gamma_{VY}, \gamma_{YV} \le 1$. Similarly,

³⁷⁷ pre-1988 infections reduced susceptibility to V by $\chi_{VA} = \chi_{VV} \cdot \gamma_{VA}$ and to Y by $\chi_{YA} = \chi_{YY} \cdot \gamma_{YA}$. Finally, $Z_V(b, y)$ was

³⁷⁸ calculated as the sum of susceptibilities in Table 1 weighted by the probabilities of the corresponding infection histories

³⁷⁹ (below). Relative susceptibility to Y, $Z_Y(b, y)$ was defined analogously.

³⁸⁰ We estimated parameters by maximum likelihood using R (version 3.4.3) and package optimParallel. We calculated the

³⁸¹ total likelihood as the product of the likelihood for each lineage in each observation year, with the number of cases in each

³⁸² combination treated as an independent multinomial draw. For plotting, we summed the observed and predicted cases for each

³⁸³ birth year across observation years.

³⁸⁴ Code implementing the analyses and figures is available at `https://github.com/cobeylab/influenza_B_distributions`.

³⁸⁵ Infection history probabilities

³⁸⁶ To calculate the probabilities in Table 1, we assumed infections occur in discrete time measured in units of annual influenza

³⁸⁷ seasons. We considered possible infections in each season between birth and the last season before observation season y.

³⁸⁸ For a person born in year b and previously unexposed to influenza B, we let $a_{b,i}$ be the probability of becoming infected in

³⁸⁹ season i. This probability was then modified by protection from previous infections as in Table 1. Given that an infection

³⁹⁰ occurred in season i, we assumed that the probability it was caused by lineages A ("ancestor"), V or Y was equal to their

³⁹¹ frequencies in that season, $f_{A,i}$, $f_{V,i}$ and $f_{Y,i}$, with $f_{A,i} = 1$ for all i before 1988, and $f_{V,i} + f_{Y,i} = 1$ since (Figure 1). For

³⁹² simplicity, we assumed that people could not be infected more than once in each season (including simultaneous infections by

³⁹³ the two lineages.)

Let $\Phi_{i,j}^B$ be the probability that no infections with influenza B occurred for a naive person born in b from seasons i to j (inclusive). It is given by:

$$\Phi_{i,j}^B = \prod_{k=i}^{j}(1 - a_{b,k}) \tag{2}$$

where k indexes years (influenza seasons). Thus the first probability in Table 1, that of being fully naive to influenza B, is given by:

$$P_0(b, y) = \Phi_{b,y-1}^B \tag{3}$$

To shorten the expressions for the remaining probabilities, we let $\Phi_{i,j}^V(h)$, $\Phi_{i,j}^Y(h)$ and $\Phi_{i,j}^{VY}(h)$ be the probability that no V infections, no Y infections, and neither V nor Y infections occurred between seasons i and j (inclusive), respectively. Unlike Φ^B, which applies to naive people, Φ^V, Φ^Y and Φ^{VY} depend on the person's infection history, h. To calculate the probability of an initial infection with A but no subsequent infections with V or Y, $P_{A,0,0}(b, y)$, we integrated, across all possible seasons i of first infection, the joint probability that the person's first influenza B infection occurred in season i with the ancestral lineage A and that no subsequent infections with V or Y occurred from $i + 1$ to $y - 1$:

$$P_{A,0,0}(b, y) = \sum_{i=b}^{y-1} \left[\Phi_{b,i-1}^B \cdot a_{b,i} \cdot f_{A,i} \cdot \Phi_{i+1,y-1}^{VY}(A) \right] \tag{4}$$

where the probability that the first infection with influenza B occurred in season i with the ancestor A is obtained by multiplying the probability of no previous infections ($\Phi_{b,i-1}^B$) by the probability of an infection in season i ($a_{b,i}$) and by the frequency of lineage A in season i ($f_{A,i}$). The probability of no infections with either V or Y after the initial infection with A in season i, $\Phi_{i+1,y-1}^{VY}(A)$, depends on protection from the A infection against V (χ_{VA}) and Y (χ_{YA}):

$$\Phi_{i+1,y-1}^{VY}(A) = \prod_{k=i+1}^{y-1} \{1 - a_{b,k}[f_{V,k}(1 - \chi_{VA}) + f_{Y,k}(1 - \chi_{YA})]\} \tag{5}$$

Similarly, the joint probability of first infection with the ancestor A and subsequent infection with V, but not with Y, is given by:

$$P_{A,V,0}(b,y) =$$
$$\sum_{i=b}^{y-1} \left[\Phi_{b,i-1}^{B} a_{b,i} f_{A,i} \cdot \sum_{j=i+1}^{y-1} \Phi_{i+1,j-1}^{VY}(A) a_{b,j} f_{V,j} (1 - \chi_{VA}) \Phi_{j+1,y+1}^{Y}(A,V) \right] \tag{6}$$

where we again integrated over all possible seasons i when the first infection with influenza B might have occurred. Given the
first infection occurred in season i with the "ancestral" lineage A, we calculated the probability of subsequent infection with
V, but not with Y, by integrating over all possible seasons j when the first infection with V may have occurred. Given the
initial infection with A in season i, the joint probability that the first V infection occurred in j is given by the probability that
neither V nor Y infections occurred from $i+1$ to $j-1$, $\Phi_{i+1,j-1}^{VY}(A)$, times the probability of a V infection in season j, given
by $a_{b,j} f_{V,j} (1 - \chi_{VA})$. The probability $\Phi_{j+1,y-1}^{Y}(A,V)$ of no subsequent Y infections after season j given the previous A and V
infections is then given by:

$$\Phi_{j+1,y-1}^{Y}(A,V) = \prod_{k=j+1}^{y-1} \{1 - a_{b,k} f_{Y,k} [1 - \max(\chi_{YA}, \chi_{YV})]\} \tag{7}$$

where we assumed that only the strongest cross-protection (from the previous A and V infections) applies. By analogy, for $P_{A,Y,0}(b,y)$ we have:

$$P_{A,Y,0}(b,y) =$$
$$\sum_{i=b}^{y-1} \left[\Phi_{b,i-1}^{B} a_{b,i} f_{A,i} \cdot \sum_{j=i+1}^{y-1} \Phi_{i+1,j-1}^{VY}(A) a_{b,j} f_{Y,j} (1 - \chi_{YA}) \Phi_{j+1,y+1}^{V}(A,Y) \right] \tag{8}$$

where

$$\Phi_{j+1,y-1}^{V}(A,Y) = \prod_{k=j+1}^{y-1} \{1 - a_{b,k} f_{V,k} [1 - \max(\chi_{VA}, \chi_{VY})]\} \tag{9}$$

To calculate $P_{A,\{VY\}}(b,y)$, we first computed the probabilities for the particular cases where either V or Y were the second infection, $P_{A,V\to Y}(b,y)$ and $P_{A,Y\to V}(b,y)$, such that $P_{A,\{VY\}}(b,y)$ is the sum of the two. The first is given by:

$$P_{A,V\to Y}(b,y) =$$
$$\sum_{i=b}^{y-1} \left[\Phi_{b,i-1}^{B} a_{b,i} f_{A,i} \cdot \sum_{j=i+1}^{y-1} \Phi_{i+1,j-1}^{VY}(A) a_{b,j} f_{V,j} (1 - \chi_{VA}) [1 - \Phi_{j+1,y+1}^{Y}(A,V)] \right] \tag{10}$$

and the second is given by:

$$P_{A,Y\to V}(b,y) =$$
$$\sum_{i=b}^{y-1} \left[\Phi_{b,i-1}^{B} a_{b,i} f_{A,i} \cdot \sum_{j=i+1}^{y-1} \Phi_{i+1,j-1}^{VY}(A) a_{b,j} f_{Y,j} (1 - \chi_{YA}) [1 - \Phi_{j+1,y+1}^{V}(A,Y)] \right] \tag{11}$$

Next we write down the probabilities of infection histories with either V or Y as first infections and no subsequent infections. For $P_{V,0}(b,y)$:

$$P_{V,0}(b,y) = \sum_{i=b}^{y-1} \Phi_{b,i-1}^{B} \cdot a_{b,i} \cdot f_{V,i} \cdot \Phi_{i+1,y-1}^{Y}(V) \tag{12}$$

where

$$\Phi_{i+1,y-1}^{Y}(V) = \prod_{k=i+1}^{y-1} [1 - a_{b,k} f_{Y,k} (1 - \chi_{YV})] \tag{13}$$

By analogy, for $P_{Y,0}(b,y)$:

$$P_{Y,0}(b,y) = \sum_{i=b}^{y-1} \Phi_{b,i-1}^{B} \cdot a_{b,i} \cdot f_{Y,i} \cdot \Phi_{i+1,y-1}^{V}(Y) \tag{14}$$

where

$$\Phi_{i+1,y-1}^V(Y) = \prod_{k=i+1}^{y-1} [1 - a_{b,k} f_{V,k}(1 - \chi_{VY})] \tag{15}$$

Finally, $P_{V,Y}(b, y)$ and $P_{Y,V}(b, y)$ are given by:

$$P_{V,Y}(b, y) = \sum_{i=b}^{y-1} \Phi_{b,i-1}^B \cdot a_{b,i} \cdot f_{V,i} \cdot [1 - \Phi_{i+1,y-1}^Y(V)] \tag{16}$$

$$P_{Y,V}(b, y) = \sum_{i=b}^{y-1} \Phi_{b,i-1}^B \cdot a_{b,i} \cdot f_{Y,i} \cdot [1 - \Phi_{i+1,y-1}^V(Y)] \tag{17}$$

Because case data had information on age and not the exact birth year, we averaged each exposure history probability across the two possible birth years given the age and the observation year (for instance, a 10-year-old in 2000 may have been born in either 1989 or 1990). Because the probabilities of different infection histories become very similar for cohorts born long before the lineages split, we used the probabilities calculated for the birth year 1970 for all previous cohorts to decrease computation time.

Season-specific attack rates

Let $P_{\text{inf}}(b, i, t)$ be the probability that a previously unexposed person born in year b has been infected with influenza B after experiencing fraction $t \in [0, 1]$ of season i. Assuming a constant instantaneous attack rate $\alpha_{b,i}$ throughout the season, $P_{\text{inf}}(b, i, t)$ is given by:

$$P_{\text{inf}}(b, i, t) = 1 - e^{-\alpha_{b,i} t} \tag{18}$$

We let the instantaneous attack rate $\alpha_{b,i}$ be equal to an age-specific baseline multiplied by an intensity score S_i representing the strength of influenza B circulation in season i relative to other seasons:

$$\alpha_{b,i} = -\ln[1 - \beta(b, y)] \cdot S_i, \tag{19}$$

where $\beta(b, y)$ takes on value β_1, if birth year b corresponds to an age of less than 5 in year y, β_2, if the corresponding age is 6-17, and β_3, for ages 18 and older. The probability of infection for an unexposed person born in year b across the entire season, $a_{b,i}$, is obtained by substituting $\alpha_{b,i}$ in Eq. [18] and setting $t = 1$:

$$a_{b,i} = 1 - e^{-\alpha_{b,i}} = 1 - [1 - \beta(b, y)]^{S_i} \tag{20}$$

The definition of $\alpha_{b,i}$ in Eq. (19) was chosen such that for a season with average influenza intensity ($S_i = 1$), the annual probability of infection for an unexposed person is equal to $\beta(b, y)$.

For the season corresponding to the first year of life ($i = b$), people are only susceptible to infections during a fraction of the season, depending on when they were born and how long they were protected by maternal antibodies. In those cases, we defined $a_{b,i}$ as the expected probability of infection across all possible weeks of birth:

$$a_{b,i} = \frac{1}{W_b} \sum_w 1 - e^{-\alpha_{b,i} \phi_i(b, w+M)}, \quad \text{for } i = b \tag{21}$$

where $\phi_i(b, w)$ is the fraction of cases in season i observed in or after week w of year b and W_b is the number of weeks in year b. Because people are assumed to be completely protected against infection by maternal antibodies for the first M weeks following birth, ϕ_i was computed for an effective birth week $w + M$. Based on the fraction of children under the age of 1 with detectable antibodies to influenza B (Bodewes et al., 2011), we set M to 26 weeks (approximately 6 months). Averaging $\phi_i(b, w + M)$ over all possible birth weeks w in year b gives the expected fraction of season i experienced by a person born in year b assuming births are distributed uniformly in time. We estimated $\phi_i(b, w)$ by fitting the incomplete beta function to the cumulative fraction of cases in the seasons for which we had case data, using R package FlexParamCurve. Following Gostic et al. (2016), we truncated season-specific infection probabilities so that they never exceed 0.75 even in years of high estimated influenza B intensity.

Intensity scores

We defined the intensity score S_i in Eq. (19) as:

$$S_i = \frac{\text{\% influenza B in ILI specimens} \times \text{ILI incidence, for season } i}{\text{mean [\% influenza B in ILI specimens} \times \text{ILI incidence] across seasons}} \tag{22}$$

Annual influenza surveillance reports from New Zealand's Institute of Environmental Science and Research (ESR) available from 2003 on (Institute of Environmental Science and Research, 2020) give the "isolation" or "detection" rate (the number of influenza-positive swabs divided by the number of swabs tested), the percentage of influenza A and B viruses among all influenza-positive isolates (both sentinel and non-sentinel), and the estimated number of ILI cases in New Zealand for each season. The reports do not directly give the fraction of ILI specimens that were influenza B-positive. Instead, we calculated the fraction of ILI isolates that were influenza B-positive in a season by multiplying the fraction of ILI isolates that were influenza-positive by the fraction of influenza-positive specimens that were influenza B. For seasons without data on the fraction of influenza-positive specimens in ILI specimens (1988 to 2000), without data on the fraction of influenza B in influenza-positive specimens (1988-89), or without estimates of the total number of ILI cases (1988-2001), we used the average values of those quantities across the remaining seasons. Although reports are not available for 1990-2002, the 2003 annual report lists the frequency of influenza B in influenza-positive specimens for those seasons.

The World Health Organization's FluNet has weekly data on the fraction of ILI specimens that were influenza B-positive in Australia from 1997 to the present. However, in data from before 2003, the number of influenza-positive specimens was usually the same as the reported number of specimens processed for that week, suggesting strong case ascertainment or reporting bias. We thus used data on the percent of influenza-positive ILI cases from 2003 on. Annual influenza reports from 1994 to 2010 are available from the Australian government's Department of Health website (Australian Government Department of Health, 2019). They report numbers of influenza A- and B-positive isolates but not the total number of specimens tested. Thus, only the fraction of influenza B in influenza isolates (but not in ILI isolates) can be estimated from those reports. We therefore used data from the WHO to calculate the fraction of influenza B-positive specimens in ILI specimens for 2003-2017. For 1994-2002, we multiplied the fraction of influenza B in influenza-positive specimens for each season (from the Department of Health reports) by the average annual fraction of influenza-positive specimens in ILI specimens from 2003-2017 (from the WHO data) to arrive at the fraction of influenza B positive ILI specimens. Finally, for seasons where data were missing altogether (1988-1993), we used the average annual fraction of influenza B positive specimens in ILI specimens for subsequent seasons (1994-2017).

To estimate ILI incidence in Australia, we used the maximum weekly number of ILI cases per 1000 consultations for each season from Department of Health annual and weekly reports (weekly reports are available for years since the last annual report in 2010). Different ILI definitions were used from 1994-2003 and from 2004-2010, and starting in 2009 reported weekly ILI rates were averaged from multiple branches of the Australian influenza surveillance system. We thus normalized values by the average value within each of those periods (1994-2003, 2004-2008 and 2009-18) to arrive at a normalized peak number of ILI cases per 1000 consultations in Australia.

Historical frequencies of influenza B lineages

To estimate historical frequencies of B/Victoria and B/Yamagata, we downloaded data on lineage and date and country of isolation for all influenza B isolates on the Global Initiative on Sharing All Influenza Data (GISAID) website collected until 09/30/2018. To complement these data, we searched the NCBI Influenza Virus Database for all protein-coding HA sequences of influenza B viruses isolated from humans and excluding laboratory strains (information on passage history for GISAID entries was scarce and non-standardized and so we did not filter out laboratory strains from the GISAID data). Because lineage information was missing for virtually all sequences retrieved from NCBI, we used BLAST to assign each sequence to either B/Victoria or B/Yamagata based on the highest bit score match with reference sequences B/Victoria/2/87 and B/Yamagata/16/1988.

We combined data from both databases to estimate the frequency of B/Yamagata and B/Victoria isolates in each season. Isolates collected in year y in Europe or North America were assigned to season $y-1/y$, if collected before October, and to season $y/y+1$, if collected in October-December. Because most European and North American isolates were collected before October of the respective year (median across years = 83% for Gisaid and 82% for NCBI), we assumed isolates with missing month of collection in those regions were collected before October (and thus belonged to the season beginning in the previous calendar year).

⁴⁶⁹ Isolates with the same name but reported for different countries or seasons were considered separately. We condensed
⁴⁷⁰ multiple occurrences of the same isolate in the same country and season (within or across datasets) into one, disregarding
⁴⁷¹ isolates for which different lineages were assigned in different countries/seasons. Using isolates present in both databases,
⁴⁷² we found that our BLAST lineage assignment matched the lineage reported on GISAID in 98% (3,159/3,217) of cases. We
⁴⁷³ disregarded isolates for which our BLAST assignment and the reported GISAID assignment disagreed. The final dataset
⁴⁷⁴ consisted of 35,158 isolates, 23 of which (0.07%) were represented more than once (in different countries or seasons). We
⁴⁷⁵ estimated the frequency of a lineage as the number of isolates belonging to that lineage divided by the total number of influenza
⁴⁷⁶ B isolates collected in a season.

⁴⁷⁷ Because all 23 isolates collected in New Zealand and Australia in the 1990s were B/Yamagata (Figure 1-Supplement 2),
⁴⁷⁸ and because previous work suggests B/Victoria circulated only in East Asia during that period (Shaw et al., 2002), we assumed
⁴⁷⁹ the frequency of B/Yamagata in New Zealand and Australia to be 100% from 1988 to 2000, even though no isolates were
⁴⁸⁰ available for several individual years within that range. In 2001 and 2003, when both lineages are known to have been
⁴⁸¹ circulating in New Zealand and Australia but fewer than 10 isolates were available for the two countries combined in the
⁴⁸² sequence databases, we used frequencies estimated from all other countries combined. Frequencies estimated from all other
⁴⁸³ countries combined were strongly correlated with estimates based on isolates from Australia and New Zealand only (Pearson's
⁴⁸⁴ correlation coefficient = 0.91, 95% CI 0.81-0.96). We also considered using frequencies estimated from isolates collected in
⁴⁸⁵ the United States, which were also correlated with frequencies in Australia and New Zealand (Figure 1-Supplement 2), but
⁴⁸⁶ the correlation was weaker (0.73, 95% CI 0.48-0.88). For the sensitivity analysis of lineage frequencies in the 1990s, we
⁴⁸⁷ re-fitted the model using frequencies from all countries combined for all years from 1988 on in which fewer than 10 isolates
⁴⁸⁸ were available for New Zealand and Australia combined, including years in the 1990s.

⁴⁸⁹ We hoped to use the sequence databases to get more reliable estimates of lineage frequencies in the 1980s than those
⁴⁹⁰ provided by early antigenic characterization (Rota et al., 1990), but fewer than 10 isolates were available for each year before
⁴⁹¹ 1988. To accommodate uncertainty, we grouped infections with B/Victoria and B/Yamagata before 1988 with infections by
⁴⁹² the ancestral influenza B strains circulating before the lineages split.

⁴⁹³ We compared our estimates of lineage frequencies based on sequence data to estimates based on antigenic characterization
⁴⁹⁴ of circulating strains from epidemiological surveillance reports (Figure 1-Supplement 2). Surveillance reports from Australia
⁴⁹⁵ are available from the Australian Government's Department of Health website (Australian Government Department of Health,
⁴⁹⁶ 2019). Surveillance reports from New Zealand are available from the website of New Zealand's Institute of Environmental
⁴⁹⁷ Science and Research (ESR) (Institute of Environmental Science and Research, 2020). Although reports from New Zealand
⁴⁹⁸ are only available from 2003 on, Fig. 27 of the 2012 report shows B/Victoria and B/Yamagata frequencies from 1990 to 2002
⁴⁹⁹ (without reporting the number of isolates used to estimate those frequencies). Annual summaries of influenza surveillance in
⁵⁰⁰ the United States are published by the Centers for Disease Control and Prevention (e.g., Garten et al. (2018)).

Model validation with independent serological data

⁵⁰² We compared the fraction of children predicted by our model to have been previously infected with B/Victoria, B/Yamagata or
⁵⁰³ either lineage with the fraction of children that had detectable antibodies against the corresponding lineage (or any influenza B
⁵⁰⁴ strain) in the Netherlands (Bodewes et al., 2011). Sera from children 0-7 years old collected between February 2006 and June
⁵⁰⁵ 2007 were tested using the hemagglutination inhibition assay against a panel of reference B/Victoria and B/Yamagata strains
⁵⁰⁶ as well of strains isolated in the Netherlands during the study period. Sera were considered positive if their hemagglutination
⁵⁰⁷ inhibition titer was ≥ 10 against at least one strain from the corresponding set (all influenza B strains, B/Victoria strains, and
⁵⁰⁸ B/Yamagata strains). We compared these data with predictions under the maximum likelihood-parameter estimates of our
⁵⁰⁹ model fitted to the complete New Zealand data.

⁵¹⁰ We also used the seroprevalence data to independently estimate the annual probability of infection for preschoolers and
⁵¹¹ school-age children, equivalent to the β_1 and β_2 parameters in our model. Assuming a constant instantaneous attack rate α, an
⁵¹² individual of age A years is still naive (and therefore seronegative) to influenza B with probability $P_N(A)$ given by:

$$P_N(A) = e^{-\alpha A} \tag{23}$$

⁵¹³ The probability of observing X seronegative individuals in a sample of n individuals of age A can be calculated assuming
⁵¹⁴ $X \sim \text{Binomial}[n, P_N(A)]$, and α thus can be estimated by maximum likelihood. The annual attack rate can then be calculated
⁵¹⁵ from the instantaneous attack rate α as $\beta_{\text{Netherlands}} = 1 - e^{-\alpha}$.

We make two modifications to Eq. (23) to account for the presence of maternal antibodies early in life and for uncertainty
in the age of individuals when their serum was collected. First, we assume individuals spend a time m (in units of years)
fully protected against influenza B due to the presence of maternal antibodies. Consistent with the fraction of children under
the age of 1 with detectable antibodies to influenza B (Bodewes et al., 2011), we assumed $m = 0.5$ year. Second, because
ages were reported at the resolution of one year (e.g. an individual 2.6 years old is reported as being 2 years old), we assume
individuals with recorded age A were sampled at a randomly distributed time $T \in [0, 1)$ during the interval between the ages
of A and $A + 1$. Thus, we let $P_N(A)$ be given by the expectation over T:

$$P_N(A) = E[e^{-\alpha(A+T-m)}] = \int_0^1 e^{-\alpha(A+t-m)} f(t) dt \tag{24}$$

where $f(t)$ is the probability density function of T. Assuming T is uniformly distributed between 0 and 1 (i.e., $f(t) = 1$), we
have:

$$
\begin{aligned}
P_N(A) &= \int_0^1 e^{-\alpha(A+t-m)} dt \\
&= e^{-\alpha(A-m)} \int_0^1 e^{-\alpha t} dt \\
&= e^{-\alpha(A-m)} \left[\frac{-e^{-\alpha t}}{\alpha} \right] \Big|_0^1 \\
&= e^{-\alpha(A-m)} \frac{(1 - e^{-\alpha})}{\alpha}
\end{aligned}
\tag{25}
$$

523 valid for $A > m$ and $\alpha > 0$. Letting α_1 and α_2 be the instantaneous attack rates for preschoolers and school-age children:

$$
P_N(A) = \begin{cases} e^{-\alpha_1(A-m)} \frac{(1-e^{-\alpha_1})}{\alpha_1}, & \text{if } A \le A_s \\ e^{-\alpha_1(A_s-m)} e^{-\alpha_2(A-A_s)} \frac{(1-e^{-\alpha_2})}{\alpha_2}, & \text{if } A > A_s \end{cases}
\tag{26}
$$

524 where A_s is the age at which children start going to school (4 years old in the Netherlands; Government of the Netherlands
525 (2020)). Note that for school-age children (the equation for $A > A_s$ on the bottom) the correction term for uncertainty in
526 sampling is not necessary for the time spent in preschool (assumed to be exactly A_s years), only for the time after preschool
527 $(A - A_s)$.

Handling cases with missing lineage information

We assumed cases with missing lineage information in 2002 ($n = 61$), 2011 ($n = 312$) and 2019 ($n = 206$) belonged to
B/Victoria, since 99% or more of identified cases in those seasons were B/Victoria (86/87 cases in 2002, 276/280 cases in
2011, 552/552 cases in 2019) as were 94%, 92% and 92% of isolates from sequence databases (for Australia and New Zealand
combined). We assumed cases with missing lineage information belonged to B/Yamagata in 2013 ($n = 37$), 2014 ($n = 77$)
and 2017 ($n = 87$), when the majority of identified cases were B/Yamagata (268/272, 131/138 and 473/489, respectively), as
were 99%, 94% and 84% of isolates in sequence databases. Unidentified cases in other seasons were disregarded because both
lineages were present at higher frequencies among identified cases. Removing unidentified cases altogether in all seasons led
to similar parameter estimates.

Sequence divergence analysis

To estimate the amount of evolution within and between lineages, we analyzed all complete HA and NA sequences from
human influenza B isolates available on GISAID in July 2019. The set of isolates used in this analysis differs from the set used
to estimate lineage frequencies because we required isolates to have complete sequences (although not all sequences listed as
complete on GISAID were in fact complete). Two isolates collected in 2000 (B/Hong Kong/548/2000 and B/Victoria/504/2000)
were deposited as B/Victoria but our BLAST assignment indicated they were in fact B/Yamagata (their low divergence from
B/Yamagata strains was a clear outlier). NA sequences from isolates B/Kanagawa/73 and B/Ann Arbor/1994 were only small
fragments (99 and 100 amino acids long) poorly aligned with other sequences and were thus excluded. We also excluded NA
sequences from B/Yamagata isolates B/Catalonia/NSVH100773835/2018 and B/Catalonia/NSVH100750997/2018 because
they were extremely diverged (60% and 38%) from the reference strain B/Yamagata/16/88 and aligned poorly with other
sequences.

To compare sequence diversity within and between lineages over time, we aligned sequences using MAFFT v. 7.310
(Katoh et al., 2002) and calculated percent amino acid differences in pairs of sequences from the same lineage and in pairs
with one sequence from each. For each year, we sampled 100 sequences from each lineage (or used all sequences if 100 or
fewer were available) to limit the number of pairwise calculations. To estimate how much B/Yamagata and B/Victoria evolved
since the late 1980s, we calculated percent amino acid differences between each B/Yamagata and B/Victoria sequence and the
corresponding HA and NA sequences of reference strains B/Yamagata/16/88 and B/Victoria/2/87. Unlike in the analysis of
pairwise divergence within each time point, we used all sequences from each lineage in each year. We excluded sites in which
one or both sequences had gaps or ambiguous amino acids.

To compare HA and NA divergence between influenza B lineages with divergence between influenza A subtypes, we
downloaded complete HA and NA sequences from H3N2 and H1N1 isolated since 1977 and available on GISAID in August
2019. Homologous sites in the HA of H3N2 and H1N1 are difficult to identify by conventional sequence alignment, and instead
we used the algorithm by Burke & Smith (Burke and Smith, 2014) implemented on the Influenza Research Database website
(Influenza Research Database, 2019). Both H3N2 and H1N1 sequences were aligned with the reference H3N2 sequence
A/Aichi/2/68. We verified that this method matched sites on the stalk and head of the H1N1 HA with sites on the stalk and head
of H3N2 HA by comparing the resulting alignment with the alignment in Fig. S2 of Kirkpatrick et al. (2018). To limit the total
number of influenza A sequences analyzed we randomly selected 100 H3N2 and 100 H1N1 sequences for years in which more
than 100 sequences were available and used all available sequences for the remaining years. Isolates A/Canterbury/58/2000,
A/Canterbury/87/2000 and A/Canterbury/55/2000 were excluded because both H1N1-like and H3N2-like sequences were
available under the same isolate name on GISAID.

Demographic data

568 We obtained annual age distributions for the general population in New Zealand from StatsNZ (StatsNZ, 2019) and in Australia
569 from Australian Bureau of Statistics (Australian Bureau of Statistics, 2019). When calculating changes in the mean age of the
570 population over time, we excluded people over 90 years old, who were pooled together in New Zealand's demographic tables.

Acknowledgments

Frank Wen helped with data collection from GISAID. This work was completed in part with resources provided by the
University of Chicago Research Computing Center. This project has been funded in part with Federal funds from the National
Institute of Allergy and Infectious Diseases, National Institutes of Health, Department of Health and Human Services under
grant DP2 AI117921 and CEIRS Contract No. HHSN272201400005C awarded to S.C. M.V. was supported by a William
Rainey Harper Dissertation Fellowship by the University of Chicago. K.K. was supported by MIDAS CIDID Center of
Excellence (U54-GM111274), V.D. was supported by NIH CEIRS Contract No. HHSN272201400006C, C.D. was supported
by an Early Carreer Fellowship (1113269) from the Australian National Health and Medical Research Council. The content is
solely the responsibility of the authors and does not necessarily represent the official views of the funding sources.

Competing interests

581 No competing interests declared.

References

Anderson RM, Crombie JA, Grenfell BT. The epidemiology of mumps in the UK: a preliminary study of virus transmission, herd immunity
and the potential impact of immunization. Epidemiology & Infection. 1987 Aug; 99(1):65–84. doi: 10.1017/S0950268800066875.

Andrews SF, Huang Y, Kaur K, Popova LI, Ho IY, Pauli NT, Henry Dunand CJ, Taylor WM, Lim S, Huang M, Qu X, Lee JH, Salgado-Ferrer
M, Krammer F, Palese P, Wrammert J, Ahmed R, Wilson PC. Immune history profoundly affects broadly protective B cell responses to
influenza. Science translational medicine. 2015 Dec; 7(316):316ra192. `http://stm.sciencemag.org/content/7/316/316ra192`,
doi: 10.1126/scitranslmed.aad0522.

Arevalo P, McLean HQ, Belongia EA, Cobey S. Earliest infections predict the age distribution of seasonal influenza A cases. eLife. 2020
Jul; 9:e50060. `https://doi.org/10.7554/eLife.50060`, doi: 10.7554/eLife.50060.

Australian Bureau of Statistics, Estimated Resident Population By Single Year Of Age, Australia. Accessed July 5, 2019.;
2019. `https://www.abs.gov.au/AUSSTATS/ABS@Archive.nsf/log?openagent&3101059.xls&3101.0&Time%20Series%`
`20Spreadsheet&AF756051EBD9FEA7CA2581470023DEE3&0&Dec%202016&27.06.2017&Latest`.

Australian Government Department of Health, National Influenza Surveillance Scheme annual reports. Accessed June 21, 2019.; 2019.
`https://www.health.gov.au/internet/main/publishing.nsf/Content/cda-pubs-annlrpt-fluannrep.htm`.

Bedford T, Riley S, Barr IG, Broor S, Chadha M, Cox NJ, Daniels RS, Gunasekaran CP, Hurt AC, Kelso A, Klimov A, Lewis NS, Li X,
McCauley JW, Odagiri T, Potdar V, Rambaut A, Shu Y, Skepner E, Smith DJ, et al. Global circulation patterns of seasonal influenza viruses
vary with antigenic drift. Nature. 2015 Jun; 523:217–220. `http://dx.doi.org/10.1038/nature14460`, doi: 10.1038/nature14460.

Bedford T, Suchard Ma, Lemey P, Dudas G, Gregory V, Hay AJ, McCauley JW, Russell Ca, Smith DJ, Rambaut A. Integrating influenza
antigenic dynamics with molecular evolution. eLife. 2014 Jan; 3:e01914. `http://www.pubmedcentral.nih.gov/articlerender.`
`fcgi?artid=3909918&tool=pmcentrez&rendertype=abstract`, doi: 10.7554/eLife.01914.

Belshe RB, Coelingh K, Ambrose CS, Woo JC, Wu X. Efficacy of live attenuated influenza vaccine in children against influenza B viruses
by lineage and antigenic similarity. Vaccine. 2010 Feb; 28(9):2149–2156. `http://www.sciencedirect.com/science/article/`
`pii/S0264410X09018714`, doi: 10.1016/j.vaccine.2009.11.068.

Bodewes R, Mutsert Gd, Klis FRMvd, Ventresca M, Wilks S, Smith DJ, Koopmans M, Fouchier RaM, Osterhaus ADME, Rimmelzwaan GF.
Prevalence of Antibodies against Seasonal Influenza A and B Viruses in Children in Netherlands. Clinical and Vaccine Immunology. 2011
Mar; 18(3):469–476. `https://cvi.asm.org/content/18/3/469`, doi: 10.1128/CVI.00396-10.

Burke DF, Smith DJ. A Recommended Numbering Scheme for Influenza A HA Subtypes. PLOS ONE. 2014 Nov; 9(11):e112302.
`https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0112302`, doi: 10.1371/journal.pone.0112302.

Bush RM, Fitch WM, Bender CA, Cox NJ. Positive selection on the H3 hemagglutinin gene of human influenza virus A. Molecular
Biology and Evolution. 1999 Nov; 16(11):1457–1465. `https://academic.oup.com/mbe/article/16/11/1457/2925391`, doi:
10.1093/oxfordjournals.molbev.a026057.

Caini S, Kusznierz G, Garate VV, Wangchuk S, Thapa B, Júnior FJdP, Almeida WAFd, Njouom R, Fasce RA, Bustos P, Feng L, Peng
Z, Araya JL, Bruno A, Mora Dd, Gámez MJBd, Pebody R, Zambon M, Higueros R, Rivera R, et al. The epidemiological signature
of influenza B virus and its B/Victoria and B/Yamagata lineages in the 21st century. PLOS ONE. 2019 Sep; 14(9):e0222381. `https:`
`//journals.plos.org/plosone/article?id=10.1371/journal.pone.0222381`, doi: 10.1371/journal.pone.0222381.

Chen R, Holmes EC. The Evolutionary Dynamics of Human Influenza B Virus. Journal of Molecular Evolution. 2008 May; 66(6):655.
`https://doi.org/10.1007/s00239-008-9119-z`, doi: 10.1007/s00239-008-9119-z.

Davies JR, Grilli EA, Smith AJ. Influenza A: infection and reinfection. Epidemiology & Infection. 1984 Feb; 92(1):125–127. `https://www.`
`cambridge.org/core/journals/epidemiology-and-infection/article/influenza-a-infection-and-reinfection/`
`AC35CB3A3029B02A6711317FEFEE5C99`, doi: 10.1017/S002217240006410X.

Dudas G, Bedford T, Lycett S, Rambaut A. Reassortment between Influenza B Lineages and the Emergence of a Coadapted PB1–PB2–HA
Gene Complex. Molecular Biology and Evolution. 2015 Jan; 32(1):162–172. `https://academic.oup.com/mbe/article/32/1/`
`162/2925578`, doi: 10.1093/molbev/msu287.

Ferguson NM, Galvani AP, Bush RM. Ecological and immunological determinants of influenza evolution. Nature. 2003 Mar; 422(6930):428–
433. `https://www.nature.com/articles/nature01509`, doi: 10.1038/nature01509, number: 6930 Publisher: Nature Publishing
Group.

Frank AL, Taber LH. Variation in frequency of natural reinfection with influenza A viruses. Journal of Medical Virology. 1983; 12(1):17–23.
`https://onlinelibrary.wiley.com/doi/abs/10.1002/jmv.1890120103`, doi: 10.1002/jmv.1890120103.

Galbraith NS, Pusey J, Young SJ, Crombie DL, Sparks JP. Mumps Sruveillance in England and Wales 1962-81. The Lancet. 1984
Jan; 323(8368):91–94. `http://www.sciencedirect.com/science/article/pii/S0140673684900151`, doi: 10.1016/S0140-
6736(84)90015-1.

Garten R, Blanton L, Elal AIA, Alabi N, Barnes J, Biggerstaff M, Brammer L, Budd AP, Burns E, Cummins CN, Davis T, Garg S, Gubareva
L, Jang Y, Kniss K, Kramer N, Lindstrom S, Mustaquim D, O'Halloran A, Sessions W, et al. Update: Influenza Activity in the United
States During the 2017–18 Season and Composition of the 2018–19 Influenza Vaccine. MMWR Morbidity and Mortality Weekly Report.
2018; 67:634–642. `https://www.cdc.gov/mmwr/volumes/67/wr/mm6722a4.htm`, doi: 10.15585/mmwr.mm6722a4.

Gostic KM, Ambrose M, Worobey M, Lloyd-Smith JO. Potent protection against H5N1 and H7N9 influenza via childhood hemagglu-
tinin imprinting. Science. 2016 Nov; 354(6313):722–726. `https://science.sciencemag.org/content/354/6313/722`, doi:
10.1126/science.aag1322.

Gostic KM, Bridge R, Brady S, Viboud C, Worobey M, Lloyd-Smith JO. Childhood immune imprinting to influenza A shapes birth
year-specific risk during seasonal H1N1 and H3N2 epidemics. PLOS Pathogens. 2019 Dec; 15(12):e1008109. `https://journals.`
`plos.org/plospathogens/article?id=10.1371/journal.ppat.1008109`, doi: 10.1371/journal.ppat.1008109.

Government of the Netherlands, Primary education in the Netherlands. Accessed January 11, 2020; 2020. `https://www.government.`
`nl/topics/primary-education`.

Grenfell BT, Anderson RM. The estimation of age-related rates of infection from case notifications and serological data. Jour-
nal of Hygiene. 1985; 95(2):419–436. `https://www.cambridge.org/core/journals/epidemiology-and-infection/`
`article/estimation-of-agerelated-rates-of-infection-from-case-notifications-and-serological-data/`
`71C6598262C607CA228B91C4EB29EA79`.

Gupta S, Ferguson N, Anderson R. Chaos, Persistence, and Evolution of Strain Structure in Antigenically Diverse Infectious Agents. Science.
1998 May; 280(5365):912–915. `https://science.sciencemag.org/content/280/5365/912`, doi: 10.1126/science.280.5365.912.

Hayward AC, Fragaszy EB, Bermingham A, Wang L, Copas A, Edmunds WJ, Ferguson N, Goonetilleke N, Harvey G, Kovar J, Lim
MSC, McMichael A, Millett ERC, Nguyen-Van-Tam JS, Nazareth I, Pebody R, Tabassum F, Watson JM, Wurie FB, Johnson AM, et al.
Comparative community burden and severity of seasonal and pandemic influenza: results of the Flu Watch cohort study. The Lancet
Respiratory Medicine. 2014 Jun; 2(6):445–454. `http://www.sciencedirect.com/science/article/pii/S2213260014700347`,
doi: 10.1016/S2213-2600(14)70034-7.

Horby P, Mai LQ, Fox A, Thai PQ, Thi Thu Yen N, Thanh LT, Le Khanh Hang N, Duong TN, Thoang DD, Farrar J, Wolbers M, Hien NT.
The Epidemiology of Interpandemic and Pandemic Influenza in Vietnam, 2007–2010The Ha Nam Household Cohort Study I. American
Journal of Epidemiology. 2012 May; 175(10):1062–1074. `https://academic.oup.com/aje/article/175/10/1062/90300`, doi:
10.1093/aje/kws121.

Huang QS, Bandaranayake D, Wood T, Newbern EC, Seeds R, Ralston J, Waite B, Bissielo A, Prasad N, Todd A, Jelley L, Gunn W,
McNicholas A, Metz T, Lawrence S, Collis E, Retter A, Wong Ss, Webby R, Bocacao J, et al. Risk Factors and Attack Rates of Seasonal
Influenza Infection: Results of the Southern Hemisphere Influenza and Vaccine Effectiveness Research and Surveillance (SHIVERS)
Seroepidemiologic Cohort Study. The Journal of Infectious Diseases. 2019 Jan; 219(3):347–357. `https://academic.oup.com/jid/`
`article/219/3/347/5054657`, doi: 10.1093/infdis/jiy443.

Huang QS, Lopez LD, McCallum L, Adlam B. Influenza surveillance and immunisation in New Zealand, 1997–2006. Influenza and Other
Respiratory Viruses. 2008; 2(4):139–145. `https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1750-2659.2008.00050.`
`x`, doi: 10.1111/j.1750-2659.2008.00050.x.

Influenza Research Database, HA Subtype Numbering Conversion (beta) - Influenza Research Database. Accessed August 27, 2019.; 2019.
`https://www.fludb.org/brc/haNumbering.spg?method=ShowCleanInputPage&decorator=influenza`.

Institute of Environmental Science and Research, Influenza surveillance in New Zealand - 2016. Accessed March 16, 2020.; 2017.
`https://surv.esr.cri.nz/PDF_surveillance/Virology/FluAnnRpt/InfluenzaAnn2016.pdf`.

Institute of Environmental Science and Research, Influenza Annual Reports. Accessed June 21, 2020.; 2020. `https://surv.esr.cri.`
`nz/virology/influenza_annual_report.php`.

Katoh K, Misawa K, Kuma Ki, Miyata T. MAFFT: a novel method for rapid multiple sequence alignment based on fast Fourier transform.
Nucleic Acids Research. 2002 Jul; 30(14):3059–3066. `https://academic.oup.com/nar/article/30/14/3059/2904316`, doi:
10.1093/nar/gkf436.

Kilbourne ED. Influenza Pandemics of the 20th Century. Emerging Infectious Diseases. 2006 Jan; 12(1):9–14. `https://www.ncbi.nlm.`
`nih.gov/pmc/articles/PMC3291411/`, doi: 10.3201/eid1201.051254.

Kirkpatrick E, Qiu X, Wilson PC, Bahl J, Krammer F. The influenza virus hemagglutinin head evolves faster than the stalk domain. Scientific
Reports. 2018 Jul; 8(1):1–14. `https://www.nature.com/articles/s41598-018-28706-1`, doi: 10.1038/s41598-018-28706-1.

Kucharski AJ, Gog JR. Age profile of immunity to influenza: Effect of original antigenic sin. Theoretical Population Biology. 2012 Mar;
81(2):102–112. `http://www.sciencedirect.com/science/article/pii/S004058091100116X`, doi: 10.1016/j.tpb.2011.12.006.

Kucharski AJ, Gog JR. The Role of Social Contacts and Original Antigenic Sin in Shaping the Age Pattern of Immunity to Seasonal
Influenza. PLOS Computational Biology. 2012 Oct; 8(10):e1002741. `https://journals.plos.org/ploscompbiol/article?id=`
`10.1371/journal.pcbi.1002741`, doi: 10.1371/journal.pcbi.1002741.

Kucharski AJ, Lessler J, Read JM, Zhu H, Jiang CQ, Guan Y, Cummings DAT, Riley S. Estimating the life course of influenza A(H3N2)
antibody responses from cross-sectional data. PLoS biology. 2015 Mar; 13(3):e1002082. `http://journals.plos.org/plosbiology/`
`article?id=10.1371/journal.pbio.1002082`, doi: 10.1371/journal.pbio.1002082.

Langat P, Raghwani J, Dudas G, Bowden TA, Edwards S, Gall A, Bedford T, Rambaut A, Daniels RS, Russell CA, Pybus OG, McCauley
J, Kellam P, Watson SJ. Genome-wide evolutionary dynamics of influenza B viruses on a global scale. PLOS Pathogens. 2017
Dec; 13(12):e1006749. `https://journals.plos.org/plospathogens/article?id=10.1371/journal.ppat.1006749`, doi:
10.1371/journal.ppat.1006749.

Laurie KL, Horman W, Carolan LA, Chan KF, Layton D, Bean A, Vijaykrishna D, Reading PC, McCaw JM, Barr IG. Evidence for Viral Interference and Cross-reactive Protective Immunity Between Influenza B Virus Lineages. The Journal of Infectious Diseases. 2018 Jan; 217(4):548–559. `https://academic.oup.com/jid/article/217/4/548/4792826`, doi: 10.1093/infdis/jix509.

Levandowski RA, Gross PA, Weksler M, Staton E, Williams MS, Bonelli J. Cross-reactive antibodies induced by a monovalent influenza B virus vaccine. Journal of Clinical Microbiology. 1991 Jul; 29(7):1530–1532. `https://www.ncbi.nlm.nih.gov/pmc/articles/PMC270149/`.

Levandowski RA, Regnery HL, Staton E, Burgess BG, Williams MS, Groothuis JR. Antibody responses to influenza B viruses in immunologically unprimed children. Pediatrics. 1991 Nov; 88(5):1031–1036.

Linderman SL, Chambers BS, Zost SJ, Parkhouse K, Li Y, Herrmann C, Ellebedy AH, Carter DM, Andrews SF, Zheng NY, Huang M, Huang Y, Strauss D, Shaz BH, Hodinka RL, Reyes-Terán G, Ross TM, Wilson PC, Ahmed R, Bloom JD, et al. Potential antigenic explanation for atypical H1N1 infections among middle-aged adults during the 2013-2014 influenza season. Proceedings of the National Academy of Sciences of the United States of America. 2014 Oct; 111(44):15798–15803. `http://www.ncbi.nlm.nih.gov/pubmed/25331901`, doi: 10.1073/pnas.1409171111.

Melegaro A, Jit M, Gay N, Zagheni E, Edmunds WJ. What types of contacts are important for the spread of infections? Using contact survey data to explore European mixing patterns. Epidemics. 2011 Sep; 3(3):143–151. `http://www.sciencedirect.com/science/article/pii/S1755436511000247`, doi: 10.1016/j.epidem.2011.04.001.

Monto AS, Koopman JS, Longini IM. Tecumseh Study of Illness XIII. Influenza infection and disease, 1976–1981. American Journal of Epidemiology. 1985 Jun; 121(6):811–822. `https://academic.oup.com/aje/article/121/6/811/92221`, doi: 10.1093/oxford-journals.aje.a114052.

Mossong J, Hens N, Jit M, Beutels P, Auranen K, Mikolajczyk R, Massari M, Salmaso S, Tomba GS, Wallinga J, Heijne J, Sadkowska-Todys M, Rosinska M, Edmunds WJ. Social Contacts and Mixing Patterns Relevant to the Spread of Infectious Diseases. PLoS Medicine. 2008 Mar; 5(3):e74. `https://dx.plos.org/10.1371/journal.pmed.0050074`, doi: 10.1371/journal.pmed.0050074.

Nachbagauer R, Choi A, Izikson R, Cox MM, Palese P, Krammer F. Age Dependence and Isotype Specificity of Influenza Virus Hemagglutinin Stalk-Reactive Antibodies in Humans. mBio. 2016 Mar; 7(1):e01996–15. `https://mbio.asm.org/content/7/1/e01996-15`, doi: 10.1128/mBio.01996-15.

Petrie JG, Parkhouse K, Ohmit SE, Malosh RE, Monto AS, Hensley SE. Antibodies Against the Current Influenza A(H1N1) Vaccine Strain Do Not Protect Some Individuals From Infection With Contemporary Circulating Influenza A(H1N1) Virus Strains. The Journal of Infectious Diseases. 2016 Dec; 214(12):1947–1951. `https://academic.oup.com/jid/article/214/12/1947/2632603`, doi: 10.1093/infdis/jiw479.

Puzelli S, Di Martino A, Facchini M, Fabiani C, Calzoletti L, Di Mario G, Palmieri A, Affanni P, Camilloni B, Chironna M, D'Agaro P, Giannecchini S, Pariani E, Serra C, Rizzo C, Bella A, Donatelli I, Castrucci MR, Ansaldi F, Arvia R, et al. Co-circulation of the two influenza B lineages during 13 consecutive influenza surveillance seasons in Italy, 2004–2017. BMC Infectious Diseases. 2019 Nov; 19(1):990. `https://doi.org/10.1186/s12879-019-4621-z`, doi: 10.1186/s12879-019-4621-z.

Rajendran M, Nachbagauer R, Ermler ME, Bunduc P, Amanat F, Izikson R, Cox M, Palese P, Eichelberger M, Krammer F. Analysis of Anti-Influenza Virus Neuraminidase Antibodies in Children, Adults, and the Elderly by ELISA and Enzyme Inhibition: Evidence for Original Antigenic Sin. mBio. 2017 May; 8(2):e02281–16. `https://mbio.asm.org/content/8/2/e02281-16`, doi: 10.1128/mBio.02281-16.

Ranjeva S, Subramanian R, Fang VJ, Leung GM, Ip DKM, Perera RAPM, Peiris JSM, Cowling BJ, Cobey S. Age-specific differences in the dynamics of protective immunity to influenza. Nature Communications. 2019 Apr; 10(1):1–11. `https://www.nature.com/articles/s41467-019-09652-6`, doi: 10.1038/s41467-019-09652-6.

Rota PA, Wallis TR, Harmon MW, Rota JS, Kendal AP, Nerome K. Cocirculation of two distinct evolutionary lineages of influenza type B virus since 1983. Virology. 1990 Mar; 175(1):59–68.

Shaw MW, Xu X, Li Y, Normand S, Ueki RT, Kunimoto GY, Hall H, Klimov A, Cox NJ, Subbarao K. Reappearance and Global Spread of Variants of Influenza B/Victoria/2/87 Lineage Viruses in the 2000–2001 and 2001–2002 Seasons. Virology. 2002 Nov; 303(1):1–8. `http://www.sciencedirect.com/science/article/pii/S0042682202917198`, doi: 10.1006/viro.2002.1719.

Skowronski DM, Chambers C, De Serres G, Sabaiduc S, Winter AL, Dickinson JA, Gubbay JB, Drews SJ, Fonseca K, Charest H, Martineau C, Hickman R, Chan T, Jassem A, Petric M, Rose C, Bastien N, Li Y, Krajden M. Vaccine Effectiveness Against Lineage-matched and -mismatched Influenza B Viruses Across 8 Seasons in Canada, 2010–2011 to 2017–2018. Clinical Infectious Diseases. 2019 May; 68(10):1754–1757. `https://academic.oup.com/cid/article/68/10/1754/5127119`, doi: 10.1093/cid/ciy876.

Skowronski DM, Chambers C, De Serres G, Sabaiduc S, Winter AL, Dickinson JA, Gubbay JB, Fonseca K, Drews SJ, Charest H, Martineau C, Krajden M, Petric M, Bastien N, Li Y. Age-Related Differences in Influenza B Infection by Lineage in a Community-Based Sentinel System, 2010–2011 to 2015–2016, Canada. The Journal of Infectious Diseases. 2017 Sep; 216(6):697–702. `https://academic.oup.com/jid/article/216/6/697/4068745`, doi: 10.1093/infdis/jix393.

Skowronski DM, Hamelin ME, Janjua NZ, De Serres G, Gardy JL, Rhéaume C, Bouhy X, Boivin G. Cross-Lineage Influenza B and
Heterologous Influenza A Antibody Responses in Vaccinated Mice: Immunologic Interactions and B/Yamagata Dominance. PLoS ONE.
2012 Jun; 7(6):e38929. `https://dx.plos.org/10.1371/journal.pone.0038929`, doi: 10.1371/journal.pone.0038929.

Smith AJ, Davies JR. Natural Infection with Influenza A (H3N2). The Development, Persistence and Effect of Antibodies to the Surface
Antigens. The Journal of Hygiene. 1976; 77(2):271–282. `https://www.jstor.org/stable/3862080`.

Smith DJ, Lapedes AS, Jong JCd, Bestebroer TM, Rimmelzwaan GF, Osterhaus ADME, Fouchier RAM. Mapping the Antigenic and Genetic
Evolution of Influenza Virus. Science. 2004 Jul; 305(5682):371–376. `https://science.sciencemag.org/content/305/5682/371`,
doi: 10.1126/science.1097211.

Sočan M, Prosenc K, Učakar V, Berginc N. A comparison of the demographic and clinical characteristics of laboratory-confirmed influenza
B Yamagata and Victoria lineage infection. Journal of Clinical Virology. 2014 Sep; 61(1):156–160. `http://www.sciencedirect.`
`com/science/article/pii/S1386653214002510`, doi: 10.1016/j.jcv.2014.06.018.

StatsNZ, StatsNZ Infoshare - Estimated Resident Population by Age and Sex (1991+) (Annual-Dec). Accessed August 1st, 2019.; 2019.
`http://archive.stats.govt.nz/infoshare/`.

Sun W, Kang DS, Zheng A, Liu STH, Broecker F, Simon V, Krammer F, Palese P. Antibody Responses toward the Major Antigenic Sites of
Influenza B Virus Hemagglutinin in Mice, Ferrets, and Humans. Journal of Virology. 2019 Jan; 93(2). doi: 10.1128/JVI.01673-18.

Tan Y, Guan W, Lam TTY, Pan S, Wu S, Zhan Y, Viboud C, Holmes EC, Yang Z. Differing Epidemiological Dynamics of Influenza B Virus
Lineages in Guangzhou, Southern China, 2009-2010. Journal of Virology. 2013 Nov; 87(22):12447–12456. `https://jvi.asm.org/`
`content/87/22/12447`, doi: 10.1128/JVI.01039-13.

Tricco AC, Chit A, Soobiah C, Hallett D, Meier G, Chen MH, Tashkandi M, Bauch CT, Loeb M. Comparing influenza vaccine efficacy
against mismatched and matched strains: a systematic review and meta-analysis. BMC Medicine. 2013 Jun; 11(1):153. `https:`
`//doi.org/10.1186/1741-7015-11-153`, doi: 10.1186/1741-7015-11-153.

Vijaykrishna D, Holmes EC, Joseph U, Su YCF, Halpin R, Lee RTC, Deng YM, Gunalan V, Lin X, Stockwell TB, Fedorova NB, Zhou B,
Spirason N, Kühnert D, Bošková V, Stadler T, Costa AM, Dwyer DE, Huang QS, Jennings LC, et al. The contrasting phylodynamics of
human influenza B viruses | eLife. eLife. 2015; 4:e05055. `https://elifesciences.org/articles/05055`, doi: 10.7554/eLife.05055.

Virk RK, Jayakumar J, Mendenhall IH, Moorthy M, Lam P, Linster M, Lim J, Lin C, Oon LLE, Lee HK, Koay ESC, Vijaykrishna D,
Smith GJD, Su YCF. Divergent evolutionary trajectories of influenza B viruses underlie their contemporaneous epidemic activity.
Proceedings of the National Academy of Sciences. 2020 Jan; 117(1):619–628. `https://www.pnas.org/content/117/1/619`, doi:
10.1073/pnas.1916585116.

773

Figure 1–Supplement 1. Distribution of medically attended influenza B cases in New Zealand (2001-2019) and Australia (2002-2013) by birth year of the patient. The fraction of cases in each birth year was calculated relative to all cases observed for each lineage (separately by type of surveillance in New Zealand).

Figure 1–Supplement 2. Comparison of lineage frequency estimates based on sequence data and surveillance reports.
Frequencies are shown by annual influenza seasons, which span multiple calendar years in the United States but are contained in a single calendar year in Australia and New Zealand (e.g., 2006 refers to the 2006/2007 influenza season in the United States and to the 2006 season in Australia and New Zealand). Numbers indicate the number of isolates collected in that season and deposited on GISAID or the NCBI Influenza Virus Database (blue) or tested by surveillance (orange; the total number of isolates tested was not reported in some of the surveillance reports). Sequence database isolates from Australia and New Zealand were grouped together to estimate lineage frequencies.

Figure 1–Supplement 3. Amino acid divergence in the hemagglutinin (HA) and neuraminidase (NA) proteins within and between influenza B lineages. The left panel shows the fraction of amino acid sites that differ between pairs of sequences in each season (each point is a pair). Up to 500 randomly chosen pairs are shown, while the solid lines represent the annual average calculated from all pairs given two samples of up to 100 sequences from each lineage (i.e, up to 4,950 pairs). The right panel shows divergence of strains circulating at different times from reference strains B/Victoria/2/87 and B/Yamagata/16/88.

Figure 1–Supplement 4. Lineage frequencies used in sensitivity analysis to test the effect of having only B/Yamagata in New Zealand in the 1990s. If fewer than 10 isolates were reported in New Zealand and Australia combined in a season, we estimated frequencies using isolates from all countries represented in the databases. The number inside each circle indicates the number of isolates used to estimate lineage frequencies in each season.

Figure 2–Supplement 1. Likelihood profiles for protection parameters estimated from the complete New Zealand data. The 95% confidence interval based on a likelihood ratio test with one degree of freedom is indicated by the vertical dashed lines.

Figure 2–Supplement 2. Observed and predicted distributions of influenza B cases in New Zealand by observation year. Vertical red bars are 95% bootstrap confidence intervals. Season/lineage combinations with fewer than 5 cases were omitted to improve visualization.

Figure 2–Supplement 3. Likelihood profiles for protection parameters estimated from sentinel cases alone. The 95% confidence interval based on a likelihood ratio test with one degree of freedom is indicated by the vertical dashed lines.

Figure 2–Supplement 4. Age distributions predicted by the model fitted to sentinel cases only. Red lines and dots show the predicted distribution under the model. Vertical bars are 95% bootstrap confidence intervals.

Figure 2–Supplement 5. Likelihood profiles for protection parameters estimated from birth years starting in 1952 instead of 1959. The 95% confidence interval based on a likelihood ratio test with one degree of freedom is indicated by the vertical dashed lines.

781

Figure 2–Supplement 6. Likelihood profiles for protection parameters estimated from birth years starting in 1966 instead of 1959. The 95% confidence interval based on a likelihood ratio test with one degree of freedom is indicated by the vertical dashed lines.

783

Figure 2–Supplement 7. Likelihood profiles for protection parameters estimated by the model assuming differential reporting of children under 5. The 95% confidence interval based on a likelihood ratio test with one degree of freedom is indicated by the vertical dashed lines.

Figure 2–Supplement 8. Likelihood profiles for protection parameters estimated from influenza B cases in Australia.
The 95% confidence interval based on a likelihood ratio test with one degree of freedom is indicated by the vertical dashed lines.

Figure 2–Supplement 9. Observed and predicted distributions of influenza B cases in Australia by observation year.
Vertical red bars are 95% bootstrap confidence intervals. Season/lineage combinations with fewer than 5 cases were omitted to improve visualization.

Figure 2–Supplement 10. Model predictions from the sensitivity analysis of the frequency of B/Yamagata in New Zealand in the 1990s. Red lines and dots show the predicted distribution under the model. Vertical bars are 95% bootstrap confidence intervals.

Figure 3–Supplement 1. Bivariate profiles of age-specific baseline probabilities of infection and protection from B/Yamagata against B/Victoria. The red circle indicates the maximum likelihood value in the profile. The contour indicates the 95% confidence interval based on a bilinear interpolation of the profiled points.

Figure 3–Supplement 2. Frequency of past influenza B infections in children predicted by the model compared with observed seroprevalence in the Netherlands. The fraction of children previously infected with influenza B in the Netherlands in 2006-2007 was estimated by Bodewes et al. (2011) as the fraction having serum antibodies against at least one influenza B strain from a panel of viruses (left panel), at least one B/Victoria strain (center panel) or at least one B/Yamagata strain (right panel). Bars show 95% binomial confidence intervals. Fractions for Australia and New Zealand were generated independently from the seroprevalence data by fitting a statistical model to medically attended influenza B infections. For the model predictions, binomial confidence intervals assume a sample size equal to the number of children with the corresponding age in the seroprelavence data.

Figure 3–Supplement 3. Amino acid divergence in the hemagglutinin and neuraminidase surface proteins between influenza A subtypes and between influenza B lineages. Divergence in hemagglutinin is shown separately for the antigenically variable head region (left) and the more conserved stalk region (center). Each point represents a pair of strains isolated in the same year and deposited on GISAID. When more than 100 isolates from a subtype or lineage were isolated in a single year, we used a random sample of 100 isolates, resulting in up to 4,950 pairs of each kind in each year. Up to 500 randomly chosen pairs are shown for each year. Average divergence values reported in the main text are based on the full set of pairs given the randomly sampled sequences.